


NEW AGE

Second Edition

PHARMACOGNOSY AND PHARMACOBIO TECHNOLOGY

Ashutosh Kar



NEW AGE INTERNATIONAL PUBLISHERS

PHARMACOGNOSY AND
PHARMACOBIOLOGY

(Revised-Expanded Second Edition)

**This page
intentionally left
blank**

PHARMACOGNOSY AND PHARMACBIOTECHNOLOGY

(Revised–Expanded Second Edition)

Ashutosh Kar

Professor and Head of Pharmacy
Shri RNS College of Pharmacy
Gormi, Bhind (M.P.), Pin-477 660

Formerly

Professor, School of Pharmacy, Addis Ababa University,
Addis Ababa (Ethiopia)
Dean, Chairman & Professor, Faculty of Pharmaceutical Sciences,
Guru Jambheshwar University, **Hisar (India)**
Professor, School of Pharmacy, Al Arab Medical University
Benghazi, (Libya)
Professor, College of Pharmacy (University of Delhi)
Delhi, (India)
Professor & Head, Department of Pharmaceutical Chemistry,
Faculty of Pharmaceutical Sciences, University of Nigeria,
Nsukka, (Nigeria)



NEW AGE INTERNATIONAL (P) LIMITED, PUBLISHERS

New Delhi • Bangalore • Chennai • Cochin • Guwahati • Hyderabad
Jalandhar • Kolkata • Lucknow • Mumbai • Ranchi

Copyright © 2007, 2003 , New Age International (P) Ltd., Publishers
Published by New Age International (P) Ltd., Publishers

All rights reserved.

No part of this ebook may be reproduced in any form, by photostat, microfilm, xerography, or any other means, or incorporated into any information retrieval system, electronic or mechanical, without the written permission of the publisher.
All inquiries should be emailed to rights@newagepublishers.com

ISBN (13) : 978-81-224-2915-2

PUBLISHING FOR ONE WORLD

NEW AGE INTERNATIONAL (P) LIMITED, PUBLISHERS

4835/24, Ansari Road, Daryaganj, New Delhi - 110002

Visit us at www.newagepublishers.com

Dedication

*Dedicated with humility and reverence
to the fond memory of beloved
parents who encouraged
and flared passion in
me to learn more
always.*

Thanks

*Wish to thank Leena, Ashish
and Abhijeet for their boundless
patience and eternal understanding
during completion of this text.*

Love

*Aditi, our grand-daughter, who
brought in an eternal saga of
love, and energised
our inspirations to
perform better.*

**This page
intentionally left
blank**

Preface to the Second Edition

Overwhelming appreciation, whole-hearted acceptance, and qualified success of the **First Edition** of the textbook entitled '**Pharmacognosy and Pharmacobiotechnology**' extended by the postgraduate students specializing in Pharmacognosy and Phytochemistry, Biotechnology, Pharmaceutical Chemistry, Chemistry of Natural Products; besides the enormous undergraduate students in Bachelor of Pharmacy programmes in all the reputed Pharmacy Degree Colleges not only in India but also abroad are quite encouraging.

Exclusively based upon the commendable comments and constructive criticisms received from various academic colleagues across the country the author has meticulously presented this entirely revised and duly expanded **Second Edition**. Moreover, the reading text material has been updated thoroughly, various biosynthetic pathways modified adequately, chemical structures and tabular contents enumerated more explicitly. Besides, the critical definitions, important statements, terminologies, names of chemical constituents have been duly highlighted so as to facilitate its readers to comprehend them accurately.

Tremendous achievement and advance in the different segments of highly sophisticated '**Research Techniques**' solely based on PC-modulated modern analytical techniques helped not only to clarify the rather complex chemical structures of unravelled chemical constituents from **Natural Products**, but also established precisely the plethora of '**Biosynthetic Pathways**' dominating the plant kingdom. The present textbook essentially comprises of nearly sixty '**biosynthetic routes**' of predominantly important natural chemical entities, such as: **alkaloids, antibiotics, glycosides, marine-derived drug substances, and terpenoids**.

It is, however, pertinent to add here that certain extremely preliminary aspects to the related pharmacognostical characteristic features of '**Natural Plant Products**', namely: morphological structures, adulterants used in herbal products, habitats, method of cultivation, geographical distribution etc., have been expunged from the text, to which the students invariably obtain sufficient exposure in the early stages of their systematic curriculum follow up.

The **Second Edition** essentially comprises of **five additional chapters**, namely: (i) **Nutraceuticals**, (ii) **Enzymes and Protein Drugs**, (iii) **Biomedicinals from Plant Tissue Cultures**, (iv) **Hi-Tech Products from Plant Sources**, and (v) **Indian Traditional Herbal Drugs**, i.e., chapter-11 through chapter-15. The judicious and thoughtful inclusion of these five chapters would certainly expose the PG/UG students of the aforesaid disciplines to an exceptionally solid platform in the scientific pursuit of their knowledge.

The author feels convinced and earnestly believes that the **Second Revised and Expanded Edition** of **Pharmacognosy and Pharmacobiotechnology** shall largely fulfil the much needed value-added substantial text materials.

It is hoped that **Pharmacognosy and Pharmacobiotechnology** will continue to enjoy its popularity amongst the august teaching fraternity, brilliant students, herbal practitioners, pharmacognosists, herbal chemists, phytochemists, biotechnologists, and above all the researchers who would like to make an illustrious career in their respective professional discipline in the **New Millennium**.

Finally, the author wishes to place on record his deep sense of gratitude to Shri Saumya Gupta M.D., and the entire professionals of New Age International (P) Limited Publishers, for their excellent support to bring out this edition in such a short-span.

Gurgaon

Dr. Ashutosh Kar

Preface to the First Edition

Etymological evidences reveal that ‘**pharmacognosy**’ refers to the knowledge (from the Greek *gnosis*) of drug (*Pharmacon*) substances. **Pharmacognosy** may also be referred to as—‘*Study of sources, and chemical and physical properties of drugs*’. In the present context *pharmacognosy*, since Dioscorides’s treatise, has spread its tentacles to investigations of a wider section of naturally occurring materials essentially comprised of plants, animals, substances originated from microorganisms and even biotechnology and genetically engineered entities. Jean Bruneton, the famous French pharmacognosist, describes pharmacognosy as—‘*Study of starting materials and substances intended for therapeutics, and of biological origin, in other words obtained from plants, animals, or by fermentation from microorganisms*’.

Since the past two centuries the identification, isolation and characterization of naturally occurring substances across the world have been accomplished by the concerted efforts through a central preoccupation of innumerable research chemists and biological scientists. In the recent past, the world has witnessed an overwhelming progress towards intensification of interest more so in natural products from the herbal-based pharmaceutical industries with the epoch-making discoveries of extremely useful new drugs, namely, taxol, artemisinin ginsengoside Rg1 ginkgolide A, doxorubicin and the like, from the nature’s natural reserves.

The other predominant aspect is ‘**pharmacobiotechnology**’, an area that encompasses the intricate production of natural-product-drug substances on the basis of the copious volumes of scientific evidences amalgamated with tremendous progress and breakthroughs particularly in the fields of ‘**biotechnology**’ and ‘**molecular biology**’. It is indeed an altogether newer frontier charged with innovative ideas and approaches in modern-drug-discovery scenario to modify and improve upon the quality of life of human beings on this planet. Therefore, in the present textbook, an earnest attempt has been made to deal with the newest drugs on one hand and the oldest ones on the other in a very systematic and lucid manner with a strong conviction that these all belong to the natural origins.

Interestingly, the last five decades have witnessed a quantum jump in relevant and useful publications especially with regard to books pertaining to medicinal plants, medicinal herbs, biologically active natural products, phytochemistry of medieval plants, alternative medicine; besides herbal and botanical remedies for commoners. It is, however, pertinent to mention about the legitimate exposure *vis-a-vis* the in depth knowledge of the various aspects of medicinal plants well within the broader limits of *pharmacognosy*—a professional discipline widely recognised not only amongst the pharmacy and medical herbalism academic programmes but also of utmost significance to non-medical professionals.

The present text essentially comprised of ten chapters, namely: introduction to phytochemistry, pharmacobiotechnology, carbohydrates, glycosides, terpenoids, phenylpropanoids, alkaloids, bitter

principles, antibiotics, and drug molecules of marine organisms. Keeping in view the intensive and remarkable progressive advances accomplished in phytochemical and technological research, it was thought worthwhile to make adequate coverage of pharmacognosy essentially needed not only for the pharmacy degree syllabuses in general but also for the professional class in particular.

The drugs have been classified on a unique broad-based, widely accepted and literature-supported manner.

These are carefully selected and arranged in each chapter, organized on the strong footing of chemical relationships, their biosynthetic approach, thereby elaborating sufficient basic fundamentals for the better advanced knowledge and vivid understanding of the wonderful natural products as 'drugs'.

Each individual drug belonging to various groups dealt with in the present textbook has been carefully selected based on its academic merit, status and relevance. It has been treated in a most scientific and methodical manner essentially consisting of the following highlights, namely: latest classification, authentic nomenclature, synonym(s), biological source(s), chemical structure, chemical name, molecular formula, isolation or preparation methods, characteristic features, identification test(s), derivatization, characteristics of derivatives, therapeutic uses, and biosynthetic pathways, wherever applicable. A number of important classes of compounds and their relevant features have been summarized in tabular forms selected figures in the text have been incorporated at appropriate places to make the ensuing subject matter more easily understandable to its readers biosynthetic pathways have been explicitly dealt with. The text contains more than one thousand chemical structures of drugs and their intermediates, more than fifty biosynthetic pathways and about fifteen figures.

Bearing in mind the extraordinary pace and appreciable momentum gathered by the global pharmaceutical market, followed by an encouraging number of newer drug companies joining the modern trend of market demands, there exists an enormous scope in phytochemical research and development efforts. The world-wide intensive quest for newer, safer and effective drugs from natural products is not confined only to the terrestrial plants from tropical rain forests and to animals, but also to the plants and microorganisms occurring in deep oceans surrounding this earth.

It is earnestly believed that in the present textbook the modern concepts of **pharmacognosy** shall fulfil the necessary requirements of undergraduate and graduate students of various universities in India and other developing nations. Those who intend to continue their research in medicinal plants and desire to establish a strong base in the production of herbal-drug industries may also find this compilation equally informative and useful.

Addis Ababa

Dr. Ashutosh Kar

Contents

<i>Preface to the Second Edition</i>	vii
<i>Preface to the First Edition</i>	ix
1. Introduction	1
1.1 Pharmacognosy—A Brief History	1
1.2 Importance of Natural Drug Substances	5
1.2.1 Serve as Extremely Useful Natural Drugs	5
1.2.2 Provide Basic Compounds Affording Less Toxic and More Effective Drug Molecules	10
1.2.3 Exploration of Biological–Active–Prototypes towards Newer and Better Synthetic Drugs	10
1.2.4 Modification of Inactive Natural Products by Suitable Biological/Chemical Means into Potent Drugs	13
1.3 Natural Drugs Substances: Cultivation and Production	15
1.3.1 Plant Products	15
1.3.2 Cell-Culture Techniques	16
1.3.3 Microbial Metabolites	16
1.3.4 Animal Derivatives	18
1.4 Phytochemistry	19
1.4.1 Constituents	21
2. Pharmacobiotechnology	41
2.1 Introduction	41
2.2 Theory	42
2.2.1 Mutation, Crossing-over and Recombinant of Meiosis	43
2.2.2 Third Revolution in Modern Medicine	43
2.2.3 Genetic Code	44
2.2.4 Specific Sets of Genes in Each Individual Organ	44
2.2.5 Reverse Transcriptase (RT)	44
2.2.6 3D-Proteins	45
2.2.7 From Nervous System to Immune System	45
2.2.8 Body's Defence Mechanism	46
2.2.9 PCR—in Forensic & Research	46
2.2.10 DNA—in Metabolic Pathways	46
2.2.11 Recombinant Vaccination Vector	46
2.2.12 OKT3—Monoclonal Antibody	47

2.3	Important Means in Biotechnology	47
2.3.1	Recombinant DNA (rDNA)	47
2.3.2	Restriction Enzymes	47
2.3.3	DNA–Ligase	48
2.3.4	Cloning Vector	48
2.3.5	Hybridization Probes	49
2.3.6	Cloning Process	50
2.4	Recombinant Proteins	58
2.4.1	Bacterial Systems	58
2.4.2	Glycosylation	59
2.4.3	Mammalian Tissue Culture Expression Systems	60
2.5	Biotechnology Vs Modern Pharmacy Practice	60
2.5.1	Human Proteins as Drugs	61
2.5.2	New Drug Classes	63
2.5.3	Vaccines	64
2.5.4	New Immunodiagnostic Agents	66
2.5.5	DNA Probes and RFLP Analysis	68
2.5.6	Enzyme Linked Immunosorbant Assay (ELISA)	69
2.6	Biotechnology Based Pharmaceuticals for the Millennium	71
2.6.1	Genetically Engineered Vaccine	72
2.6.2	Gene Splicing and DNA Recombinant Procedures	73
2.6.3	Antibodies in Biotechnology	75
2.6.4	Gene Therapy	77
2.6.5	3D Picture of the ‘Lock’ and ‘Keys’	78
2.7	Biotechnology and Modern Drug Discovery	79
2.7.1	Approved Medicines	79
2.7.2	Medicines Under Development	80
2.7.3	Human Clone	80
2.8	Biotechnology: Some Thought Provoking Newer Ideas	82
2.8.1	Potato Vaccine	82
2.8.2	Functional Food Revolution	82
3.	Carbohydrates	84
3.1	Introduction	84
3.2	Classification	86
3.2.1	Homoglycans	86
3.2.2	Heteroglycans	100
3.3	Carbohydrate Biogenesis	119
4.	Glycosides	122
4.1	Introduction	122
4.1.1	O-Glycosides	125
4.1.2	S-Glycosides	125
4.1.3	N-Glycosides	125
4.1.4	C-Glycosides	126

4.2	Classification	127
4.2.1	Anthracene Glycosides (or Anthraquinone Glycosides)	127
4.2.2	Phenol Glycosides	139
4.2.3	Steroid Glycosides	144
4.2.4	Flavonoid Glycosides	157
4.2.5	Coumarin and Furanocoumarin Glycosides	168
4.2.6	Cyanogenetic Glycosides	173
4.2.7	Thioglycosides	179
4.2.8	Saponin Glycosides	182
4.2.9	Aldehyde Glycosides	195
4.2.10	Bitter Glycosides	196
4.2.11	Miscellaneous Glycosides	200
4.3	Biosynthesis of Glycosides	202
4.3.1	Biosynthesis of Anthracene Glycosides	203
4.3.2	Biosynthesis of Phenol Glycosides	204
4.3.3	Biosynthesis of Steroid Glycosides	204
4.3.4	Biosynthesis of Flavonoid Glycosides	204
4.3.5	Biosynthesis of Coumarin and Furanocoumarin Glycosides	205
4.3.6	Biosynthesis of Cyanogenetic Glycosides	207
4.3.7	Biosynthesis of Thioglycosides	209
4.3.8	Biosynthesis of Saponin Glycosides	209
4.3.9	Biosynthesis of Aldehyde Glycosides	210
4.4	Profile of Glycosides in Natural Plant Sources	210
5.	Terpenoids	215
5.1	Introduction	215
5.2	Classification	218
5.2.1	Monoterpenoids	218
5.2.2	Sesquiterpenoids	228
5.2.3	Diterpenoids	233
5.2.4	Triterpenoids	236
5.2.5	Tetraterpenoids and Carotenoids	238
5.2.6	Volatile Oils (or Essential Oils)	240
5.2.7	Resins and Resin Combinations	306
5.2.8	Oleoresins	323
5.2.9	Oleo-Gum-Resins	328
6.	Phenylpropanoids	340
6.1	Introduction	340
6.2	Classification	340
6.2.1	Hydroxycinnamic Acids	341
6.2.2	Phenylpropenes	344
6.2.3	Coumarins	345
6.2.4	Abridged Phenylpropanoids	354

6.2.5	Biphenylpropanoid Derivatives	361
6.2.6	High Molecular Weight Phenylpropanoids	366
6.3	Biosynthesis of Phenylpropanoids	369
7.	Alkaloids	372
7.1	Introduction	372
7.1.1	Nomenclature	374
7.1.2	Occurrence and Distribution in Different Organ's of Plant	374
7.1.3	Site of Formation of Alkaloids in Plants	377
7.1.4	Function of Alkaloids in Plants	377
7.1.5	Isomerism	378
7.1.6	General Characteristics of Alkaloids	380
7.1.7	General Methods of Extraction and Isolation of Alkaloids	389
7.2	Classification of Alkaloids	395
7.2.1	Alkaloids Derived from Amination Reactions	401
7.2.2	Alkaloids Derived from Anthranilic Acid	427
7.2.3	Alkaloids Derived from Histidine	436
7.2.4	Alkaloids Derived from Lysine	441
7.2.5	Alkaloids Derived from Nicotinic Acid	454
7.2.6	Alkaloids Derived from Ornithine	461
7.2.7	Alkaloids Derived from Tyrosine	475
7.2.8	Alkaloids Derived from Tryptophan	495
7.3	Alkaloids in Tissue Cultures	542
7.4	Alkaloids in Chemosystematics	543
8.	Bitter Principles	547
8.1	Introduction	547
8.2	Classification of Bitter Principles	547
8.2.1	Phenolic Bitter Principles	548
8.2.2	Lactone Bitter Principles	550
8.2.3	Chromone Bitter Principles	553
8.2.4	Coumarin Bitter Principles	556
8.2.5	Coumarone Bitter Principles	559
8.2.6	Miscellaneous Bitter Principles	561
9.	Antibiotics	568
9.1	Introduction	568
9.2	Antibiotic Development	569
9.2.1	Quest for New Antibiotics	569
9.2.2	Large-Scale Production	571
9.3	Classification of Antibiotics	579
9.3.1	Aminoglycosides	579
9.3.2	Anthracyclines	590
9.3.3	Cephalosporins	596
9.3.4	β -Lactams	611

9.3.5 Lincosamides	616
9.3.6 Macrolides	618
9.3.7 Penicillins	625
9.3.8 Polypeptide Antibiotics	641
9.3.9 Tetracyclines	649
9.3.10 Miscellaneous Antibiotics	657
10. Drug Molecules of Marine Organisms	695
10.1 Introduction	695
10.2 Classification of Drug Molecules of Marine Organisms	696
10.2.1 Cytotoxic/Antineoplastic Agents	696
10.2.2 Cardiovascular Active Drugs	701
10.2.3 Marine Toxins	709
10.2.4 Antimicrobial Drugs	716
10.2.5 Antibiotic Substances	718
10.2.6 Antiinflammatory and Antispasmodic Agents	720
10.2.7 Miscellaneous Pharmacologically Active Substances	721
10.3 Marine Natural Products: An Upgradation Profile	726
10.3.1 Microbial Transformations	726
10.3.2 Puupehenone: Semisynthetic Analogues	729
10.4 Summary	733
11. Nutraceuticals	735
11.1 Introduction	735
11.2 Phytochemicals as Nutraceuticals	737
11.2.1 Terpenoids (or Isoprenoids)	738
11.2.2 Non-Carotenoid Terpenoids	742
11.2.3 Polyphenolics [or Polyphenol Extracts]	744
11.2.4 Phenolic Acids	751
11.2.5 Non-Flavonoid Polyphenolics	753
11.2.6 Glucosinolates [or Thioglucosides]	755
11.2.7 Thiosulphinates [or Cysteine Sulphoxides]	759
11.2.8 Phytosterols	760
11.2.9 Anthraquinones	761
11.2.10 Glucosamine [Synonym: Chitosamine;]	763
11.2.11 Octacosanol [Synonym: Octacosyl Alcohol]	764
11.2.12 Carnitine [Synonym: g-Trimethyl-b-hydroxybutyrobetaine;]	765
11.2.13 Capsaicin [Synonyms: Axsain; Mioton; Zacin; Zostrix;]	767
11.2.14 Piperine	768
11.2.15 Chlorophyll	768
11.2.16 Pectin	770
11.2.17 Dominant Phytochemical Pigments	770
11.2.18 Tocotrienols and Tocopherols	770
11.2.19 α -Lipoic Acid and Ubiquinones	771

11.3 Contemporary Nutraceuticals	772
11.3.1 Spiruline	773
11.3.2 Broccoli	773
11.3.3 Aloe Vera Gel and Aloe Juice	774
11.3.4 Soyfoods	776
11.3.5 Omega-3 Fatty Acids	776
11.3.6 Pomegranate Juice	777
11.3.7 Walnuts	777
11.3.8 Certified Organic Mushroom Nutrace	777
12. Enzyme and Protein Drug Substances	779
<i>A. Enzyme as Drug Substances</i>	779
12.1 Introduction	779
12.2 Enzyme Variants	782
12.3 Enzymes of Pharmaceutical Relevance and Utility	783
12.4 Brief Description of Enzymes Used as Drugs	784
12.4.1 Bromelain	784
12.4.2 Chymotrypsin	784
12.4.3 Collagenase	784
12.4.4 Deoxyribonuclease [DNase]	785
12.4.5 Fibrinolysin	785
12.4.6 Hyaluronidase	785
12.4.7 Muramidase	785
12.4.8 Papain	786
12.4.9 Pancreatin	786
12.4.10 Pancrealipase	787
12.4.11 Pepsin [Greek: Pepsis = digestion]	787
12.4.12 Rennin [or Chymosin]	787
12.4.13 Seratiopeptidase	787
12.4.14 Streptokinase	788
12.4.15 Urokinase	788
12.4.16 L-Asparaginase	789
<i>B. Protein as Drug Substances</i>	789
12.5 Introduction	789
12.6 Protein Variants	790
12.7 Brief Description of Proteins Used as Drugs	791
12.7.1 Complement Protein (Complement Factor C-3) [Latin; Complere = to Complete]	791
12.7.2 Gelatin [Latin: Gelatina = Gelatin]	792
12.7.3 Collagen [Synonym: Ossien]: (Greek: kolla = glue, + gennan = to produce)	792
12.7.4 Casein [Latin: caseus = cheese]	793
12.7.5 Lectins [Synonyms: Agglutinins; Affinitins; Phasins; Protectins;]	794

12.7.6	Yeast	794
12.7.7	Thaumatococcus [Synonym: Talin;]	795
13.	Biomedicinals From Plant-Tissue Cultures	797
13.1	Introduction	797
13.2	Profile of Plant-Tissue Cultures	800
13.2.1	Type of Cultures	800
13.2.2	Composition of Culture Medium	801
13.2.3	Surface Sterilization of Explants	803
13.2.4	Preparation of Tissue Cultures	804
13.3	Biomedicinals in Plant-Tissue Cultures	805
13.3.1	Secondary Metabolites	806
13.3.2	Usefulness of Secondary Metabolites	808
13.3.3	Secondary Metabolites in Chemosystematics	809
13.3.4	Newer Products Developed	810
13.4	Bioproduction of Commendable Secondary Metabolites	811
14.	Hi-Tech Products from Plant Sources	814
14.1	Introduction	814
14.2	High Throughput Screening (HTS)	815
14.2.1	HTS and Bioassays	816
14.2.2	Access to Plants <i>vis-a-vis</i> Natural Source Materials	817
14.2.3	HTS and Selection for Plant Materials	818
14.2.4	Identification Process of Plants for Targeted Sets	819
14.2.5	Dereplication and Isolation of Active Compounds	820
14.3	Success of HTS of Plant Source Materials for New Lead Chemical Entities	820
14.3.1	Use of MS for Identification of Potent Biologically Active and Important Drug Molecules	821
14.4	Hi-Tech Products	822
14.4.1	Genistein [Syn. Genisteol; Prunetol;]	822
14.4.2	Camptothecin	823
14.4.3	Rhein [Syn: Monorhein; Rheic Acid; Cassic Acid; Parietic Acid; Rhubarb Yellow]	823
14.4.4	Taxanes	824
14.4.5	Homoharringtonine (HHT)	824
15.	Indian Traditional Herbal Drugs	827
15.1	Introduction	827
15.2	Indian Traditional Herbal Drugs	828
15.2.1	Cardiovascular Drugs	828
15.2.2	Immunomodulators and Adaptogens	830
15.2.3	Antidiabetic Drugs	830
15.2.4	Antineoplastic Drugs	831
15.2.5	Antiviral Drugs	832
	Index	835

**This page
intentionally left
blank**

1

Introduction

- Pharmacognosy—A Brief History
- Importance of Natural Drug Substances
- Natural Drugs Substances: Cultivation and Production
- Phytochemistry
- Further Reading References

1.1 PHARMACOGNOSY—A BRIEF HISTORY


‘**Pharmacognosy**’—has been coined by the merger of two Greek words **Pharmakon** (drug) and **Gnosis** (knowledge) *i.e.*, the **knowledge of drugs**. The nomenclature—‘**Pharmacognosy**’ was used first and foremost by C.A. Seydler, a medical student in Halle/Saale, Germany, who emphatically employed **Analetica Pharmacognostica** as the main title of his thesis in the year 1815. Besides, further investigations have revealed that Schmidt has made use of the terminology ‘*Pharmacognosis*’ in the monograph entitled **Lehrbuch der Materia Medica** (*i.e.*, Lecture Notes on Medical Matter) which dates back to 1811, in Vienna. This compilation exclusively deals with the medicinal plants and their corresponding characteristics.

It is indeed quite interesting to observe that our **ancients** were duly equipped with a vast, in-depth and elaborated knowledge of plethora of drugs from the vegetable origin but unfortunately they possessed a scanty knowledge with regard to the presence of chemically pure compounds in most of them.

Camphor found its enormous use in the treatment and cure of many ailments, for instance: **internally** as—*a stimulant and carminative*; **externally** as—an **antipruritic**, *counterirritant* and *antiseptic* by the ancient Egyptians, Chinese, Indians, Greeks and Romans.

Earlier it was obtained by mere cooling of volatile oils from—*ssasafras*, *rosemary*, *lavender*, *sage*; while the Ancient Greeks and Romans derived it as a by product in the manufacture of wine. Nowadays, camphor is obtained on a large-scale synthetically (racemic mixture) from the α -pinene present in the turpentine oil (Chapter 5).

African natives used plant extracts in their ritual ceremonies whereby the subject would lose his/her complete body movements but shall remain mentally alert for 2 or 3 days. Later on, the earlier civilization also discovered a number of fermented drinks solely derived from carbohydrate—rich plant substances invariably containing **alcohols** and **vinegar**. With the passage



CAMPHOR
(A Bicyclic Ketone)

of time they also recognised certain plant products exclusively used for poisoning their spears and arrows in killing their preys and enemies as well. Interestingly, they found that some plant extracts have the unique property of keeping the new meat fresh and also to mask its unpleasant taste and flavour.

The human beings belonging to the ancient era in different parts of the globe independently discovered the inherent stimulating characteristics of a wide variety of drinks exclusively prepared from the vegetative source as stated below in Table 1.1.

Table 1.1 Stimulating Characteristics from Vegetative Sources

S. No.	Common Name	Biological origin (Family)	Part used	Active Ingredient	Distribution
1.	Guarana	<i>Paullinia cupana</i> Kunth (<i>Sapindaceae</i>)	Seed	Caffeine (2.5-5.0%) Tannin (Catheochutannic acid) 25%	Brazil, Uruguay
2.	Paraguay Tea or Mate	<i>Ilex paraguariensis</i> St. Hill. (<i>Aquifoliaceae</i>)	Leave	Caffeine (upto 2%)	South America
3.	Coffee Bean or Coffee Seed	<i>Coffea arabica</i> Linne' or <i>C. liberica</i> (<i>Rubiaceae</i>)	Seed	Caffeine (1-2%) Trigonelline (0.25%); Tannin (3-5%) Glucose & Dextrin (15%); Fatty Oil (trioleoglycerol) and palmitoglycerol (10-13%) Protein (13%)	Ethiopia, Indonasia, Sri Lanka, Brazil
4.	Coca Kola or Kolanuts	<i>Coca nitida</i> (Ventenat) Schott et Endlicher (<i>Sterculiaceae</i>)	Seed	Caffeine anhydrous ($\leq 1\%$)	Sierra Leone, Congo, Nigeria Sri Lanka, Ghana Brazil, Indonesia Jamaica
5.	Tea or Thea	<i>Camellia sinensis</i> Linne's O. Kuntze (<i>Theaceae</i>)	Leave or Leaf bud	Caffeine (1-4%) Gallotannic acid (15%) Volatile oil (Yellow) 0.75%	China, Japan, India, Indonesia Sri Lanka
6.	Cacao Beans or Cacao seeds	<i>Theobrome cacao</i> Linne' (<i>Sterculiaceae</i>)	Seed	Fixed Oil (35-50%) Starch (15%) Protein (15%) Theobromine (1-5%); Caffeine (0.07-0.36%)	Ecuador, Columbia Malasia, Curacao, Mexico, Trinidad Brazil, Nigeria Camerrons, Ghana Philippines, Sri Lanka

Figure 1 shows the basic nucleus of 'Xanthine' and 'Purine'; besides the three well-known members of the Xanthine family viz., **Caffeine**, **Theophylline** and **Theobromine**.




Fig. 1.1 The Xanthine and Purine Structures

Figure 1.2, illustrates the mode of synthesis of caffeine essentially from the same precursors present in *Coffea arabica* as the three purine alkaloids (see Fig 1.1) found in order biological systems which have been studied so far at length, either from a compound which may afford an active 1-carbon fragment (e.g., serine, methanol, glycine and formalin) or from formic acid.




Fig. 1.2 Mode of Synthesis of Caffeine

- Methionine along with the said four compounds act as active precursors of the three 'Methyl Groups' at N_1 , N_3 and N_7 positions respectively.
- Glycine is responsible for the contribution of C-4, C-5 and C-7,
- Carbon dioxide contributes C-6,
- N-1 is provided from aspartate, and
- N-3 and N-9 are derived from the amide nitrogen of glutamate.

Such elaborated and intensive studies of chemical constituents present in 'Natural Products' could only be feasible with the advent of various advancement in the field of 'Phytochemistry'.

However, it is pertinent to mention here that the scientific reasonings for the various age-old established characteristic medicinal properties have been adequately ascertained and determined in the past two centuries. A critical survey of literatures would reveal that a few chemical entities were not only identified but also known to the therapeutic armamentarium between the said era. A few typical examples are enumerated below in a chronological order, as stated in Table 1.2.

Table 1.2 Examples of Plant Constituents in Use from 1627 to 1830

S. No.	Period	Researcher	Chemical Entity	Remarks
1.	1627-1691	R. Boyle	Alkaloid(s) (probably)	Present in Opium
2.	1645-1715	N. Lemery (French Apothecary)	Alcohol	As a solvent in extraction processes
3.	1709-1780	A.S. Marggraf (German Apothecary)	Sugar	isolated from many plant sources including Sugar-Beet
4.	1742-1786	K.W. Scheele	Organic acids oxalic, malic, citric, gallic, tartaric and prussic (HCN)	isolated from natural sources.
5.	1805	Serturner (German Chemist)	Meconic acid	Present in Opium
6.	1811	Gomeriz (Portugese Chemist)	Cinchonine	Isolated from Cinchona Barks
7.	1817	Serturner (Geman Chemist)	Morphine	An alkaloid present in Opium
8.	1817	Pelletier and Caventou (French Chemist)	Strychnine	An alkaloid from Strychnos Nux Vomica
9.	1819	- do -	Brucine	- do -
10.	1820	Meissner	Veratramine	An alkamine from Green Hellebore.
11.	1830	–	Amygdalin	A cyanophore glycoside from Bitter Almond.

Considerable progress has been made in the nineteenth century when chemists seriously took up the challenge of synthesizing a plethora of organic compounds based on **‘biologically-active-prototypes’**. Some of these purely ‘synthesized compounds’ essentially possessed structures of ever increasing complexity; and later on, after systematic pharmacological and microbiological evaluations proved to be yielding excellent useful therapeutic results. Evidently, as most of these **‘tailor-made’** compounds having marked and pronounced therapeutic indices were found to be existing beyond the realm of **‘pharmacognosy’** or more specifically **‘phytochemistry’**—an altogether new discipline under the banner of **‘medicinal chemistry’** came into existence. However, this particular discipline almost remained dormant since the era of Paracelsus. But now, the **‘medicinal chemistry’** has acclaimed deserving wide recognition across the globe due to its own legitimate merit and advantages.

In short, *three* major basic disciplines became largely prevalent with regard to the development of drugs, namely:

- **Pharmacognosy:** embracing relevant information(s) with regard to medicines exclusively derived from natural sources, for instance: plants, animals and microorganisms,
- **Medicinal Chemistry:** covering entirely the specific knowledge not only confined to the science of ‘synthetic drugs’ but also the basic fundamentals of **‘drug-design’**, and
- **Pharmacology:** dealing particularly the actions of **‘drugs’** and their respective effects on the cardiovascular system and the CNS-activities.

Over the years, with the tremendous growth of scientific knowledge and valuable informations the three aforesaid disciplines have fully-emerged as '**complete sciences**' within their own spheres.

Though copious volumes of ancient literatures in Chinese, Egyptian, Greek, Unani and Indian (Ayurvedic) systems of herbal medicines were found to contain factual and invariably exaggerated claims of their therapeutic efficacies, yet when they are evaluated intensively on a scientific basis with the advent of latest analytical techniques, such as: FT-IR, NMR, MS, GLS, HPLC, HPTLC, X-Ray Diffraction, ORD, CD and UV-spectroscopy—it has adequately and promptly provided an elaborated structure of various **complex chemical constituents**. A few select typical examples of known compounds are given in Table 1.3.

1.2 IMPORTANCE OF NATURAL DRUG SUBSTANCES

In general, natural drug substances offer *four* vital and appreciable roles in the modern system of medicine thereby adequately justifying their legitimate presence in the prevailing therapeutic arsenal, namely:

- (i) Serve as extremely useful natural drugs.
- (ii) Provide basic compounds affording less toxic and more effective drug molecules.
- (iii) Exploration of biologically active prototypes towards newer and better synthetic drugs.
- (iv) Modification of inactive natural products by suitable biological/chemical means into potent drugs.

The aforesaid aspects shall be briefly dealt with in the sections that follows:

1.2.1 Serve as Extremely Useful Natural Drugs




On a recent survey conducted by the World Health Organisation (WHO) globally, around 20,000 medicinal plants are being used profusely either in **pharmaceutical industry** or in **folk medicines**. Interestingly, about 1.4% do possess well-established, widely—proven and broadly accepted unequivocally active constituents.

De Souza *et al.** in 1982 opined on a serious note that—“*the usual success rate of discovering new drugs from natural sources is solely based not only on the conception but also on the implementation of ingenious comprehensive strategies which invariably explore and exploit the untrapped potential of the natural sources*”. In fact, there are *four* ways by which the above objectives may be accomplished reasonably and legitimately, such as:

- (a) Isolation of novel genotypes from marine and terrestrial ecosystems,
- (b) Genetic engineering: creating novel and altered genotypes,
- (c) Biochemical manipulation of selected pathways, and
- (d) Supersensitive and specific selection techniques and evaluation for varied bioactivities.

* De Souza, NJ *et al.*, *Annu. Rep. Med. Chem.*, **17**, 301, 1982.


Table 1.3 Examples of Chemical Constituents Present in Herbal Plants

S.No.	Common name(s)	Biological origin (Family)	Chemical constituents	Distribution	Uses
1.	Artemisinin or Qinghaosu	Artemisia annua Linne, Linne, (Asteraceae)	 <p style="text-align: center;">ARTEMISININ</p>	China	Treatment of cerebral malaria; Active against both chloroquine sensitive and chloroquine-resistant strains of <i>Plasmodium falciparum</i>
2.	Doxorubicin or Adriamycin and Daunorubicin or Cerubidine	<i>Streptomyces coeruleorubidus</i> <i>Streptomyces peucetius</i> var <i>caesius</i>	 <p style="text-align: center;">DOXORUBICIN : R = OH; DAUNORUBICIN : R = H;</p>	—	Treatment of breast cancer, various types of carcinomas, acute Leukemia; Daunorubicin Treats acute Lymphocytic Leukmias.
3.	Ginkgo	<i>Ginkgo biloba</i> Linne (<i>Ginkgoaceae</i>)	 <p style="text-align: center;">GINKGOLIDE-A</p>	Eastern Asia Southeastern United States.	Ginkgolides A, B, C, and M inhibit platelet-activating factor (PAF); reduces capillary fragility and blood loss from the capillary vessels that may ultimately check ischemic brain damage.

(Contd.)

4. **Ginseng**

Panax quinquefolius
Linne and
Panax ginseng C.A. Mey
(Araliaceae)




American ginseng
(p.q.) in Eastern
United States and
Canada;
Asian ginseng (P.g.)
In Eastern Asia
presently cultivated
profusely in Korea,
Japan and the former
Soviet Union.

Known to possess
tonic stimulant
diuretic and
carminative
properties;
reported to act
significantly on
metabolism, CNS
7 endocrines;
Exhibits adaptogenic
(antistress) activity.

5. **Gum opium or
opium or
Poppy Seed or
Maw Seed**

Papaver somniferum
Linne or
Papaver album
Deendolle
(Papaveraceae)




Turkish Anatolian
plain extends to
Northern border of
Laos; India; China,
Democratic peoples
republic of Korea.

Strongly narcotic
and hypnotic;
Centrally acting
analgesic.




6. **Rauwolfia
seperntina**

Rauwolfia serpentina
Linne Bentham
(Apocynaceae)



India, Myanmar,
Sri Lanka; Vietnam;
Malaysia; Indonesia;
The Phillipines.


Treatment control
of hypertension;
As an antipsychotic
agent.

<p>7. Taxol or Pacitaxel or Pacific Yew</p>	<p><i>Taxol brevifolia</i> Nutt (<i>Taxaceae</i>)</p>		<p>Northwestern United States.</p>	<p>Treatment of metastatic carcinoma of the ovary after failure of first line or followup chemotherapy; Treatment of breast cancer after failure of combination chemotherapy for metastatic disease.</p>
<p>8. Curare or South American Arrow Poison</p>	<p><i>Strychnos castenaei</i> Weddell; <i>S. toxifera</i> Bentham; <i>S. crevauxii</i> G. Planchon (<i>Logniaceae</i>); and <i>Chondendron tomentosum</i> Ruiz et Pavon (<i>Menispermaceae</i>)</p>		<p>Orinpoco basin; Upper Amazon regions; Eastern Ecuadorian plateau.</p>	<p>As a disgnostic aid in myasthenia gravis; As an adjunct to electroshock treatment in neuropsychiatry to control convulsion caused due to tetanus and strychnine poisoning.</p>
<p>9. Yohimbine</p>	<p><i>Pausinystalia yohimbe</i> (K. Schum) Pierre (<i>Rubiaceae</i>)</p>		<p>West Africa.</p>	<p>Treatment of impotence in patients with vascular or diabetic problems.</p>

(Contd.)

10. **Cantharanthus**
or
Vinca

Cantharanthus roseus G. Don
(*Apocynaceae*)
Formerly designated as
Vinca rosea Linne'
It has a close resemblance to
Vinca minor Linne', commonly known as Periwinkle.



Madagascar,
India, South
America, Austria,
South Africa,
Europe, The
West Indies,
Southern
United States.

Treatment of
lymphocytic
lymphoma,
advanced testicular
carcinoma,
histiocytic
lymphoma,
myosis
fungoids,
Hodgkin's disease,
Kaposi's sarcoma
choriocarcinoma
and breast cancer
Unresponsive to
other diagnosis.

A few typical examples of drugs derived from natural sources and their respective uses are given in Table 1.4.

Table 1.4 Examples of Drugs from Natural Products













S.No.	Name	Biological Origin(s)	Isolation	Synthesis	Uses
1.	Atropine	<i>Atropa beladona</i> (Linne)	1831	1883	Spastic colitis, gastro enteritis, peptic ulcer; antispasmodic.
2.	Ergotamine	<i>Claviceps purpurea</i> (Fries)	1918	1961	To prevent or abort vascular headaches, (Migraine and cluster headache)
3.	Morphine	<i>Papaver somniferum</i> (Linne)	1805	1956	As narcotic analgesics strongly hypnotic.
4.	Prostaglandins (PGE ₁ & PGE ₂)	C-20 Lipid Metabolites <i>in vitro</i> from essential unsaturated fatty acids of food (Linoleic acid)	1962	1969	PGE ₁ -certain congenital heart defects as a gastric antisecretory and gastroprotective agent; PGE ₂ -for termination of second trimester pregnancies.
5.	Physostigmine	<i>Physostigma venosum</i> (Balfour)	1864	1935	In ophthalmology to treat glaucoma; decreases intraocular pressure.
6.	Quinine	<i>Cinchona succirubra</i> (Pavon et Kloyzsch)	1820	1944	For treatment of malarial fever.
7.	Scopolamine (Hyscyamine)	<i>Atropa belladonna</i> , <i>Datura stramonium</i> , <i>Hyoscyamus niger</i> ; (or Egyptian henbane)	1881	1956	As CNS-depressant; in motion sickness; in preanaesthetic sedation; in obstetric amnesia along with other analgesic to calm delirium.

1.2.2 Provide Basic Compounds Affording Less Toxic and More Effective Drug Molecules


The numerous examples of naturally occurring plant products that serve as prototypes for other medicinally potent compounds either having closely related structures prepared exclusively by semisynthetic routes or possessing relatively simpler (less complex) purely synthetic structural analogues have been adequately described in various literatures. A few interesting examples of such compounds shall be described in this context as under:

1.2.3 Exploration of Biological-Active-Prototypes towards Newer and Better Synthetic Drugs

A plethora of better synthetic drug models gained considerable recognition in the therapeutic arsenal that were solely derived on the **biologically-active-prototypes**. It is, however, pertinent to mention here that these synthetic models not only possessed similar and better therapeutic index but also

S. No.	Natural	Semisynthetic	Synthetic
1.	Morphine (Narcotic Analgesic) 	Hydromorphone 	Methadone  Propoxyphene  Ibuprofen 
2.	Salicin and salicylic acid (Analgesic)  	Acetyl salicylic acid (Aspirin) 	Ibuprofen 
3.	Ephedrine (Adrenomimetic) 	Phenylpropanolamine 	Tetrahydrozoline 


(Contd.)



Note: The dotted areas in the above chemical structures show the presence of essential characteristic features in the natural (biological-active-prototype), semisynthetic and synthetic models.

exhibited fewer side effects than their corresponding naturally occurring constituents. A few typical examples are enumerated below:


(a) *Procaine from Cocaine—As Local Anaesthetic:*



(b) *trans-Diethylstilbestrol from—Estradiol—as Estrogenic Hormone:*



(c) *Chloroquine from Quinine—as Antimalarial:*




1.2.4 Modification of Inactive Natural Products by Suitable Biological/Chemical Means into Potent Drugs

This particular role of natural products is not only distinctly different from the rest, as discussed in section 1.2.1 through section 1.2.3, but also has its prime importance by virtue of the fact that certain constituents present in them do not exhibit any significant biological activity or chemical means surprisingly give rise to quite effective and potent drugs that are not easily obtainable by other known methods.


The following examples will expatiate the above facts squarely:

Examples:

1. Vitamin* A from—carotene (isolation from carrots *i.e.*, *Dacus carota*)




β -Carotene




Vitamin A (Retinol)


2. Taxol by conversion of 10-Dasacetyl**baccatin III** (Isolated from the needles of *Taxus baccata*):



3. Progesterone and Pregnenolone by conversion of Diosgenin (aglycone of Saponin Dioscin from *Dioscorea tokoro* Makino):




* 2 Moles of RETINOL are produced from 1 mole of *d*- β -Carotene which acts as a precursor of the former.



Pregnenolone

4. Hydrocortisone and Corticosterone from Stigmasterol (occurs abundantly as Phytosterol Mixture from Soyabeans and Calabar Beans)



In addition to the Third World nations, the technologically advanced countries like the United States have experienced a phenomenal change towards the acceptance of herbal medicines over the expanded OTC usages of such drugs. It is believed that in the Twenty First Century a quantum leap forward would be distinctly seen in the **world pharmaceutical market**. A few such projection of pharmaceutical products by the year 2001 may be glanced as detailed below:

S. No.	Name of Product(s)	Uses	Estimated Sales (USD)* /Year
1.	Plantago Seed or Psyllium Seed or Plantain Seed	Cathartic, Purgative	300 Million
2.	Scopolamine and Nicotine Patches	Motion sickness, Calming delirium, As anticholinergic agent	1 Billion
3.	Taxol	Antineoplastic agent	1 Billion
4.	Vinblastine and Vincristine	-do-	100 Billion

* USD: United State Dollar.

It may appear to be quite realistic and amazing that in the near future about **50% of the healthy-market-share** would be captured legitimately by **drugs belonging to the natural origin**.

It is not out of place to assert that on one hand science is advancing in a tremendous logarithmic progression towards gene-synthesis, rocket-fuels, super computers, electronic cash-transaction across the globe, fax machine, paperless offices, modern analytical computer-aided instruments, auto analysers for routine industrial analysis for on-going chemical and biological processes and final products metioulously designed and skillfully formulated life-saving drugs; while on the other hand the confidence of the people being restored at a steady pace towards the ancient herbal drugs right from the treatment of constipation to management and control of malignancies in human beings. Of course, the so-called '**Crude-Drugs**' are presently available in well refined and latest state-of-the-art packings as **over the counter (OTC) drugs** through chemists and druggists and super markets across the world. Perhaps that day is not too far when a common person will be tempted to grow medicinal plant in the kitchen garden rather than growing spring onions, lettuce, cucumber and french beans for their daily needs. It is a pity that the inhabitants of the modern society is virtually over-loaded by the usage of tonnes of chemicals used in the form of medicines for the cure of various ailments.

1.3 NATURAL DRUGS SUBSTANCES: CULTIVATION AND PRODUCTION

It is pertinent to mention here that the actual production of '**natural drug substances**' invariably adopts a number of different routes and methods based on the fact that their diversified origin as present in **plant, microbial and animal kingdom**. These *three* sources shall be discussed individually in the sections that follow:

1.3.1 Plant Products

Many countries in the world have a 'God-gifted' natural reserve of medicinal plants. Because of their judicious and cautious administration by the expertise of indigenous-systems of medicine people could survive and thus explore and conquer the world as per the historical evidence. In the past, lack of knowledge, non-availability of adequate storage facilities and proper scientific means and methods of cultivation and collection a good number of useful medicinal plants almost reached a point of not only depletion but also extinction. With the advent of scientific knowledge abundantly available these medicinal plants are now grown in an organised fashion whereby proper identification, right cultivation, due harvesting in the correct time of the year to yield maximum desired chemical constituents, and adequate prevention from spoilage and infestation due to improper storage. Nowadays, plant-extracts are available commercially across the globe so that these may be incorporated duly in several tried and tested herbal preparations. Various advanced '**analytical methods**' help a long-way in establishing the true picture of their quality, for instance: percentage of **Eugenol** present in *Clove oil* determines its quality; percentage of **Cineol** in *Eucalyptus oil* shows its purity; percentage of **Total Alkaloids** in *Datura stramonium* depicts its medicinal value.

A few countries in the world are noted for their supply of certain specialized plant extracts, namely:

India	:	Opium extracts;
China	:	Extract of <i>Artemisia annua</i> ;
United States	:	<i>Ginkgo biloba</i> extracts (GBE)
Korea, Japan	:	<i>Panax ginseng</i> extracts;
Madagascar	:	<i>Catharthus roseus</i> extracts;
Eastern Europe	:	Ergot produced by mechanical inoculation of rye plants with spores of a selected fungus.

1.3.2 Cell-Culture Techniques

It essentially involves the production of the ‘**desired secondary constituents**’ that caters for a viable alternative means of **drug-plant-cultivation**. Extensive studies have revealed that under the influence of ‘**stress-conditions**’, for instance: reacting with a suitable pathogen-may ultimately help in simulating the yield of some specific highly desired constituents in plant-cell suspension cultures. However, the actual slow growth of the *cell-biomass* possess a serious obstacle in the wide acceptance of this innovated technique. Perhaps the day is not too far when the **plant genes** which are responsible for coding enzymes catalyzing the desired biosynthetic routes may be converted to rather **more swiftly growing bacterial or fungal cells**.

1.3.3 Microbial Metabolites

A number of ‘**microbial metabolites**’ produced by well-defined process of fermentation give rise to certain very useful therapeutically potent drugs, especially the antibiotics and related antineoplastic agents as exemplified below:

(a) **As Antibiotics:** for instance:

- (i) **Chloromycetin** – from *Streptomyces venezualae* Bartz,
- (ii) **Erythromycin** – from *Streptomyces erythreus* (Walksman) Walksan & Henrici,
- (iii) **Gentamycin** – from *Micromonospora purpurea* MJ Weinstein *et al.*
- (iv) **Penicillin O** – from *Penicillium chrysogenum*,
- (v) **Streptomycin** – from *Streptomyces griseus* (Krainsky) Walksman et Henrici,
- (vi) **Tetracycline** – from *Streptomyces viridifaciens*.

(b) **As Antineoplastic Agents:** for examples:

- (i) **Dactinomycin** – from several *Streptomyces spp.*
- (ii) **Daunorubicin** – from *Streptomyces peucetius* G. Cassinelli; *P. orezzi*.
- (iii) **Mitomycin C** – from *Streptomyces caespitosus* (griseovinaceus)
- (iv) **Pilcamycin** (or **Mithramycin**) – from *Streptomyces argillaceus* n. sp. and *S. tanashiensis*

Figure 1.3, illustrates the outline of the fermentation process usually accomplished in a pharmaceutical industry whereby dried drugs are produced in a large scale. However, in certain specific instances, *per se* **cephalosporins**, the end product obtained by the fermentation process is routed through semisynthetic means to yield the desired pharmaceutical substance.




Fig. 1.3 Out-line of Fermentation Process of Drugs

It is pertinent to mention here that the production of '**genetically-engineered-drugs**', bears fundamentally a close resemblance to the various fermentation processes normally employed for the antibiotics. The major noteworthy difference in this specific instance lies in the fact that a gene controlling the production of the desired constituent is virtually transferred from its basic source to a fast-growing microbial cell-line whereby permitting the large-scale production in comparatively a much shorter duration.

However, it is rather a '**difficult task**' to isolate a **gene coding** for a particular antibiotic. Interestingly, in the **actinomycetin fungi**, the '*gene*' was separated conveniently from the **chromosomal genes** and cloned on naturally occurring plasmids. It has been observed that though plasmids are found in streptomycetes, only in the specific case of **methylenomycin*** *biosynthesis*, the extrachromosomal element essentially consists of several structural gene absolutely necessary for the production of antibiotic.

In general, a number of methods are employed to identify clones that usually harbour the plasmids carrying antibiotic-biosynthetic genes, namely:

- (a) Mutants that are found to be blocked at different steps in the **aminoglycosidic-production-pathways** are known and also available. These 'blocked mutants' may be employed as recipients for the prediction of respective genes from **shotgun-cloning-experiments**. Shotgun cloning is the isolation of a specific DNA sequence and subsequent screening for the desired phenotype. The plasmids eventually isolated from the transformants, wherein antibiotic-biosynthesis is restored by the cloned genes, would ultimately be introduced for maximizing the final yield.
- (b) The latest technique of **insertional mutagenesis** may be used effectively to obtain not only the mutant but also the cloned DNA in a single experiment.

* Member of a family of cyclopentenoid antibiotic related structurally to **sarkomycins**, and having *in vitro* activity Vs Gram positive and Gram negative organism.

- (c) As the enzymes that are intimately involved in the **biosynthesis of aminology** essentially possess relatively wider substrate specificities, the transfer of genes between such species that cause the production of various **aminoglycosides** were invariably utilised to generate **newer antibiotics**.

If genes that code for the synthesis of chosen precursors are duly cloned interspecifically many existing aminoglycosides may be produced by just a one-step-fermentation process. **Mutasynthesis*** has paved the way for the introduction of a plethora of interesting hybrids, for instance; **mutamicins**, **hybrimycins** and **hydroxygentamicin**, and

- (d) Conversion of **Amikacin (I)** from **Kanamycin (II)**:

Amikacin (I) is one of the most effective aminoglycosides. It may be produced chemically from **Kanamycin (II)** but this route is rather expensive and not cost-effective. However, an **aminoglycoside producing** strain of *Bacillus circularis* is capable of converting (II) into (I) by the addition of **hydroxyaminobutyric acid**. Thus, the interspecific transfer of this gene may be used to persuade successfully a kanamycin producing streptomycetes to afford (I) and this **recombinant DNA** route could prove to be an economical one.

Figure 1.4, summarizes the conversion of **Amikacin (I)** from **Kanamycin (II)** *via* **chemical and recombinant DNA routes**.




Fig. 1.4 Conversion of Amikacin(I) from Kanamycin(II)

1.3.4 Animal Derivatives

Animal Derivatives are also referred to as the **biologics** in literatures. They are invariably categorized into *two* groups, namely:

- (a) **Prepared from the blood of animals:** Such as: *serum*, *antitoxins* and *globulins*. These are usually obtained by the aid of certain specific treatment particularly carved to enhance the strength of desired constituents.

* Here, a mutationally blocked producing organism is able to incorporate a precursor analogue to produce a modified form of antibiotic.

(b) **Prepared from inoculation of suitable culture medium:** For instance: *vaccines, toxins, tuberculins* collectively termed as the '**microbial products**'. These products afford protection against a host of *causative pathogenic microorganisms*. They are produced by inoculating an appropriate culture medium that may consist of living tissue along the right pathogen. The resulting product is purified suitably, and may be used as a '**drug**'.

In conclusion, it may be emphasised that even in developed countries a variety of natural products enjoy their well-deserved recognition in the therapeutic arsenal. However, their actual and precise method of production is more or less an extremely individualized aspect. Many advanced countries like United States, Germany, France, Great Britain and most parts of Europe, where medical practice is found to be oriented toward the utilisation of **preferred-single chemical entities**, a major portion of natural drugs are treated to afford either one or more active components, such as:

Ginsengoside Rg ₁	from :	Ginseng;
Morphine	from :	Opium;
Reserpine	from :	Rauwolfia;
Taxol	from :	Pacific Yew;
Ergotamine	from :	Ergot;
Vincristine & Vinblastine	from :	Catharanthus;
Digoxin	from :	Digitalis;
Ginkgolide-A	from :	Ginkgo;
Artemisinin	from :	Qinghaosu, etc.

It is worthwhile to state that in **technologically-advanced nation** like China, India, Korea, Japan make use of a good number of herbal medicines having multicomponent entities with proven and advantageous therapeutic values. Mostly such preparations are available in the form of film coated tablets, capsules, syrups, powdered mixtures and dispensed under modern packing norms. Of course, there is a visible-upward tendency to adopt these preparations, from reputed manufactures in the Western World for the cure of a number of human diseases.

It is pertinent to mention here that the age-old practice of using **hydroalcoholic tinctures** and **fluid-extracts** have become more or less **obsolete nowadays**.

Various official compendia-like USP, NF, BP, Eur. P, Int. P, IP have duly incorporated the **standards of some purified natural products**; and hence, the quality in such cases may not be a significant concern at all.

1.4 PHYTOCHEMISTRY

'**Phytochemistry**' or the '**Chemistry of Natural Products**' may be strategically placed somewhere in between *natural product organic chemistry* and *plant biochemistry*. In fact, it is intimately related to the above two disciplines. However, in a broader sense phytochemistry essentially deals with the enormous different types of organic substances that are not only elaborated but also accumulated by plants. It is solely concerned with the following various aspects namely:

- Natural distribution.
- Chemical structure,
- Biosynthesis structure,
- Biosyntheses (or biogenesis),
- Metabolism, and
- Biochemical function.

Importantly, with the advent of most up-to-date analytical procedures the detailed phytochemical study of an unknown plant may be accomplished right from elucidation of chemical structure of pure constituents to the elaborated study of their biological characteristics.

Figure 1.5, illustrates the schematic development of a 'drug' from a 'medicinal plant' that may serve as a fruitful guide for various phytochemistry studies:




Fig. 1.5 Schematic Development of a 'Drug' from a 'Medicinal Plant'

It is, however, important to mention here that the ‘**living organism**’ of the universe (*i.e.*; plants, microbes and animals) may be regarded as mother nature’s splendid and huge BIOSYNTHETIC LABORATORY. It not only caters for the survival of the so called ‘living creatures’ of the earth in terms of providing a broad spectrum of **essential chemical constituents**, for instance: *proteins, fats, carbohydrates* and *vitamins* but also meticulously bring forth an enormous quantum of **physiologically active chemical entities**, such as: *alkaloids, glycosides, volatile oil (terpenoids), steroids, antibiotics, bitter principles, tannins* and the like.

The ‘**living organisms**’ give rise to a number of interesting phytochemical aspects over the years that may be viewed closely under the following *three* heads, namely:

- (i) Constituents,
- (ii) Drug Biosynthesis (or Biogenesis) and
- (iii) Classification

1.4.1 Constituents

The huge number of chemical substances that are present in the plant-kingdom and animal kingdom in one form or the other are termed as ‘**constituents**’. These **constituents** may be further divided into **two** main categories, namely:

- (a) Active Constituents, and
- (b) Inert Constituents.

1.4.1.1 Active Constituents

The chemical entities that are solely responsible for existing pharmacological, microbial or in a broader-sense therapeutic activities are usually termed as **active constituents**. Most drugs like: *alkaloids, glycosides, steroids, terpenoids, bitter principles* are the bonafide members of this particular category.

1.4.1.2 Inert Constituents

The chemical compounds, though present in plant and animal kingdom, which do not possess any definite therapeutic values as such but are useful as an adjunct either in the formulation of a ‘**drug**’ or in surgery are collectively known as **inert constituents**.

Examples:

- (a) **Plant Drugs:** The following **inert constituents** are invariably present in plants, namely:

Cellulose	: Microcrystalline forms of cellulose are used as combination binder-disintegrants in tableting. Colloidal cellulose particles aid in stabilization and emulsification of liquid;
Lignin	: To precipitate proteins, and to stabilise asphalt emulsions;
Suberin	: Esters of higher monohydric alcohols and fatty acids;
Cutin	: -do-
Starch	: As pharmaceutic aid <i>i.e.</i> ; tablet filler, binder and disintegrant;
Albumin	: Soyabean albumins—as emulsifiers;
Colouring Matters	: Cochineal for colouring food products and pharmaceuticals.

(b) **Animal Drugs:** The under mentioned **inert constituents** are mostly present in animals, namely:

- Keratin** : For coating “**enteric pills**” that are unaffected in the stomach but dissolved by the alkaline into intestinal secretions;
- Chitin** : **Deacylated chitin (chitosan)**—for treatment of water; sulphated chitin—as anticoagulant in laboratory animals.

It has been observed that the very presence of ‘*Inert Constituents*’ either act towards modifying or check the absorbance and the therapeutic index of the ‘*active constituents*’.

Obviously, to get at the right active constituents one has to get rid of the host of ‘inert constituents’ by adopting various known methods of separation, purification and crystallization. Therefore, most literatures invariably refer to the former as ‘secondary’ plant products.

The presence of these **secondary plant products (active constituents)** are governed by **two** school of thoughts, namely:

- (a) **Superfluous Metabolites:** *i.e.*, substances that have no value as such and perhaps their presence are due to the lack of excretory mechanism in them and ultimately result as the ‘**residual lock-up**’ *superfluous metabolites*, and
- (b) **Characteristic Survival Substances:** *i.e.*, substances which exert a positive survival value on the plant wherein they are actually present. They offer more or less a ‘**natural defence-mechanism**’ whereby these host plants are survived from destruction owing to their astringent, odorous and unpalatable features.

Examples: Poisonous alkaloidal containing plants; astringent containing shrubs; and pungent volatile oil-containing trees etc.

1. Genetic Composition (or Heredity): In reality, **genetic effects** exert both qualitative and quantitative alterations of the active constituents in medicinal plants.

Examples:

(i) **Eugenol:** It is naturally present in two different species in varying quantities as follows:

Eugenia caryophyllus (Sprengel) Bullock et Harrison: 70–95%

Syzygium aromaticum (L.) Merr et L.M. Perry: Not less than 85%.

(ii) **Reserpine-rescinamine group of Alkaloids:**

Rauvolfia serpentina (Linné) Bentahm: NLT* 0.15%;

Rauvolfia vomitoria Afzelius (from Africa): NLT 0.20%;

(*NLT: Not less than)

(iii) **Rutin:**

Fagopyrum esculatum Moznch : 3-8%;

Sophora japonica Linné : 20%;

(iv) **Menthol:**

Mentha piperita L. : 50-60%

Mentha arvensis Linnévar : 75-90%

(Japanese Mint Oil)

2. Environment Factors:

The **environment factors** largely contribute to the quantitative aspect of secondary constituents *i.e.*, active constituents. It is pertinent to mention here that medicinal plants belonging to the same species which are phenotypically identical *i.e.*, they essentially bear a close resemblance with regard to their form and structure, may not, however, genotypically be the same *i.e.*, possessing the same genetic composition. This particular natural phenomenon evidently gives rise to an altogether marked and pronounced difference in their chemical composition, specially with reference to active constituents. In a more logical and scientific manner it may be said that these plants categorically belong to different chemical races.

Example:

- (i) **Ergotamine:** Modified strains of *Claviceps purpurea* (Fries) have been developed, exclusively for field cultivation, that are capable of producing nearly 0.35% of **ergotamine** (in comparison to the normal one producing NLT 0.15% of total **ergot alkaloids**).
- (ii) **Eucalyptol** (*Syn: Cineole, Cajeputol*): It is present in the fresh leaves of *Eucalyptus globus* Libillardiere to the extent of 70–85%. It has been observed that the chemical races of some species of **Eucalyptus invariably** display significant variations in the content of eucalyptus and related components present in the essential oils.

There are a number of environmental factors which may afford considerable changes in active plant constituents, for instance: *composition of soil* (mineral contents); *climate* (dry, humid, cold); *associated flora* (*Rauwolfia serpentina* and *R. vomitoria*) and lastly the methods of cultivation (using modified strains, manual and mechanical cultivation). For a specific instance it may be recalled that a soil rich-in-nitrogen content evidently gives rise to a relatively higher yield of alkaloids in the medicinal plants; whereas a soil not so abundant in nitrogen content and grown in comparatively dry zones may yield an enhanced quantum of volatile oil.

- 3. **Ontogeny (or Ageing of Plant):** The age of a medicinal plant has a direct impact on the concentration of the '*active constituent*'. It is, however, not always true that older the plant greater would be the active principal.

Example:

- (i) **Cannabidiol:** It is present in *Cannbis sativa* L. (*C sativa* var. Indica Auth), possessing euphoric activity; and its content attains a maximum level in the growing season and subsequently the decline commences gradually. Interestingly, the concentration of **dronabinol** (or **tetrahydrocannabinol**) starts to enhance reciprocally till the plants gets fully matured.
- (ii) **Morphine:** The well-known narcotic-analgesic present in the air-dried milky exudate collected by incising the capsules of *Papaver somniferum* Linné or *P. album* Decandolle is found to be the highest peak just 2 to 3 week after flowering. An undue delay in harvesting from this '*critical-period*' would ultimately result into the decomposition of morphine. It is worth to be noted that a prematured harvesting of latex would certainly enhance the content of allied alkaloid like **codeine** and **thebaine**.

In short, it is a prime importance to affect the harvesting of medicinal plant at the right time so as to maximise the yield of the active principal.

1.4.1.3 Drug Biosynthesis (or Biogenesis)

In the recent past, a good deal of well-deserved importance and recognition have been attributed to the exclusive study of the biochemical pathways that precisely lead to the formulation of ‘*active constituents*’ otherwise referred to as the **secondary constituents** mostly employed as *drugs*. This specific study is normally termed as **Drug Biosynthesis or Biogenesis**.

As a ‘**medicinal chemist**’ is required to know the synthesis of **chloroquine**—an antimalarial drug from pure synthetic compounds, a ‘**phytochemist**’ is supposed to know the biogenesis of **quinine** in the cinchona bark. With the advent of isotopically labelled organic compounds known in the early fifties it was quite possible to establish scientifically that the host of amino acids along with their corresponding derivatives more or less acted as precursors of complex alkaloids. However, these logical studies confirmed the earlier hypothesis stated above by Trier in 1912.

Figure 1.6, summarizes the various *biosynthetic pathways and their inter-relationships* that ultimately lead to the formation of different kinds of **secondary constituents** (*i.e.*, **active constituents**) belonging to the plant kingdom which are invariably employed as drugs having *potent therapeutic index*.




Fig. 1.6 Biosynthetic Pathways and their Interrelationships

1.4.1.4 Classification

A plethora of pharmacologically active naturally occurring substances derived from ‘**medicinal plants**’ essentially comprise of rather large and complex molecules that invariably possess one or more than one of the chemical functional moieties which are responsible for attributing characteristic

features for alcohol, phenols, esters, aldehydes, ketones, oxides and organic acids. In fact, the aforesaid chemical groups are often attached to the molecular skeletons (*e.g.* **aromatic, heterocyclic compounds**) of noticeable diversified nature and complexity.

In the light of the following *two* observations, the phytochemical classification is eventually done on a more rational and broader perspective:

- (a) **Morphine** and **salicylic acid** has one phenolic—OH group in their molecule but structurally they are world-apart, and
- (b) **Essential (or volatile) oils** mostly contain a mixture of substances, such as: *hydrocarbons, ketones, aldehydes, and terpenes.*

Therefore, ideally the phytochemical classification is solely based on the types of plant constituents present in the natural products, namely:


- (i) Comprising of C and H only,
- (ii) Comprising of C, H and O only,
- (iii) Comprising 'O' into heterocyclic rings,
- (iv) Comprising of N, S and P,
- (v) Mostly containing Nitrogen
- (vi) Comprising of diversified chemical entity, and
- (vii) Mixtures.

The above phytochemical classification will be further expatiated with the help of some typical examples from the domain '**pharmacognosy**' along with their structures, wherever possible, as under:


1.4.1.4.1 Comprising of C and H only: They essentially consist of hydrocarbon present in the natural products.

Examples:


- (a) **β -Myrcene:** It is an **unsaturated acyclic hydrocarbon** found in *oil of bay, verbena, hop* and others.




- (b) **Ocimene:** It is also an **unsaturated acyclic hydrocarbon** found in the essential oil distilled from the fresh leaves of *Ocimum basilicum* L. and from the fruits of *Evodia rutaecarpa* (Juss.) Hook & Thoms. It exists in two modifications and forms. The *cis*- and *trans*-refers to the stereochemistry at the double-bond between C-3 and C-4.




- (c) ***p*-Cymene (Dolcymene)**: It is an **aromatic hydrocarbon** and occurs in a number of essential oils *e.g.*, sage, lemon, thyme, nutmeg, coriander, origanum, cinnamon.



- (d) **Limonene (Syn: Cinene, Cajeputene, Kautschin)**: It is an **alicyclic hydrocarbon** and further classified into *monocyclic terpene*. It occurs in various ethereal oils particularly in oils of lemon, orange, caraway, dill and bergamot.



- (e) **α -Pinene**: It is also an **alicyclic hydrocarbon** and further classified into *bicyclic monoterpene*. *d*- α -pinene obtained from Port Oxford Cedar Wood Oil (*Chamaecyparis lawsorliana* Parl.). *l*- α -Pinene obtained from Mandarin Peel Oil (*Citrus reticulata* Blando).




1.4.1.4.2 Comprising of C, H, and O only: A wide spectrum of plant constituents containing C, H and O have been identified.


Examples:

(A) Alcohols:

- (a) **Geraniol (or Lemonol)**: It is an **olephenic terpene alcohol** constituting the major portion of oil of rose and oil of palmarosea. It is also found in many volatile oils, for instance: citronella, lemon grass *etc.*





- (b) **Menthol (or Peppermint Camphor):** It is a **monocyclic terpene alcohol** obtained from **peppermint oil** or other **mint oils** or prepared synthetically on large scale by carrying out the hydrogenation of thymol.




(B) Aldehydes:

- (a) **Citral:** It is an aliphatic terpene aldehyde present in oil of lemon grass, lemon, lime, ginger root and in the oils of several *Citrus species etc.* The **citral** from natural sources is a mixture of *two* isomers **geraniol** and **neral**.



- (b) **Vanillin:** It is a **cyclic terpene aldehyde**. It occurs in vanilla in potato parings, in Siam benzoin, Peru balsam, clove oil etc. It is made synthetically either from **guaiacol** or **eugenol**; also from waste (**lignin**) of the wood pulp industry.




(C) Ketones:

- (a) **Carvone:** It is a **monocyclic terpene ketone**. *dl*-Carvone is found in *gingergrass oil*; *d*-carvone is found in caraway seed and dill seed oils, *l*-carvone is found in *spearmint and kuromoji oils*.
- (b) **Camphor:** It is a **bicyclic terpene ketone**. It naturally occurs in all parts of the camphor tree, *Cinnamomum camphora* T. Nees & Ebermeier; while 3/4th of the camphor consumed in USA is manufactured from **pinene** as the racemic form.


***d*-camphor:** is found in oil of sassafras, rosemary, lavender and sage;

***l*-camphor:** is found in lavender and artemisia;

***dl*-camphor:** is found in oil of sage and in oil of *Chrysanthemum sinense* var. japonicum.




Carvone



Camphor


(D) Phenols:

- (a) **Thymol:** It is a **monocyclic phenol**. It is obtained from the volatile oil of *Thymus vulgaris* L. and *Monarda punctata* L. and several spices of *Ocimum*. Commercially it is synthesized from *p*-cymene, *m*-cresol and piperitone.




Thymol

- (b) **Eugenol (or Allyguaicol):** It is a **dihydric phenol** and is the main constituent of several important essential oils, such as: **oil of clove, oil of cinnamon leaf, oil pimenta.**




Eugenol

- (c) **Myristicin:** It is a **trihydric phenol** occurs in **oils of nutmeg, mace, French parsley, dill oils and carrot.**



Myristicin

(d) **Apiole:** It is a **tetrahydic phenol** which occurs in **Dill oil** (*Anethum graveolus L.*) and known as Apiole (Dill); and also in **Parsley oil** (*Petroselinum sativum* Blanchet, Sell) and termed as Apiole (Parsley).




(E) Quinones:

Examples:


Anthraquinone Glycosides: A plethora of glycosides having aglycone moieties related to anthracene are present in such drugs as aloe, rhubarb, senna, frangula and cascara sagrada. In general, the glycosides on hydrolysis give rise to corresponding aglycones which are di-, tri-, or tetra-hydroxyanthraquinones or invariably structural modifications of these compounds.

Examples: Frangulin-A upon hydrolysis yields **emodin** and **rhamnose** as shown below:




(F) Acids:

(a) **Caffeic acid:** It is the constituent of plant and isolated from green **coffee beans**. It probably occurs in plants only in conjugated forms *e.g.* **chlorogenic acid**.




- (b) **Ferulic acid:** It is widely distributed in small amounts in various plants species. It is isolated from *Ferula foetida* Reg.



Ferulic acid


(G) Esters:

- (a) **Pyrethrins (Pyrethrin I & Pyrethrin II):** It is the active insecticidal constituents of pyrethrum flowers.



Pyrethrins


- (b) **Methyl Salicylate:** It is present in a number of oils, namely: **wintergreen oil, betula oil, sweet birch oil, teaberry oil.**



Methyl Salicylate


(H) Lactones:

- (a) **Podophyllotoxin (Syn: Condyline, Podofilox, Martec):** It is an *antineoplastic glycoside* found in the rhizomes of North American *Podophyllum peltatum* L.




Podophyllotoxin

- (b) **α -Santonin:** It is an *anthelmintic* isolated from the dried unexpanded flower heads of *Artemesia maritima* L., sens lat.

 α -Santonin


(I) Terpenoids:

- (a) **Gibberellins:** It represents a class of *plant growth hormones* first isolated from the cultures of *Gibberella fujikuroi* (Sawada) Wollenweber.



Gibberelane


- (b) **Primaric Acid:** It is obtained from American rosin, French galipot and from *Pinus maritima* Mill.



Pimaric Acid


(J) Carotenoids:

- (a) **Xanthophyll (Syn: Vegetable lutein; Vegetable lutenol; Bo-Xan):** It is one of the most widespread **carotenoid alcohol** present in nature. It occurs in egg-yolk, nettles, algae, and petals of many yellow flowers. It also occurs in the coloured feathers of birds.



Xanthophyll


- (b) **β -Carotene:** It is most abundantly distributed in the plant and animal kingdom. In plants it occurs invariably with chlorophyll. It acts as a **precursor of Vitamin A**. It was first isolated from carrots and hence bears its name. It usually represents the nature's 'red' colouration in plant kingdom.



β -Carotene


(K) Steroids:

- (a) **Cevadine:** It is one of the **steroidal alkaloids** obtained from *Veratrum viride*. American or Green hellebore from its dried rhizome and roots.




Cevadine

- (b) **Digitoxin:** It is a **cardiotonic steroidal-glycoside** obtained from *Digitalis purpurea L*; *D. lanata* and other species of Digitalis. About 10 Kilo leaves yield only 6 Grams of pure **digitoxin**.



Digitoxin

- (c) **Ergosterol:** It is usually obtained from yeast that synthesizes it from simple sugars such as glucose. The damp yeast yield about 2.5 g ergosterol; however, the particular variety of yeast is very important.




Ergosterol

1.4.1.4.3 Comprising of 'O' into Heterocyclic Rings: There are a number of natural plant constituents that essentially possess an oxygen atom into the heterocyclic ring-system. A few typical example are cited below to initiate some commendable interest in the domain of "**phytochemistry**":

1.4.1.4.3A Furan-based Constituents: These constituents are derived from *five-membered heterocyclic ring 'furan'*, namely:


- (a) **Furfural (or 2-Furfuraldehyde):** It is a **heterocyclic aldehyde** that usually occurs in the first fraction of many essential oils, belonging to the natural order-**Pinaceae**, for instance: *Pinus palustris* (**Pine oil**) and **cade oil**. It is also present in **oil of orris rhizome, clove oil, petit-grain, lavender and cinnamon oils**.



Furfural


1.4.1.4.3B Pyran-based Constituents: They are derived from *six-membered heterocyclic ring 'pyran'*, namely:

- (a) **Dicoumarol (Syn: Dicoumarin; Dufalone; Melitoxin):** It was originally isolated from **sweet clover** (Improperly cured Mililotus hay).




Dicoumarol

- (b) **Umbelliferone (Syn: Hydrangin; Skimmetin):** It is present in many plants and obtained by the distillation of resins belonging to the natural order **Umbelliferae**. It is the **aglucon** of *skimmin*.




Umbelliferone

- (c) **Meconic Acid (Syn: Oxchelidonic acid):** It is obtained from **opium** *i.e.*, *Papaver somniferum* which contains 4 to 6% of **meconic acid**.



Meconic Acid

- (d) **Coumestrol:** An *estrogenic factor* occurring naturally in forage crops, especially in **ladino clover** (*Trifolium repens* L.), **strawberry clover** (*T. fragiferum*) and **alfalfa** (*Medicago sativa* L.).




Coumestrol


1.4.1.4.3C Flavan based Constituents:

- (a) **Caechin (Syn: Catechol; Cyanidol):** It is a flavonid found primarily in **higher woody plants** as (+)-**catechin** along with (-)-**epicatechin** (*cis*-form).

Source: From **mahogany wood** and **catechu** (gambir and acacia).

*d*-Catechin


- (b) **Leucocyanidin (Syn: Flavan; Leucocyanidol):** It is obtained from the petals of **Asistic cotton flower** (*Gossipum spp.*) Stephens; *Butea frondosa* Koen ex Roxb. and **taxifolin**.



Leucocyanidin

1.4.1.4.3D Phenylbenzopyrylium based Constituents:


(a) **Cyanidin Chloride:** It is isolated from **bananas** and prepared by the reduction of **quercitin**.



Cyanidin Chloride

1.4.1.4.3E Carbohydrates: There are several examples of well known compounds belonging to the **carbohydrates**:

(a) **Glucose:** It occurs naturally in the free state in fruits and other parts of plants, it is also found in the combined form in **glucosidase**, in **di- and oligosaccharides**, in the **polysaccharides** (cellulose and starch) and in **glycogen**.



Glucose


(b) **Algin (Syn: Kelgin; Allose; Protanol):** It is a *gelling polysaccharide* extracted from **giant brown seaweed** [**giant kelp** (*Macrocystis pyrifera*) (L.) Ag]; **horsetail kelp** [*Laminaria digitata*) (L.) (Lamour)]; and **sugar kelp** [*Laminaria saccharina*) (L.) (Lamour)].

(c) **Pectin:** It is a **polysaccharide** substance present in cell walls of all plant tissues that functions as an *intercellular cementing materials*. Orange and lemon rind serve as the richest source of pectin and contains about 30% of the same.

1.4.1.4.4 Comprising of N, S and P: It can be divided into *three* groups, namely:


(A) Comprising of N:

(a) **Amygdalin:** The name **amygdalin** is currently used interchangeably with '**lactrile**'. It is a *cyanogenic glycoside* that occurs in seeds of *Rosacea*, principally in bitter almonds and also in peaches and apricots.




Capsaicin

- (b) **Trigonelline (Syn: Coffearin; Gynesine; Trigenolline):** It occurs in the seeds of *Trigonella foenumgraecum* L., in coffee beans, in the seeds of *Strophanthus spp.* and the *Cannabis sativa* L., Besides in seeds of many other plants. It is also found in **jellyfish** and in **sea urchin**.



Trigonelline


- (c) **Trimethylamine:** It occurs as a degradation product of nitrogenous plant and animal substances. It is widely distributed in animal tissue and especially in fish.



Trimethylamine


1.4.1.4.6 Comprising of Diversified Chemical Entity: The naturally occurring plant products invariably represent a class of entirely diversified chemical entity and nature. A few typical examples are cited below:

- (a) **Thiamine Hydrochloride (Syn: Vitamin B₁; Aneurine Hydrochloride; Bivatin; Metabolin; Bedome; Bewon):** It occurs abundantly in plant and animal tissues, notably in rice husk, cereal grains, eggs, milk, green leaves, yeast, liver, tubers and roots.




Thiamine Hydrochloride


- (b) **Ascorbic Acid (Syn: Vitamin C; Cantaxin; Cevalin):** It is widely distributed in the plant and animal kingdom. However, the good sources are **fresh tea-leaves, citrus fruits, hip berries** and **acerola**. It was isolated from **lemons** and **paparika**.



- (c) **Chloramphenicol (Syn: Chloromycetin; Levomycetin; Klorita):** It is a broad-spectrum antibiotic obtained from cultures of the soil bacterium *Streptomyces venezualae*.



- (d) **Penicillium O (Syn: Panicillium AT):** It is an antibiotic produced by *Penicillium chrysogenum*.



1.4.1.4.7 Mixtures: A good number of naturally occurring plant substances contain a mixture of constituents, namely:

(A) Tannins:

(a) **Hydrolyzable Tannins** : **Examples:**

Chest Nut	:	Bark and wood;
Oak	:	-do-
Sumac	:	Leaves;
Turkish Tannins	:	Galls of <i>Cynips tinctoria</i>

(b) **Consensed Tannins** : **Examples:**

Eucalyptus	:	Bark;
Black catechu	:	Hearthwood;
Spruce	:	Bark;
Gambier	:	Leaves and Twigs

(c) **Pseudotannins** : **Examples of drug containing pseudotannins are as follows:**

Gallic acid	:	Rhubarb;
Catechins	:	Acacia; catechu; cocoa; guarana;
Chlorogenic acid	:	Nux-Vomica; coffee; mate;
Ipecacuanhic acid	:	Ipecacuanha
(B) Volatile Oils	:	Examples
(Syn: Essential Oils)		
Oil of Chenopodium	:	<i>p</i> -Cymene; α -terpinene; <i>l</i> -limonene; methadiene;
Oil of Cinnamon	:	Cinnamaldehyde; eugenol; cinnamyl acetate;
Oil of Cloves	:	Eugenol; acetyleugenol; caryophyllene; vanillin; furfural;
Oil of Bitter Orange	:	<i>d</i> -Limonene; citral; decyl aldehyde; linalool; terpineol; methyl anthranilate;
Oil of Juniper	:	Pinene; cadinene, camphene; terpineol; juniper; camphor

(C) Resins:**Examples:**

- (a) **Rosin (Syn: Colophony: Yellow resin):** It is obtained as a residue left over after distilling off the essential oil from the oleoresin obtained from *Pinus palustris* and other species of *Pinus*.
- (b) **Guaic (Syn: Guaiacum; Resin guaic) : It contains**
- | | | |
|---|---|----------------|
| α -and β -Guaiaconic acid | : | \approx 70%; |
| Guaiac acid + Guaiaretic acid + Related compounds | : | 11%; |
| Vanillin + Guaiac Yellow + Guaiac | : | |
| Saponin (Guaiacin) | : | 1.5% |

(D) Latex:**Examples:**

- (a) **Opium Latex:** It is the air-dried milky exudation from incised unripe capsules of *Papaver somniferum L.*, or *P. album* Mill. It contains about 20 alkaloids, constituting about 25% of the opium, meconic acid, sulphuric acid and lactic acid, sugar, resinous and wax-like materials.
- (b) **Euphorbia (Syn: Cat's hair; Snake weed; Queensland; Asthma weed; Pill bearing Spurge):** It is obtained from the dried herb of *Euphorbia hirta L.*, or *E. pilifera L.* It contains several resins and an unstable glucoside.

FURTHER READING REFERENCES

1. **'Biochemical Evolution of Plants'** in *Comprehensive Biochemistry*, Vol. 29A, M. Florkin. and E.H. Stotz (Eds.), Elsevier, Netherland, 1974.
2. **'Biochemical Systematics'**, in *Plant Taxonomy*, 2nd ed., V.H. Heywood, Edward Arnold, London, 1976.

3. Grifo F *et al.* in *Bio-diversity and Human Health*, Grifo F and Rosenthal J (Eds) Island Press, Washington D.C., 1997.
4. Hamburger M and Hostettmann K: **Bioactivity in Plants—The Link between Phytochemistry and Medicine**, In: *Thirty Years of Phytochemistry (1961-1991) Phytochemistry*, **30**(12), pp. 3849-3874, 1991.
5. **'Handbook of Medicinal Herbs'**, J.A. Duke, CRC-Press, London, 1st Edn., 2001.
6. Jack T: **Molecular and Genetic Mechanisms of Floral Control**, *Plant Cell*, **16**, S1-S17, 2004.
7. Jeevaratnam K *et al.*: **Biological Preservative of Foods—Bacteriocins of Lactic Acid Bacteria**, *Ind. J. of Biotech.*, **4**(4), 446-454, 2005.
8. Miller JS: in *Sampling the Green World*, Stuessy TF and Sohmer SH (Eds). Columbia University Press, New York, 1996.
9. Paul L Huang *et al.*: **Developing Drugs from Traditional Medicinal Plants: Chemistry & Industry**, pp. 290-293, 1992.
10. Radenbaugh K: **Syn Seeds: Applications of Synthetic Seeds to Crop Improvement**, CRC Press, Boca Raton (USA), 1993.
11. Topsell KBG: **Natural Product Chemistry: A Mechanistic, Biosynthetic, and Ecological Approach**, Apotekarsocieteten, Stockholm, 1997.
12. **'Toxicology and Clinical Pharmacology of Herbal Products'**. M.J. Cupp (Ed.), Humana Press, New Jersey, 1st Edn, 2000.

2

Pharmacobiotechnology

- Introduction
- Theory
- Important Means in Biotechnology
- Recombinant Proteins
- Biotechnology Vs Modern Pharmacy Practice
- Biotechnology Based Pharmaceuticals for the Millennium
- Biotechnology and Modern Drug Discovery
- Further Reading References

2.1 INTRODUCTION

In a broader-sense “**biotechnology**” may be defined as—‘*the use of organism or enzymes for the large-scale production of useful substances ranging from not only agricultural products, food products and environmental science but also in the field of medicinal compounds, vaccines and diagnostics*’.

Interestingly, Robbers *et al.** (1996) coined an altogether newer terminology called the **Pharmacobiotechnology** so as to specifically refer to the wide application of “**biotechnology**” to the development of *pharmaceuticals* or *pharmaceutically active substances*.

The 1980s proved to be a golden–era in the field of **biotechnology**. In this particular decade science has virtually conquered the peak of “**Everest**” when it really became absolutely possible to detect, isolate and decipher the large congregation and wide–spectrum of natural proteins that invariably play a major role in coordinating the various functions extremely critical and vital to human life and health. Thus for the first time the numerous complicated processes that were mostly responsible as the root-cause of a plethora of mysterious major and dreadful ailments have been unearthed successfully and hence, could be modulated duly.

In short, this upcoming, innovating and fool proof comparatively newer developing methodology has made an ever-lasting impact that more or less embraced the unique-top-notch blending of meaningful discoveries from various diversified fields, such as:

- Recombinant DNA–technology
- Molecular biology,
- DNA-alteration

* Robbers, J.E.; Speedie, M.K. and Tyler, E., ‘*Pharmacognosy and Pharmacobiotechnology*’, Williams & Wilkins, London, 1996.

- Gene-splicing,
- Genetic engineering,
- Immunology, and
- Immunopharmacology

into an wonderful perfect scientific blend of high-tech-state-of-the-art industry.

Though, at present the viability and potential of newer range of pharmacobiotechnologically developed products is almost facing a staggering fate *vis-à-vis* the stringent US-FDA regulation and Drug Laws in the other countries of the world, it is expected earnestly with a great hope and wishful desire that in the new millenium quite a few of them , which are at present under critical phases of trials, would see the light-of-the-day with a thunderous bang to improve the quality of life of the human suffering due to the host of dreadful ailments.

It is, however, pertinent to mention here that in the recent past a number '**biotechnology-based products**' have already made a gainful entry into the Western World. A few typical examples are mentioned below:

Actimmune^(R)	:	For chronic granulomatous disease,
Betaseron^(R)	:	For relapsing ,remitting multiple sclerosis,
Epogen^(R)	:	For treatment of anemia associated with chronic renal failure,
KoGENate^(R)	:	For treatment of haemophilia A,
Leupine^(R)	:	For autologus bone marrow transplantation,
OncoScint^(R) CR/OV	:	For detection,staging and follow up of colorectal and ovarian cancer, and
Proleukin^(R)	:	For renal cell carcinoma.

2.2 THEORY

Since 1953, the epoch making discovery of a 3D structure of DNA by the help of exclusive X-ray diffraction studies carried out by Watson and Crick* (1953) there has been a tremendous quantum growth in the development and application of **biotechnology**. These advancements shall be briefly highlighted at this juncture so as to ascertain the much–deserved-acclaim of this nescent technology not only in the field of drugs but also in various aspects of our day-to-day life, namely:

- (i) Mutation, Crossing-over and Recombination at Meiosis:
- (ii) Third revolution in modern medicine,
- (iii) Genetic code,
- (iv) Collection of specialized cells,
- (v) Reverse transcriptase,
- (vi) 3D-Proteins,
- (vii) From nervous systems to immune systems,
- (viii) Body's defence mechanism,
- (ix) PCR-in Forensic and Research,

* James Watson and Francis Crick, 1953 (Nobel Prize, 1962).

- (x) DNA-Metabolic Pathways,
- (xi) Recombinant vaccinia vector, and
- (xii) OKT3-Monoclonal Antibody.

2.2.1 Mutation, Crossing-over and Recombinant of Meiosis

‘**Mother nature**’ has been actively engaged in carrying out silently an wonderful task of ‘natural’ genetic experiments since the past several billion years. The outcome of these splendid and meticulous experiments are as follows:

Mutation (or Random Heredity Alteration)

It has been exploited judiciously in enhancing the yield of antibiotic and strain selection. Thus the initial yield of penicillin from **Penicillium notatum** Westling, amounting to 4 mg L⁻¹ and from **Penicillium chrysogenum** Thom, amounting to 40 mg L⁻¹ has been increased to a phenomenal extent of 21,000 mg L⁻¹ by the **utilisation of mutation and strain selection**.

Crossing Over

It essentially refers to the simultaneous breakage and exchange of corresponding segments of the homologous chromosomes. The relentless efforts of inbreeding (hyberdization) experiments invariably give rise to a better new species, such as:

- (i) *Improved strains of cereals*: wheat, corn, rice producing much more yield per acre,
- (ii) *Improved versions of fruits*: seedless-grapes, seedless-watermelons, larger and sweeter oranges, larger and bulky tomatoes,
- (iii) *Improved versions of vegetables*: larger potatoes, pumpkins, tubers, cucumbers, etc.,
- (iv) *Improved hyberdization*: Tangelo due to crossing the tangerine with the grapefruit,
- (v) *Improved hyberdization in animals*: Alsetian Dogs-due to cross breeding of a German shephard dog with a wild-wolf; mule-due to crossing a donkey and a horse.

Recombination at Meiosis (or Fertilization): The innovation of such processes have brought about a revolution to the status of present diversity of life on various parts on this globe.

2.2.2 Third Revolution in Modern Medicine

According to Sadee* (1987) not only the physical characteristic features but also the gross-structures of each and every organism basically originated by virtue of the genetic code inherited and actually present within the nucleus of each cell. The major building blocks responsible for the architectural design of the cell are due to the essential components like carbohydrates, lipids, proteins and nucleic acids. Eventually the **enzymes**, which are nothing but a special class of proteins, invariably build and use such types of molecules in the course of a cell-life, *viz.*, maturation, maintenance and finally reproduction. Deoxyribonucleic acid (DNA) is the central point wherein the code for building the proteins in a cell takes place. In fact, DNA is the **genetic blueprint** of an organism. It essentially

* Sadee, W. ‘A Third Revolution in Modern Medicine. The World and I, Washington Times, Washington D.C., Pt I (Nov), 178; Pt II (Dec), 162 (1987).

comprises of nucleotides that are appropriately connected together in a sufficiently lengthy structure which more or less looks like a ladder.

2.2.3 Genetic Code

The human *genome** contain appropriately three billion nucleotide units of which four different nucleotides essentially comprise of the bases, namely: *adenimine*, *cytosine*, *guanine* and *thymidine* that are compactly stored into the **chromosomes**. Importantly, these possess the genetic code for about one million different type of gene. Now, each of these genes controls and regulates the synthesis of a protein that is made up of a long chain-like-sequence of amino acids ranging in number from 50 to 1000. Nirenberg *et al.* (1966) and Khorana (1966) for the first time in the world accomplished successfully the determination of the **genetic code**, In fact, they proved scientifically the manner whereby the inherent nucleotide sequence present in a gene actually regulates the particular sequence wherein the requisite number (say 20) of individual amino acids shall be combined to produce a specific protein molecule. It has been observed that one single codon is comprised of three nucleotides placed in a series; and every codon represents one specific amino acid. Thus, the very sequential arrangement of a codons in the DNA, following transcription into messenger RNA (*i.e.*, mRNA), eventually establishes the particular sequence of amino acids that will ultimately give rise to a specific protein. In short, both **molecular biology** and **biotechnology** has one common goal to achieve, that is to decipher the manner by which the genes and their corresponding proteins help in the management and control of basic cellular processes.

2.2.4 Specific Sets of Genes in Each Individual Organ


It has been observed that each organ, which is nothing but a collection of specialized cells, possesses some very specific sets of genes. Since, essential organs of the body, such as: brain, heart, liver, tissue, blood etc is tantamount to carry out a certain exclusively particular set of jobs, they are supposed to be triggered or for activation and followed up by deactivation *i.e.*, turned “on” and “off” as and when required by the system.

Thus, under the supreme command of the genetic code of DNA and subsequently mediated by mRNA a broad-spectrum of proteins are duly generated by every individual cell in a continuous fashion. A good number of these highly specific and specialized proteins are consumed within the cell itself, whereas the remaining are secreted directly into the extra cellular fluid. It is assumed justifiably that over a certain long duration and bearing in mind the enormous possible permutation and combination of nearly 500 amino acids, that each organism would have evolved a reasonably large excess of unique proteins having optimized characteristic functions.

2.2.5 Reverse Transcriptase (RT)

The basic underlying principle of ‘**modern biology**’ rests on the fact stated below:

* **Genome:** The entire DNA capable of expressing all the genetic information in the cell.



In 1970, another landmark was established with the advent **reverse transcriptase (RT)** which was responsible for the actual conversion of its own genomic RNA into a double-stranded RNA. It is, however, pertinent to mention here that the latest trends in biotechnology profusely bank upon this particular enzyme (RT). A few typical and glaring examples of RT, sometimes also referred to as **cellular catalyst's** are enumerated as under:

- Produce the chemical building blocks of cell life, for instance: fats and carbohydrates.
- To carry out the digestion of food.
- Generate hormones which in turn regulate and monitor organism,
- Serve as fuel for energy production, and
- Production of important molecules like DNA.

2.2.6 3D-Proteins

In true sense, the proteins are responsible for the creation of the cell cytoskeleton which ultimately gives rise to an organised three-dimensional (3D) structure. Generally speaking, the proteins not only help in the transport but also regulate the movement of various molecules throughout the cellular structure. They are strategically located in the outer-cell-membrane and aid in the transportation of ions as well as nutrients across the cell-membranes. Proteins play various vital roles as described below briefly:

- (a) Regulate gene-activity by binding to DNA.
- (b) Both proteins and peptides (smaller fragments of proteins) are usually secreted by cells as hormones like **insulin** and as **neurotransmitters**.
- (c) Caters as carrier molecule such as hemoglobin to act as body's oxygen transport mechanism, and
- (d) Serve as receptor sites for various hormones which in turn tune up the cell-function as per the varying pattern of body's requirements.

2.2.7 From Nervous System to Immune System

It is virtually an open-secret that '**hormones**' not only influence directly a good number of specific-cell-surface-receptors but also more or less on all functions of the body right from the **nervous system to the immune system**. By virtue of the fact that these hormones are capable of exerting highly selective, potent and above all hitting the bullseye by affording a specific action on the selective target cells have gained a wide recognition as a most promising and viable candidate for a new generation-drug in terms of the '**magic-bullet**' hypothesis laid down by Ehrlich. These hormones have the following **merits**, namely:

- When given parenterally they approach the target receptors just on the outer-periphery of the cells without even penetrating the membranes, and

- Capable of binding to **cell-surface-receptors** intimately thereby activating the cell's specific function instantly;

Example: Interleukin-2, which is still an experimental anticancer substance, may trigger stimulation to certain immune cells so as to exert a direct influence on cancerous cell-growth.

2.2.8 Body's Defence Mechanism

The so called '**immune system**' of the human body actually governs its defence mechanism against all sorts of pathogenic invading organisms. The *two* vital factors that have a direct influence on the body's defence mechanism (immune system) are, namely:

- (a) **Memory**: the capability of recognizing and responding immediately to the previously encountered infections; and
- (b) **Specificity**: the capability of concentrating directly on the specific pathogens.

The various cells which are solely responsible to the spectrum of immune responses are – **Phagocytic cells (macrophages); B-cells; Suppressor T-cells; Helper T-cells; Killer T-cells** and **Natural Killer (NK) cells**; and **Memory cells**.

2.2.9 PCR-in Forensic & Research

The DNA research on the **polymerase chain reaction (PCR)** has turned out to be a very efficacious and useful technique in the specialised field of forensic as well as research applications. Many mysteries from the scene of murder, hitherto impossible to detect precisely, has now been made easy with the blood-strains collected from the victim. Likewise, the authenticity of fraternity may be established beyond any reasonable doubt with the help of DNA studies.

2.2.10 DNA-in Metabolic Pathways

The recent developments in the field of **metabolic engineering** has made a tremendous impact on the intermediary metabolism with a big bang. A few typical examples whereby DNA helps in the metabolic pathways are enumerated below:

- To facilitate and solve the branch-point-control problems,
- Introduction of identical enzymes obtained from different sources into the studied organism has brought to light not only improved and newer degree of flexibility but also introduced much better and acceptable metabolic characteristics into the older mechanism.
- To increase copies of a gene at a rate-controlling point,
- To remove a poisonous product by the addition of a single specific gene
- To accomplish an altogether new pathway into an organism that otherwise ceases short of the desired product by addition several genes.

2.2.11 Recombinant Vaccination Vector

It is now known that the **vaccinia virus** may be used as a **molecular vehicle** for transporting **foreign genes** into other organism. With the advent of both extensive and intensive research it is quite

possible to exploit this **recombinant vaccinia vector** to act as a vehicle for the production of **live vaccines** for otherwise difficult ones to produce.

2.2.12 OKT3–Monoclonal Antibody

Antibodies are regarded as the body's **missile defence mechanism**. These are large protein molecules produced by WBC which seek out and destroy dreadful foreign substances. They are used gainfully in matching donors and recipients in organ transplantation, in blood typing, in the measurement and identification of hormones, toxins and various antigens in blood and fluids. The remarkable use of monoclonals as a pinpoint attack on a cancerous cell or to eradicate cells left after a conventional **chemotherapy**. They may also be utilized to transport drugs, radioactive particles and toxins to such cells. Recently, **OKT3 monoclonal antibody** has been duly cleared and approved by the US-FDA for the treatment of acute renal allograft rejection.

2.3 IMPORTANT MEANS IN BIOTECHNOLOGY

There are a number of ways and means whereby this wonderful ever-expanding field of **biotechnology** has gained enough recognition to be identified as a top-brass zone of research. A few such important tools of technology shall be discussed in the sections that follows:

2.3.1 Recombinant DNA (rDNA)

It may be defined as—“**the hybrid DNA produced by joining pieces of DNA from different sources**”. It is usually designated as **rDNA**.

Recombinant DNA is normally formed outside the living cell with the aid of highly critical enzymes termed as ‘**restriction enzymes**’ which will essentially cleave the DNA at particular sites. Subsequently, another type of enzymes known as ‘**ligases**’ will help to insert the cut piece of DNA (called the ‘**insert**’) directly into the **Vector DNA** (*i.e.*, **viral DNA** or **plasmid**). The resulting vector DNA shall be capable of entering conveniently into a ‘**host**’ cell or otherwise called as **microorganism**.

It is pertinent to mention here that once the **foreign DNA** (or **vector DNA**) gains an entry into the microorganism (*i.e.*, host organism), it is known as a “**recombinant organism**”.

2.3.2 Restriction Enzymes

Restriction enzymes are also known as ‘**restriction endonucleases**’. They are found to ‘**digest**’ DNA into corresponding short strands at very particular sites which is exclusively governed by the following *two* factors, namely:

- (a) By a four-base-system *i.e.* four different nucleotides containing the bases – **adenine, cytosine, guanine** and **thymidine**; and
- (b) By a relatively larger stretch of **specific DNA sequence**.

Restriction enzymes, in fact just act as a pair of “**molecular scissors**”. Evidently, the longer the number of bases that command a particular cleavage-site for a specific enzymes, the less frequently it will afford a cut to a long strand of DNA.

Table 2.1 illustrates the examples of a few **restriction enzymes*** and their cleavage sites respectively:

Table 2.1 Examples of Restriction Enzymes

Name of Enzymes	Generating Organism	Cleavage Site
<i>Pvu1</i>	<i>Proteius vulgaris</i>	CG ATCG GCTA GC
Hpa1	Haemophilus <i>parainfluenzae</i>	GTT AAC CAA TTG
<i>Eco R1</i>	<i>Escherichia coli</i>	G AATTC CTTAA G
<i>Sau 3A1</i>	<i>Staphylococcus aureus</i>	GATCZ CTAG

Most restriction enzymes have either ‘**sticky ends**’ or ‘**blunt ends**’.

2.3.3 DNA–Ligase

It has been observed that the DNA pieces cut with the same enzyme shall possess ‘**sticky ends**’ which would anneal under the appropriate and favourable conditions when such ‘pieces’ are pooled together. The resulting annealed pieces usually exhibit only single strands (termed as ‘**nicks**’) at the specific sites where they were actually cleaved. When such mixed pieces are treated with an enzymes, termed as **DNA – ligase**, it will ultimately give rise to an **intact piece of recombinant DNA** by the combination of phosphodiester bond (at the ends of the DNA strands) with the complementary bases (**adenine, cytosine, guanine and thymidine**).

2.3.4 Cloning Vector

A **cloning vector** may be a **plasmid**** or a **bacteriophage*****. In **genetic engineering** (*i.e.*, **gene cloning techniques**) – a gene having a close resemblance to a particular protein shall be joined together with a **cloning vector** so as to enable it get transferred into a host cell.

* (a) **Restriction enzymes** are named by using the **first letter** of the **genus name** combined with the **first two letters** of the **species name**;

(b) **Strain** is designated by a capital letter (*e.g.* Eco R1) or an arabic numeral and a capital letter (eg Sau 3A1) and Roman numerals follow immediately to specify individual enzymes when a given microbe has more than one.

** These are self replicating, double-stranded, circular DNA molecules which are invariably found and maintained in the respective host-cell as an independent extrachromosomal moiety. They are normally characterised by specificity of host, size, and copy number.

*** They are viruses that infect bacteria. These are employed as cloning vectors. The foreign DNA is usually inserted into the viral genome in an analogous fashion. At times the genes could be eliminated from the bacteriophage to result a vector that might infect cells but not kill the infected cell; thus, the DNA may be expressed within the recipient bacterial cell.




Fig. 2.1 Genetic Map of Plasmid puc19

Figure 2.1 represents the genetic map for a **plasmid puc 19** that has been used in cloning where,

- A = Ampicillin,
- B = *lac Z* gene,
- C = Multiple cloning site
- D = *lac I* gene, and
- E = Origin of replication.

Here, the **plasmid** contains an ampicillin resistance gene (A) that permits only the selection of plasmid – bearing organism. The multiple cloning site (C) possesses a member of restriction enzymes sites. It essentially exists very much within the ***lac Z* gene (B)** which in turn encodes **β -galactosidase**. In fact, **isopropylethiogalactoside (IPTG)** induces B that eventually interferes with the binding process of the repressor proteins encoded by (D) to the (B) promoter. In this manner (B) undergoes depression ultimately leading to its subsequent *transcription* and *translation*. At this juncture, any foreign genes strategically inserted into the **multiple cloning site (C)** shall positively interfere with the production of **β -galactosidase activity**. However, these would be expressed by themselves either from the promoter belonging to the gene or from the ***lac Z* gene (B)** promoter.

2.3.5 Hybridization Probes

Hybridization probes constitute another extremely important and vital technique exclusively meant for **genetic engineering**. In reality, probe is nothing but a complementary **sequence of DNA** that is specifically labelled with anyone of the **three** different means namely:

- (a) A radioactive substance,
- (b) A fluorescent material, and
- (c) A chromagenic material.

Probe DNA or probes may be either **larger segments of DNA** or **synthetic oligonucleotides** or even **whole plasmids**. In usual practice, the DNA which is subjected to probe shall be conveniently bound to a solid support that could be either nitrocellulose or nylon. The product thus obtained is heated in a mixture along with the **probe DNA**. Subsequently, the strands would separate and reanneal thereby allowing the probe to bind with its complementary sequence. Consequently, the resulting product shall determine the very presence of that segment of DNA by rendering the **hyberdized DNA** either **chromogenic** or **fluorescent** or **radioactive**.

2.3.6 Cloning Process

Literally it refers to – ‘*a cutting used for propagation*’. In the present context cloning means, to make identical copies. The recent advances accomplished in the field of “*Biotechnology*” the **cloning process** has been exploited in the following *six* aspects, namely:

- (i) DNA–cloning,
- (ii) Cloning larger DNA fragments in specified cloning vectors,
- (iii) Cloning Eukaryotic DNAs in bacterial plasmids,
- (iv) Cloning Eukaryotic DNAs in phage genomes,
- (v) Cloning cDNAs, and
- (vi) Expression cloning.

The above diversified **cloning processes** shall be dealt individually as under:

2.3.6.1 DNA–Cloning

The **DNA cloning** is nothing but a broad based technique whereby large quantum of a particular DNA-segment are produced. Usually, the resulting DNA segment which is to be cloned is first linked to a vector DNA, that serves as a vehicle for carrying foreign DNA into a suitable host cell, such as the bacterium *Escherichia coli*. The **vector** (*i.e.*, *E. coli*) essentially contains sequences which in turn permits to be replicated within the host cell. In order to clone DNAs within bacterial hosts two types of vectors are commonly employed, namely:

- (a) The DNA segment to be cloned is introduced into the bacterial cell by **first** joining it to a plasmid and **secondly**, causing the bacterial cells to take up the plasmid from the medium, and
- (b) The DNA segment is joined to a portion of the **genome** of the bacterial virus lambda (λ) which is subsequently allowed to infect a culture of bacterial cells. Thus, a huge quantum of viral progeny are produced, each of which contains the foreign DNA segment.

It is, however, pertinent to mention here that by following either of the two methods stated above—the DNA segment once gets inside a bacterium, it will undergo the replication process with the bacterial (or viral) DNA and partitioned to the daughter cells (or progeny viral particles). In this manner, the actual number of bacterial cells which are actually formed.

Besides, **cloning** may also be employed as a versatile method to **isolate in a pure form any specific DNA fragment** amongst a relatively large heterogeneous population of DNA molecules.

2.3.6.2 Cloning Larger DNA Fragments in Specified Cloning Vectors

It has been observed that neither plasmid or **lambda phage** (λ) vectors are adequately suitable for **cloning DNAs** whose length is more than 20-25 kb*. This specific lacuna has revitalized the interest of researchers to look into the development several other vectors which might facilitate to clone much larger **segments of DNA**. However, the most important of these vectors are termed as **yeast artificial chromosomes (YACs)**

* [Kilobase (kb)]: A 1000 – base fragment of nucleic acid. A *kilo base pair* is a fragment containing 1000 base pairs.

YACs are nothing but artificial versions of a normal yeast chromosome. They normally comprise of all the elements of a yeast chromosome which are absolutely necessary for the specific structure to be replicated during **S-phase** and subsequently segregated to **daughter cells** during **mitosis**, including:

- One or more origins of replication,
- Having telomers at the ends of the chromosomes, and
- A centromere to which the spindle fibers may get attached during chromosome separation.

Invariably, the **YACs** are designed in such a fashion so as to provide essentially:

- (a) A gene whose encoded product permits those particular cells having the **YACs** to be selected from those that lack the element, and
- (b) The **DNA fragment** to be cloned like other cells, subsequently the yeast cells shall pick up DNA from their respective medium that caters for the path whereby **YACs** are introduced directly into the cells.

It has been observed that **DNA fragments** cloned in **YACs** range typically from 100kb to 1,000 kb in length. *Example:*

‘The **restriction enzyme** usually recognizes the eight-nucleotide sequence GCGGCCGC, which in turn specifically cleaves mammalian DNA into fragments approximately one million base pairs long’.

Fragments of this length can now be introduced conveniently into **YACs** and subsequently cloned within host yeast cells.

2.3.6.3 Cloning Eukaryotic* DNAs in Bacterial Plasmids**

A **foreign DNA** intended to be cloned is strategically inserted into the plasmid to give birth to a **recombinant DNA** molecule. However, the plasmid used for DNA cloning are exclusively the modified versions of those occurring in the bacterial cells. Consequently, the bacterial cells are able to take up DNA from their medium. This particular phenomenon is termed as ‘*transformation*’ and forms the basis for **cloning plasmid in bacterial cells**.

Figure 2.2 represents the DNA cloning using bacterial plasmids. First of all the recombinant plasmids each containing a different foreign DNA insert are added to a bacterial culture (*E. coli*) which has been previously treated with Ca^{+2} ions. These bacteria are gainfully stimulated to take up DNA from their respective surrounding medium upon exposure to a brief thermal-shock treatment yielding **plasmid DNA** (purified). Secondly, **human DNA** are also obtained in the purified form. Subsequent treatment of **human DNA** and **plasmid DNA** with **EcoRI***** result into the cleavage of human and bacterial DNA into various sized fragments. Now, these small fragments join together to yield recombinant DNAs with DNA ligase and thus give rise to the *plasmids*. These population of plasmids invariably contain various segments of **human DNA**. Incubation of these plasmids with

* **Eukaryote:** A cell or organism having a unit membrane–enclosed (true) nucleus and has no extracellular form.

** **Plasmid:** An extrachromosomal genetic element that is not essential for growth and has no extracellular form.

*** **EcoRI:** Enzymes designation for *E. coli* with recognition sequence G A A* T T C [arrow indicate the sites of enzymatic attack; indicate the site of methylation].




Fig. 2.2 DNA Cloning Using Bacterial Plasmids

E. coli cells under controlled experimental parameters ultimately yield plasmid that are free from *E. coli*. It has been observed that only a very small percentage of the cells are competent to pick up and retain one of the combinant replicate molecules. Once it is taken up the plasmid undergoes replication autonomously within the recipient and is subsequently passed on to its progeny during cell division. The isolated **recombinant plasmids** can then be treated with the same restriction enzymes used in their formation, that releases the **cloned DNA segments** from the remainder of the DNA which served as the vector. Thus, the **cloned DNA** can be separated from the **plasmid**.

2.3.6.4 Cloning Eukaryotic DNAs Phase Genomes

A *bacteriophage*, or more commonly a *phage* is a virus particle which infects a bacterial cell. In fact, a phage particle normally comprises of *two* essential components: first, a *phage head* that contains the genetic material and secondly, a *tail* through which the genetic material is injected into the host cell.

Interestingly, one of the most broadly explored of these phage particles, termed *Bacteriophage Lambda* [or *bacteriophage* (λ)], has more or less turned out to be a commonly employed **cloning vector***.

The **genome**** of lambda is a linear and double-stranded DNA molecule having 50kb length.

Figure 2.3 depicts the protocol for **cloning eukaryotic DNA** fragment in lambda (λ) phase.

In usual practice, the modified strain (**mutant***** employed in most cloning experiments contains two cleavage sites for the enzymes EcoRI that ultimately fragments the genome into *three* large segments. However, the two outer segments essentially contain all informations required for the infectious growth, whereas the middle fragment could be rejected conveniently and replaced suitably by a piece of DNA upto 25 kb in length.

It has been observed that the genes of eukaryotes are often split, with non-coding intervening sequences—known as *introns*, thereby separating the coding regions—termed as *exons*. The two outer segments undergo *splicing*[†] with **eukaryotic fragment** to result into the formation of recombinant DNA. Consequently, the recombinant DNA molecules can be packaged into phage heads *in vitro* and in turn these genetically engineered phage particle may be employed to infect host bacteria. Once gaining entry into the bacteria, the **eukaryotic DNA segment** is adequately amplified along with the viral DNA and subsequently packaged into an altogether new generation of virus particle that are released when the cell undergoes *lysis*^{††}. The released particle thus obtained infect new cells, and without any loss of time either a *plaque*^{†††} or a clear spot in the ‘*bacterial lawn*’ is visible distinctly at the site of infection. Each **plaque**, which is nothing but a **zone of lysis**, possesses millions of phage particle, each carrying a single copy of the same **eukaryotic DNA segment**. Interestingly, a single pertridish may accommodate more than 10, 000 different plaques.

2.3.6.5 Cloning cDNAs

It is pertinent to mention that the explanation of **cloning cDNAs** has been specifically restricted to cloning DNA fragment isolated from extracted DNA *i.e.*, genomic fragments. In other words, the isolation of a genomic DNA means the eventual isolation of a particular gene or a family of genes out of a pool of hundreds of thousands of unrelated sequences. Besides, it becomes more or less necessary to study the following different aspects during the course of isolation of **genomic fragment**, namely:

* **Vector**: A genetic element able to incorporate DNA and cause it to replicate in another cell.

** **Genome**: The complete set of genes present in an organism.

*** **Mutant** : A strain differing from its parent because of mutation.

† **Splicing** : The processing step whereby introns are removed and exons are joined.

†† **Lysis**: Rupture of a cell, resulting in a loss of cell contents.

††† **Plaque**: A zone of lysis or cell inhibition caused by virus infection on a lawn of cells.




Fig. 2.3 Sequence for Cloning DNA Fragments in Lambda (λ) Phage.

- Non-coding intervening sequences,
- Regulatory sequences flanking on either sides the coding portion of a gene,
- Different members of a multigene family that invariably lie very close in the genome,
- Evolution of DNA sequences, such as: duplication and DNA of various species *vis-a-vis* their rearrangement, and
- Interspersion of transposable genetic elements'.

There are **two** aspects which are very important with **cloning cDNAs**, namely:

- (a) Analysis of gene structure, and
 (b) Analysis of gene expression.

Figure 2.4 illustrates the manner by which cDNAs are synthesized for cloning in a plasmid.

In order to **clone cDNAs**, *first* of all a sizable population of mRNA is isolated; *secondly*, it is employed as a template to provide a single-stranded DNA complement; *thirdly*, the resulting product (single stranded) is duly converted to a double stranded population with the help of a DNA polymerase; and *fourthly*, they are finally combined with the desired vector. It is quite evident that essentially




Fig. 2.4 Synthesis whereby cDNAs get Cloned in a Plasmid.

mRNA populations typically consist thousands of altogether different species, and as with experiments employing genomic DNA fragments, the clones should be invariably screened to isolate only one particular sequence from a heterogeneous population of recombinant molecule.

From Figure 2.3, it may be observed that when polypeptide (A) and mRNA are annealed, it provides a small segment of primer attached to poly(A) to the tail of mRNA. Now, with the help of **reverse transcriptase** the primer to poly(A) gets fully developed. Alkali helps in the separation of DNA and RNA strands to give rise to fully developed primer alone, which on treatment with RNA polymerase 1 yields the combined product. The resulting product when digested with S1 nuclease two separate strands of the primer and poly(A) are obtained. Lastly, integrate cDNA into the **plasmid vector** that will produce a bacterium wherein DNA can be cloned.

2.3.6.6 Expression Cloning

For practical applications it is quite important that such systems must be available wherein the cloned genes may be expressed. In other words, **expression cloning** is an alternative method for identifying a *phage plaque* which essentially contains a particular cDNA. In this specific method the cDNA being cloned is inserted directly in the downstream region from a strong bacterial promoter, which adequately ensures that the foreign DNA is not only transcribed but also translated in the course of the infections process. Interestingly, those phage which has originally incorporated the gene being sought must form plaques that essentially possess the protein encoded by the gene. Further identification of the plaque is performed on replica plates of employing a labeled probe which binds particularly to the encoded protein. The antibodies serve invariably as the most commonly used probe to identify the desired cloned genes which have been critically located on the replica plate whereas the genes may be subsequently isolated from the viruses on the original plate.

2.3.6.7 Amplifying DNA: The Polymerase Chain Reaction (PCR)

The *conventional molecular cloning techniques* may be considered *in vivo* DNA-amplifying tools. Interestingly, the latest development in the field of **synthetic DNA*** has evolved an altogether new method for the rapid amplification of DNA *in vitro*, broadly termed as the **Polymerase Chain Reaction (PCR)**. In reality, PCR is capable of multiplying DNA molecules to the extent of a billion fold *in vitro*, thereby giving rise to huge amounts of very specific genes employed for various purposes, such as: **cloning, sequencing or mutagenesis**. In short, PCR utilizes the enzyme DNA polymerase, which eventually copies DNA molecules.

The polymerase chain reaction (PCR) for amplifying specific DNA sequences have been shown in Figure 2.5 [Stage–A through Stage–F]. These six stages have been duly explained here under:

Stage–A: The target genes (DNA–combined form) if *first* heated to affect the separation of the strands of DNA; *secondly*, a reasonably excess amount of two **oligonucleotide primers****, of which one is complementary strand, is added along with DNA-polymerase;

Stage–B: As the resulting mixture attains the ambient temperature, the excess of primers relative to the target DNA makes sure that most target strands anneal to a primer exclusively and not to each other. In this way, the primer extension ultimately gives rise to a copy of the original double-stranded DNA.

* **Synthetic DNA**—short fragments of DNA of specified base sequence and widely used in molecular genetics.

** **Primers:** A molecule (usually a polynucleotide) to which DNA polymerase can attach the first nucleotide during DNA replication.




Fig. 2.5 PCR for Amplifying Specific DNA Sequences (Stage-A through Stage-F)

Stage-C: Further follow up of *three* above mentioned steps sequentially viz; *heating*, *primer annealing* and *primer extension* results into the formation of a copy of the original double-stranded DNA. In other words, **DNA polymerase extends the primers employing the target strands as a template.**

Stage-D: Another primer extension of the resulting product yields the second double-stranded DNA.

Stage-E: The end product obtained from the previous step is subjected to incubation for a suitable duration; and the resulting mixture is heated once again so as to separate the strands. Subsequently, the mixture is brought to the room temperature whereby the primers aptly get hybridized with the complementary regions of newly synthesized DNA. Thus, the whole process is repeated. In this particular instance, the two additional PCR-cycles give rise to 8 and 16 copies, respectively, of the original DNA sequence.

Stage-F: It represents a plot between the number of PCR cycles (along the X-axis) and the copies of the target gene (along the Y axis). The graphical illustration depicts the effect of carrying out 20 PCR cycles on a DNA preparation initially having only 10 copies of a target gene. The resulting graph is **semilogarithmic** in nature.

Advantages of PCR–Technique

PCR-technique has **two** cardinal advantages, namely:

- (a) Each and every cycle virtually doubles the content of the original target DNA, and
- (b) A 10^6 to 10^8 fold increase in the target sequence is actually achieved after a 20-30 PCR cycle run.

2.4 RECOMBINANT PROTEINS

After accomplishing the isolation, cloning and sequencing of a gene, it becomes absolutely necessary to express it in an appropriate *expression** system which could be either **fungal, bacterial mammalian tissue culture, or insect tissue culture**. The inclusion of a host organism together with the *specific vector*** gives rise to an expression system. At this juncture a *promoter**** not only augments relatively high yields of protein but also yields a secreted protein by sheer manipulation. It is, however, pertinent to mention here that the criteria of selecting an expression system are namely: economic factors and the structure of proteins.

The production of **recombinant proteins** and peptides can be accomplished to a fairly reasonable extent by the following *three* techniques, namely:

- (a) Bacterial systems,
- (b) Glycosylation, and
- (c) Mammalian tissue culture expression systems.

These techniques shall be discussed briefly in the sections that follows:

2.4.1 Bacterial Systems

It essentially makes use of the microorganism *E. coli* which being considerably inexpensive, but unfortunately they fail to yield an active protein invariably. Ideally, a constant endeavour is always

* *Expression:* The ability of a gene to function within a cell in such a way that the gene product is formed.


** *Vector:* An agent normally an insect or other animal, able to carry pathogens from one host to another. OR A genetic element able to incorporate DNA and cause it to be replicated in another cell.

*** *Promoter:* The site on DNA where the RNA polymerase begins transcription.

made to look for cheaper, dependable and reproducible methods whereby newer active protein is produced first on a smaller scale and then scaled up to a pilot and finally to the commercial scale without any compromise on the quality of protein.

It has been observed that the proteins produced by bacterial systems essentially have an **N-formylmethionine** (designated as '**f-Met**') at the N terminus because that is considered to be a critical signal for the exact initiation of *translation*,* to a stage whether either a cleavage site is introduced or the resulting protein is cleaved and secreted in the process.

It has been observed that when proteins are overexpressed in a bacterial system of *E.coli*, they exhibit a tendency to aggregate in intracellular bodies usually termed as *inclusion bodies*. They may prove to be either beneficial or harmful for the subsequent protein recovery that could be accomplished by anyone of the following *two* methods usually adopted.



2.4.2 Glycosylation

Glycosylation mainly contributes in enhancing the molecular weight of several mammalian glycoproteins. It also helps in modifying activity of the protein of choice. It has been observed that the eukaryotic proteins which are usually either secreted or inserted in membranes enter through the golgi apparatus and endoplasmic reticulum right into the eukaryotic cell and are duly glycosylated during this particular phenomenon. *O-glycosylation* takes place upon threonines or serines in the protein sequence, whereas *N-glycosylation* occurs exclusively at the asparagine residues which are essentially part of the sequence.

Precisely, a recombinant protein which has duly undergone the process of glycosylation for improving its activity should be produced only in the eukaryotic cells. Thus, **glycosylation** may be accomplished by using specific yeast cells, such as:

* **Translation:** The synthesis of protein using the genetic information in a messenger RNA as a template.

** **Periplasmic space:** The area between the cytoplasmic membrane and the cell wall in gram negative bacteria (e.g. *E. coli*).

Yeast	Characteristics
<i>Saccharomyces cerevisiae</i>	: Attached oligosaccharides having more than 100 mannose residues.
<i>Pichia pastoris</i>	: Transfection* with a baculovirus expression system.

2.4.3 Mammalian Tissue Culture Expression Systems

The **mammalian tissue culture expression systems**, for instance: *Chinese hamster ovary cells*, are invariably transfected with a *viral vector*. In reality, they require a very complex media for the production of protein, besides the entire process shows slower growth and very poor yields thereby making it an expression system. Of course, there exists an ample genuine scope whereby the exorbitant cost may be brought down considerably by the advent of more stable and more efficient mammalian tissue culture systems.

Once a protein has been developed by the help of biotechnology it is essential that its identification and determination of its purity is established by known modern–analytical techniques, such as:

Characteristics	Techniques used
Molecular Weight of Protein	: Polyacrylamide electrophoresis to assess any proteolytic, cleavages
Bioactivity	: Bioassay;
Charge	: Isoelectric focussing electrophoresis to assess any amino acid substitution;
Hydrophobicity or charges	: High performance liquid chromatography
In charge	: to assess folding errors;
Purity	: Capillary electrophoresis and pyrogenicity;
Confirmation of identity	: X-ray crystallography with respect to initially identified protein.

2.5 BIOTECHNOLOGY VS MODERN PHARMACY PRACTICE

In the recent past, ‘**biotechnology**’ has already gained a tremendous magnitude of commendable success in the most upcoming field of modern pharmacy practice. These multifold plus points may be further expatiated with the help of the following *four* cardinal aspects, namely:

- (a) A host of *modern drug substances* available in the therapeutic armamentarium belong to the class of **protein** and **peptides**. Therefore, they essentially require prime consideration towards their stability, dosage forms, storage, administration and lastly the probable side-effects as compared to the relatively much smaller synthetic drug molecules;
- (b) The proper selection of the recombinant products would heavily depend upon not only the correct choice of the expression system but also the small differences existing in the structure which may result from the use of various known expression systems;
- (c) A plethora of new drug class have evolved as a result of the advent of more in-depth knowledge of pathophysiology besides newer modes of therapies for such diseases that could not be successfully treated earlier; and
- (d) A good number of ‘**diagnostic products**’, available in the form of ‘**kits**’, are solely based on *monoclonal antibody technology*. They have been exclusively developed for the proper

* **Transfection:** The transformation of a prokaryotic cell by DNA or RNA from a virus.

diagnosis of *thyroid disorders, anaemia, fertility for pregnancy, allergies, cancer*; and finally the *management of several hormone-related imbalances (i.e., disorders)*. In short, *diagnostic products* represent one of the **largest biotech startups**.

It is worth while to mention here that since the world-wide recognition of **biotechnology** an enormous progress has already been accomplished in modern pharmacy practice. A few typical and classical instances shall be cited as under, namely:

- (i) Human proteins as drugs,
- (ii) New Drug Classes,
- (iii) Vaccines,
- (iv) New Diagnostic Agents, and
- (v) DNA probes and RELP analysis.

The abovesaid various aspects shall now be treated individually in the sections that follows:

2.5.1 Human Proteins as Drugs

It has been established beyond any reasonable doubt that DNA, the genetic material, not only directs the production of the proteins which essentially comprise the structure of, but also regulate the processes in the living system.

Genetic engineering fundamentally comprises of picking up a particular gene from a chromosome of one brand of organism and then inserting the same into the chromosome of another. In other words, **genetic engineering** is nothing but a sheer manipulation of DNA in a deliberate and controlled manner. It really allows the wisdom and skill of biotechnologists to lay hand on specific individual components of complex living systems and subsequently produce them on a large-scale either in comparatively much simple micro-organisms or mammalian cells inside fermentation tanks.

However, the production of human proteins exclusively by the aid of **genetic engineering** exclusively guided by the following *three* factors, namely:

- (a) Ability to isolate the DNA of interest,
- (b) Selection of an appropriate organism (*e.g. E. coli; B. subtilis*) wherein the selected DNA is first inserted and then produced and
- (c) Ability to extract and purify the resulting product after fermentation.

Various categories of human proteins are invariably used as drugs, such as: **hormones, blood products** and **lymphokines**. These specialized proteins shall be discussed briefly as under:

2.5.1.1 Hormones

Interestingly, hormones are relatively simple protein molecules that usually serve as a medium organ-to-organ communication in the body. Examples:

1. **Human Insulin: Insulin** is a protein essential for correct sugar metabolism and is produced in the body by the pancreas. *Humulin^(R)* (Eli Lilly) was the first approved and marketed clinical product produced by recombinant DNA technology in *E. coli* (1983). Soonafter in 1987, NOVO–

a Danish company evolved a continuous process for the production of genetically engineered human insulin from *yeast cells*.

2. **Calcitonin gene-related peptide (CGRP): Calcitonin** being a thyroid hormone which helps in the regulation of calcium retention in bones. It has been observed that patients either suffering from **Paget's disease*** or post menopausal condition thereby rendering the bones structurally weakened. **Cloned CGRP** was developed by Sandoz (Basel, Switzerland) with a view that the peptide could be useful in the treatment of cardiovascular diseases.

2.5.1.2 Blood Products

Interestingly, more than fifty proteins naturally occurring in human blood have been successfully cloned either in mammalian cells or in *E.coli*, but only a handful of these have gained access for detailed clinical studies. A few typical examples will be cited below:

- (a) **Tissue Plasminogen Activator (tPA):** It is a protein generated by the body cells that is exclusively involved in the dissolution of unnecessary and unwanted blood clots. However, tPA is formed by natural cells of the body exactly in a similar manner as the interferons, but the former is produced in such a small quantity that its isolation in sizable quantities poses a difficult problem for its ultimate utility either in research or medicinal purposes. With the advent of biotechnology it is now possible to *clone* the **human tPA gene** into bacterial cells and thus produce any amount required. When tPA is administered intravenously to a patient suffered from heart attack, it reaches directly to the culprit blood clot that blocks the cardiac vessel thereby acting as a '*Cardiac-drano*' i.e. it helps in clearing off the blocked passage quickly and swiftly.
- (b) **Factor VIII:** The **genetically engineered Factor-VIII** is used exclusively for the treatment of haemophilia.

2.5.1.3 Lymphokines

Lymphokines represent a class of small proteins, which essentially include substance like: **interleukins, interferons** and **tumour-necroses factor**, and are usually secreted most naturally by cells of the immune system. They exert a *hormone-like action* through cell-to-cell interactions rather than spread out over the entire body. Perhaps this aspect offers a clear line of demarcation with regard to their usage as pharmaceutical substances as most drugs need to be administered systemically.

Examples: A few typical examples of **lymphokines** are enumerated below along with their applications in therapeutic domain:

- (i) **α -Interferon:** for rare cancer *hairy cell leukemia*;
- (ii) **Interleukin-2:** for cancer
- (iii) **Tumour Necrosis Factor (TNF)**—for killing tumour cells and causing a generalized wasting syndrome.

* **Paget's disease:** A chronic form of osteitis with thickening and hypertrophy of the long bones and deforming of the flat bones.

2.5.2 New Drug Classes

The **recombinant DNA technology** has paved the way in the management and control of a number of diseases related to body's immune system. In the past two decades the US-FDA approved quite a few recombinant DNA products that are being used as drugs across the world. Recently, the exact role of **cytokines** has been revealed in order to understand the intricacies of the relatively complex immune system.

The most recent and epoch-making developments with regard to the new drug classes are the *novel vaccines, proteins, hormones, glycoprotein, blood-clotting proteins* and *immunoactive drugs*. A few typical examples have been enumerated in Table 2.1, to stress upon the versatility of newer recombinant DNA products.

Table 2.1 Approved Biotechnology Medicines

S. No.	Name of the Product	Brand Name (Mfg. Co.)	Indication
Vaccines:			
1.	Hepatitis B vaccine	Engerix B ^(R) (Smith Kilne Beecham)	Hepatitis B prevention. Comprise of highly specific antibodies that more or less act as 'magic bullets'.
2.	Human Insulin	Humulin ^(R) (Eli-Lilly); Novalin ^(R) (Nova Nordisk)	To combat insulin dependant diabetes; it is the first human health product used in medicine.
Proteins			
3.	Interleukin- 2 Recombinant	Proleukin ^(R) (Chiron/ Cetus)	For metastatic renal cancer. A protein drug through which immune cells communicate with each other.
4.	Interferon Alfa- 2a, Recombinant	Roferon-A (Hoffman-LaRoche)	For hairy cell leukemia; AIDS related Kaposi's sarcoma; Antiviral activity, especially against RNA Viruses; Enhances the targetting of monoclonal antibody-tethered cytotoxic drugs to cancer cells.
5.	Interferon alfa – 2b, Recombinant	Intron-A (Schering Alferon) (Interferon Sciences)	For hairy cell leukemia; AIDS related Kaposi's sarcoma, chronic hepatitis types B & C; <i>Condylomata acuminata</i> .
Hormones:			
6.	Somatropin	Humatrope ^(R) (Eli-Lilly); Protropin ^(R) (Genentach)	Identical to human pituitary derived somatropin; for human growth hormone deficiency in children.
7.	Calcitonin	Cibacalcin ^(R) (Ciba); Miacalcin ^(R) (Sandoz) Calcimar ^(R) (Rhône-Poulenc Porer)	To decrease osteoelastic activity thereby inhibiting the movement of bone salts from bone to the blood; also decreases renal tubular secretion of calcium.
Glycoprotein (Purified)			
8.	Alteplase Recombinant	Actirase ^(R) (Genentech)	For acute myocardial infarction; Pulmonary embolism.

(Contd.)

(Table 2.1 contd.)

S. No.	Name of the Product	Brand Name (Mfg. Co.)	Indication
Blood Clotting Proteins:			
9.	Antihemophilic Factor	Profilate ^(R) (Alpha) Hemofil T ^(R) (Hyland); KoGENate ^(R) (Miles); Recombinant ^(R) (Baxter)	For the management and control of severe hemorrhage in the patient with hemophilia A.
Immunoactive Drugs:			
10.	Muromonab CD3	Orthoclone OKT3 (Ortho)	For acute allograft rejection in renal transplant patients; and cardiac transplant patients.

2.5.3 Vaccines

Vaccine is a suspension of infectious agents, or some part of them given for the purpose of establishing resistance to an infectious disease. Traditionally, vaccines comprise of *four* general classes, namely:

- Those containing living attenuated infectious organisms, for example, **vaccine for poliomyelitis**;
- Those containing infectious agents either by chemical or physical means, for instance: *vaccines used to protect human beings against typhoid fever, rabies and whooping cough.*
- Those containing soluble toxins of microorganisms sometimes used as such, but usually forming toxoids such as : *vaccine used in the prevention of diphtheria and tetanus,* and
- Those containing substances extracted from infectious agents, for examples: *capsular polysaccharides extracted from pneumococci.*

At this juncture, it is worthwhile to draw a line between the **traditional vaccines** as stated above and the **recombinant vaccines** (or **subunit vaccines**) obtained exclusively by the aid of genetic engineering techniques known so far are described below:

S. No.	Traditional Vaccines	S. No.	Recombinant Vaccines (or Subunit Vaccines)
1.	Prepared either by attenuating or killing the pathogens with a view to disabling the disease causing function of the pathogens.	1.	By the help of genetic engineering techniques it is quite possible to eliminate the potential vaccine induced illness by separating effectively the <i>immune activation function of pathogens</i> from the disease producing functions.
2.	Immune activating molecules remained unharmed.	2.	First, all the gene(s) solely dedicated for encoding the cell surface molecules that eventually trigger the immune activation function is isolated.
3.	Modified pathogens on being injected into a subject induce immunity to the modified pathogen thereby preventing the disease.	3.	Secondly, these genes are removed from the pathogens genome by endonuclease action and subsequently inserted into a cloning vehicle that later on employed to transform either a bacteria (<i>E. coli</i>) or yeast (<i>Saccharomyces cerevisiae</i>) host cell.
Disadvantages:			
(a) In certain instances, the <i>whole virus vaccine (i.e., the attenuated pathogens)</i> invariably revert back to their virulent state in the produced disease in the vaccinated subject rather than preventing it,			

(Contd.)

(Table contd.)

S. No.	Traditional Vaccines	S. No.	Recombinant Vaccines (or Subunit Vaccines)
	<p>(b) Possibility of certain pathogens being escaped from attenuation (or being killed).</p> <p>(c) Impurities may be incorporated through the method of production based on different <i>animal species</i>. They may be removed by very tedious and extensive procedures.</p> <p>Advantages: In spite of various serious adverse reactions they have been used successfully in combating diseases like; Polio, Small pox, Yellow fever, Measles, Mumps, Tetanus and Diphtheria.</p>	<p>4. A series of antigenic molecules are synthesized by the host cells that may be suitably isolated, purified and made into a vaccine.</p> <p>5. When this recombinant vaccine is injected into a subject, the individual affords an immune response against the antigen.</p> <p>6. The synthesis of suitable memory B cells is the net result of the immune response.</p> <p>7. The immunized subject on being exposed to a pathogen that essentially bears that specific antigen on its cell surface, the resulting immune response swiftly eradicates the pathogen totally.</p> <p>Advantage: As it contains only the immune-activating subunit* of the pathogen, it is incapable of inducing the disease state in the subject.</p>	

* **Recombinant vaccines** only contain a subunit of the original pathogen, hence they are also termed as '**Subunit Vaccines**'.

2.5.3.1 Specificity of Vaccines

Whether it is a *whole virus vaccine* or a *subunit vaccine* both exert their action by including an immune response against a specific subset of molecules that could be the reason why a vaccine specifically prepared against one influenza strain fails to respond against an altogether different strain. Based on this observation the health authorities in hospitals and OPDs always advise the youngsters, the elderly ones and above all the chronic ones to receive the shot against influenza every year periodically. It has been duly observed that each outbreak of flu is invariably caused by a different and newer influenza virus which means logically to produce new vaccines with each outbreak of the disease. The same logical explanation holds good for polio whereby three different strains give rise to three types of vaccines to control and prevent the same disease.

2.5.3.2 Search for AIDS Vaccines

Human Immunodeficiency Virus (HIV) belongs to the class of viruses termed as *retroviruses* which differ from most of the cell types. Here they usually comprise of RNA as their only major genetic material in comparison to the more common DNA. It has been observed that whenever a HIV virus infects a cell it not only injects its RNA but also the specific genetic material usually an enzyme termed as *reverse transcriptase*. Now, the reverse transcriptase affords the synthesis of a DNA molecule complementary to the viral RNA inside the host cell. Thus the virally derived DNA exerts the '*infective assault*' upon the host cell and ultimately leads to a condition commonly known as **Acquired Immunodeficiency Syndrome (AIDS)**.

It is pertinent to mention here that HIV is not adequately suitable for production of vaccine by virtue of the following *four* aspects, namely:

- (a) Being an extremely complex virus it may conveniently get lost in the inner portion of the infected cells thereby leaving absolutely no trace of viral antigens on the cell surface. At the end, the overall immune system may appear to behave as if no outside virus were ever present,

- (b) Unfortunately, till date no appropriate animal model has been established for human HIV infection, which has slowed down the active research and subsequent screening through an elaborated experimental laid-down procedure.
- (c) Consideration of ethical aspects with regard to human screening of any probable and potential AIDS vaccines, and
- (d) HIV possesses its unique ability to insert pieces of its self-genome right in the genome of the infected host, thereby rendering the genetic material invisible to the respective host human-system.

Keeping in view the above serious limitations encountered a good number of **AIDS vaccines** have been prepared meticulously and tested adequately. A few such examples are illustrated below:

- (i) **Thymosin Alpha 1** : For AIDS in combination with
(Immunomodulator) PEF-II-2 and Retrovir^(R)
Manufactured by (under Phase I/II Clinical Trial)
Alpha 1 Biomedicals
(Bethesda, MD)
- (ii) **Soluble CD4** : This protein interacts with HIV
(Product of Recombinant and thereby blocks its attachment
DNA technology) to intact CD4 T cells.

2.5.4 New Immunodiagnostic Agents

Immunodiagnostic agents are regarded as stranded tools in the physicians diagnostic regimens. They afford an improved testing technology whereby diagnosis could be faster, cheaper and easier.

2.5.4.1 Monoclonal Antibodies (MABs)

Antigen recognition by antibodies, that more or less acts like an anchor in body's defence mechanism system to combat dreadful diseases, has been judiciously explored for quite sometime towards the logical development of newer diagnostic tests for various ailments making use of antibodies from animal sources.

Hybridoma Technology The **hybridoma technology** is the process whereby the fusion of an immortal cell with a lymphocyte to produce an immortal lymphocyte. In other words, the antibody producing cells are duly fused with the respective tumour cells which results into the production of a hybrid that could be grown in an appropriate laboratory culture and which reproduces antibodies of a single specificity (**monoclonal antibodies**).

The tremendous growth in the field of **pharmacbiotechnology** in the recent past has broadened the scope of monoclonal antibodies (MABs) in *two* vital aspects of immunodiagnostics, namely:

- (i) MABs in diagnostics, and
- (ii) MABs in Imaging and Therapy.

2.5.4.2 MABs in Diagnostics

More recently MABs have gained rapid and wide recognition into the ever-expanding field of health-care diagnostics. There are usually *four* important techniques that find their utility in diagnostics, for instance:

- (a) **Immunoassays:** Most immunoassays make use of radioactive antibodies [*i.e.* **radio-immunoassays (RIA)**] whereby the sample showing positive test which allow the antibody to get bound to it and thus the radioactivity will be retained onto the sample. However, the stringency and authenticity of RIA tests mostly restrict it to centralized specialist diagnostic facilities exclusively;
- (b) **Enzyme Immunoassays (EAI):** In this particular instance a specific colour-producing enzyme is coupled to the antibody. Thus, the outcome of the results may be read either directly by a naked eye or spectrophotometrically;
- (c) **Enzymes- Cascade Technique*:** Here, several enzyme reactions taking place are coupled to generate an appreciable amplification of the original binding signal which is either read by a naked eye or spectrophotometrically; and
- (d) **Fluorescence Immunoassays (FIA) and Luminescence Immunoassays (LIA):** Precisely, these are nothing but related techniques wherein the 'table' either exhibits fluorescence or light respectively.

Examples:

1. **Pregnancy Dipstick Test:** Based on MABs the pregnancy dipstick test determines the pregnancy either at home or in a clinical laboratory.
2. **Ovulation Dipstick Test:** Another dipstick test based on MABs ascertains the positive or negative ovulation in a subject, and
3. **AIDS Test:** MABs based AIDS test kit is available to identify its presence in donated blood sample(s).

2.5.4.3 MABs in Imaging and Therapy

Ironically, the most acute and severe hinderances ever encountered in the treatment of cancer virtually lies in the fact that *malignant cells* have a very close resemblance to the *normal cells*. It is, therefore, quite possible that the therapeutic agents that are actually intended to destroy (kill) malignant cells also destroy normal cells perhaps due to their close similarity. However, it has been established beyond any reasonable doubt that the surfaces of malignant cells do differ in certain respects from those of normal cells. As we have seen earlier that MABs only recognise specific antigens on cells, they are being fully exploited to image cancerous tumours particularly in an intense on-going clinical research programme and in therapy against a variety of malignancies, such as: lymphomas, melanomas, colon and breast cancer.

A few typical examples have been duly expatiated below, namely:

- (a) **Gastrointestinal Cancer**:** MABs is employed singly to combat gastrointestinal cancer. The underlying principle being that when the antibodies opt to bind to the turnover, they invariably exhibit a tendency to attract the cells of the immune system to act against the prevailing cancerous tissue.
- (b) **Lung, Breast, Prostate and Pancreas Cancer:** It is however, pertinent to mention here that enough research activities have triggered off in the recent past towards the development of monoclonal conjugates of *two* important class of drugs namely:

* Developed by I Q (BIO) in Great Britain.

** A collaborative research by Centocor (USA) & Hoffman – LaRoche (Switzerland).

- (i) **Anthracycline Drugs***: Such as antibodies having quinones and related structures, eg. Adriamycin^(R) (Adrio); Bufex^(R) (Bristol);
- (ii) **Desacetyl Vinblastine****: When desacetyl vinblastin *ie.* a chemical entity obtained either from the plant source or produced by plant cell culture, is conjugated to a monoclonal which consequently acts specifically on lung, prostate, breast and pancreas malignant cells.
- (c) **Drug Targetting and Tumour Imaging**: Another theory of **drug targetting** *vis-s-vis* **tumour imaging** is based on the use of highly specific and relatively *small molecular size toxins* [*e.g.*, castor seed) and toxin from bacteria (microorganism source)] conjugated with monoclonals. It has been observed that the same monoclonals that are being employed to target may also be used to image tumours via conjugation to radioactive elements. Once the conjugate has been injected, whole-body-scanning methods may be carried out to localize and quantify the malignant thereby helping the physicians either to initiate a preliminary diagnosis or to ascertain if the patient is giving due response to the conventional therapy.

2.5.5 DNA Probes and RFLP Analysis

Probe that are specific to different *genes* as well as *DNA fragments* have paved the way for the diagnosis of a number of prenatal problems. Besides, the genetic probes makes it possible to establish future outset of some ailments for instance: emphysema (*i.e.*, pathological distention of interstitial tissues by gas or air) and Huntington's disease (*i.e.*, an inherited disease of the CNS which usually has its outset between ages 25 and 55).

The cross section of **recombinant DNA** and medicine invariably occurs when genetically engineered probes help in the cure and management of human ailments. DNA probe is a radioactive labeled DNA fragment that acts as a complementary to a particular gene or gene segment. Thus, suitable probes may be employed to analyse not only the abnormal genes but also the human genome. Interestingly, prenatal medicine makes use of probes to ascertain the presence of the genes in some ailments, namely: *Cystic fibrosis**** and *Tay-Sachs disease*****.

In addition, **DNA probe** is considered to be the most versatile and powerful technique for the identification of individuals.

The application of **DNA probe** as diagnostics has been extended to test for AIDS, understand the course and causes of cancer, genetic disease (*e.g.*, *muscular dystrophy*, *sickle cell anaemia*) and bacterial infections (*e.g.*, *walking pneumonia*, *Legionnaire's disease*).

Restriction Length Fragment Polymorphism (RELP) Analysis Each and every human being born on this earth essentially inherits one set of genes from its father and another one from its

* A collaborative research of Carbo-Erba (Italy) and Cytogen (New Jersey, (USA).

** Eli – Lilly (USA).

*** A single gene defect manifesting in multiple body systems as chronic obstructive pulmonary disease, pancreatic exocrine deficiencies, urogenital dysfunction and abnormally high electrolyte concentration in the sweat.

**** An inherited disease transmitted as an autosomal recessive trait. Because of the lack of the enzymes hexosaminidase A, which is important in sphingolipid metabolism, sphingolipids accumulate in the cells, especially those of the nerves and the brain. Death normally occurs before the age of 4. There is no specific therapy.

mother which in most cases have been found to be identical. Invariably, mutation of genes that one inherits from one or other of one's parents ultimately leads to diseases like muscular dystrophy and haemophilia. Besides, these distinct differences occurring between the two sets of DNA there exists many thousands of relatively 'Silent' differences usually termed as **restriction length fragment polymorphisms (or RELP)**.

The enormous progress made in the field of *biotechnology* has made it possible to afford the diagnosis of genetic defects prenatally. The following steps are performed sequentially:

- (a) Cells from the growing foetus are first drawn by sophisticated methods, namely: **Chorionic vili sampling and aminocentesis**,
- (b) DNA preparations are subsequently made by standard methods,
- (c) Specific enzymes that chop DNA in strategic positions are usually employed so as to trace 'silent' genetic markers and also the **RELPS**, associated with the prevailing ailments,
- (d) Carrying out a close comparison between the 'pattern of silent genetic markers' on the foetal DNA and those on the paternal and maternal DNA, it is now relatively possible to ascertain with a high degree of probability, whether the foetus under investigation is either normal or diseased.

In addition to the above mentioned diseases **RELP analysis** has also been judiciously and effectively extended to some more gruesome diseases such as: *schizophrenia, Alzheimer's disease and complications of heart*.

2.5.6 Enzyme Linked Immunosorbant Assay (ELISA)

The unique covalent attachment of enzymes to antibody molecules provided an immunological method that not only affords high specificity but also high sensitivity. **ELISA** utilizes antibodies to which enzymes (*eg: peroxidase, alkaline phosphatase and 3-galactosidase* have been covalently bound in such a fashion that neither the enzyme's catalytic characteristics nor the antibody's specificity are changed.

It is however, pertinent to state that **ELISA** may prove to play a major role in pin-pointing the early diagnosis of plethora of human ailments or medical conditions that are of major public health significance, for instance:

Diseased Condition	Primary(P)/Confirmatory (C)
• Pregnancy	P
• Leprosy (<i>Mycobacterium leprae</i>)	P
• Syphilis (<i>Treponema pallidum</i>)	P
• Tuberculosis (<i>Mycobacterium tuberculosis</i>)	C
• Leishmaniasis	C
• Hepatitis	P
• AIDS	P
• T3-T4, TSH (Thyroid hormones)	P

In fact *two* fundamental **ELISA methodologies** are prevalent, namely:

(a) **Direct ELISA:** It is meant for detecting *antigen* exclusively, for instance:

- (i) Virus particles from faecal sample, and
- (ii) Virus particles from a blood sample.

Method of Testing:

- (i) Specimen is first added to the wells of a microtitre plate that has been duly precoated with antibodies specific for the antigen to be detected. If the antigen (*i.e.* **virus particulates**) is present in the sample, it would be trapped by the available antigen binding sites present on the antibodies.
- (ii) The unbound material is washed out and a second antibody (containing a **conjugated enzyme**) is added,
- (iii) As the second antibody is also specific for the antigen it gets bound to any available remaining exposed determinants,
- (iv) After giving a proper wash, the enzymes activity of the bound material present in every microtiter well is estimated by adding the substrate of the enzymes, and
- (v) The characteristic colour thus produced is found to be directly proportional to the amount of antigen present.

(b) **Indirect ELISA:** It is meant for detecting *antibodies* present in human serum. It is invariably employed to specifically detect antibodies to **human immunodeficiency virus (HIV)**.

Method of Testing (HIV- ELISA):

- (i) The microtiter plates are first and foremost coated with a disrupted preparation of HIV particles (approximately 200ng of disrupted HIV particle is needed in each and every well)
- (ii) The resulting plates are subjected to a brief incubation period so as to ensure plates of the antigens to the surface of the microtiter wells.
- (iii) Now, a diluted serum sample is added and the mixture thus obtained is again incubated so as to permit **HIV–antibodies** to bind to **HIV antigens**.
- (iv) In order to detect the presence of antigen–antibody complexes, a *second antibody* is now introduced, which is essentially an enzyme conjugated anti-human IgG preparation,
- (v) Again after a short incubation (with the second antibody) followed by a washing step to ensure complete removal of any unbound second antibody, and
- (vi) The enzyme activity is now determined with the production of a colour which is directly proportional to the quantum of anti-human TgG antibody bound.

Note:








1. It is a positive indication by virtue of the fact that the binding of the second antibody and the antibodies from the patients serum ultimately recognised the HIV antigens *i.e.*, the patient possesses antibodies HIV, and the same has been exposed to HIV.
2. Always the control sera* are also assayed simultaneously along with any sample(s) so as to measure the extent of background absorbance in the particular assay.

* **Control sera:** which is known to be HIV–negative

However, the ‘**Indirect ELISA-Test**’ for the detection of antibodies to HIV, the casual agent of AIDS, has been depicted in Figure 2.5 along with the various sequential procedural details for both positive and negative test *vis-à-vis* the graphic representation between the quantum of antibody and the intensity of the colour produced.

2.6 BIOTECHNOLOGY BASED PHARMACEUTICALS FOR THE MILLENNIUM

From a recent survey (1993)* it has been broadly accepted that *biotechnology* based pharmaceuticals (or **Biotechnology Drugs**) will enhance tremendously to combat today’s complex ailments. In the last two decades American Pharmaceutical concerns have grossly shifted their interest towards life-saving medicinals based solely on biotechnology that is shown by the quantum jump of their on-going research projects from a meager 2% (in 1980) through 33% (in 1993) to a maximum of upto 70% (in 1993) for larger pharmaceuticals**.

S. Methodology No.	Positive Test	Negative Test
1. Miceotitre wells are coated with antigen preparation from disrupted HIV particles(*)		
2. Patient serum sample is added. HIV specific antibodies bind to HIV antigen		
3. Washing is done with buffer		
4. Human anti-IgG antibodies conjugate to enzymes (E+E) is added		
5. Washing is done with buffer		
6. Substrate for enzyme is added and quantity of colored product(.) is measured, which is directly proportional to the antibody concentration.		
Colour Intensity	+ + + +	-

* Mossionghoff G: Biotechncology medicines in Development (1993).

** Study by the Boston Consulting Group (BCG); April, 1993.




Fig. 2.6 Indirect ELISA Test for Detecting Antibodies to HIV, the Casual Agent of AIDS.

Interestingly, the **biotechnology based pharmaceuticals** for the millenium may be broadly categorised under the following *five* heads; namely:

- (i) Genetically engineered vaccine,
- (ii) Gene-splicing and DNA recombinant procedures,
- (iii) Antibodies in biotechnology,
- (iv) Gene Therapy
- (v) 3-D Picture of the 'Lock' and 'Keys'

These different categories shall be discussed briefly in the sections that follows:

2.6.1 Genetically Engineered Vaccine

In general, **vaccines** preparations made of either *dead organisms or living attenuated microorganisms* which may be administered to man or animal to specifically stimulate their immunity to infection by the same or closely related organisms. It has been observed that once an individual's immune system has positively responded to a particular antigen, it gives rise to a state of resistance that will remain for a reasonably long period of time. Thus, **vaccines** are regarded as important **immunizing agents**.

Interestingly, many viruses invariably possess an outer-coat made of protein. When human subjects or animals become immune to virus diseases, it is achieved by virtue of the fact that their immune systems to recognize the particular *virus protein-coat* and subsequently make corresponding antibodies to it. Now, mostly the pharmaceutical concerns actively engaged in **gene-cloning** maintains a keen interest in '*yeast*' as possible host for the expression of cloned gene.

Following are the examples of some **genetically engineered vaccines**, namely:

- (a) **Hepatitis B Surface Antigen (HBSAG):** It is prepared by cloning a coated protein gene in yeast, extracting the protein formed by the genetically altered yeast, and obtaining from it the desired vaccine. It confers immunity to **Hepatitis B** without exposure to the virus itself.
- (b) **Viral Vaccines:** In reality, **viral vaccines** are of *two* types, namely:
 - (i) those consisting of inactivated viruses that are incapable of mutiplication in the body, for instance, *Salk Poliovaccine*; and
 - (ii) those consisting of viruses that multiply at low rates in the body, but do not exhibit the symptoms of the disease, such as: *Sabin Oral Polio Vaccine*
- (c) **Whooping Cough Vaccine:** It is essentially a **bacterial vaccine** that consists of dead cells and provides a long-term immunity.
- (d) **Vaccines from Live Attenuated Viruses:** Both *Small Pox Vaccine and Measles Vaccine* obtained from live attenuated viruses induce a high degree of immunity in the vaccinated subjects, effective for a long duration and thus prove to be highly effective in the control and management of the respective diseases.

- (e) **Influenza Vaccine:** The vaccine of influenza viruses that have been grown in chick embryos and subsequently rendered non-infectious either by UV irradiation or by formalin. It prevents influenza by its parenteral injection and the duration of immunity rarely exceeds a year because of the fact that the causative agent (virus) of influenza is capable of undergoing mutation rather frequently.

It is pertinent to mention here that by the turn of the century the development of **synthetic vaccines** via **genetic combination** and **gene cloning** would likely to promote legitimately their medicinal utility and availability.

Figure 2.7 illustrates the various steps involved for the preparation of **vaccine** from hepatitis virus by the method of **genetic engineering**.




Fig. 2.7 Preparation of Vaccine from Hepatitis Virus by the Technique of Genetic Engineering

2.6.2 Gene Splicing and DNA Recombinant Procedures

The techniques invariably referred to as '**gene-splicing**' are well defined means of judiciously and constructively rearranging the genetic code to give rise to an organism with an altogether new, altered and most desirable features. It essentially comprise of the following *five* cardinal aspects namely:

- (i) A simple microbe may be empowered to generate a specific chemical,

- (ii) This desired characteristic informative feature or product is strategically located on the DNA molecule,
- (iii) **'Engineering'** aspect is mainly concerned with **cutting out** only that particular section of the string and subsequently grafting this into another organism.
- (iv) **'Splicing'** is generally carried out with a specific set of enzymes termed as '*ligation enzymes*', that would help in sticking the fragments once again, and
- (v) In fact, **'cutting'** and **'splicing'** involves more or less a very delicate surgery and these functions are actually accomplished *chemically* in solution only.

A brief description of **recombinant DNA (rDNA)** has already been made in section 3.1 in this chapter.

Figure 2.8 vividly depicts the basic **gene-splicing** and **recombinant DNA** procedures in an elaborated sequential manner.




Fig. 2.8 Basic Gene Splicing and Recombinant DNA Procedure in Sequential Order.

First of all the bacterial cells are removed from the mixture of *E.coli* with plasmids and the resulting chromosomes are retained. Secondly, the large chromosomes are removed by centrifugation and the resulting plasmid DNA is treated with restriction enzymes and incubated accordingly. Thirdly, add to it fragments of **eukaryote DNA** cleaved by the same restriction enzymes and add ligase; incubate to join tails. Fourthly, the circular molecules are removed by centrifugation and retained, whereas the linear fragments are rejected. Fifthly, the plasmids are resuspended and the ones with eukaryotes gene-inserts being heavier are removed by centrifugation. Sixthly, these segregated heavier plasmids are treated with bacterial protoplasts, Seventhly, the plasmids infected protoplasts are duly cultured on agar plates, grown into colonies of identically infected bacteria. Finally, the resulting clones may be incubated into large volumes of media and produced in large quantity.

2.6.3 Antibodies in Biotechnology

An extremely important and versatile aspect of **recombinant DNA technology** is the application of **antibodies** in biotechnology. In fact, **antibodies** are produced by the **B-cells** (or **plasma cells**) exclusively. They are virtually composed of four protein chains that one are interconnected by 'disulphide bonds'. It has been observed that the surface of the antibody essentially bears a highly *specific marking* (or 'lock') that would promptly recognize the particular foreign particle (or 'key') with which it readily undergoes complexation or binding. It has now been well established that different antibodies are actually generated in each individual person for their characteristic and highly specific interaction with antigens. In other words, antibodies play the role of the **immune surveillance** in the living system. Certain antibodies have the job of surveying foreign cells or molecules and mark the invading antigenic entity distinctly for destruction by other immune system cells. Consequently, the marked **antigens** are actually removed from the living system thereby leaving the unmarked ones *in situ*.

It is understood that there exists approximately 4000 strategic locations in the human genome which are responsible for various genetic disorders (diseases). Interestingly, out of this lot only 30 (*i.e.* 1200) have been studied in sufficient details. It has been observed that certain abnormalities found on the genes are usually termed as 'point mutations'. In all these instance, a single nucleic acid base present in a gene is normally replaced by a different one. Therefore, in the encoded protein the net result would be visualised by the exchange of a single amino acid.

Example: The genes encoding the **hemoglobin protein sequences** invariably possess a minimum of forty 'point mutations'.

It is pertinent to state here that the **molecular probing**, with regard to the characteristic alteration in the shape of blood cells in *sickle-cell anemia**, would go a long way in revealing these types of abnormalities parentally, or in the early stage of life so as to make it possible to institute a suitable remedial measure or a preventive action adequately (*i.e.*; '**gene therapy**').

Figure 2.9 clearly illustrates that the manner by which genetic defects could be detected in a logical way in *foetal cells*.

It has been noticed that quite often variance do take place in the genetic code on account of error into the *chemical sequence*. The defects of this sort are invariably caused by chemical change or by inheritance and ultimately results into specific disease condition. At this particular juncture the

* **Sickle Cell Anemia:** The gene for sickle cell anemia is hemoglobin S. Thus each RBC has both normal hemoglobin A and abnormal hemoglobin S. These will not become sickled until extremely low concentration of oxygen occur.

prenatal examination of the *foetal cells* would help in locating the **faulty genes** and thus, identify particular subjects who may develop certain disease. The DNA is carefully removed from the *foetal cells* and with the help of an appropriate enzyme the DNA is cleaved whereby the same is unwound into respective single strands. In this particular instance, one nucleotide base (say, 'A') is defective giving rise to a situation termed as '**point mutation**'. Now, a purely synthetic DNA is manufactured in the laboratory whose genetic sequence will allow it to bind to the '**point mutation**' i.e; its 'T' with a defective 'A'. A radioactive tag (-) is then attached to the probe which is subsequently mixed with the **foetal DNA**. This specific radioactive tag makes it possible for investigators to know whether the bases get bound or not. In this particular instance (Figure 2.9), they do, thereby suggest that the foetus under investigation may develop the blood disease.

Occasionally variance do occur in the genetic code due to error into the chemical sequence. Defects of this by chemical change or by inheritance and results into disease condition. Timely prenatal screening may locate faulty genes & thus, identify particular subjects who may contract certain diseases.

The DNA is carefully removed from the foetal cell.

An enzyme is employed to cleave the DNA. It is unwound into respective single strands. In this instance, one nucleotide base is defective giving rise to a situation termed as a "Point mutation".

- A purely synthetic DNA is prepared in the lab. Its genetic sequence will allow it to bind to the 'Point mutation'. (i.e. its 'T' with a defective 'A').
- This probe is mixed with the foetal DNA. The radioactive tag makes it possible for researchers to know whether the bases get bound. In this instance, they do, thereby indicating that the foetus may develop the blood disease.




Fig. 2.9 Searching for Genetic Defects in Foetal Cell.

2.6.4 Gene Therapy

It has been adequately proved that there exists two probable cellular targets essentially required for carrying out the process of 'gene replacement therapy' namely: the **Somatic Cell*** and the **Germ Cell****.

In the light of the ethical issues inherent in 'gene therapy', five narrowly cardinal objectives have been put forward by the scientific review committees, namely:

- (i) Research should be limited to somatic cells only, so that a treated individual cannot pass genetic alterations to offspring,
- (ii) Risk to the patient must be outweighed by the potential therapeutic benefit,
- (iii) Target diseases must be limited to those that involve a known defect in a single gene, and the normal gene must be cloned and available,
- (iv) Disease must involve cells that can be isolated from a patient, altered in tissue culture and subsequently reimplanted in the patient, and
- (v) Planned well-defined procedures must meet strict safety standards in animal trials, before attempts are made with human beings.

Targets of human **gene therapy** essentially include several serious ailments such as:

- (a) **Lesch-Nyhan Syndrome:** [M. Lesch, b: 1939, William Leo Nyhan, b: 1926, U.S. **Pediatricians**] A functional lack of a single enzymes *i.e. hypoxanthin – guanine phosphoridbosyl transferase*, produced by a single gene, which is essential for purine metabolism, gives rise to an inherited metabolic disease that affects only males, in whom mental retardation, aggressive behaviour, self-mutation, and renal failure are exhibited, and
- (b) **Tumour Necrosis Factor (TNF):** Another effort is exclusively focussed on new approaches in the treatment of cancer, whereby immune system cells known to be associated with tumours are suitably modified to produce a protein with appreciable anti-tumour activity termed as TNF. Alternatively, TNF is a protein mediator or cytokine released primarily by **macrophages** and **T lymphocytes** that helps regulate the immune response and some **hematopoietic functions**.

There are *two* factors, namely:

- (i) **Alpha (TNF-a)** : [Syn: Cachectin – produced by macrophages, and
- (ii) **Beta (TNF- b)** : [Syn: Lymphotoxin]- produced by activated CD4*** + T cells.

Interestingly, **gene therapy** is a process by which a patient is cured by altering his or her *genotype*****. **Severe Combined Immunodeficiency (SCID)******* is treated by **gene therapy** for a variety of valid reasons, such as:

-
- * **Somatic Cell:** [Gr. Soma = body] Pertaining to nonproductive cells or tissues;
 - ** **Germ Cell:** A cell whose function is to reproduce the organism. It usually has a single set of chromosomes. Germ cells are called 'ova' in females and 'spermatozoa' in males.
 - *** **CD4:** A protein on the surface of cells that normally helps the body's immune system combat disease. HIV attaches itself to the protein to attack WBC, causing a failure of host defence.
 - **** **Genotype:** The precise genetic constitution of an organism (Phenotype the observable characteristics of an organism).
 - ***** **SCID:** A syndrome marked by gross functional impairment of both humoral and cell-mediated immunity and by susceptibility to fungal, bacterial and viral infections.

- (i) There is no possible cure for the disease and is fatal at an early age,
- (ii) **SCID**, results from an alteration in a specific gene that has been isolated and cloned and thus is available for use in treatment.
- (iii) **SCID**, results from the hereditary absence of a single enzymes **adenosine deaminase (ADA)**, and
- (iv) The cells that usually express a gene are a type of WBC that can be easily removed from a patient, cultured *in vitro*, genetically modified and then reintroduced into the patient by transfusion.

In the recent past, the somatic cells have been regarded as ‘**targets**’ for **gene therapy**. It has been observed that **adenosine deaminase deficiency (ADD)**, a very serious type of disorder of the immune system, takes place once in each 200, 000 newly born babies. **Adenosine deaminase (ADA)** plays a vital role for the production as well as maintenance of *two* of the most important cell types that are found to be active in the immune response, namely: **B-lymphocytes** and **T-lymphocytes**. In the absence of these aforesaid cell types the children are susceptible to every possible infection to which they are exposed.

Examples:

- (i) The case of David, the ‘**Bubble Boy**’ from Texas (USA) in 1971, represents a typical example wherein he suffered from ADA deficiency causing severe immune dysfunction. Eventually, David had SCID and died at the age of 13 years probably due to a series of infections spread out through his entire body as a result of the *total collapse of prevailing immune systems*.
- (ii) Blaese and Anderson (1990) isolated **T-cells** from the blood of a 4-year old girl suffering with ADD. They were successful in the transformation of these isolated T-cells with a tailor made, **genetically nonmutant ADA gene**. The resulting **engineered T-cells**, having the inserted **ADA gene**, were subsequently inducted to the same child through blood transfusion. The investigators had a strong prediction that the newly administered gene will not only function properly but also help in the synthesis of enough ADA to restore normal immune function required by the affected child.

2.6.5 3D Picture of the ‘Lock’ and ‘Keys’

The explanation of ‘**lock**’ and ‘**keys**’ has been made in Section 2.6.3 of this chapter. With the advent of latest developments in the fields of science and technology and the unique combination of X-ray crystallographic studies, molecular mechanic calculations and supercomputers have tremendously helped in revealing the three-dimensional (3D) arrangement vividly. Based on this 3D picture of the ‘**lock**’ scientists are in a position to design specifically shaped drug molecules (‘**keys**’) that would conveniently fit into the active sites of the 3D protein (or the folded protein). In this way, it has really paved the way towards a quantum jump for a realistic and rational approach to drug design. It is earnestly expected that in the near future this 3D-picture of the ‘**lock**’ and ‘**keys**’ would certainly prove to be a great asset.

2.7 BIOTECHNOLOGY AND MODERN DRUG DISCOVERY

In the past two decades, an exponential growth has emerged in the field of modern drug discovery exclusively based on **'biotechnology'**. An unique blend of wisdom, skill, knowledge and an enormous strength of perseverance with regard to the most advanced and sophisticated techniques of **recombinant DNA technology, gene splicing, cloning of genes, genetically engineered vaccines** and the like have given birth to the discovery of a plethora of drug substances. Interestingly, these newer types of drugs would certainly prove to be beneficial in treating some of the most dreadful human ailments besides improving the quality of life.

These **'biotechnology medicines'** may be broadly categorised into *two* heads, namely:

- (a) Approved medicines, and
- (b) Medicines under development.

The two different categories of biotechnology medicines shall be exemplified in the sections that follow.

2.7.1 Approved Medicines

A number of medicines that have been approved by several drug authorities are enumerated below:

S.No	Name of the Product	Company	Indication (s)
1.	Activase^(R) (Alteplase)	Genentech (S. San Francisco CA)	Acute myocardial infarction
2.	Actimmune^(R) (Interferon gamma –1b)	Genentech (S. San Francisco CA)	Pulmonary embolism; Chronic granulomatous disease
3.	Beta seron^(R) (Interferon beta – 1b)	Berlex Laboratories (Wayne, NJ)	Multiple sclerosis
4.	Cerezyme^(R) (Imiglucerase)	G-enzyme	Type-1 caucher's disease
5.	Epogen^(R) (Epoctin alfa)	Amgen (Thousand Oaks, CA)	Anemias of chronic renal disease; AIDS, cancer; chemotherapy
6.	Engerix-B^(R) (Hepatitis B vaccine)	Smith Kline Beecham (Philadelphia, PA)	Hepatitis –B prevention
7.	Humatrope^(R) (Somatropin)	Eli-Lilly (Indianapolis, IN)	Human growth hormones deficiency in children
8.	Humulin^(R) (Human insulin)	Eli-Lilly (Indianapolis, IN)	Diabetes;
9.	Intron-A^(R) (Interferon alfa-2b)	Schering Plough (Madison, NJ)	Hairy-cell leukemia; genital warts; AIDS-related Kaposi's sarcoma; hepatitis – C; hepatitis B
10.	Leukine^(R) (Sargramostim)	Immunex (Seattle, WA)	Myeloid recognition after bone marrow trans-plantation
11.	Neopogen^(R) (Filgrastim)	Amgen (Thousand Oaks, CA)	Neutropenias due to myelosuppressive chemotherapy; myeloid reconstitution after bone marrow transplantation
12.	Orthoclone OKT3 (Muromonab CD-3)	Ortho Biotech (Raritan, NJ)	Reversal of acute kidney transplant rejection; reversal of heart and liver transplant rejection
13.	Proleukin^(R) (Aldesleukin)	Chiron (Emeryville, CA)	Renal cell carcinoma;
14.	Recombivax HB^(R) (Hepatitis B vaccine)	Merck (Rahway, NJ)	Hepatitis B prevention
15.	Recombinate^(R) (Antihemophilic factor)	Baxter Healthcare/ Hyland Division	Hemophilic A

2.7.2 Medicines Under Development

A good number of medicines are in the last phase of clinical trial *i.e.*, Phase-III with regard to their development status in USA. A few typical examples of drugs under this class are mentioned below:

S. No.	Name of Product	Company	Indication(s)	(Class)
1.	P1XY 321	Immunex (Seattle, WA)	Chemotherapy-induced neutropenia and thrombocytopenia	Colony-stimulating factors
2.	Procrit^(R) (Epoetin alfa)	Ortho Biotech (Raritan, NJ)	Prevention of anemia associated with surgical blood loss; autologous blood donation adjuvant	Erythropoietin
3.	Epidermal growth factor	Intraptics (Inrvine, CA)	Corneal and cataract surgeries	Growth factor
4.	Recombinant human platelet derived growth factor-BB(PDGF)	Chiron (Emeryville, CA)	Wound healing	Growth factor
5.	Nutropin^(R) (Somatropin for injection)	Genentech (S. San Francisco, CA)	Turner's Syndrome*	Human Growth Hormones
6.	Beta Interferon	Biogen (Cambridge, MA)	Multiple sclerosis	Interferon
7.	Immuneron (Interferon gamma)	Biogen (Cambridge, MA)	Rheumatoid arthritis	Interferon
8.	Roferon-A^(R) (Interferon alfa 2a)	Hoffman Roche (Nutley, NJ)	Chronic myelogenous leukemia; Hepatitis C	Interferon
9.	Recombinant human interleukin-3	Sandoz Pharmaceuticals (East Hanover, NJ)	Adjuvant to chemotherapy autologous bone marrow transplants	Interferon
10.	Anti-LPS Mab	Chiron (Emeryville, CA)	Sepsis	Monoclonal Antibodies
11.	CentorX (Mab)	Centocor (Malvern, PA)	Anti-platelet prevention of blood clots	Monoclonal Antibodies
12.	ESTM (Mab)	Pfizer (New York, NY)	Gram-negative sepsis	Monoclonal Antibodies
13.	Myoscint (Mifarmonab)	Centocor (Malvern, PA)	Cardiac imaging agents	Monoclonal Antibodies
14.	Antitumor necrosis factor	Miles Inc; (West Haven, CT)	Sepsis syndrome	Tumor necrosis factors
15.	Melacine (Therapeutic vaccine)	Ribi Immunotherm (Hamilton, MT)	Stage III-IV Melanoma	Vaccines
16.	VaxSyn[®] HIV-1 (gp 160)	MicroGene Sys (Menden, CT)	AIDS	Vaccines
17.	Antril (Interleukin I receptor antagonist)	Synergen (Boulder, CO)	Severe sepsis	Others

* A congenital endocrine disorder caused by failure of the ovaries to respond to pituitary hormone stimulation;

2.7.3 Human Clone

Severino Antinori and **Robert Edwards** jointly produced the first test-tube baby and created a history in the world. Now, Antinori, a 53 year old Italian embryologist wants to repeat the history by creating the world's first **human clone**. In spite of severe glaring ethical challenges, Antinori with the help of first human ***in vitro fertilization (IVF)*** technique enabled a 62 year old woman in 1994 to become the oldest to have a baby. He advocates and argues strongly that the technology of cloning is nothing but a logical and legitimate extension of IVF that may help in fertile couple to bear children.

After decades of constant dedicated efforts the scientists have developed the key techniques to “reset” the DNA of living cells that essentially possessed specialist functions so they behaved as though they were a *newly fertilized embryo* which grew into a **clone of the adult**. In early 1998, the experts at the University of Hawaii succeeded the **cloning of mice**.

Thus, the prospect of **‘human cloning’** is now perceived as an absolute reality in the near future. In view of the above rapid development in biotechnology the US government and the European parliament have made strict legislations to outlaw its practice on human beings. Although, it has been entirely left upto individual American and European states to accept or reject it outright. In Britain, the strict control of the **Human Fertilization and Embryology Authority (HFEA)** which has strongly pronounced its obvious intension to block and reject any requests to do work related to human cloning.

Antinori argues and seeks support from the world body for him to go ahead with human clone—

“What about the man who does not produce any sperm at all? What should he do? If he cannot reproduce himself why should he not reproduce his ‘genes’ in this way—this is one of the few cases where it is acceptable to clone”.

our medicines and how we keep healthy.

‘Functional-food’ was first developed by the US Institute of Medicine as food wherein the concentration of one or more ingredients have been duly manipulated to enhance their all-out contribution towards a healthy and complete diet.

Since 1969, when the very idea of healthy foods was first conceived , a host of such products slowly gained entry for the human consumption, for instance:

- Bran and soy; low-fat, low-salt and no-sugar products
- Ginseng—based products
- Fish oil supplement products,
- Extra-strong cabbage,
- Specially designed pasta, and
- Genetically altered porridge

Besides, some specially produced vegetables and other eatables also took due cognizance of their added therapeutic value such as:

Name of Product	Indication
Special broccoli	For cancer
Tomatoes	To prevent prostate malignancies
Spagetti	For arthritis
Egg sandwich	To prevent heart disease
Cornflakes	To minimise the risk of breast cancer

Likewise, the genetically altered rapessed oil plant to result hybrids that produce a substantial amount of **β-carotene**, the well known precursor of Vitamin-A, commonly found in rather very small concentrations in carrots, palm oil etc. It has also been proved logically* that if one consumes

* According to Dr. G. Kishore, the St. Louis based Chief Biotechnologist at Monsanto (USA).

a tablespoon of this oil daily, one would have an adequate amount of Vitamin-A required by the body. Perhaps, it would help a long way in providing between 250m to 1 billion people around the world, having various stages of Vitamin-A deficiency, and saving 10m children dying each year globally as a result of this deficiency, as an alternative '*functional food*'.

2.8 BIOTECHNOLOGY: SOME THOUGHT PROVOKING NEWER IDEAS

Since, the intensive and extensive research has gained a surmountable momentum in '**biotechnology**' a number of newer thought provoking ideas have been put forward by the scientists, such as: *potato vaccine*, *functional food revolution* and *human clone*. These different aspects shall be discussed briefly in the sections that follows:

2.8.1 Potato Vaccine

A researcher in US has recently developed a **genetically engineered potato** that could afford protection against food poisoning and diarrhoea. Charles Arntzen* has successfully transferred the antibiotic gene of *E.coli* into the genetic material of potato. As the potato could not be tricked into producing enough *E.coli* proteins to elicit an immune response, an '*artificial gene*' identical to the bacterium was duly synthesized. It has been observed that small chunks of raw potato incorporated with the **synthetic gene** were administered orally by 14 volunteers, antibodies against *E.coli* appeared in the gut. According to the researcher this '**vegetable vaccine**' is going to be a very effective strategy for delivering oral vaccines as it goes straight into the gut. It is earnestly believed that this **engineered potato** may prove to be the forerunner of many edible vaccines against a wide range of diseases, including *cholera* and *hepatitis B*.

2.8.2 Functional Food Revolution

The time is not far when it is all set to convert our kitchens into culinary pharmacies;. Interestingly, it is termed as **functional food revolution**. This new concept of 'functional food' bears an attitude which is poised to revolutionise the manner we normally take.

FURTHER READING REFERENCES

1. Cleveland J *et al.* Bacteriocins: Safe, natural antimicrobials for Food Preservation, *Int. J. Food Microbiol.*, **71**, 1-20, 2001.
2. De Souza NJ, *Coleus Forskohlii* Briq.: **The Indian Plant Source for Froskolin**, In SP Chaudhuri (Ed) *Recent Advances in Medicinal, Aromatic, and Spice Crops*, Today and Tomorrow's Printers and Publishers, New Delhi, Vol. I, pp. 83-91, 1991.
3. Ellaiash *et al.*: Optimization of process Parameters for Alkaline Protease Production under solid state fermentation by alkalophilic *Bacillus* sp., *Asian J. Microbiol Biotechnol Environ Sci.*, **5**, 49-54, 2003.
4. Gennaro, A.R., '**Remington : The Science and Practice of Pharmacy**', Mac Publishing Company, Easton, Pennsylvania Vol. I & II 19th edn. 1995.

* A plant biologist from the Boyce Thompson Institute for Plant Research in New York.

5. Karp, G., '**Cell and Molecular Biology : Concept and Experiments**', John Wiley and Sons Inc., New York, 1996.
6. Madian, M.T., J.M., Martinko and J.Parker, '**Brock's: Biology of Microorganism**', Prentice Hall, International, Inc., New Jersey, 8th ed., 1997.
7. Misawa M and Nakemishi TM, **Antitumour Compounds Production by Plant Cell Cultures**: In Bajaj VPS (Ed) *Medicinal and Aromatic Plants II*, Springer Verlag, New York, pp. 192-207, 1988.
8. Ota SK and Kurmasagar I, *Detection of monodon baculovirus and white spot syndrome virus in apparently healthy Penaeus monodon postlarvae from India by polymerase chain reaction*, *Aquaculture*, 220, 59-67, 2003.
9. Peters, P. '**Biotechnology**', Wm C. Brown Publishers, Dubuque, 1993.
10. Sandmann G: Genetic **Manipulation of Carotenoid Biosynthesis: Strategies, Problems and Achievements**, *Travels Plant Sci.*, 6, 14-17, 2001.
11. '**The International Biotechnology Handbook**', Euromonitor Publishers Limited, London, 1988.
12. Wilson, M., and S.S. Lindow : '**Release of Recombinant Microorganisms**, *Ann. Rev. of Microbiol.* 47, 913-944, 1993.

3

Carbohydrates



- Introduction
- Classification
- Carbohydrate Biogenesis
- Further Reading References

3.1 INTRODUCTION

The Germans first and foremost introduced the word '**kohlenhydrates**' which was later on coined to **carbohydrates**. The name obviously suggests that these compounds are essentially the hydrates of carbon. In reality, all carbohydrates comprise of carbon, hydrogen and oxygen; whereas, the last two elements are found to exist in the same proportions as in water (*i.e.*, H₂O – 2:1). However, it has been observed that there are certain compounds that do conform to the said '**hydrate rule**' *i.e.*, maintain the ratio of H and O (2:1) but do not belong to the category of carbohydrates, for instance:





- (i) Formaldehyde [HCHO] 2:1
- (ii) Acetic Acid [CH₃COOH] 2:1
- (iii) Lactic Acid [C₃H₆O₃] 2:1

Besides, there exist such compounds that evidently show the chemical properties of carbohydrates but do not necessarily abide by the above mentioned '**hydrate rule**', for example:

S.No.	Name	Emperical Formula	Structure	Remarks
1.	Cymarose	C ₇ H ₁₄ O ₄		<i>Apocynum cannabinum</i> L. <i>A. androsaemifolium</i> L., <i>A. venetum</i> L.
2.	Digitoxose	C ₆ H ₁₂ O ₄		Obtained by mild hydrolysis of the glycosides eg; digitoxin, gitoxin and digoxin

(Contd.)

(Table contd.)

3.	Rhamrose	$C_6H_{12}O_5$		<i>Rhus toxicodendron</i> L.
4.	Sarmentose	$C_7H_{14}O_4$		Hydrolysis of sarmentocymarins, a glycoside isolated from seeds of <i>Strophanthus sarmentosus</i> DC by the enzymatic method.
5.	Oleandrose	$C_7H_{14}O_4$		<i>Nerium oleander</i> L. (Laurier Rose)
6.	Digitalose	$C_7H_{14}O_5$		Obtained by hydrolysis of the glycoside from the seeds of <i>Strophanthus eminii</i> Aschers & Pax.

In view of the above cited glaring examples with regard to various anomalies the terminology ‘**Carbohydrates**’ has still been retained to represent not only the sugars but also those substances that are related to them basically in structure and other characteristic features.

Invariably, the carbohydrates belong to the chemical class of the aldehydes, ketone alcohols, and also the condensation polymers of these partially oxidized polyalcohols collectively known as ‘**Polysaccharides**’ or ‘**Oligosaccharides**’.

Glycan is the generic term for polysaccharide and in the systematic nomenclature the latter is assigned a suffix “-an”. Generally, the polysaccharide may be classified into *two* broad heads, namely:

- (a) **Homoglycan:** The polysaccharide is termed as homoglycan when it contains only one type of monosaccharide unit, and
- (b) **Heteroglycan:** The polysaccharide is known as heteroglycan when it involves more than one kind of monosaccharide unit.

However, a more accurate and precise demarcation of polysaccharides essentially makes use of nomenclature that includes, *first* the type of monosaccharide building unit, and *secondly*, the exact position and configuration of the glycosidic linkage involved.

Examples:

- (i) **Homoglycan:** *e.g.*, *Cellulose*. It may also be expressed as β -1, 4 -D-glycan by virtue of the following reasons, namely:
 - Prevailing attached unit is D-glucose,

- D-glucose bears the β -configuration at the anomeric C-atom (*i.e.*, C-1),
 - C-1 is linked to C-4 of the next identical unit of D-glucose.
- (ii) **Heteroglycan:** *e.g.*, *D-gluco-D-mannose*. It is made up of D-glucose and D-mannose. The two altogether different monosaccharides usually show up in an orderly manner. In this particular instance, the diheteroglycan is composed of two different types of monosaccharides that has been arranged in an alternating and regular fashion.

It is worthwhile to mention at this juncture that plant kingdom provides a variety of complex polysaccharides, such as: **cellulose, starch, dextran, inulin** and the like. These complex polysaccharides yield the respective sugar residues upon hydrolysis, for example:

Pentosans	Hydrolysis →	Pentoses, Arabinose, Xylose, Ribose;
Hexosans	Hydrolysis →	Hexoses, Glucose, Fructose;
Fructan	Hydrolysis →	Inulin that results Fructose;
Glucan	Hydrolysis →	Starch that gives Glucose.

Nevertheless, the starch and sugars find their abundant applications not only as food or food supplements, but also as indispensable adjuvants in the formulation of a wide range of pharmaceutical products all over the globe.

3.2 CLASSIFICATION

In broader sense the **polysaccharides** or **glycan** may be classified into *two* major groups, namely:

- (a) Homoglycans, and
- (b) Heteroglycans

These two major groups would be described in details with the help of important representative members in the sections that follows:

3.2.1 Homoglycans

A large number of plant products belonging to this particular category are namely: *honey, starch, hetastarch, inulin, lichenin, dextran, cyclodextrins, cellulose, cotton, and dextrin*.

3.2.1.1 Honey

Synonyms Madhu, Madh, Mel, Honey (English);

Biological Source Honey is a viscid and sweet secretion stored in the honey comb by various species of bees, such as: *Apis dorsata, Apis florea, Apis indica, Apis mellifica*, belonging the natural order *Hymenoptera* (Family: *Apidae*).

Geographical Source Honey is available in abundance in Africa, India, Jamaica, Australia, California, Chili, Great Britain and New Zealand.

Preparation Generally, honey bees are matched with social insects that reside in colonies and produce honey and beeswax. Every colony essentially has one 'queen' or 'mother bee', under whose

command a huge number of 'employees' exist which could be mostly sterile females and in certain seasons male bees. The 'employees' are entrusted to collect nectar from sweet smelling flowers from far and near that mostly contains aqueous solution of sucrose (ie; approximately 25% sucrose and 75% water) and pollens. Invertase, an enzyme present in the saliva of bees converts the nectar into the invert sugar, which is partly consumed by the bee for its survival and the balance is carefully stored into the honey comb. With the passage of time the water gets evaporated thereby producing honey (ie; approximately 80% invert sugar and 20% water). As soon as the cell is filled up completely, the bees seal it with wax to preserve it for off-season utility.

The honey is collected by removing the wax-seal by the help of a sterilized sharp knife. The pure honey is obtained by centrifugation and filtering through a moistened cheese-cloth. Invariably, the professional honey collectors smoke away the bees at night, drain-out honey, and warm the separated combs to recover the beeswax.

Description

Appearances	: Pale yellow to reddish brown viscid fluid,
Odour	: Pleasant and characteristic,
Taste:	: Sweet, Slightly acrid,
Specific gravity	: 1.35-1.36
Specific rotation	: +3° to -15°
Total Ash	: 0.1-0.8%

However, the taste and odour of honey solely depends upon the availability of surrounding flowers from which nectar is collected. On prolonged storage it usually turns opaque and granular due to the crystallisation of dextrose and is termed as '**granular honey**'.

Chemical Constituents The average composition of honey ranges as follows: Moisture 14-24%, Dextrose 23-36%, Levulose (Fructose) 30-47%, Sucrose 0.4-6%, Dextrin and Gums 0-7% and Ash 0.1-0.8%. Besides, it is found to contain small amounts of essential oil, beeswax, pollen grains, formic acid, acetic acid, succinic acid, maltose, dextrin, colouring pigments, vitamins and an admixture of enzymes eg; diastase, invertase and inulase. Interestingly, the sugar contents in honey varies widely from one country to another as it is exclusively governed by the source of the nectar (availability of fragment flowers in the region) and also the enzymatic activity solely controlling the conversion of nectar into honey.

Substituents/Adulterants Due to the relatively high price of pure honey, it is invariably adulterated either with artificial invert sugar or simply with cane-sugar syrup. These adulterants or cheaper substituents not only alter the optical property of honey but also its natural aroma and fragrance.

Uses

1. It is used as a sweetening agent in confectionaries.
2. Being a demulsent, it helps to relieve dryness and is, therefore, recommended for coughs, colds, sore-throats and constipation.
3. Because of its natural content of easily assimilable simple sugars, it is globally employed as a good source of nutrient for infants, elderly persons and convalescing patients.

3.2.1.2 Starch

(Corn starch, Potato Starch, Rice Starch, Wheat Starch)

Synonym Amylum

Biological Source Starch comprises of mostly polysaccharide granules usually separated from the fully grown grains of Corn [*Zea mays* Linn.]; Rice [*Oryza sativa* Linn.] ; and Wheat [*Triticum aestivum* Linn.] belonging to the family *Gramineae* and also from the tubers of Potato [*Solanum tuberosum* Linn.] family *Solanaceae*.

Geographical Source USA, Canada, Australia, China, India, CIS – countries (Russia), Thailand, Indonesia, Vietnam, Pakistan and many other tropical and sub-tropical countries are the major producers of starch in the world.

Preparation In general, cereal grains *e.g.*, corn, rice and wheat mostly comprise of starch bundles, oil, soluble protein and the insoluble protein termed as '*gluten*'; whereas the potato contains starch, mineral salts (inorganic), soluble proteins and vegetable tissues. Obviously, various specific methods are normally employed to separate starch either from cereal grains or from potato. These methods are briefly enumerated below, namely:

(a) Methods for Maize (Corn) Starch

Maize grains are first washed with running water to get rid of dust particles and adhered organic matters. They are now softened by soaking in warm water (40-60°C) for 48 to 72 hrs charged with a 0.2-0.3% solution of SO₂ to check the fermentation. The swollen grains are passed through '**Attrition Mill**' to split and partly crush them to separate the embryo and the epicarp. It is extremely important to isolate the germ (embryo) which may be accomplished by addition of water, whereby the germs float and are segregated by skimming off promptly. The corn oil, a rich source of Vitamin E, is recovered from the germ by the process of expression. After removal of the germ the resultant liquid mass is subsequently freed from the accompanying **cell debris** and **gluten** (insoluble protein) by passing through a number of fine sieves. The milky slurry thus obtained is a mixture of starch and gluten particles which is then subjected to centrifugation by custom-designed **starch purification centrifuges**. Thus, the starch which being relatively heavier settles at the bottom and the gluten being lighter floats on the surface and removed quickly by a jet of water. Consequently, the starch is washed thoroughly with successive treatment of fresh water, centrifuged or filter pressed and ultimately dried either on a *moving belt dryer* or *flash dryer*.

(b) Method for Rice Starch

The rice* is adequately soaked in a solution of NaOH (0.5% w/v) till such time when the gluten is softened and dissolved partially. The resulting grains are wet-milled and taken up with water. The suspension is purified by repeatedly passing through sieves and the starch is recovered by centrifugation. Finally, the starch is duly washed, dried, powdered and stored in HDPE** bags.

(c) Method for Wheat Starch

Wheat being an extensively used common staple food, therefore, its utility for making starch is restricted by many government authorities. First of all the wheat flour is made into a stiff ball of

* Broken pieces of rice obtained during the polishing are mostly used for preparation of rice starch.

** HDPE : High density polyethylene.

dough which are kept for a short duration. The gluten present in the dough swells up and are shifted to grooved-rollers that move forward and backward slowly. Constant sprinkling of water is done which carries off the starch along with it whereas gluten remains as a soft elastic mass. The slurry of starch is purified by centrifugation, washed, dried, powdered and packed in HDPE bags.

(d) Method for Potato Starch

The tubers of potato are thoroughly washed to get rid of the sticking soil. These are subsequently chopped into small pieces and made into a fine pulp by crushing in a **Rasp Machine**. The resulting slurry is passed through metallic sieves to remove the cellular matter as completely as possible. The starch suspension (slurry) is purified by centrifugation, washed, dried and the stocked in HDPE bags.

Description

Starch occurs in nature as irregular, angular, white masses that may be easily reduced to power.

- Appearance** : White – rice and maize starch,
 Creamy white – Wheat starch,
 Pale yellow – potato starch,
- Odour** : Odourless
- Taste** : Bland and mucilaginous.

Nevertheless, all the four types of starch mentioned above do possess a definite shape and characteristic features as illustrated in Fig. 3.1





Fig. 3.1 Characteristic Features of (A) Maize Starch, (B) Rice Starch, (C) Wheat Starch, (D) Potato Starch.

Chemical Constituents

In general, under ideal experimental parameters hydrolysis of starch in acidic medium yields glucose in theoretical proportion that essentially represent the main building block of the starch molecule. It has been established that starch molecule is essentially made up of *two complex polysaccharides*, namely:

(a) Amylopectin: (α -Amylose)

Amylopectin is insoluble in water and swells in it thereby giving rise to a thick paste upon boiling with water. It produce a distinct violet or bluish red colouration with iodine* solution (0.1 N). It has a highly branched structure that is composed of several hundred short chains of about 20-25 **D-Glucose units** each. Interestingly, one terminal of each of these chains is joined through C-1 to a C-6 with the next chain and so on and so forth as shown below:




Amylopectin (Chair – Conformations Anticipated)

(b) Amylose: (β -Amylose)

Amylose is water soluble and gives an instant bright blue colour with iodine solution (0.1 N). Based on the fact that amylose upon hydrolysis yields the only **disaccharide (+) – Maltose** and the only **monosaccharide D-(+) – Glucose**, it has been suggested that amylose is comprised of chains of a number of D-(+) – glucose units, whereby each unit is strategically linked by an **alpha-glycoside bondage** to C-4 of the next unit as depicted below:

* Iodine colour reaction is influenced by the starch-chain ie; longer the branching the colour varies from Blue→Blue violet→Red→Brown.



Amylose (Chair Conformations Anticipated)

Amylose invariably constitutes upto 25% of the total starch content; however, the proportion varies with the particular species under consideration. Amylose is found to be either absent or present to a very small extent ($\leq 6\%$) in some glutinous or waxy starches available in the plant kingdom.

Substituents and Adulterants

A number of biological species containing starch is generally employed to substitute (adulterate) the conventional commercially available starch used as **food** and as **pharmaceutical adjuvants**, namely:

S.No.	Name	Biological Source	
1.	Topioca Starch or Cassava Starch	<i>Manihot esculenta</i> Pohl., <i>Manihot aipi</i> Pohl., <i>Manihot utilissima</i> Pohl.	Family: Euphorbiaceae
2.	Sago Starch	<i>Metroxylon sagu</i>	
3.	Brazilian Arrowroot Starch or Sweet-Potato Starch	<i>Ipomoea batatas</i> Lam.	
4.	Nuts Starch	<i>Tapa bispinosa</i> Roxb.	Family: Onagraceae

Uses


1. It possesses both absorbent and demulcent properties.
2. It is employed in dusting powder because of its unique protective and absorbent property.
3. It is used in the formulation of tablets and pills as a vital disintegrating agent and a binder.
4. It is utilized as a diagnostic aid for the proper identification of crude drugs.
5. It is employed as a diluent (or filler) and lubricant in the preparation of capsules and tablets.
6. It is used as an indicator in iodimetric analyses.
7. It is an antidote of choice for iodine poisoning.
8. Dietetic grades of corn starch are marked as '**Maizena**' and '**Mondamin**'.
9. '**Glycerine of starch**' is used not only as an emolient but also as a base for the suppositories.
10. It is the starting material for the large scale production of liquid glucose, glucose syrup, dextrose and dextrin.
11. It finds its extensive industrial application for the sizing of paper and textile.
12. It possesses nutrient properties as a food and in cereal based weaning foods for babies e.g., **Farex**^(R) (Glaxo) and **Cerelac**^(R) (Nestle).

13. It is used topically and externally to allay itching.
14. It is used profusely in laundry starching.

3.2.1.3 Hetastarch

Hetastarch is a semisynthetic material that essentially comprises of more than 90% amylopectin, which has been treated with *pyridine* and *ethylene chlorohydrin*, so as to give rise to 7 to 8 hydroxyethyl substituents present for every **10 D-glucopyranose units** of the starch polymer. The molecular weight is approximately 450, 000 daltons.

Amylose Derivative



Where R or R' = H or CH₂ CH₂ OH

Amylopectine Derivative It specifically differs from the amylose derivative in that the sequence is frequently interrupted by a unit in which R is the residue of an additional ***o*-hydroxyethylated α-D-glucopyranosyl** moiety that essentially constitutes the first unit in a branch or sub-branch of the polymer.

Uses

1. It serves as a '**Plasma Volume Expander**'. A 6% solution is osmotically equivalent to a 5% albumin solution. But in the blood, it draws up certain quantum of water either from the intestinal or intracellular fluids, thus expanding the blood volume slightly in excess of the volume infused. This acquired expansion lasts for 24 to 36 hours.
2. It is employed in the management and treatment of **hypovolemic shock***.
3. It is also used as a suspension medium for **leukapheresis****.
4. It is employed as a **cryoprotective***** agent for erythrocytes.

3.2.1.4 Inulin

It is found to bear a close resemblance to starch except that it is a *levulan* rather than a **dextran**. The following characteristic features make it altogether different from starch, namely:

- Gives yellow colouration by iodine.
- Does not gelatinize with water.

* Shock caused due to diminished blood volume.

** The separation of leukocytes from blood, which are then transfused back into the patient.

*** A chemical that protects cells from the effect of cold.

- Not commonly found in plants in the form of granules having concentric layers, and
- Upon hydrolysis in acidic medium yields fructoses.

Synonyms Dahlin; Alantin; Alant starch.

Biological Source It occurs in certain plants of the *Compositae* family, such as:

<i>Inula helenium</i> Linn.	: Roots contain inulin;
<i>Eupatorium cannabinum</i> Linn.	: Plant contains inulin;
<i>Cynara scolymus</i> Linn.	: Flower heads contain inulin;
<i>Carpesium cernuum</i> Linn.	: Roots contain inulin;
<i>Calendula officinalis</i> Linn.	: Roots contain inulin;
<i>Aretium lappa</i> Linn.	: Roots contain 45% inulin.

Geographical Source

A. lappa – Western Himalayas from Kashmir to Simla;

C. officinalis – India, Pakistan;

C. cernuum – Temperate Himalayas and Nilgiri Hills (India);


S. scolymus – Throughout India; *E. cannabinum*—Temperate Himalayas;

I. helenium – Europe and Asia.

Preparation Isolated from *dahlia tubers* and from other members of the family *Compositae*.

Description The crystals are spherical in shape when prepared from water.


Chemical Constituents




Inulin

[Structure of INULIN showing arrangement of *Fructofuranose* residues in chain]

Inulin is quickly hydrolyzed by acids to D-fructose by the enzymes inulase but does not undergo hydrolysis by the amylases. However, methylated inulin upon hydrolysis gives rise to 3,4,6-trimethylfructose as a major product and 1,3,4,6-tertamethylfructose as a minor product, thereby suggesting that the fructose residues are present in the furanose form and the adjacent units are joined through C-1 and C-2 (*i.e.*, the glycosidic hydroxyl moiety).



3,4,6-Trimethyl Fructose
(Furanose-Form)
(Major Product)



1,3,4,6 – Tetramethyl Fructose
(Furanose-Form)
(Minor Product)

Uses

1. It is used in **culture media** as a fermentative identifying agent for certain bacterial species.
2. It is filtered exclusively by the glomeruli and is neither secreted nor reabsorbed by the tubules. Hence, it is employed as a **diagnostic agent** for evaluation of glomerular filtration *i.e.*, renal-function test (or kidney function test).
3. It is considered to be valuable in the diet of the diabetic patients.

3.2.1.5 Lichenin

Synonym Moss starch; Lichen starch;

Biological Source *Cetraria islandica* (L.) Ach., Family: *Parmeliaceae*. It is known as Iceland Moss.

Description It is a cellulose like polysaccharide which occurs as a cell wall component in lichens. It is readily soluble in hot water to give rise to a colloidal solution. It is more rapidly hydrolyzed than cellulose. It produces cellobiose upon acetylation with acetic anhydride and sulphuric acid. It is a white powder.

On methylation followed by hydrolysis it yields 2,3,6-trimethylglucose as a major component and tetramethylglucose as minor one thereby suggesting that the chain present in lichenin is not branched at all as in cellulose.

Chemical Constituents The exact chemical structure of lichenin molecule is yet to be established; however, it has been indicated that it may contain both β -1, 4 and β -1, 6 linkages.

3.2.1.6 Dextran

Dextran is a carbohydrate substance made up predominantly of D-glucose units *i.e.* $(C_6H_{10}O_5)_n$. It is α -1, 6 linked polyglucan.

Synonyms Dextraven; Expandex; Gentran; Hemodex; Intradex; Macrose; Onkotin; Plavolex; Polyglucin.

Biological Source A number of organisms produce dextrans; however, only *two* of them, namely; *Leuconostoc mesenteroides* and *L. dextranicum*, belonging to the family *Lactobacteriaceae*, have been used commercially.


Preparation Commercially, dextrans are manufactured by the process of fermentation of sucrose (a disaccharide) either by cell-free enzymatic fermentation technique or by whole culture technique.

The enzymes that are responsible for producing dextrans from sucrose as a substrate are collectively known as ‘**dextran-sucrases**’.

Native dextran possesses a very high molecular weight, whereas the clinical dextrans have lower molecular weights, for instance; **Dextran 40** [Gentran 40^(R) (Baxter); Rheomacrodex^(R) (Pharmacia)]; **Dextran 70** [Hyskon^(R) & Macrodex^(R) (Kabi Pharmacia)]; **Dextran 75** [Gentran 75^(R) (Baxter)]; These may be accomplished by controlled depolymerization *e.g.*, ultrasonic vibration, fungal dextranase and acid hydrolysis.

Description Dextrans obtained by the precipitation from methanol vary considerably with regard to their characteristic physical and chemical properties which solely depends upon the individual method of preparation.

Chemical Constituents The interaction between ‘**n**’ molecules of sucrose and ‘**x**’ number of glucose moieties yields dextran together with ‘**n**’ molecules of fructose as shown in the following equation:



Uses

1. Dextran 40 is employed as an isotonic solution either to prime pumps or to improve flow in surgery concerned with *cardiopulmonary bypass*. Thus, it exerts its effect by lowering the viscosity of blood and improving flow; and the latter is caused due to hemodilution.
2. Dextrans as a whole helps in minimising platelet adhesiveness, which property is gainfully exploited in their usage for *prophylaxis of thrombosis and thromboembolism both during and after surgery*.
3. Both Dextran 70 and Dextran 75 find their extensive use as plasma extender for the control and management of *hypovolemic shock*. Hypertonic solutions usually afford the dehydration of tissues, whereby the abstracted water being added to the plasma causing increase in its volume. For this reason they are quite useful in the prevention and treatment of *toxemia of pregnancy* and *nephrosis*.
4. Dextran 70 and Dextran 75 are used in 6% solutions to prevent pending shock caused by hemorrhage, trauma, and severe burns.
5. Dextran 40 (10% solution) is not only used to lower blood viscosity but also to improve microcirculation at low flow rates.
6. Dextrans are employed in the formulation of fat-soluble vitamins (*viz.* Vitamin A, D, E, K).
7. It is also used in preparing sustained released tablets.
8. Dextrans find their abundant applications in various types of confectionaries, for instance: ice-creams, candies, jellies, syrups and cake-topings.
9. It is employed as an adjunct in cosmetic preparation exclusively meant for soothing wrinkles.

3.2.1.7 Cyclodextrins

Cyclodextrins invariably consist of 6, 7 or 8 molecules (*viz.* α , β and γ cyclodextrin) in a 1, 4-configuration to result into the formation of rings having various diameters. In fact, based on the geometry of the chiral isomer, only one would possibly gain entry into the cavity in the ring while the other is excluded evidently.

Synonyms Cyloamyloses; Cycloglycans; Scharinger dextrans.

Biological Source Starch on being treated with the amylase of *Bacillus macerans*, a specific enzyme, gives rise to a mixture of cyclodextrins. They are naturally occurring carbohydrates.

Preparation It is obtained from the action of *B. macerans* amylase on starch to yield homogeneous cyclic α – (1 \rightarrow 4) linked D-gluco-pyranose units.

Description The various rings constituting the cyclodextrins appear to be as doughnut shaped. However, α -cyclodextrin *i.e.*, the smallest of the lot, has a diameter about two times that of 18-crown-6 (*viz.*, as crown ethers) and its hole (4.5 \AA across) is approximately two times as broad.

Chemical Constituents Cyclodextrins mainly are comprised of *three* different types, as detailed below:

S.No.	Name/Mol. Formula	Chemical Name	Shape
1.	α -Cyclodextrin (C ₃₆ H ₆₀ O ₃₀)	Cyclohexamylose	Hexagonal plates, or blood-shaped needles
2.	β -Cyclodextrin (C ₄₂ H ₇₀ O ₃₅)	Cycloheptaamylose	Parallelogram shaped crystals
3.	γ -Cyclodextrin (C ₄₈ H ₈₀ O ₄₀)	Cyclooctamylose	Square plates or rectangular rods.

The structure of α -cyclodextrin may be represented in Figure 3.2, in *two* different manners, namely:




Fig. 3.2 A Schematic Representation of α -Cyclodextrin:
(a) Chair-conformation Based (b) Doughnut Shaped.

- (a) Chair-conformation based cyclic structure, and
(b) Doughnut shaped or like a tiny-pail with the bottom knocked out.

Uses

1. As enzyme models based on the fact that, like enzymes, they first and foremost *bind* the substrate and then, through substituent groups, *act on it*.
2. As a complexing agent to explore the various types of enzyme action.
3. It may be employed as an additive to the mobile phase (in HPLC*), but it invariably gets bound to an inert support material.

3.2.1.8 Cellulose

Cellulose represents one of the most widely distributed and abundantly available organic matter on this planet. It is, in-fact, the most important structural element of *higher-plant-cell walls*. In nature, wood (40-50% cellulose) caters as the major source of cellulose for industrial utilities, whereas cotton (98% cellulose) provides the balance requirement globally.

Geographical Source It has been observed that nearly thirty billion MT of carbon is transformed annually into organic compounds by higher plants and out of this approximately 1/3rd is converted into cellulose. As cellulose is profusely utilized in the form of wood to build houses, paper industry and textile industry, a considerable amount of research has been duly conducted on this well-known polysaccharide.

Preparation The scientific and large-scale methods for preparing cellulose essentially involves the removal of excess of the non-cellulose substances *e.g. Lignin*. In fact, there are *three* well defined and established procedures whereby the undesired '*lignin content*' present in wood shavings are removed exhaustively, namely:

- (a) **Treatment with Sodium Bisulphite [Sulphonate Process]:** The small wood chips are boiled with a solution of sodium bisulphite whereupon the lignin is removed as *lignosulphonate*,
- (b) **Treatment with Sodium Hydroxide [Soda Process]:** The wood chips on being boiled with sodium hydroxide solution removes the lignin content as soluble products, and
- (c) **Treatment with NaOH and Na₂SO₄ [Sulphate Process]:** The sodium sulphide (Na₂S) obtained by the interaction of NaOH and Na₂SO₄ will remove most of the lignin component from the wood shavings.


However, the traces of lignin may be removed by bleaching with chlorine. The remaining mixture of hemicellulose and cellulose are subsequently extracted by subjecting it to alkaline treatment. The readily soluble hemicellulose are removed by treatment with higher concentration of NaOH solution, whereas the cellusans (Xylans) may be removed by treatment with a 5% solution of NaOH.

Description **Cellulose** has molecular weights ranging from 250,000 to 1,000,000 or even more. It is assumed that at least 1500 glucose units may be present in each molecule. Based on the findings by X-ray analysis and electron microscopy it is revealed that these long chains lie side by-side in bundles, held together by H-bonds available between the huge number of adjoining –OH moieties. Further, these bundles are twisted together to give rise to rope-like structures, that ultimately are clubbed together to yield the normal apparently visible fibers. Interestingly, in the case of wood

* **HPLC:** High performance liquid chromatography.

these cellulose “ropes” are meticulously embedded in lignin to afford a structure that resembles to concrete reinforced structures used for making buildings.

Chemical Constituents Cellulose is comprised of chains of D-glucose units, whereby each unit is joined by a glycosidic linkage to C-4 of the next unit.



Cellulose

Cellulose Cellulose derived from various sources and also from different modes of preparations usually display great differences not only in their mean chain length but also in their degree of homogeneity. Generally, the cellulose that are distinctly more homogenous are the most suitable for industrial utilities.

Uses

1. The viscose when forced through a spinnerette into an acid-bath, it gives rise to the generation of cellulose as fine filaments that yield threads of a substance termed as RAYON.
2. Cellulose undergoes an analogous reaction to produce cellulose xanthate, that is made to dissolve in alkali to yield a viscous colloidal dispersion known as VISCOSE.
3. Methyl, ethyl and benzyl ethers of cellulose are proved to be important in the commercial production of films, textiles and various types plastic materials.

3.2.1.9 Absorbent Cotton

Synonyms Purified cotton; Cotton wool; Surgical cotton.

Biological Source Cotton comprises of the *epidermal trichomes* (or hairs) from the seeds of different species of *Gossypium*, such as : *G. herbaccum*; *G. hirsutum*; *G. barbedense*; belonging to the family *Malvaceae*. In fact, absorbent cotton or purified cotton consists exclusively of the trichomes that are completely freed from adhering impurities, fat, properly bleached and finally sterilized.

Geographical Source Cotton is produced on large scale in USA, Egypt, China, India, South America and certain parts of Africa. Egyptian cotton–yarn enjoys a world-wide reputation. Both India and China are not only self-sufficient in the production of absorbent cotton but also exports a substantial quantity to various countries.

Preparation The cotton plant after flowering bears fruits which are also called ‘capsules’ or ‘balls’. These are usually 3-5 celled. Once the fruit ripens they open-up widely that contains a number of seeds per loculus. The brown coloured seeds are normally surrounded with a thick mass of white hairs. The *long-lint hairs* are known as ‘staple’ or ‘floss’; whereas the *short-fuzz hairs* are called ‘linters’. The cotton fibres (*i.e.*, mass of white hairs) along with their seeds are collected manually by hand picking. The raw cotton is subjected to a mechanical process called ‘ginning’ whereby only the hairy substance is collected separately and the undesired substances, such as: dirt, leaf-fragments and other foreign materials are removed separately. ‘Delinting’ is the mechanical

process which discards the short hairs that eventually passed along with the cotton fibres obtained from the 'ginning' process. The raw segregated long-sized cotton hairs are subsequently freed from colouring matters and traces of wax and oil coating the hairs which render them non-absorbent. The treated absorbent cotton obtained above is processed through the 'carding machine' so as to arrange the fibres in parallel direction and also to get rid of immature fibres completely. Short fibres are once again removed by 'combing' mechanically. Finally, the processed cotton fibres are defatted (with alkali) washed, bleached (with chlorinated soda) and then washed (with diluted mineral acid). It is again washed, dried, recarded and sterilized.

Description White, soft, fine, filament like hairs appearing under the microscope as hollow, flattened and twisted bands, striate and slightly thickened at the edges, practically odorless and tasteless. Cotton fibres are usually 2.5 to 4.5 cm in length and 25 to 35 μ in diameter.

Chemical Constituents Absorbent cotton is mostly cellulose 93-94% and moisture 6-7% (for structure see under section 3.2.1.8)

Uses

1. It is employed as surgical dressings.
2. It is mostly used in the textile industry to prepare a wide range of fibres.
3. It is invariably employed as its derivatives to be recognized as the most versatile adjunct in pharmaceutical formulations, for instance:

Microcrystalline cellulose	–	as Tablet Disintegrant
Carboxymethyl Cellulose (CMC)	–	as Binder and thickening agent;
Cellulose acetate phthalate	–	as an Enteric coating material;
Ethyl Cellulose	}	– as Binder and Film coating substance;
Methyl Cellulose		
Hydroxypropyl methyl Cellulose		
Oxidised Cellulose	–	as local Haemostatic;
Purified 'Rayon'	–	as Surgical aid;
Pyroxylin	–	as an ingredient in the preparation of Collodian and nail polishes.

4. It is used as a filtering medium and also as an insulating material.
5. Pharmaceutical grade cotton seed oil is used as an emolient and in the preparation of *Steroid Hormone Injections*.
6. It is used for making explosives.

3.2.1.10 Dextrin

Synonyms British Gum; Starch Gum; Leicom; Pyrodextrin; Torrefaction dextrin; Canary dextrin; Yellow dextrin; White dextrin.

Preparation Dextrin is prepared by carrying out the incomplete hydrolysis of starch with dilute acid or by heating dry starch.

Various types of dextrin are prepared as detailed below namely:

- (a) **British Gum, Starch Gum:** It is produced at high temperature in the absence of acid.

Characteristic features:

- (i) Dark brown colour, odourous,
- (ii) High viscosity, very soluble in cold water,
- (iii) Does not reduce Fehling's Solution, and
- (iv) Gives reddish-brown colour with iodine.

(b) **Canary Dextrin, Yellow Dextrin:** It is prepared by hydrolyzing starch at high temperature for a longer duration but in the presence of small quantum of acid.

Characteristic features:

- (i) Light brown to yellow colour, slight odour, and
- (ii) Low viscosity, very soluble in cold water.

(c) **White Dextrin:** It is prepared by hydrolysis at low temperature for a shorter duration but in the presence of large quantum of acid.

Characteristic features:

- (i) White colour, odourless,
- (ii) Slightly soluble in cold water, and gives a red colour with iodine, and
- (iii) Very soluble in hot water and gives a blue colour with iodine.

Uses

1. As an excipient for dry extracts and pills.
2. It is used for preparing emulsions and dry bandages.
3. It is employed for thickening of dye-pastes and mordants used in printing fabrics in fast colours.
4. It is used for sizing paper and fabrics.
5. It is employed for preparing felt and printing tapestries.
6. It is used for preparing printer's inks, glues and mucilage.
7. It is employed for polishing cereals.
8. It is extensively used in making matches, fireworks and explosives.

3.2.2 Heteroglycans

In general, **Gums** represent a heterogenous group of acidic substances, that essentially possess in common the characteristic property of swelling in water to form either gels or viscous, sticky, solutions. It has also been advocated that gums are the resulting products obtained from normal plant metabolism. In addition, it is also believed that gums may have been produced from starch or cellulose through hydrolysis, followed by oxidation to uronic acids and finally undergoing the process of esterification or formation of salt accordingly.

In actual practice, the **natural gums** may be classified into *four* different groups, namely:

- (a) Exudate Gums
- (b) Seed Gums
- (c) Marine Gums, and
- (d) Microbial Gums

These different types of gums would be discussed along with some typical examples in the sections that follows:

3.2.2.1 Exudate Gums

It has been observed that a large number of plants which grow in a semiarid environment generate **exudate gums** in sufficient amount when either an incision is made on their bark or they get damaged that invariably helps to seal off the cracked wound thereby preventing dehydration of the plant.

A plethora of exudate gums find their abundant applications as a pharmaceutical aid, namely: **Acacia, Tragacanth and Karaya Gum.**

3.2.2.1.1 Acacia

Synonyms Indian Gum; Gum Acacia; Gum Arabic.

Biological Source According to the USP, **acacia** is the dried gummy exudation from the stems and branches of *Acacia senegal* (L.) Willd; family; *Leguminosae*, or other African species of Acacia. It is also found in the stems and branches of *Acacia arabica*, Willd.

Geographical Source The plant is extensively found in India, Arabia, Sudan and Kordofan (North-East Africa), Sri Lanka, Morocco, and Senegal (West Africa). Sudan is the major producer of this gum and caters for about 85% of the world supply.

Cultivation and Collection **Acacia** is recovered from wild as well as duly cultivated plants in the following manner, such as:

- (a) **From Wild Plants:** The Gum after collection is freed from small bits of bark and other foreign organic matter, dried in the sun directly that helps in the bleaching of the natural gum to a certain extent, and
- (b) **From Cultivated Plants:** Usually, transverse incisions are inflicted on the bark which is subsequently peeled both above and below the incision to a distance 2-3 feet in length and 2-3 inches in breadth. Upon oxidation, the gum gets solidified in the form small translucent beads, sometimes referred to as 'tears'. Tears of gum normally become apparent in 2-3 weeks, which is subsequently hand picked, bleached in the sun, garbled, graded and packed.

Description

Colour: Tears are usually white, pale-yellow and sometimes creamish-brown to red in colour. The powder has an off-white, pale-yellow or light-brown in appearance.


Odour: Odourless (There is a close relationship between colour and flavour due to the presence of tannins).

Taste: Bland and mucillagenous.


Shape & Size: Tears are mostly spheroidal or ovoid in shape and having a diameter of about 2.5-3.0 cm.

Appearance: Tears are invariably opaque either due to the presence of cracks or fissures produced on the outer surface during the process or ripening. The fracture is usually very brittle in nature and the exposed surface appears to be glossy.

Chemical Constituents Acacia was originally thought to be composed only of *four* chemical constituents, namely : (–) **arabinose**; (+) – **galactose**; (–)–**rhamnose** and (+) **glucuronic acid**.



On subjecting the gum acacia to hydrolysis with 0.01 N H_2SO_4 helps in removing the combined product of (-) – arabinose and (+) – galactose, whereas the residue consists of the product (+) – galactose and (+) – glucuronic acid. These two products are formed in the ratio of 3:1.



It also contains a **peroxidase enzyme**.

Chemical Tests

1. **Lead Acetate Test:** An aqueous solution of acacia when treated with lead-acetate solution it yields a heavy white precipitate.
2. **Borax Test:** An aqueous solution of acacia affords a stiff translucent mass on treatment with borax.

3. **Blue Colouration due to Enzyme:** When the aqueous solution of acacia is treated with benzidine in alcohol together with a few drops of hydrogen peroxide (H_2O_2), it gives rise to a distinct-blue colour indicating the presence of enzyme.
4. **Reducing Sugars Test:** Hydrolysis of an aqueous solution of acacia with dilute HCl yields reducing sugars whose presence are ascertained by boiling with Fehling's solution to give a brick-red precipitate of cuprous oxide.
5. **Specific Test:** A 10% aqueous solution of acacia fails to produce any precipitate with dilute solution of lead acetate (a clear distinction from Agar and Tragacanth); it does not give any colour change with Iodine solution (a marked distinction from starch and dextrin); and it never produces a bluish-black colour with $FeCl_3$ solution (an apparent distinction from tannins).

Uses

1. The mucilage of acacia is employed as a demulscent.
2. It is used extensively as a vital pharmaceutical aid for emulsification and to serve as a thickening agent.
3. It finds its enormous application as a binding agent for tablets *e.g.*, cough lozenges.
4. It is used in the process of '*granulation*' for the manufacture of tablets. It is considered to be the gum of choice by virtue of the fact that it is quite compatible with other plant hydrocolloids as well as starches, carbohydrates and proteins.
5. It is used in conjunction with gelatin to form conservates for microencapsulation of drugs.
6. It is employed as colloidal stabilizer.
7. It is used extensively in making of candy and other food products.
8. It is skillfully used in the manufacture of spray – dried '*fixed*' flavours – stable, powdered flavours employed in packaged dry-mix products (puddings, desserts, cake mixes) where flavour stability and long shelf-life are important.

3.2.2.1.2 Tragacanth

Synonym Gum Tragacanth

Biological Source The dried gummy exudation from *Astragalus gummifer* Labill. (white gavan) or other Asiatic species of *Astragalus* belonging to the family of *Leguminosae*.

Geographical Source It is naturally found in various countries, *viz.*, Iran, Iraq, Armenia, Syria, Greece and Turkey. A few species of *Astragalous* are located in India, *viz.*, Kumaon, Garhwal and Punjab. *Persian tragacanth* are exported from Iran and North Syria, whereas the *Smyrna tragacanth* from the Smyrna port in Asiatic Turkey.

Collection The thorny shrubs of **tragacanth** normally grow at an altitude of 1000-3000 meters. As an usual practice transverse incisions are inflicted just at the base of the stem, whereby the gum is given out both in the pith and medullary rays. Thus, the absorption of water helps the gum to swell-up and subsequently exude through the incisions. The gummy exudates are duly collected and dried rapidly to yield the best quality white product. It usually takes about a week to collect the gum exudates right from the day the incisions are made; and this process continues thereafter periodically.

Description

Colour: White or pale-yellowish white

Odour: Odourless

Taste: Tasteless

Shape: Curved or twisted ribbon-like flakes marked with concentric ridges that is indicative of successive exudation and solidification. Fracture is normally short and horny.

Size: Flakes are usually $25 \times 12 \times 12$ mm.

Appearance: Translucent

Chemical Constituents Interestingly, tragacanth comprises of two vital fractions: *first*, being water-soluble and is termed as '**tragacanthin**' and the *second*, being water-insoluble and is known as '**bassorin**'. Both are not soluble in alcohol. The said two components may be separated by carrying out the simple filtration of a very dilute mucilage of tragacanth and are found to be present in concentrations ranging from 60-70% for bassorin and 30-40% for tragacanthin. Bassorin actually gets swelled up in water to form a gel, whereas tragacanthin forms an instant colloidal solution. It has been established that no methoxyl groups are present in the tragacanthin fraction, whereas the bassorin fraction comprised of approximately 5.38% methoxyl moieties. Rowson (1937) suggested that the gums having higher methoxyl content *i.e.*, possessing higher bassorin contents, yielded the most viscous mucilages.

Chemical Test

1. An aqueous solution of tragacanth on boiling with conc. HCl does not develop a red colour.
2. Ruthenim Red* solution (0.1% in H₂O) on being added to powdered gum tragacanth whereby the particles will not either acquire a pink colour or are merely stained lightly.
3. When a solution of tragacanth is boiled with few drops of FeCl₃ [aqueous 10% (w/v)] it produces a deep-yellow precipitate.
4. It gives a heavy precipitate with lead acetate.
5. When tragacanth and precipitated copper oxide are made to dissolve in conc. NH₄OH it yields a meagre precipitate.

Substituents/Adulterants **Karaya gum** which is sometimes known as **sterculia gum** or **Indian tragacanth** and is invariably used as a substitute for gum tragacanth.

Uses

1. It is used as a demulcent in throat preparations.
2. It is employed as an emolient in cosmetics (*e.g.*, hand lotions).
3. It is used as a pharmaceutical aid as a suspending agent for insoluble and heavy powders in mixtures.
4. It is effectively employed as a binding agent for the preparation of tablets and pills.
5. It is also used as an emulsifying agent for oils and waxes.
6. A substantial amount find its application in calico printing and in confectionary.
7. It is used in making medicinal jellies *e.g.*, spermicidal jelly.
8. A 0.2-0.3% concentration is frequently used as a stabilizer for making ice-creams and various types of sauces *e.g.*, tomato sauce, mustard sauce.
9. It is used to impart consistence to troches.
10. The mucilages and pastes find their usage as adhesives.

* Ruthenium oxychloride ammoniated, C₁₆H₄₂N₁₄O₂R_{u3}, soluble in water and used in microscopy as reagent for pectin and gum.

3.2.2.1.3 Karaya Gum

Synonyms Gum Karaya; Kaday; Katilo; Kullo; Kuteera; Sterculia; Indian Tragacanth; Mucara.

Biological Source **Karaya Gum** is the dried exudate of the tree *Sterculia urens* Roxb; *Sterculia villosa* Roxb; *Sterculia tragacantha* Lindley and other species of *Sterculia*, belonging to the family: *Sterculeaceae*. It is obtained from *Cochlospermum* *Geographical Source: gossypium, De Candolle or other species of cochlospermum Kunth* –family: *Bixaceae*.

Geographical Source The *S. urens* is found in India especially in the *Gujarat* region and in the central provinces.

Preparation The gum is obtained from the *Sterculia* species by making incisions and, thereafter, collecting the plant exudates usually after a gap of 24 hours. The large irregular mass of gums (tears) which weigh between 250 g to 1 kg approximately are hand picked and despatched to the various collecting centres. The gum is usually tapped during the dry season spreading over from March to June. Each healthy fully grown tree yields from 1 to 5 kg of gum per year; and such operations may be performed about five times during its lifetime. In short, the large bulky lumps (tears) are broken to small pieces to cause effective drying. The foreign particles *e.g.*, pieces of bark, sand particles, leaves are removed. Thus, purified gum is available in *two* varieties, namely:

- (a) **Granular or Crystal Gum:** Having a particle size ranging between 6 to 30 mesh, and
- (b) **Powdered Gum:** Having particle size of 150 mesh

Description

Colour : White, pink or brown in colour

Odour : Slight odour resembling acetic acid

Taste : Bland and mucilageous taste

Shape : Irregular tears or vermiform pieces.

It is water insoluble but yields a translucent colloidal solution.

Chemical Constituents **Karaya gum** is partially acetylated polysaccharide containing about 8% acetyl groups and about 37% uronic acid residues. It undergoes hydrolysis in an acidic medium to produce (+)-galactose, (-)-rhamnose, (+)-galacturonic acid and a trisaccharide acidic substance. It contains a branched heteropolysaccharide moiety having a major chain of 1, 4-linked α -(+)-galacturonic acid along with 1, 2-linked (-)-rhamnopyranose units with a short (+)-glucopyranosyluronic acid containing the side chains attached 1→3 to the main chain *i.e.*, (+)-galactouronic acid moieties.

Chemical Test It readily produces a pink colour with a solution of Ruthenium Red.

Substituent/Adulterant It is used as a substitute for **gum tragacanth**.

Uses

1. It is employed as a denture adhesive.
2. It is used as a 'binder' in the paper industry.
3. It is also employed as a thickening agent for dyes in the textile industry.
4. It is widely used as a stabilizer, thickner, texturizer and emulsifier in foods.

5. It is used as a bulk laxative.
6. It finds its usage in lozenges.
7. It is employed extensively in wave set solution and in skin lotions.
8. It is used in preparations concerned with composite building materials.

3.2.2.2 Seed Gums

Seed Gums are hydrocolloids contained in some seed embryos where they actually play the role as polysaccharide food reserves.

A few typical examples of seed gums are described below, such as: Plantago seed; Pectin; Locust bean gum; and Guar gum.

3.2.2.2.1 Plantago Seed The origin of the word '**Plantago**' is from the Latin and means sole of the foot, referring to the shape of the leaf. Likewise, '*Psyllium*' is from the Greek and means *flea*-describing the seed.

Synonyms Psyllium seed; Plantain seed; Flea seed; Ispaghula; Isapgol; Isabgul.

Biological Source It is the dried ripe seeds of *Plantago psyllium* L., or *Plantago arenaria* Waldst & Kit (*P. ramosa* Asch.) (Spanish or French psyllium seed) or of *Plantago ovata* (blond or Indian plantago seed) or of *Plantago amplexicaulis* belonging to the family: *Plantaginaceae*.

Geographical Source *P. amplexicaulis* is grown on the Panjab plains, Malwa and Sind and extending to Southern Europe.

P. psyllium is an annual pubescent herb practically native to the Mediterranean countries. It is grown in France and constitutes the main bulk of the American imported psyllium seed.

P. ovata is extensively grown in Pakistan; besides it is found to be native to Mediterranean countries and Asia.

Preparation The crops are grown usually on light, well drained sandy loamy soils; and during their entire growth period the climate must be cool and dry. The ripe and matured fruits are normally collected after a span of about three months. The seeds are separated by thrashing lightly on a solid support. The dust and foreign particles are removed by sieving and against a current of moderate air-blast.

Description

Colour : Pinkish grey to brown


Odour : No characteristic odour

Taste : Bland and mucilaginous

Weight : 100 seeds weigh between 0.15-0.19 g

Figure 3.3 gives an account of the dorsal surface as well as the ventral surface of *Ispaghula seed* and *Psyllium seed* along with their overall shape, size and outersurface.

Chemical Constituents **Plantago** seeds generally comprise of approximately 10% of mucilage invariably located in the epidermis of the testa together with proteins and fixed oil. The mucilage essentially consists of **pentosan** and **aldobionic acid**.




Shape : Ovate or boat shaped

Size : Length = 1.8-3.5 mm, Width = 1.0-1.7 mm.

Outersurface: The *Convex surface* has a central brown oval spot, whereas the *Concave surface* bears a deep furrow having its hilum covered with a thin whitish membrane.

Fig. 3.3 Dorsal and Ventral Surfaces of *Ispaghula Seed* and *Psyllium Seed*.



The various products of hydrolysis are, namely: **xylose, arabinose, rhamnose and galacturonic acid.**

Chemical Tests

1. Its mucilage gives a distinct red colouration on treatment with Ruthenium Red solution.
2. **Swelling Factor***: It establishes the purity of the drug and ranges between 10 to 14. It is easily determined by transferring accurately 1.0g of the drug in a 25 ml measuring cylinder duly filled with 20 ml of water with intermittent shaking. The exact volume occupied by the seeds after a duration of 24 hours of wetting is noted carefully which represents the *swelling factor* of the seeds under investigation.

Substituents/Adulterants A number of species of **Plantago** have been studied extensively for their mucilage contents. Interestingly, *Plantago rhodosperma* which is particularly habitated in Missouri and Louisiana (USA) and *Plantago wrightiana* are worth mentioning. The former species

* **Swelling Factor**: It represents the quantitative swelling due to the presence of mucilage present in the drug substance.

contains mucilage to the extent of 17.5% whereas the latter contains about 23%. However, these two species compare favourably with the official drug.

In addition to the above, a few species like *P. purshii*, *P. aristata* and *P. asiatica* are also employed as a substitute for **plantago seeds**.

Uses

1. Plantago seeds are mostly employed as demulscent and in the treatment of chronic constipation.
2. It is also used in amoebic and bacillary dysentery.
3. Mucilage of the isapgol is invariably employed in the preparation of tablets (*e.g.*, granulation)
4. It is used as a stabilizer in the ice-cream industry
5. The crushed seeds are employed as a poultice for rheumatic pain
6. The *acid form of polysaccharide* is obtained by carefully removing the cations from the mucilage by treatment with cation-exchange resins and spray drying the resultant products. This '*specialized product*' finds its enormous applications as a tablet disintegrator, as enteric coating substance and finally employed in the sustained release drug formulations.

3.2.2.2.2 Pectin **Pectin**, in general, is a group of polysaccharides found in nature in the primary cell walls of all seed bearing plants and are invariably located in the middle lamella. It has been observed that these specific polysaccharides actually function in combination with both cellulose and hemicellulose as an intercellular cementing substance. One of the richest sources of pectin is lemon or orange rind which contains about 30% of this polysaccharide.

Pectin is naturally found in a number of plants namely: lemon peel, orange peel, apple pomace, carrots, sunflower-heads, guava, mangoes and papaya. The European countries, Switzerland and USA largely produce pectin either from apple pomace or peels of citrus fruits. Evaluation and standardization of pectin is based on its '*Gelly-Grade*' that is, its setting capacity by the addition of sugar. Usually, pectin having '*gelly grade*' of 100, 150 and 200 are recommended for medicinal and food usages.

Biological Sources **Pectin** is the purified admixture of polysaccharides, obtained by carrying out the hydrolysis in an acidic medium of the inner part of the rind of citrus peels, for instance: *Citrus limon* (or Lemon) and *Citrus aurantium* belonging to the family *Rutaceae*, or from apple pomace *Malus sylvestris* Mill (*Syn: Pyrus malus* Linn, family: *Rosaceae*).

Geographical Source Lemon and oranges are mostly grown in India, Africa and other tropical countries. Apple is grown in the Himalayas, California, many European countries and the countries located in the Mediterranean climatic zone.

Preparation The specific method of preparation of **pectin** is solely guided by the source of raw material *i.e.*, lemon/orange rind or apple pomace; besides the attempt to prepare either low methoxy group or high methoxy group *pectins*.

In general, the preserved or freshly obtained lemon peels are gently boiled with approximately 20 times its weight of fresh water maintained duly at 90°C for a duration of 30 minutes. The effective pH (3.5 to 4.0) must be maintained with food grade lactic acid/citric acid/tartaric acid to achieve maximum extraction. Once the boiling is completed the peels are mildly squeezed to obtain the liquid portion which is then subjected to centrifugation to result into a clear solution. From this

resulting solution both proteins and starch contents are suitably removed by enzymatic hydrolysis. The remaining solution is warmed to deactivate the added enzymes. The slightly coloured solution is effectively decolourized with activated carbon or bone charcoal. Finally, the pectin in its purest form is obtained by precipitation with water-miscible organic solvents (*e.g.*, methanol, ethanol, acetone), washed with small quantities of solvent and dried in a vacuum oven and stored in air-tight containers or polybags.


Note: As Pectin is fairly incompatible with Ca^{2+} , hence due precautions must be taken to avoid the contact of any metallic salts in the course of its preparation.

Description

- Appearance** : Coarse or fine- powder
Colour : Yellowish white
Odour : Practically odourless
Taste : Mucilaginous taste
Solubility : 1. Completely soluble in 20 parts of water forming a solution containing negatively charged and very much hydrated particles.
 2. Dissolves more swiftly in water, if previously moistened with sugar syrup, alcohol, glycerol or if first mixed with 3 or more parts of sucrose.

Chemical Constituents Pectin occurs naturally as the partial methyl ester of α (1 \rightarrow 4) linked (+) – polygalacturonate sequences interrupted with (1–2) – (–) – rhamnose residues. The neutral sugars that essentially form the side chains on the pectin molecules are namely: (+) – galactose, (–) – arabinose, (+) – xylose, and (–) – fructose.

Schneider and Bock (1938) put forward the following probable structure for **pectin galacturonan**:



Pectin Galacturonan

Chemical Tests

1. A 10% (w/v) solution gives rise to a solid gel on cooling.
2. A *transparent gel* or *semigel* results by the interaction of 5 ml of 1% solution of pectin with 1 ml of 2% solution of KOH and subsequently setting aside the mixture at an ambient temperature for 15 minutes. The resulting gel on acidification with dilute HCl and brisk shaking yields a voluminous and gelatinous colourless precipitate which on warming turns into white and flocculent.

Uses

1. It is employed mostly as an intestinal demulscent. It is believed that the unchanged molecules of polygalacturonic acids may exert an adsorbent action in the internal layers of the intestine, thereby producing a protective action along with *Kaolin* to prevent and control diarrhoea.

2. As a pharmaceutical aid pectin is used frequently as an emulsifying agent and also as a gelling agent preferably in an acidic medium.
3. It is employed extensively in the preparation of jellies and similar food products *e.g.*, jams, sauces, ketchups.
4. Pectin in the form of pastes exerts a bacteriostatic activity and hence, is used frequently in the treatment of indolent ulcers and deep wounds.
5. A combination of *pectin* and *gelatin* find its application as an encapsulating agent in various pharmaceutical formulations to afford sustained-release characteristics.

3.2.2.2.3 Locust Bean Gum

Synonym Carob Flour; Arobon; Carob Gum; Ceratonia; Johannisbrotmehl;

Biological Source The Gum essentially consists of the hydrocolloid from the powdered endosperm of tree pods of *Ceratonia siliqua* Linn, belonging to the family *Leguminosae* (**St. John's bread**). It normally takes about 15 long years for a full-grown tree to yield seeds which, therefore, restricts the provision of a regular production of the gum to cater for the ever-expanding needs for the hydrocolloids.

Geographical Source The tree is found in abundance in Egypt, Cyprus and Sicily. It is very sensitive to low temperature. It is also commercially grown in countries like: Algeria, Greece, Israel, Italy, Morocco, Portugal and Spain.

Preparation The locust bean pods comprise of about 90% pulp and 8% kernels. The kernels are separated from the pods mechanically by means of **Kibbling Machine**. The kernels comprise of mainly the endosperm (42-46%), husk (30-33%) and germ to the extent of 25%. First of all the seeds are duly dehusked and splitted lengthwise the seeds are duly dehusked and splitted lengthwise to facilitate the separation of the endosperm from the **embryo*** (*i.e.*, the yellow germ). The dried gum is pulverised and graded as per the mesh-size *e.g.*, 150, 175 and 200 mesh sizes available in the European market.

Description

Colour : Translucent white, yellow green

Odour : Odourless

Taste : Tasteless and mucilaginous. Acquires a leguminous taste when boiled with water

Solubility : Insoluble in alcohol. Dispersable in concentration upto 5%

Viscosity : As it is a neutral polysaccharide, hence pH has no effect on viscosity between 3-11.

Chemical Constituents **Locust bean gum** comprises of *proteins*, for instance: albumins, globulins and glutelins; *carbohydrates*, such as: reducing sugar, sucrose, dextrans, and pentosans; besides ash, fat, crude fiber and moisture.

* Embryo enhances the rate of fermentation of gum solutions and hence it must be removed as completely as possible.

Chemical Tests The mucilage of this gum when gently boiled with 5% KOH solution it yields a clear solution; but agar and tragacanth gives rise to a yellow colour, whereas karaya gum produces a brown colour.

Substituent Adulterant In food industry it is employed as a substitute for strach.

Uses

1. It is used as a stabilizer, thickner and binder in foods and cosmetics.
2. It is widely employed as a sizing and finishing agent in textiles.
3. It finds its abundant use as fiber - bonding in paper manufacture.
4. It is used as an adsorbent - demulcent therapeutically.
5. It is employed as drilling mud additive.

3.2.2.2.4 Guar Gum

Synonyms Guar flour; Decorpa; Jaguar; Gum cyamopsis; Cyamopsis gum; Burtonite V-7-E.

Biological Source **Guar gum** is the ground endosperms of *Cyamopsis tetragonolobus* (L.) Taub; belonging to family *Leguminosae*.

Geographical Source It grows abundantly in tropical countries like: Indonesia, India, Pakistan and Africa. In USA, southern western regions it was introduced in the year 1900 and its large-scale production commenced in early 1950's.

Preparation First of all the fully developed white seeds of Guar gum are collected and freed from any foreign substances.

The sorted seeds are fed to a mechanical '**splitter**' to obtain the *bifurcated guar seeds* which are then separated into husk and the respective cotyledons having the '*embryo*'. The gum is found into the endosperm. Generally, the guar seeds comprise of the following:

Endosperm	: 35 to 40%
Germ (or Embryo)	: 45 to 50%, and
Husk	: 14 to 17%


The **cotyledons**, having a distinct bitter taste are separated from the endosperm by the process called '**winnowing**'. The crude guar gum *i.e.*, the endosperms is subsequently pulverised by means of a '*micro-pulveriser*' followed by grinding. The relatively softer cotyledons sticking to the endosperms are separated by mechanical '*sifting*' process. Thus, the crude guar-gum is converted to a purified form (*i.e.*, devoid of cotyledons), which is then repeatedly pulverized and shifted for several hours till a final white powder or gramular product is obtained.

Description

Colour	: Colourless; Pale-yellowish white powder
Odour	: Characteristic smell
Taste	: Mucilagenous
Solubility	: Insoluble in alcohol with water it gives a thick transparent suspension

Chemical Constituents It has been found that the water soluble fraction constitutes 85% of Guar gum and is commonly known as **Guaran**. It essentially consists of linear chains of (1 → 4) -β-D-

mannopyranosyl units with α -D-galactopyranosyl units attached by (1 \rightarrow 6) linkages. However, the ratio of D-galactose to D-mannose is 1:2.



Guaran

Chemical Tests

1. On being treated with iodine solution (0.1 N) it fails to give olive-green colouration.
2. It does not produce pink colour when treated with Ruthenium Red solution (distinction from sterculia gum and agar)
3. A 2% solution of lead acetate gives an instant white precipitate with guar gum (distinction from sterculia gum and acacia)
4. A solution of guar gum (0.25 g in 10 ml of water) when mixed with 0.5 ml of benzidine (1% in ethanol) and 0.5 ml of hydrogen peroxide produces no blue colouration (distinction from gum acacia).

Uses

1. It is used therapeutically as a bulk laxative.
2. It is employed as a protective colloid.
3. It is also used as a thickener and its thickening property is 5 to 8 times more than starch.
4. It finds its use in peptic ulcer therapy.
5. It is used as an anorectic substance *i.e.*, it acts as an appetite depressant.
6. It is employed both as a binding and a disintegrating agent in tablet formulations.
7. It is used in paper sizing.
8. It is abundantly employed as film forming agent for cheese, salad dressing, ice-cream and soups.
9. It is used in pharmaceutical jelly formulations.
10. It is widely used in suspensions, emulsions, lotions, creams and toothpastes.
11. It is largely used in mining industry as a flocculant and also as a filtering agent.
12. It is also employed in water treatment plants as a coagulant aid.

3.2.2.3 Marine Gums

A variety of algae and seaweeds comprise of **marine gums** as components of cell walls and membranes or present in the intracellular regions where they actually serve as reserve food material.

A few typical examples of marine gums would be discussed in the sections that follow.

3.2.2.3.1 Algin

Synonyms Sodium alginate; Alginic acid sodium; Sodium polymannuronate; Kelgin; Minus; Protanal;

Biological Source **Algin** is a gelling polysaccharide extracted from the giant brown seaweed (*giant kelp. Macrocystis pyrifera* (L.) Ag., *Lessoniaceae*) or from *horsetail kelp (Laminaria digitata* (L.) Lamour, *Laminariaceae*) or from *sugar kelp (Laminaria saccharina* (L.) Lamour). Some other common species are *Laminaria hyperborea* and *Ascophyllum nodosum*

Geographical Source The different varieties of seaweeds are invariably found in the Pacific and Atlantic Oceans, more specifically along the coastal lines of USA, Canada, Scotland, Japan and Australia. In India the Western coast of Saurashtra is also a potential source of algin. However, USA, and UK are the largest producers of algin in the world.

Preparation The **algin** (or sodium alginate) is the sodium salt of alginic acid which is a purified carbohydrate extracted from brown seaweed (algae) by the careful treatment with dilute sodium hydroxide. The brown colour of the crude algin is due to the presence of a carotenoid pigment associated with it which may be eliminated by treating the aqueous solution with activated carbon and spray drying the powder.

Description

Colour : Yellowish-white, cream coloured, buff coloured


Odour : Odourless

Taste : Tasteless

Solubility : Insoluble in alcohol, ether, chloroform and strong acids, freely soluble in water

Viscosity : A 1% (w/v) aqueous solution at 20°C may show a viscosity ranging between 20-400 *centipoises*.

Chemical Constituents Alginic acid is mainly comprised of D-mannuronic acid residues which on methylation and hydrolysis gives rise to the formation of 2,3 dimethyl-D- mannuronide. Therefore, the ring as well as bridge oxygen atoms involve C-4 and C-5 and the carboxyl groups are absolutely free to react (to form sodium salts), whereas the aldehydic moieties are duly utilized by the respective glycosidic linkages. It has been observed that these mannuronic acid entities are joined by β -1, 4-glycosidic linkages. The resulting structure could be either linear or very slightly branched.



Alginic Acid

Chemical Tests

1. The aqueous solution of algin gives an instant white copious precipitate with calcium chloride solution.

2. A 1% (w/v) aqueous solution of algin yields a heavy gelatinous precipitate with diluted sulphuric acid.
3. It is not precipitated by saturated ammonium sulphate solution (distinction from agar and tragacanth)
4. It gives effervescence (liberates CO₂) with carbonates.
5. It readily reacts with compounds having ions of alkali metals (*e.g.*, Na⁺, K⁺, Li⁺) or ammonium (NH₄⁺) or magnesium (Mg²⁺) to produce their respective alginates (salts) that are *water soluble* and forms thick and viscous solutions characteristic of *hydrophillic colloids*.

Uses

1. It is extensively used in the manufacture of ice-creams where it serves as a stabilizing colloid, ensuring creamy texture thereby checking the growth of ice-flakes (or crystals).
2. It is also used in the flocculation of suspended solids in most water treatment plants.
3. It is employed as a stabilizing and thickening agent in food and pharmaceutical industry.
4. It is used as a film and film-forming agent in the rubber and paint industry.
5. It is widely used in the textile industry as absorbable haemostatic dressings.
6. It is employed as a binding and disintegrating agent for tablets and lozenges.

3.2.2.3.2 Agar

Synonyms Agar-agar; Gelose; Japan-agar; Chinese-isinglass; Bengal isinglass; Ceylon isinglass; Layor carang; Vegetable gelatin.

Biological Source **Agar** is the dried hydrophilic colloidal polysaccharide complex extracted from the agarocytes of algae belonging to the class *Rhodophyceae*. It is also obtained as the dried gelatinous substance from *Gelidium amansii* belonging to the family *Gelidaceae* and several other species of red algae, such as *Gracilaria* (family: *Gracilariaceae*) and *Pterocladia* (*Gelidaceae*). The predominant agar-producing genera are, namely; *Gelidium*; *Gracilaria*; *Acanthopeltis*; *Ceramium* and *Pterocladia*.

Geographical Source **Agar** is largely produced in Japan, Australia, India, New Zealand, and USA. It is also found in Korea, Spain, South Africa and in the Coastal regions of Bay of Bengal (India) together with Atlantic and Specific Coast of USA.

Preparation It is an usual practice in Japan where the **red-algae** is cultivated by placing poles or bamboos spread in the ocean which will serve as a support and shall augment the growth of algae on them. During the months of May and October the poles are removed and the algae are carefully stripped off from them. The fresh seaweed thus collected is washed thoroughly in water and subsequently extracted in digestors containing hot solution of dilute acid (1 portion of algae to 60 portions of diluted acid). The mucilagenous extract is filtered through linen while hot and collected in large wooden troughs to cool down to ambient temperature so as to form solid gel. The gel is mechanically cut into bars and passed through a wire netting to form strips. The moisture from the strips is removed by successive **freezing and thawing*** and finally sun dried and stored as thin agar strips.

Alternatively, the mass of gel if frozen and subsequently thawed and the dried agar is obtained by vacuum filtration. The crude agar is usually formed as flakes which can be powdered and stored accordingly.

* To bring down to room temperature from -20 to -30°C.

Description

- Colour** : Yellowish white or Yellowish grey
Odour : Odourless
Taste : Bland and mucilaginous
Shape : It is available in different shapes, such as: bands, strips, flakes, sheets and coarse powder
Size : Bands: width = 4cm; Length = 40 to 50 cms
 : Sheets: Width = 10-15cm; Length = 45 to 60 cms
 : Strips: Width = 4mm; Length = 12 to 15 cms

India produces about 250 MT of good quality agar using *Galidiella accrosa* as the raw material. It is insoluble in cold water in organic solvents. It readily dissolves in hot solutions and it forms a translucent solid mass which characteristic is very useful in microbiology for carrying out the *Standard Plate Count*.

Chemical Constituents **Agar** can be separated into two major fractions, namely: (a) *Agarose*—a neutral gelling fraction; and (b) *Agaropectin*—a sulphated non-gelling fraction. The former is solely responsible for the *gel-strength of agar* and consists of (+)-galactose and 3,6-anhydro-(−)-galactose moieties; whereas the latter is responsible for the *viscosity of agar solutions* and comprises of sulphated polysaccharide wherein both uronic acid and galactose moieties are partially esterified with sulphuric acid. In short, it is believed to be a complex range of polysaccharide chains having alternating α -(1→3) and β -(1-4) linkages and varying *total charge content*.

Chemical Tests

1. It gives a pink colouration with Ruthenium Red solution.
2. A 1.5-2.0% (w/v) solution of agar when boiled and cooled produces a *stiff-jelly*.
3. Prepare a 0.5%(w/v) solution of agar and add to 5 ml of it 0.5 ml of HCl, boil gently for 30 minutes and divide into two equal portions:
 - (a) To one portion add BaCl_2 solution and observe a slight whitish precipitate due to the formation of BaSO_4 (distinction from Tragacanth), and
 - (b) To the other portion add dilute KOH solution for neutralization, add 2 ml of Fehling's solution and heat on a water bath. The appearance of a brick red precipitate confirms the presence of galactose.

Substituents/Adulterants **Gelatin** and **isinglass** are usually used as substituents for **agar**.

Uses

1. It is used in making photographic emulsions.
2. It is also employed as a bulk laxative.
3. It is extensively used in preparing gels in cosmetics.
4. It is widely used as thickening agent in confectionaries and dairy products.
5. It is used in the production of ointments and medicinal encapsulations.
6. In microbiology, it is employed in the preparation of bacteriological culture media.
7. It is used for sizing silks and paper.
8. It finds its enormous usage in the dyeing and printing of fabrics and textiles.
9. It is also used as dental impression mould base.
10. It is employed as corrosion inhibitor.

3.2.2.3.3 Carrageenan

Synonyms Irish moss; Chondrus.

Biological Source Carrageenan refers to closely associated hydrocolloids which are obtained from different red algae or seaweeds. The most important sources of carrageenan are namely: *Chondrus crispus* (Linn.) stockhouse and *Gigartina mamillosa* (Goodnough and Woodward) J. Agardh belonging to the family *Gigartinaceae*.

Geographical Source The plants are abundantly found along the north-western coast of France, the coast of Nova Scotia, and the British Isles.

Preparation In general, the plants are collected mostly during June and July, and spread out on the bench for natural drying. They are then exposed to the sun rays directly whereby bleaching takes place. Now, they are treated with a brine solution, and ultimately dried and stored.

Description The *Chondrus* is more or less an allusion to the cartilage-like characteristic features of the dry thallus; whereas *Gigartina* is an absolute allusion specifically to the fruit bodies which appear as raised tubercles on the thallus.

Chemical Constituents The carrageenan bears a close physical resemblance to agar. However, its hydrocolloids are mostly galactans having sulphate esters, which are present in excess amount in comparison to agar. Carrageenan polysaccharides essentially comprise of chains of 1, 3 linked β – (+) – galactose and 1,4-linked α - (+) – galactose moieties that are invariably substituted and later on modified to the 3, 6- anhydro derivative. In fact, carrageenans may be further separated into three major components, namely: **k-carrageenan**; **i-carrageenan**; and **λ -carrageenan**.

Uses

1. Both **k-and i-carrageenans** proved to be good gelating agents because of the fact that they tend to orient in stable helices when in solution.
2. The λ -carrageenan does not form stable helices and hence represent the nongelling portion of the carrageenans which serves as a more useful thickener.
3. The fairly stable texture and supported by excellent rinsability of the hydrocolloids these are immensely useful in the formulations of toothpastes.
4. They are used as bulk laxative.
5. They are employed as a demulscent.
6. They constitute as an important ingredient in a large number of food preparations.

3.2.2.4 Microbial Gums

Microbial Gums are produced by certain selected microorganisms in the course of fermentation. The resulting exopolysaccharides are usually isolated from the fermentation broth by appropriate procedures.

A few typical examples are described below, for instance: Xanthan Gum; Chitin.

3.2.2.4.1 Xanthan Gum

Synonyms Polysaccharide B-1459; Keltrol F; Kelzan.

Biological Source The Polysaccharide Gum is produced by the bacterium *Xanthomonas campestris* on certain suitable carbohydrates.

Preparation One of the latest techniques of 'biotechnology' i.e., 'recombinant DNA technology' has been duly exploited for the commercial production of xanthan gum.

Methodology: First of all the genomic banks of *Xanthomonas campestris* are meticulously made in *Escherichia coli* by strategically mobilizing the broad-host-range cosmids being used as the vectors. Subsequently, the conjugal transfer of the genes take place from *E. coli* into the nonmucoid *X. campestris*. Consequently, the wild type genes are duly separated by virtue of their unique ability to restore mucoid phenotype. As a result, a few of the cloned plasmids incorporated in the wild type strains of *X. campestris* shall afford an increased production xanthan gum.

Interestingly, the commercial xanthan gums are available with different genetically controlled composition, molecular weights and as their respective sodium, potassium or calcium salts.

Description It is a cream coloured, odourless and free flowing powder. It dissolves swiftly in water on shaking and yields a highly viscous solution at relatively low concentrations. The aqueous solutions are extremely pseudoplastic in character. It gives rise to a strong film on evaporation of its aqueous solutions. It is fairly stable and resistant to thermal degradation. The viscosity is independent of temperature between 10 to 70°C. It is fairly compatible with a variety of salts.

Chemical Constituents **Xanthan gum** is composed of chiefly D-glucosyl, D-mannosyl and D-glucosyluronic acid residues along with variant quantum of O-acetyl and pyruvic acid acetal. The primary structure essentially comprises of a cellulose backbone with trisaccharide side chains and the repeating moiety being a pentasaccharide.

Uses

1. Its potential in chemically enhanced oil recovery is well known.
2. The inherent pseudoplastic property of its aqueous solutions rendered both toothpastes and ointments in enabling them to hold their shape and also to spread readily.
3. It is extensively employed in pharmaceuticals due to its superb suspending and emulsifying characteristic features.

3.2.2.4.2 Chitin **Chitin**, is the nitrogen containing polysaccharide which invariably occurs in certain fungi e.g., ergot. It is also commonly found in some invertebrate animals eg, crab, shrimp, lobster-specifically located in the exoskeleton of the body. Besides, it is located in the appendages of insects and crustaceans.

Biological Source The mycelia of *Penicillium* species contain approximately 20% of chitin. The relatively hard crustacean shells of crab and lobster are reported to contain 15 to 20% chitin, whereas the rather soft crustacean shells of shrimp contain between 15 to 30% chitin. It is found in the spores of many fungi and yeasts.


Preparation The hard or soft **crustacean shells** are first ground to fine powder and then treated with dilute HCl (5%) for a duration of about 24 hours whereby most of the calcium* and other impurities are eliminated completely as soluble CaCl₂. The above extract containing the proteins derived from the shells are eliminated by treating it with proteolytic enzymes like *pepsin* or *trypsin*.

* Crustacean shell contain approx. 60-75% of calcium carbonate.

The resulting pink coloured liquid extract is bleached by H_2O_2 in an acidic medium for 5-6 hours at room temperature. The bleached product is subjected to deacetylation at $120^\circ C$ with a mixture of 3 parts of KOH, 1 part EtOH and 1 part ethylene glycol. The deacetylation process is repeated several times till the 'acetyl content' reaches a minimum level. **Chitin** is obtained as an amorphous solid substance.

Description It is an amorphous solid. It is practically insoluble in water, dilute acids, dilute and concentrated alkalis, alcohol and other organic solvents. It is soluble in concentrated HCl, H_2SO_4 , anhydrous HCOOH and H_3PO_4 (78-97%). There exists a wide difference with regard to the solubility, molecular weight, acetyl value and specific rotation amongst chitins of different origin and obtained from different methods.

Chemical Constituents **Chitin** may be regarded as a derivative of cellulose, wherein the C-2 hydroxyl groups have been duly replaced by acetamido residues. In fact, it is more or less a cellulose like biopolymer mainly consisting of unbranched chains of β -(1 \rightarrow 4) -2- acetamido -2-deoxy-D-glucose. It is also termed as N-acetyl-D-glucosamine. It contains about 6.5% of nitrogen.



Chemical Tests

1. **Chitin** affords a brown colour with Iodine solution which turns into red violet on acidification with sulphuric acid.
2. **Chitin sulphate** gives rise to a characteristic strain with acidic dyes, such as: picric acid and fuchsin.
3. When chitin is heated with strong KOH solution under pressure it fails to dissolve, but undergoes deacetylation to form acetic acid and other products collectively termed as *chitosans*.
4. Hydrolysis of **chitin** in the presence of strong mineral acids forms acetic acid and glucosamine.
5. When **chitin** is dissolved in dilute nitric acid (50%) and allowed to crystallise overnight it forms beautiful spherocrystals of chitosan. These crystals on being examined under polarised light, by making use of crossed nicols, a distinct cross is observed.

Uses


1. **Chitosan** *i.e.*, **deacetylated chitin**, finds its application in water treatment operations.
2. It is used in pholographic emulsions.
3. It is used in improving the dyeability of synthetic fibers and fabrics.
4. Therapeutically it is used in wound healing preparations.
5. It shows considerable adhesivity to plastics and glass.
6. It is used as a sizing agent for cotton, wool, rayon and for synthetic fibers.

3.3 CARBOHYDRATE BIOGENESIS

One of the most vital aspects of pharmacognosy which has gained paramount importance and legitimate recognition in the recent past is the intensive and extensive studies involving the various biochemical pathways that has ultimately led to the production of ‘**secondary constituents**’ invariably employed as ‘**drugs**’. This type of specific and elaborated study is frequently termed as **biogenesis** or **drug biosynthesis**. It is quite pertinent to mention here that as it is absolutely necessary for a medicinal chemist to understand the intricacies of chemical synthesis of a potent drug substance, such as: naproxen, chloramphenicol etc., exactly in the same vein a pharmacognosist must possess a thorough knowledge of the biogenesis of drugs of natural origin.

A Swiss chemist G. Trier, as far back in 1912, not only predicted but also postulated that amino acids and their corresponding derivatives invariably act as the precursors of relatively complex naturally occurring alkaloids mostly used as potent therapeutic agents. Interestingly, soon after the second half of the twentieth century, there had been a tremendous progress in the era of **isotopically labelled organic compounds** that facilitated the affirmation as well as confirmation of the earlier speculated theories.

With the advent of most advanced knowledge in sciences it has been established that the carbohydrate biogenesis usually takes place due to the *Photosynthesis* from carbon dioxide (CO₂) as the starting material occurring abundantly both in all plants and in certain purple bacteria as depicted below:



However, the general pattern of ‘**Carbohydrate Biogenesis**’ may be shown explicitly in the following Fig. 3.4.




Fig. 3.4 Carbohydrate Biogenesis

13. Whistler, R.L., and J.N. Bemiller: (Eds), '**Industrial Gums, Polysaccharides and their Derivatives**', Academic Press, New York, 1959.
14. Weymouth-Wilson AC: **The Role of Carbohydrates in Biologically Active Natural Products**, *Nat. Prod. Rep.* **14**, 99–110, 1997.
15. Yalpani, M.: (Ed), '**Progress in Biotechnology**, Vol. 3, **Industrial Polysaccharides**, Elsevier Science Publishers, B.V. Amsterdam, 1987.

4

Glycosides

- | | |
|--|---|
| <ul style="list-style-type: none">• Introduction• Classification• Biosynthesis of Glycosides | <ul style="list-style-type: none">• Profile of Glycosides in Natural Plant Sources• Further Reading References |
|--|---|

4.1 INTRODUCTION

Glycosides, in general, are defined as the condensation products of sugars with a host of different varieties of **organic hydroxy** (occasionally *thiol*) **compounds** (invariably monohydrate in character), in such a manner that the hemiacetal entity of the carbohydrate must essentially take part in the condensation. It is, however, pertinent to state here that the *polysaccharides* are also encompassed in this broad-based overall definition of **glycosides**. The noncarbohydrate moiety is usually termed as **aglycone** (or **aglycon**), or a **genin**.

The rather older or trivial names of glycosides usually has a suffix 'in' and the names essentially included the source of the glycoside, for instance: **strophanthidin** from *Strophanthus*, **digitoxin** from *Digitalis*, **barbaloin** from *Aloes*, **salicin** from *Salix*, **cantharidin** from *Cantharides*, and **prunasin** from *Prunus*. However, the systematic names are invariably coined by replacing the "ose" suffix of the parent sugar with "oside". The *stereochemical anomeric prefix* α or β and the *configurational prefix* (*D- or L-*) immediately precede the sugar nomenclature, and lastly the *chemical name of the aglycone* precedes the name of the sugar. It may be expatiated with the help of the following examples:

- Aloin (or Barbaloin):** 10-Glucopyranosyl-1, 8-dihydroxy-3-(hydroxymethyl)-9 (10H)-anthracenone;
- Salicin:** 2-(Hydroxymethyl) phenyl- -D-glucopyranoside;
- Amygdalin:** D-Mandelonitrile- β -D-glycosido-6- β -D-glucoside;
- Digitoxin:** 3-[0-2, 6-Dideoxy- β -D-ribo-hexopyranosyl -(1 \rightarrow 4)-O-2, 6-dideoxy- β -D-ribo-hexopyranosyl (1 \rightarrow 4), 2, 6-dideoxy- β -D-ribo hexopyranosyl oxy]-14-hydroxycard- 20(22)-enolide.

Interestingly, the **glycosides** may be regarded as internal acetate. The two series of stereoisomeric glycosides are usually termed as **α and β glycosides**. Thus, taking into consideration the simple example of **methyl D-glucosides**, these α and β structures may be represented as shown below:




Fig. 4.1 Methyl- α -D-Glucoside






Fig. 4.2 Methyl- β -D-Glucoside

Figures 4.1 and 4.2 represent the **open-chain structure, cyclic structure and boat configuration of methyl- α -D-glucoside and methyl- β -D-glucoside** respectively. In this particular instance the glycosidic linkage is established by dehydration involving a *hydroxy group* of the *aglycone portion* (*i.e.*, methyl alcohol) and the *hydroxyl group* on the *hemiacetal carbon of the carbohydrate* (in question), thereby ultimately resulting into the formation of an *acetal type of structure*.

α -Configuration If the —OCH_3 moiety (generalized as OR') is in opposite steric sense as the $\text{—CH}_2\text{OH}$ moiety on C-5 (for D-family sugars), the glycosidic structure is designated as **α -configuration**.

β -Configuration If the —OCH_3 moiety is in the same steric sense as the CH_2OH group on C-5, the glycosidic structure is designated as **β -configuration**.

It has been observed that the substantial quantum of naturally occurring glycosides essentially possess the **stereo-configuration**. However, this observation may be further expatiated with the help of the following typical examples of **β -amygdalin** and **β -salicin**:



Sailent Features (β -Amygdalin):

- (i) Glycosidic linkage is β -because it is hydrolysed by **emulsin** (an enzyme),
- (ii) The linking oxygen is on the same side of the plane of the ring as the CH_2OH moiety on C-5.
- (iii) It contains several asymmetric C atoms *i.e.*, **chiral centres**, and
- (iv) It is **optically active**.

Sailent Features (β -Salicin):

1. Hydrolysed by emulsin, hence it has **β -configuration**,
2. The linking oxygen is on the same side of the plane of the ring as the CH_2OH moiety on C-5,
3. It has several **chiral centres**, and
4. It is **optically active**.


Glycosides, are found to exert a wide spectrum of therapeutic actions, both in modern medicines and in traditional medicaments, ranging from *cardiotonic, analgesic, purgative, and anti-rheumatic, demulcent* and host of other useful actions.

The Glycosidic Linkages The exact point of linkage between the carbohydrate (sugar) and non carbohydrate (aglycone) moieties is an '**oxygen bridge**' that essentially connects the reducing group present in carbohydrate to either an alcoholic or a phenolic group present in the non carbohydrate. Such glycosides are collectively termed as **O-glycosides**. However, if the 'O' is replaced by 'S' it is called **S-glycosides**; if replaced by 'N' is known as **N-glycosides**; and if replaced by 'C' is termed as **C-glycosides**.

These **four** types of glycosides shall be described briefly at this juncture with appropriate examples from the domain of medicinal plants.


4.1.1 O-Glycosides

The **O-glycosides** are usually represented as follows:



These are most abundantly found in nature in the higher plants, such as: senna, rhubarb and frangula.


Examples: Rhein-8-glycoside obtained from rhubarb.




Rhein-8-Glycoside

4.1.2 S-Glycosides

The **S-glycosides** are normally designated as below:




The presence of S-glycosides is more or less restricted to isothiocyanate glycosides, such as: **Sinigrin**, obtained from black mustard seeds. (*i.e.*, **Brassica campestris** Family: *Cruciferae*)



Sinigrin


4.1.3 N-Glycosides

The **N-glycosides** may be represented as shown below:




The most typical example of N-glycosides is the nucleosides, wherein the respective amino group of the base ultimately reacts with —OH group ribose/deoxyribose.

Examples: Adenosine: It is widely distributed in nature *e.g.*; from yeast nucleic acid.




4.1.4 C-Glycosides

The C-glycosides may be designated as shown under:




C-Glycosides are present in a variety of plant substances, such as: **aloin** in Aloe; **cascaroside** in Cascara.

(i) **Aloin or Barbaloin:**



Aloin (or Barbaloin)

(ii) **Cascarosides (A, B, C and D):**



Cascarosides

4.2 CLASSIFICATION

In reality, the most befitting classification of glycosides is rather a hard-nut-to-crack. In case, the classification is to be governed by the presence of sugar moiety, a good number of rare sugars are involved, if the aglycone function forms the basis of classification, one may come across groups from probably all major categories of plant constituents identified and reported. Of course, a therapeutic classification offers not only a positive edge over the stated classification but also affords an excellent means from a pharmaceutical viewpoint, but it grossly lacks a host of glycosides which are of great pharmacognostic interest.

The most acceptable **classification of glycosides** is based on the chemical nature of the **aglycone moiety** present in them, namely:

- (i) Anthracene glycosides
- (ii) Phenol glycosides
- (iii) Steroid glycosides
- (iv) Flavonoid glycosides
- (v) Coumarin and Furanocoumarin glycosides
- (vi) Cyonogenetic glycosides
- (vii) Thioglycosides
- (viii) Saponin glycosides
- (ix) Aldehyde glycosides
- (x) Bitter glycosides
- (xi) Miscellaneous glycosides

All these **different categories of glycosides** would be discussed individually with appropriate examples in the sections that follows:

4.2.1 Anthracene Glycosides (or Anthraquinone Glycosides)

Anthracene glycosides represent a major class of glycosides. They are abundantly found in various dicot plant families, such as: *Ericaceae*, *Euphorbiaceae*, *Leguminosae*, *Lythreaceae*, *Polygonaceae*, *Rhamnaceae*, *Rubiaceae* and *Verbenaceae* to name a few. Interestingly, some monocots belonging to the family *Liliaceae* also exhibits the presence of anthracene glycosides. Besides, they are also present in certain varieties of **fungi** and **lichens**.

A plethora of glycosides having their aglycone moieties closely related to anthracene are present in noticeable amounts in a variety of drug substances, for instance: *aloe*, *casacara*, *frangula*, *sagrada* and *senna*. These drugs are invariably employed as **cathartics**.

4.2.1.1 Aloes

Synonym: Aloe

Biological Source: Aloe is the dried latex of leaves of various species of Aloes, namely:

- Aloe barbadensis* Miller (or **Curacao Aloe**);
- Aloe ferox* Miller (or **Cape Aloe**);

Aloe perryi Baker (or **Socotrine Aloe**);

Aloe africana Miller and *Aloe spicata* Baker (or **Cape Aloe**).

All these species belong to the family *Liliaceae*.

Geographical Source

Curacao, Barbados, Aruba : Curacao Aloes or Barbados Aloes
and Bonaire (West Indian Islands)

Cape Town (South Africa) : Cape Aloes

Socotra and Zanzibar Islands : Socotrine or Zanzibar Aloes

It is also cultivated in Europe and the North West Himmalayan region in India.


Preparation

General Method The leaves are transversely cut at the base and the incised ends placed downwards in a 'V' shaped trough having a hole at its bottom. The latex drains down the trough and is collected in individual receptacles placed beneath. The latex is evaporated in a kettle made of copper till it attains such a consistency that it may be poured into metallic ingots where it gets solidified. When the latex is concentrated gradually and then cooled slowly, it gives rise to an opaque product. The aloe thus obtained is termed as '*hepatic*' or '*livery*' aloe. If the latex is concentrated rapidly, followed by sudden cooling the resulting product appears to be transparent and relatively brittle in nature. The broken surface has a vitreous or glassy surface. Such a product is commonly known as '*vitreous*', '*lucid*' or '*glassy*' aloe.

Description

S. No.	Properties	Curacao Aloes	Cape Aloes	Socotrine Aloes	Zanzibar Aloes
1.	Colour	Brownish black opaque mass	Dark brown or greenish brown to olive brown mass	Brownish yellow opaque mass	Liver brown colour
2.	Odour	Strong odour resembles with <i>Iodoform</i>	Sour and distinct odour	Unpleasant odour	Characteristic but agreeable odour
3.	Taste	Intense bitter taste	Nauseating and bitter taste	Extremely bitter and nauseous taste	Bitter taste
4.	Texture	Waxy and somewhat resinous	Breaks with a glassy fraction	Fractured surface looks conchoidal	A dull, waxy, smooth and even fracture

Chemical Constituents **Aloe-emodin** occurs in the free state and as a glycosides in various species of Aloe and also in *Rheum* (Rhubrb). **Curacao aloes** contains about two and half times the amount of aloe emodin when compared to **cape-aloes**.



Interestingly, the glycosides of anthranols, dianthrone, and oxanthrones *i.e.*, the reduced derivatives of anthraquinones, invariably found in various plant substances. These plant products do make an appreciable contribution to the inherent therapeutic values of the naturally occurring substances. The structural relationships of emodin are represented as shown in Figure 4.3.





Fig. 4.3 Emodin: Structural Relationships of its Derivatives.


Both **anthrones** and **anthranols** mostly occur either as free or combined as glycosides. From a close look at their respective structures it may be observed that they are reduced anthraquinone derivatives. Both anthrone and anthranol are isomeric in nature; however, the latter may be partially converted to the former, which is essentially a non-fluorescent substance and is not soluble in alkaline solutions. Generally, the anthrones are converted on oxidation into their corresponding anthraquinones, namely: oxanthrone and dianthrone. Hence, it has been observed that prompt oxidation usually takes place in the powdered crude drug rather than the rhizomes itself.

Besides, **aloin** (or **barbaloin**) the aloes also contain **isobarbaloin** (Curacao aloes), **β -barbaloin** = (Cape aloes), aloe emodin and resins. The principal resin present in the aloes is known as **aloesin**.



Aloesin

γ -Coniceine, which is a piperidine alkaloid is found in *Aloe gililandii*, *A. ballyi*, and *A. ruspoliana* (*Liliaceae*)

 γ -Coniceine

Aloe yields not less than 50% of water soluble extractives. It also contains volatile oil to some extent that imparts a characteristic odour to it.

Chemical Tests The overall chemical tests for *aloes* may be divided into *two* separate heads, namely: (a) General Tests, and (b) Special Tests

(a) General Tests: For this prepare a 0.1% (w/v) aqueous solution of aloes by gentle heating, add to it 0.5g of Kiesulgur and filter through. Whatman Filter Paper No. 42 and preserve the filtrate for the following tests:

1. **Borax Test (or Schoenteten's Reaction):** To 5 ml of the above test solution add 0.2 g of pure borax and heat gently till it gets dissolved. Transfer a few drops of the resulting solution into a test tube filled with distilled water, the appearance of a green coloured fluorescence due to the formation of *aloe emodin anthranol* shows its presence.
2. **Bromine Test:** When equal volumes of the test solution and bromine solution are mixed together, it yields a pale-yellow precipitate due to the production of tetrabromaloin.
3. **Modified Borntrager's Test:** It is known that aloin (or barbaloin) belongs to the class of **C-glycoside** which does not undergo hydrolysis either by heating with dilute acid or alkali, but it may be decomposed with ferric chloride due to oxidative hydrolysis. Hence, the **Modified Borntrager's test** employing FeCl_3 and HCl is used as stated below:

First of all heat together 0.1 g of powdered aloe with about 2 ml of FeCl_3 solution (5% w/v) and 2 ml of dilute HCl (6N) in a test tube over a pre-heated water bath for 5 minutes. Cool the contents and extract the liberated anthraquinone with carbon tetrachloride. Now carefully separate the lower layer of CCl_4 and add to it ammonia solution. The appearance of a rose-pink to cherry red colour confirms its presence.

(b) Special Tests

1. **Nitrous Acid Test:** Crystals of sodium nitrite together with small quantity of acetic acid when added to 5 ml of the above test solution of aloe, the following observations are noted:
 - (a) **Curacao Aloes:** A sharp pink to carmine colour due to the presence of isobarbaloin.

(b) **Cape Aloes:** A faint pink colour due to isobarbaloin.

(c) **Socotrine and Zanzibar aloes:** Colour comparatively lesser change in colour.

2. **Nitric Acid Test:** The Test solution of aloes when made to react with nitric acid, it gives rise to various shades of colour due to different types of aloes available commercially as shown below:

Caracao Aloe : Deep brownish red

Cape Aloes : Initial brownish colour changing to green

Socotrine Aloes : Pale brownish yellow

Zanzibar Aloes : Yellowish brown

3. **Cupraloin Test (or Klunge's Isobarbaloin Test):** To 10 ml of a 0.4% (w/v) aqueous solution of aloe add a drop of the saturated solution of copper sulphate, immediately followed by 1 g of NaCl and 20 drops of ethanol (90% v/v). It produces different shades of colours depending on the variety of aloes used:

Caracao Aloes : A wine red colour lasting for few hours,

Cape Aloes : A faint colouration changing to yellow quickly,

Socotrine Aloes : No colouration

Zanzibar Aloes : No colouration

Uses

1. Though, both aloes and aloin are official drugs, the former is mostly used as a purgative by exerting its action mainly on colon, whereas the latter is generally prepared over the former now-a-days.
2. Aloes find its usefulness as an external aid to painful inflammatory manifestations.
3. It constitutes an important ingredient in the '**Compound Tincture of Benzoin**' (or **Friar's Balsam**).
4. Aloe gel made from the mucilaginous latex of *A. vera* is frequently employed in the treatment and cure of radiation burns to get immediate relief from itchings and pains.
5. Aloe usually causes griping and is, therefore, administered along with carminatives.

4.2.1.2 *Rhubarb*

Synonyms: Rheum; Radix rhei; Rhubarb rhizome.

Biological Source: **Rhubarb** is the rhizome and roots of *Rheum officinale* Bail., *R. palmatum* L., *Rheum emodi* Wall ; *R. webbianum* Royle, belonging to the family *Polygonaceae*. The rhizome and roots are mostly collected from 6-7 year old plants just prior to the following season. They are commercially available either with intact cortex or partially decorticated.

Geographical Source It is obtained largely from cultivated as well as wild species of *Rhubarb* grown in regions extending from Tibet to South East China. It is also found in Germany and several European countries. In India it is grown extensively in Kashmir, Kullu, Sikkim, Uttar Pradesh, Panjab. It is also found in Nepal. It is cultivated in Southern Siberia and North America.


Preparation The rhizomes are collected either in spring or in autumn from 6 to 10 year old plants., grown at an altitude of more tha 3, 000 meters. These are duly cleaned, decordicated and dried. The relatively larger rhizomes are cut into small pieces either longitudinally or transversely. The cut

fragments are threaded and dried in the shade. They are also dried artificially in an atmosphere of hot wooden boxes and exported for commercial consumption.


Description Rhubarb is usually found to be compact, rigid, cylindrical conical or barrel shaped with 8-10 cm length and 3-4 cm thickness. They appear to be mostly longitudinally wrinkled, ridged or furrowed; whereas a few of them do exhibit transverse annulations or wrinkles. Interestingly, the flat pieces are prepared from large rhizomes that are normally cut longitudinally and, therefore, they appear to be largely as plano-convex with tapering at both ends. These two varieties of pieces possess a sharp characteristic odour and a bitter astringent taste. The surface is often smeared with a bitter yellowish powdery substance, which on being removed gives rise to a rather smooth surface that appears to be pale brown to red in colour.

Chemical Constituents Rhubarb essentially contains mainly the anthraquinone glycosides and the astringent components. The former range between 2 to 4.5% and are broadly classified into *four* categories as stated below:


- (a) **Anthraquinones with —COOH moiety**—*e.g.*, Rhein; Glucorhein;



- (b) **Anthraquinones without —COOH moiety**—*e.g.*, Emodin; Aloe-Emodin; Chrysophanol; Physcion;




- (c) **Anthrones and Dianthrones of Emodin**—as shown below:



- (d) **Heterodianthrones**—*e.g.* Palmidin A, B, and C, which are produced from two different anthrone molecules, as stated under:

- Palmidin A : Aloe-emodin anthrone + Emodin anthrone
 Palmidin B : Aloe-emodin anthrone + Chrysophanol anthrone
 Palmidin C : Emodin anthrone + Chrysophanol anthrone

However, the astringent portion of rhubarb chiefly comprises with the following components, namely: **gallic acid** as α - and β -glucogallin; **tannin** as *d*-catechin and epicatechin.



Rhubarb in addition to the above constituents, consists of **rheinolic acid**, **pectin**, **starch**, **fat** and **calcium oxalate**. The calcium oxalate content ranges between 3-40% in various species of rhubarb which reflects directly on the corresponding **ash values** (*i.e.*, total inorganic contents).

Chemical Tests

1. The Rhubarb powder on being treated with ammonia gives rise to a pink colouration.
2. Rhubarb gives a blood-red colouration with 5% potassium hydroxide.
3. It gives a positive indication with modified Borntrager's test (see under Aloes).

Uses

1. It is used mainly in the form of an ointment in the treatment and cure of chronic eczema, psoriasis and trichophytosis—as a potent *keratolytic agent*.
2. It is employed as a bitter stomachic in the treatment of diarrhoea.
3. It is also used as a purgative.

4.2.1.3 Cascara Sagrada

Interestingly, the very name '**cascara sagrada**' is *Spanish* for the sacred bark; *Rhamnus* is the ancient classical name for buckthorn, and *Purshianus* was attributed as a mark of honour and respect to the great German botanist Friedrich Pursh.

Synonyms Sacred bark; Chitten bark; Chittin bark; Purshiana bark; Persian bark; Bearberry bark; Bearwood; Cascara bark; Cortex *Rhamni purshianae*.

Biological Source *Cascara sagrada* is the dried bark of *Rhamnus purshiana* DC., belonging to family *Rhamnaceae*, from which a naturally occurring cathartic is extracted. It is usually collected at least one year prior to its use.

Geographical Source It is invariably obtained from cultivated as well as wild shrubs and small trees grown in Northern Idaho, West to Northern California, North Carolina, Oregon, in Kenya and Western Canada.

Preparation The bark is collected, during the dry season (April to August) from the 8 to 9 years old trees that have gained a height of 16-18 meters with their stems having a diameter of 8 to 10 cm, by inflicting longitudinal incisions on the fully developed stems. In usual practice, the *coppicing technique* is employed for the collection of bark. The bark is carefully stripped off from the branches and the stems. They are subsequently allowed to dry in shade by putting their inner-surface facing the ground so as to permit the completion in the enzymatic conversion of the anthranol derivative *i.e.*, glycosides (an emetic principle) to its anthraquinone derivative usually present in the fresh drug, thereby exerting a milder cathartic activity. During this span of one year the drug must be duly protected from rain or humid environment so as to check the growth of mould.

Description


- Colour** : Outside-purplish brown; Inside reddish brown.
Odour : A typically nauseatic odour.
Taste : Persistently bitter.
Size : Occurs in varying sizes of thickness between 1 to 4 mm.
Shape : Mostly occurs in quills or channels. Also available in small, flat and broken segments.

However, internally the bark exhibits longitudinal striations; whereas externally the bark appears to be quite smooth and usually displays the presence of scattered lenticles, lichens and cork. Besides, mostly insects and liveworts are found on the exterior surface of the bark.


Chemical Constituents The *cascara sagrada* bark is found to contain **two** major types of anthracene compounds, namely:

- (a) **Normal O-Glycosides** These are based on emodin like structures and constitute about 10 to 20% of the total glycosides, and
 (b) **Aloin-like C-Glycosides** These comprise of about 80 to 90% of the total glycosides.

The two **C-glycosides** are known as **barbaloin** and **deoxybarbaloin** (or **chrysaloin**) as given below:




Barbaloin (Aloin)



Deoxybarbaloin (Chrysaloin)

The main active constituents are *four* glycosides usually designed as Cascarosides A, B, C and D. From extensive and intensive studies of these cascariosides by *optical rotary dispersion* (ORD) technique it has been established that the *cascariosides A and B* are solely based on optical isomers of **barbaloin** ; whereas *cascariosides C and D* on optical isomers of **deoxybarbaloin**. However, from a close inspection of all the four basically primary glycosides of **barbaloin** and **deoxybarbaloin** it may be revealed that they possess the characteristic features of **O-glycosides** as well as **C-glycosides**.



Glycosides	Type	R	Configuration
CASCAROSIDE	A	OH	(10 S)
CASCAROSIDE	B	OH	(10 R)
CASCAROSIDE	C	H	(10 S)
CASCAROSIDE	D	H	(10 R)

Salient Features The **salient features** of the various **glycosides** are as follows:

- (i) About six anthracene derivatives isolated and identified in the drug belong to the category of **O-glycosides** which are solely based on **emodin**,
- (ii) Dried cascara bark normally produces not less than 7% of the total hydroxyanthracene derivatives, calculated as **cascarioside A**, and
- (iii) The remaining cascariosides must make up at least 60% of this total quantum.

Perhaps the presence of a '*lactone*' in the drug attributes a bitter taste to it.

Casanthranol is the purified version of a mixture of **anthranol glycosides** highly water-soluble and duly extracted from **cascara sagrada**. It has been reported that each gramme of casanthanol contains not less than 200 mg of the entire hydroxyanthracene derivative, calculated as cascarioside A, out of which not less than 80% of the respective derivatives mainly consists of cascariosides.

Chemical Test It gives a positive indication with **Modified Borntrager's test** because of the presence of **C-glycosides**.

Substituents/Adulterants The barks of *Rhamnus californica* and *R. fallax* are generally used as a substitute for **cascara sagrada** bark. Sometimes the *frangula bark* is also used as a substitute for this drug. However, the former types of barks (*Rhamnus species*) exhibit a more uniform coat of lichens along with broader medullary rays when compared to the original drug species.

4.2.1.4 *Frangula*


Synonyms Buckthorn bark; Alder buckthorn; Black dogwood; Berry alder; Arrow wood; Persian berries;

Biological Source *Frangula bark* is the dried bark of *Rhamnus frangula* Linne belonging to the family *Rhamnaceae*.

Geographical Source The plant is a shrub which grows abundantly in Europe, the Mediterranean coast of Africa and Western Asia.

Preparation The preparation of *frangula bark* resembles to that of cascara bark (see section 4.2.1.3). Just like the cascara bark, the frangula bark must be aged for at least a period of one year before it is used therapeutically so as to permit the reduced forms of the glycosides with harsh action to be oxidised to comparatively milder forms.

Chemical Constituents The seed, bark and rootbark of *Rhamnus species*, specifically in *Alder Buckthorn* (*Rhamnus alnifolia* L' Her.); in *Rhamnus carthartica* L., in *Rhamnus purshiana* DC (*Cascara sagrada*) consists of the *two* important glycosides *Frangulins A and B*, which were initially thought to be isomeric compounds. Later on *two* more glycosides known as **glucofrangulins A and B** have also been reported. However, the structures of **Frangulins A* and B**** along with **Glucofrangulins A and B** are given below:




Frangulin A : R=H;

Frangulin B : R = H;

Glucofrangulin A : R = β -D-Glucopyranose

Glucofrangulin B : R = β -D-Glucopyranose

Besides, *frangulin* the *frangula bark* contains emodin (see Section 4.2.1.2) and chrysophanic acid as shown below:



Chrysophanic Acid

* Horhammer, Wagner, *Z. Naturforsch.*, 27B, 959, 1972.

** Wagner, Demuth, *Tetrahedron Letters*, 5013, 1972.

Substituents/Adulterants As the activity of **Frangula Bark** corresponds to that of *cascara sagrada*, it finds a good substitute and comparable usage in Europe and the Near East. Interestingly, the drug substances obtained from the ripe and duly dried fruits of *Rhamnus catharticus* Linn., are invariably employed in Europe and the Near East for their recognised cathartic therapeutic activity.

Uses It is mostly used as a cathartic.

4.2.1.5 Senna

Senna was first used in the European medicine as early as the 9th or 10th century by the Arabs. An Egyptian native Issae Judaeus (850 to 900 A.D.) was reported to be the pioneer in bringing and introducing the drug to Egypt from Mecca,

Synonyms Senna leaf; Sennae folium; Tinnevelley Senna; Indian Senna;

Biological Sources Senna is the dried leaflets of *Cassia senna* L. (*Cassia acutifolia* Delile, (Alexandria senna), or of *Cassia angustifolia* Vahl (Indian or Tinnevelley Senna) belonging to the family *Leguminosae*. However, the modern taxonomists recommend to club together both the commonly available species of senna, namely: *Alexandria senna* and *Tinnevelley senna*—under one name as *Senna alexandria* Mill.

Geographical Source *C. acutifolia* grows wild in the vicinity of Nile River (Egypt) extending from Aswan to Kordofan; whereas, *C. angustifolia* grows wild in the Arabian Peninsula, Somalia, India, and Northwest Pakistan. In India the drug is cultivated in Southern part- Tinnevelly, Mysore and Madurai; Northern part- Jammu, Western part Pune and Kutch region of Gujrat.

Preparation After a duration of 2-3 months of sowing the Alexandria senna is harvested both in April and in September by cutting off the tops of the plants approximately 15 cm from the ground level and subsequently allowing them to dry in the sun. Later on, the unwanted stems and pods are segregated from the leaflets with the help of sieves using mechanical vibrators. The portion that passes through the sieves, is now 'tossed' carefully, whereby the leaves are collected on the surface and the relatively heavier stalk fragments at the bottom. The dried leaves are now graded, packed in bags and stored in dry place. The commercial drug at present is distributed through Port Sudan located on the Red Sea, and from the Port of Tuticorin, in India.


Description

Features	Alexandria senna	Tinnevelley senna
Colour	: Pale greyish green	Yellowish green
Odour	: Slight	Slight
Taste	: Mucilagenous slightly bitter and characteristic	Mucilagenous, bitter and characteristic
Size	: Length = 2-4 cm, Width = 7-12 mm;	Length = 2.5-5cm Width = 3-8 mm
Shape	: Ovate -lanceolate;	Lanceolate
Texture	: Thin and brittle	Thin and flexible

Chemical Constituents The principle active constituents of senna are *four* sennosides A, B, C and D, which are the dimeric glycosides having their aglycones composed of either *rhein* and/or

aloe-emodin moieties *i.e.*; 10, 10' -bis (9, 10-dihydro-1, 8-dihydroxy-9-oxoanthracene-3-carboxylic acid).

The structure of the above *four* glycosides are as given below:



Glycosides	R	C-10 & C-10'	Characteristics
Sennoside-A	—COOH	<i>trans</i> -	Optically active, a levoratory isomer present in large concs; water insoluble
Sennoside-B	—COOH	<i>meso</i> -	Intramolecularly compensated present in large concs; more water soluble
Sennoside-C	—CH ₂ OH	<i>trans</i> -	Aglycone (-) isomer; present in small concs;
Sennoside-D	—CH ₂ OH	<i>meso</i> -	Aglycone (+) isomer; present in small concs.

Besides, relatively small quantities of monomeric glycosides and free anthraquinones are also present in senna pods, such as: **rhein-8-glucosides, rhein-8-diglucoside, aloe emodin-8-glucoside, aloe-emodin anthrone diglycoside, rhein, aloe-emodin and isorhamnetin.**

It also contains **kaempterol** (a phytosterol), mucilage, resins, myricyl alcohol, chrysophanic acid, calcium oxalate and salicylic acid.

Specific method of extraction for the sennosides: Exclusively for commercial purposes, the sennosides are extracted as their corresponding calcium sennosides in varying strengths because of its enhanced stability.

Methodology: The drug powder (about 80-100 mesh size) is duly macerated with either 80% acetone or 90% methanol for a period of 6 hours, followed by 2 hours with cold water. This process helps to achieve an extract that contains between 17-18% sennosides and enables to extract about 65% of sennosides from the crude drug.

The sennosides and other anthracene derivatives may be extracted by the help of a mixture of polyethylene glycols (in 70% v/v ethanol) and solutions of non-ionic surfactants.

However, the isolation of individual sennosides may be achieved by employing non-polar synthetic resins having porous structural features.

Alternatively, the drug powder is macerated with citric acid in methanol which is followed by a repeated extraction with a mixture of methanol, toluene and ammonia. The resulting extract is treated with a concentrated solution of calcium chloride to salt out the sennosides as their respective calcium salts.

Chemical Tests

1. **Modified Borntrager's Test:** It gives a pink to red colouration for the presence of anthraquinone glycosides (see under section 4.2.1.1).
2. The mucilage of senna gives a distinct red colouration with Ruthenium Red solution.

Substituents and Adulterants Tinnevelly senna is invariably found to be adulterated with the following *three* cheaper varieties of senna namely:

- (a) Dog senna ie; *Cassia abovata*,
 - (b) Palthe senna ie; *Cassia auriculata*, and
 - (c) Arabian Senna or Mecea senna or Bombay senna *i.e.*; wild variety of *Cassia angustifolia* Vahl. from Southern Arabia.
- Dog senna* : It contains approximately 1% of anthraquinone derivatives.
Palthe senna : It contains no anthraquinone glycosides
Arabian senna : It is brownish-green in appearance.

Uses

1. Senna and its branded preparations, for instance: Glaxenna^R (Glaxo); Pursennid^(R) (Sandoz); Helmacid with senna^(R) (Allenburrys); are usually employed as purgative in habitual constipation. The glycosides are first absorbed in the small intestinal canal after which the aglycone portion gets separated and ultimately excreted in the large intestine (colon). The released anthraquinones irritate and stimulate the colon thereby enhancing its peristaltic movements causing bulky and soft excretion of faeces.
2. The inherent action of senna is associated with appreciable griping, and therefore, it is generally dispensed along with carminatives so as to counteract the undesired effect.

4.2.2 Phenol Glycosides

A variety of **phenol glycosides** are widely distributed in nature. It has been found that quite a few simple phenol glycosides have their aglycone portion loaded with either phenolic moieties or more often with alcoholic moieties or carboxylic acid functions. Invariably, the natural vegetative plant products, such as: Willow bark (containing *Salicin*) and Bearberry leaves (containing *arbutin*) have been employed therapeutically since ages, the former as antipyretic and the latter both as urinary antiseptic and as diuretic.

A few frequently used **phenol glycosides** commonly found in natural plant products are described below; such as: **Arbutin; Gaultherin; Salicin; Populin; Glucovanillin.**

4.2.2.1 Arbutin

Synonyms Ursin; Arbutoside; Uvasol; Uvaursi; Bearberry leaves; Busserole.


Biological source It occurs in the dried leaves of *Bergenia crassifolia* (L.) Fritsch, belonging to the family *Saxifragaceae*.

It is also obtained from the dried leaves Uva-Ursi or Bearberry *Arctostaphylos uva-ursi* (Linne') Sprengel, belonging to family (*Ericaceae*) and other related plants e.g., **coactylis** or adenotricha Fernald and McBride (family *Ericaceae*). Besides, it is extracted from the leaves of blueberry, cranberry and pear trees (*Pyrus communis* L., family; *Rosaceae*).

Geographical Source Bearberry is mostly grown in various parts of North and Central Europe, North America, Canada and Scotland.


Description **Arbutin** occurs in white needles that are promptly soluble in water and ethanol. It is very hygroscopic in nature.

Chemical Constituents The structure of **arbutin** is given below:




It has a β -D-glucopyranoside function attached to the *para* position of a phenol.

It yields upon hydrolysis, either by dilute acids or by emulsin, one mole each of D-glucose and hydroquinone.



Besides, the leaves also contain **methyl arbutin**, **quercetin**, **gallic acid**, **ursolic acid** and **tannin**. However, **arbutin** forms a complex with hexamethylenetetramine that may be used as a means to separate it from **methylarbutin**.



Chemical Tests

1. **Arbutin** yields a blue colouration with ferric chloride solution.
2. Its presence in crude drug may be detected by first moistening the powdered tissues with dilute HCl, warming cautiously over a watch glass on a low flame and carefully collecting the sublimate as crystals of hydroquinone that forms on another watch glass.

Important Features The presence of **gallotannin** usually helps in preventing certain specific enzyme, for instance: **β -glucosidase** from splitting **arbutin** that justifiably explains why the crude plant extracts are more effective therapeutically, as compared to **pure arbutin**.

Uses

1. It is used as a diuretic.
2. It finds its application as an antiseptic agent on the urinary tract.
3. It also exerts astringent actions.

4.2.2.2 Gaultherin


Synonyms Monotropitoid; Monotropitin; Methyl salicylate 2-glucoxyloside.

Biological Source It occurs in the leaves of the Canadian Wintergreen plant *Gaultheria procumbens* L., in *Monotropa hypopitys* L., belonging to family *Ericaceae*. It is also found in the bark of *Betula lenta* L., family *Betulaceae*; in *Spiraea ulmaria* L., and *S. filipendula* L., family *Rosaceae*.


Geographical Source It grows in the hills of India, Burma and Ceylon.

Description It has a needle-shaped star formation look when crystallised from acetone (99%). It is soluble in water and alcohol.

Chemical Constituents When **gaultherin** is hydrolysed with 3% H_2SO_4 , it forms one mole each of **methyl salicylate**, **D-glucose** and **D-xylose** as shown below:



However, **gaultherin** (or **monotropitin**) on being subjected to hydrolysis by the enzyme *gaultherase* gives rise to the production of one mole each of **primeverose** [*i.e.*; **6-(β -D-xyloside)-D-glucose**] a disaccharide and methyl salicylate.



4.2.2.3 Salicin

Synonyms Salicoside; Salicyl alcohol glucoside; Saligenin β -D-glucopyranoside.


Biological Source It is obtained from the bark of poplar (*Populus*) and willow (*Salix*) and also found in the leaves and female flowers of the willow. It is specifically found in *two* species of *Salix*, namely: *Salix fragilis* and *Salix purpurea*, belonging to the family *Salicaceae*. It is also found in the root bark of *Viburnum prunifolium* L., family : *Caprifoliacea* and in *Spiraea ulmaria*, family: *Rosaceae*.

Geographical Source It grows in China, Europe and in India.


Preparation The powdered bark is macerated with hot water for several hours whereby the **glucoside (salicin)** and tannin are extracted collectively. The resulting liquid extract is filtered, concentrated under vacuum and treated with lead acetate to remove the tannins as a precipitate. It is subsequently treated with hydrogen sulphite to remove the excess of lead. The clear filtrate is neutralized with ammonia, allowed to concentrate, chilled to obtain the crystals of salicin. The crude **salicin** may be further purified by treating its solution with animal charcoal and concentrating followed by cooling.

Description It occurs as colourless crystals or prisms or scales. It has a very bitter taste. It is highly soluble in hot water and practically insoluble in ether. It is a levorotatory substance [$(\alpha)_D$: -63°].

Chemical Constituents It is hydrolysed in the presence of the enzyme **emulsin** by yielding one mole each of **saligenin** (aglycone) and **D-glucose** as stated below:



When hydrolysis is done in an acidic medium by boiling for a prolonged duration, two moles of **saligenin** combine together to provide **saliretin** (water insoluble) with the loss of a mole of water, which may be summarised as shown under:



Chemical Tests

1. It gives an instant bright red colour with concentrated sulphuric acid that fades out on the addition of water.
2. Its hydrolysed product saligenin gives a blue colour with ferric chloride.
3. On oxidation with potassium dichromate and sulphuric acid and heating yields salicylaldehyde having a characteristic odour.

4. It gives specific colours with the following reagents:

- Frachde's Reagent** : Violet colour
Mandelin's Reagent* : Purple red colour
Erdmann's Reagent** : Bright red colour

Uses


1. It is used as an analgesic
2. It has been employed as a bitter stomachic.
3. It is also used as an antirheumatic agent.
4. It is used as a standard substrate in evaluating enzymes preparations containing β -glycosidase.

4.2.2.4 Populin

Synonyms Populoside; Salicin benzoate.

Biological Source It occurs in the bark leaves of *Populus tremula* L., *P. nigra* L., *P. nigral.* L. var *italica* Duroi, *P. canadensis* Moench., *P. grandidentate* Michx., and *P. tremuloides* Miehx., belonging to the family *Salicaceae*. It is perhaps also found in *Salix helix*, *Salix purpureae* L. var *helix* (L.) Koch.

Preparation It may be prepared either from *salicin* by melting with benzoic anhydride, or from *salicin* and benzoyl chloride in the presence of KOH as shown below:



Description It occurs as white needles having a sweet taste like licorice. It is readily soluble in alcohol and hot water, but sparingly soluble in cold water and almost insoluble in solvent ether.

Chemical Constituents **Populin** on hydrolysis with alkalis [eg., $\text{Ba}(\text{OH})_2$] yields salicin and benzoic acid, whereas, its hydrolysis in an acidic medium gives benzoic acid, saligenin and glucose. However, its enzymatic hydrolysis with **taka-diastase** provides **salicin** and **monobenzoyl glucose**.

Chemical Tests It gives exactly identical reactions with conc. H_2SO_4 and Frachde's Reagent as those with salicin.


* *Mandelin's Reagent*: Dissolve 0.5 g of ammonium vanadate in 15 ml of water and dilute to 100 ml with sulphuric acid. Filter the solution through glass wool.

** *Erdmann's Reagent*: A 1% (w/v) aqueous solution of ammonium diamminetetraakis-(nitrito-N) cobaltate (1-).

4.2.2.5 Glucovanillin

Synonyms Vanilloside; Avenein; Vanillin -D-glucoside. It is obtained from the green fruit of vanilla.

Preparation It is prepared from **coniferin** by oxidation with chromium-6-oxide (CrO_3) as follows:



Description It has a needle-like appearance having a bitter taste. It is readily soluble in hot water and alcohol; but almost insoluble in ether.

Chemical Constituents **Glucovanillin** is the chemical constituent present in the green fruit of vanilla.

Uses

It is mostly used in pharmaceutical preparations as a flavoring agent.

4.2.3 Steroid Glycosides

Steroid glycosides are also referred to as '**Cardiac glycosides**' in many available books on phytochemistry. In fact, there exists enough evidence in literatures to reveal that a host of medicinal plants comprise of cardiac or cardiotoxic glycosides, collectively known as '**steroid glycosides**', and they have since been employed as *arrow poisons* or *cardiac drugs*. Interestingly, from a therapeutic perspective this particular group of compounds may be regarded as one of the most important of all naturally occurring plant products.

The **cardiac glycosides** are basically steroids with an inherent ability to afford a very specific and powerful action mainly on the cardiac muscle when administered through injection into man or animal. As a word of caution, a small amount would exhibit a much needed stimulation on a diseased heart, whereas an excessive dose may cause even death.

Generally, the **steroid glycosides** are invariably employed in the therapeutic domain primarily for *two* vital reasons, namely: (a) to enhance the tone, excitability and above all the contractibility of the cardiac muscle; and (b) to increase the diuretic action, due principally to the enhanced renal circulation (an inherent secondary action).

A few important plant products belonging to this category are discussed in the sections that follows, namely:

4.2.3.1 *Digitalis*

Synonyms Foxglove; Purple foxglove; Fairy gloves; Digifortis; Digitora; Pil-Digis; Neodigitalis.

Biological Source *Digitalis* comprises of the dried leaves of *Digitalis purpurea* L., belonging to the family *Scrophulariaceae*. [The word *purpurea* has been derived from the purple colour of flowers]. It is pertinent to mention here that the fresh leaves must be dried immediately in the dark at a temperature not exceeding 60°C so that the final dried leaves should not contain more than 5% of inherent moisture. This is, however, extremely important to retain the glycosides in a good undecomposed condition.

Geographical Source It is grown in Southern and Central Europe, England, Holland, Germany, United States and India. In India, it is grown in Kashmir and Nilgiri Hills.

Preparation Good quality **digitalis** is grown specifically from the seeds of selected strains that will invariably yield only leafy plants enriched with glycoside contents. Even the soil is usually sterilized by steam before commencement of sowing. Mostly it grows both appreciably and luxuriantly at an altitude ranging between 1600-300 meters preferably in a shady environment. In actual practice, the sowing of seeds is performed in autumn (October/November) , and the seedlings are virtually transplanted in the fields in the following springs (March/ April). The leaves are normally hand picked in the *afternoon* during August / September in the *first* and *second* year, when almost 2/3rd of the flowers have fully bloomed. The leaves collected in the first year are found to contain the highest percentage of glycosides. The basal leaves and the ones located at the top are collected at the end. The discoloured leaves are sorted out and rejected outright. The selected leaves are duly spread on perforated trays (usually a thin bed), the trays are stacked one above the other in a well-closed dark drying shed heated by a stream of hot air maintained strictly at a temperature not more than 60°C. The dried leaves having a moisture content not more than 5% are carefully packed in suitable air-tight containers, charged with appropriate dehydrating agents and shipped for export.

Note The therapeutic potency *vis-à-vis* the activity of the leaves is solely due to the glycoside content. Surely, the presence of moisture and certain enzymes, namely; *oxidase* and *digipurpuridase*, are chiefly responsible for the ultimate deterioration of the glycosides of the leaves. In case, the leaves are made to dry at a temperature beyond 60°C, it has been observed that there is a drastic loss in potency on account of *chemical degradation* (irreversible).

Description

- Colour** : Dark greyish green
- Odour** : Slight
- Taste** : Bitter
- Size** : Length: 10-14 cm; width: 4-15 cm
- Shape** : Orate-lanceolate to broadly ovate

Special Features The digitalis leaves are more or less pubescent venation together with pronounced and marked veinlets on the under surface. The leaves are invariably crumpled and broken.

Chemical Constituents Nativelle (1868), Kiliani (1891), and Stoll (1938) were the pioneers who contributed valuable informations with regard to the chemical constituents present in *digitalis* through their extensive and intensive studies.

It has been reported that **digitalis** essentially contains *three* important *primary glycosides* namely: *Purpurea glycoside A*, *Purpurea glycoside B*, and *Purpurea glycoside C*, which upon hydrolysis give rise to *digitoxin*, *gitoxin* and *gitalin* respectively. These **secondary glycosides** on further hydrolysis yields noncarbohydrate moieties (called *aglycones or genins*) *digitoxigenin*, *gitoxigenin* and *gitaligenin* or *gitaloxigenin* respectively. The series of all these hydrolysed products and their structures are summarised below.

Besides, the crude drug also contains a good number of other *glycosides* (e.g.; **digitalin**, **diginin**); *saponins* (e.g.; **digitonin**, **gitin** and **digitosaponin**); tannins, gallic, formic, acetic, succinic and benzoic acids; fatty acids and enzyme *digipuridase* solely responsible for hydrolysis of purpurea glycosides.

Chemical Tests A plethora of chemical colour reactions have been evolved to be used as the qualitative tests either for the various glycosides or their corresponding aglycones in the chemical laboratory. However, the exact positions of the respective **glycosides** or their **aglycones** may be detected either on the paper chromatograms or on the thin layer chromatographic plates by virtue of the production of specific colours or by exposing the chromatograms under UV light so that the components would be detected by their fluorescence. All these specific tests are summarised in Table 4.1.




Table 4.1 Colour Tests of Glycosides

S.No	Specific Tests	Experimental Procedures	Inferences
1.	Kellar-Kiliani Test	The glycoside is dissolved in glacial acetic acid containing a trace of FeCl ₃ , add the same amount of FeCl ₃ dissolved in conc. H ₂ SO ₄ along the side of the test tube to settle at the bottom (for α-deoxy sugars eg; digitoxose)	A reddish brown colour changing to bluish green colour appears at the junction of two reagents within 2-5 minutes spreading slowly into the acetic acid layer.
2.	Legal Test	A few mg of the glycoside (except scillaren) is dissolved in a few drops of pyridine. To this is added a drop of sodium nitroprusside soln. (2% w/v), add a drop of NaOH soln. (20% w/v)	Appearance of a pink or deep red colour indicates its presence.
3.	Baljet Test	The aglycone portion of the glycoside is mixed with Baljet reagent*	Appearance of an orange or red colour shows its presence.
4.	Raymond Test	Dissolve a small quantity of the glycoside in 1 ml of ethanol (50%) Add to it 0.1 ml of Raymonds Reagent** and 2/3 drops of NaOH solution (20% w/v) (for activated methylene group at C-21 in the lactone ring)	Appearance of a violet colour slowly changing to blue gives an affirmative test.
5.	Tollen's Test	Dissolve a small amount of the glycoside in a few drops of pyridine. To this is added a few ml of Tollen's Reagent*** & heated gently (if required).	Appearance of a silver mirror shows a positive test.
6.	Xanthydro Test	Add to the glycoside solution 0.5 ml of xanthydro solution**** (for 2-desoxy sugars only)	Development of red colour shows its presence.
7.	Antimony Trichloride Test	To a solution of glycoside add a solution of antimony trichloride and trichloroacetic acid, and then heat the mixture (for <i>Cardenolides</i> & <i>Bufadienolides</i>)	Appearance of a blue or violet colour shows its presence.
8.	Kedde Test	A solution of glycosides is treated with a small amount of Kedde's Reagent***** (for <i>Cardenolides</i> & <i>Cardinolide aglycones</i>)	Development of a blue or violet colour that fades out in 1 to 2 hours shows its presence.

* **Baljet Reagent:** Aqueous solution of picric acid (1% w/v) and NaOH soln. (10% w/v). Both solutions mixed immediately before use and filtered.

** **Raymond's Reagent:** A 1% (w/v) solution of m-dinitrobenzene in ethanol (or methanol)

*** **Tollen's Reagent:** To a 0.1 N soln. of AgNO₃ is added dilute NH₄OH till the white precipitate initially formed gets dissolved after further addition of NH₄OH.

**** **Xanthydro Reagent:** A solution of 0.125% (w/v) xanthydro in glacial acetic acid containing 1% HCl.

***** **Kedde's Reagent:** Mix equal volumes of a 2% (w/v) soln. of 3, 5 dinitrobenzoic acid in methanol and a 7.5% (w/v) aqueous soln. of KOH

Substituents/Adulterants The **digitalis** leaves have been frequently adulterated with the following varieties of leaves, namely:

1. *Great mullein leaves:* The leaves of *Verbascum thapsus* belonging to the family *Scrophularineae* are usually mixed with the genuine drug leaves which may be identified and distinguished microscopically by the abundant presence of huge woolly and branched candelabra trichomes.
2. *Primose leaves:* The leaves of *Primula vulgaris* belonging to the family *Primulaceae* are invariably mixed to **digitalis**, that may be identified microscopically as follows:

- (a) *P. vulgaris* has uniseriate covering trichomes, that are normally 8-9 celled long, and
- (b) *P. vulgaris* leaves have straight lateral veins.

3. *Comfrey leaves*: The leaves of *Symphytum officinale* belonging to the family *Boraginaceae* are mixed with **digitalis**, which may be distinguished by the presence of multicellular trichomes seen at the top in the shape of a hook.

Uses

1. **Digitalis** enhances the force of contraction of heart muscle which ultimately affords an increased cardiac output, decreased size of heart, decreased venous pressure and above all the decreased blood volume. Hence, digitalis together with its various marketed preparations are employed profusely as vital cardiotonics in the management and control of different kinds of congestive heart failure, atrial flutter, atrial fibrillation, supraventricular tachycardia and premature extra systoles.
2. **Digitalis** has a tendency to exert an overall cumulative effect in the body, and hence it gets eliminated rather gradually. Therefore, it is extremely important to monitor the dosage regimen by a physician whether he relies on branded products or natural drug preparations eg., Digitoxin injection, Lanoxin, Prepared digitalis and Digitalis tincture.

4.2.3.2 Allied Drugs of Digitalis

Three important allied drugs of **digitalis** which shall be discussed in this section are, namely:

- (a) *Digitalis lanata*,
- (b) *Digitalis lutea*, and
- (c) *Digitalis thapsi*.

4.2.3.2.1 Digitalis Lanata The drug is almost three times more potent in comparison to *Digitalis purpurea* discussed earlier.

Synonyms Woolly Foxglove leaf, Grecian Foxglove; Austrian Digitalis.

Biological Source The dried leaves of *Digitalis lanata* Ehrhart, belonging to family *Scrophulariaceae*.

Geographical Source It is a biennial or perennial herb which being indigenous to Southern and Central Europe. It is also cultivated abundantly in India, Holland, United States and Ecuador.


Preparation The preparation of *D. lanata* leaves is quite similar to the one described under section 4.2.3.1 earlier.

Description The leaves are usually 5-15 cm in length, sessile, linear lanceolate to oblong lanceolate; margin- entire apex-acute; veins leave the mid-rib at an acute angle.

Chemical Constituents Stoll and Jucker* (1955) first isolated from its leaves *three* chemically pure *primary glycosides* usually termed as *Lanatoside A*, *Lanatoside B*, and *Lanatoside C*. These primary glycosides are also known as **Digilanid A**, **Digilanid B**, and **Digilanid C** respectively.

However, the inter-relationship of digitalis glycosides found in *Digitalis lanata* may be represented in the following flowchart.

* Stoll, A.; and E. Jucker, 'Modern Methods in Plant Analysis', Springer Verlag, Berlin, 1955



4.2.3.2.2 Digitalis lutea

Synonym Straw Foxglove.

Biological Source The dried leaves of *Digitalis lutea* Linn. belonging to family *Scrophulariaceae*.

Geographical Source It is found in Europe and USA.

Description The leaves have a length of 28 cm and width of 6 cm, but they usually attain only half of their size.

Chemical Constituents The chemical constituents of *D. lutea* have not been fully reported, but it does not contain calcium oxalate.

Uses

1. It is used as a common substitute for the *official drug*.
2. It is potent as *D. purpurea*.
3. It is mostly used for the same purpose as that of *D. purpurea*, but it is found to exert much lesser irritation.

4.2.3.2.3 Digitalis Thapsi

Synonyms Spanish Foxglove; Spanish digitalis.

Biological Source The dried leaves of *Digitalis thapsi* Linn. belonging to the family *Scrophulariaceae*.

Geographical Source It is grown in Italy and Spain.

Description The leaves are mostly yellowish green in colour; and have lanceolate with decurrent lamina and crenate margin. It also contains calcium oxalate crystals.

Uses Its therapeutic efficacy is almost 1.25 to 3 times more potent than *D. purpurea* and its actions are similar to those of the latter.

4.2.3.3 Squill

A survey of literature reveals that the *Squill bulbs* was thoroughly and repeatedly investigated since 1879. However, Stoll in 1933 was first able to separate and isolate two glycosides in their purest form, namely: **Scillaren A** and **Scillaren B**. These *two* naturally occurring glycosides are usually present in the crude drug in the ratio 2:1 (*i.e.*, 2 parts of Scillaren A and 1 part of Scillaren B).

Generally, the squill is available in *three* varieties, namely:

- (a) European Squill
- (b) Indian Squill, and
- (c) Red Squill.

These *three* varieties would be described in the sections that follows:

4.2.3.3.1 European Squill

Synonyms Sea, onion, Bulbus Scillae; Meerzweibel, White Squill, Squill bulb; Scila.

Biological Source **European squill** is the fleshy inner bulb scales of the white variety of *Urginea maritima* (L.) Baker (*Scilla maritima* L.) belonging to family *Liliaceae*.

Geographical Source It is found to be indigenous to those countries located near the Mediterranean region, such as: France, Malta, Italy, Greece, Spain, Algeria and Morocco.

Preparation Normally the white squill yields fully grown and healthy bulbs that have a height ranging between 18-20 cm and a diameter varying between 12-15 cm. These bulbs are grown in partially submerged condition in sandy soil in the mediterranean coastal region. The bulbs are usually collected in late August soon after the flowering season. The roots and the thin external scaly layers are removed and discarded. While the central fleshy bulbs are collected separately. These bulbs are then cut into transverse slices and subsequently dried either in sun rays or by artificial heating devices.

Description

Colour : White; Whitish yellow;

Taste : Bitter and gummy;

Size : Length = 3.5-5 cm; Width = 5-8 mm; Thickness = 2-5 mm;

Shape : Available as strips with tapering both ends.

Chemical Constituents Squill has the following glycosides, namely:


Glucoscillaren A = Scillarenin + Rhamnose + Glucose + Glucose;

Scillaren A = Scillarenin + Rhamnose + Glucose;

Proscillaridin A = Scillarenin + Rhamnose.

Scillaridin A; Scilliglaucoside; Scillipheoside; Glucoscillipheoside; Scillicyanoside.

The structures of these glycosides are as shown below:




In addition to the above cardiac glycosides, the drug also comprises of **flavonoids, calcium oxalate, xanthoscillide, sinistrin** (an *inulin* like carbohydrate) and an irritation causing volatile component.

The following flowchart evidently illustrates the various steps involved in the *acidic* and *enzymatic hydrolysis* of *Scillaren A* as shown below.

Stoll was first to separate the *two* glycosides from **squill bulbs** and named them as **Scillaren A** and **Scillaren B**. They have the following characteristic features.

Chemical Test **Scillaren A** on interaction with acetic anhydride and H₂SO₄, it gives rise to a red colour which changes gradually first to blue and finally to bluish green colour.



Characteristics of Scillaren A and Scillaren B

S.No.	Properties	Scillaren A	Scillaren B
1.	Nature	Crystalline	Amorphous
2.	Colour	Colourless	White/ Yellowish white
3.	Taste	Bitter	Very bitter
4.	Odour	Odourless	Odourless
5.	Solubility	Comparatively less water soluble	More soluble in water ethanol and methanol
6.	Activity (in Frog assay)	Less active (3 times to reference drug)	More active (5 times to reference drug)
7.	Stability	Less stable	More stable

Uses

1. It is a potent cardiotoxic without having any cumulative effect (unlike *Digitalis*).
2. It is mostly employed in small doses as an effective expectorant specially in chronic bronchitis.
3. It causes emesis in relatively higher doses.
4. The **squill glycosides** usually produce copious diuresis.
5. By virtue of the fact that the **squill glycosides** possess high therapeutic index and rapid elimination they invariably maintain compensation in such patients where a prolonged treatment is required.

4.2.3.3.2 Indian Squill

Synonyms Scilla; Sea onion; Jangli Pyaj; Urginae.

Biological Source **Indian squill** comprises of the dried slices of the bulbs of *Urginbea indica* Kunth; belonging to the family *Liliaceae*.

Geographical Source It is grown in India along the sea coasts of Konkan and Saurashtra; and also on the dry hills of the lower Himalayan range located at an altitude of 1500 meters.

Preparation The method of preparation of dried slices of **Indian squill** is very much alike the **European squill**. It loses approximately 80% of its weight after sun drying.

Description

Colour : Yellowish to White

Odour : Slight and characteristic

Taste : Acrid, bitter and mucilaginous

Size : Length = 30-60 cm; Breadth = 3-8 mm

Shape : Usually 4 to 8 slices are placed one on the top of other and gives it a curved shape.

Characteristic Features

1. It has an overall diameter of 15 cm
2. The dried slices are translucent in appearance which become flexible and rather tough soon after gaining moisture.

Chemical Constituents **Indian Squill** essentially comprises of cardiac glycosides (0.3%), alcohol soluble extractives (20-40%), mucilages (40%) and calcium oxalate.

The two major **cardiac glycosides** present in the drug are **Scillaren A** and **Scillaren B** (see Section 4.2.3.3.1).

Substituents/Adulterants The bulbs of different species of *Ledebouria* (*Scilla*, Linn) are sold in the Indian bazars, under vernacular names which are equivalent to 'small squill'.


Ledebouria hyacinthoides, is used as a substitute for squill. It has a scaly bulb, about the size and shape of a small pear, composed of very smooth and fleshy scales, which are so *imbricated* that they might be mistaken for entire coats if not carefully examined.

Uses

1. It is largely employed as a cardiotoxic, stimulant and also an expectorant.
2. It is used as a very effective expectorant both in asthma and chronic bronchitis.
3. It possesses anticancer activity against human epidermoid carcinoma of the nasopharynx in tissue culture.
4. It is in no way a perfect replacement for *Digitalis* since it possesses not only irritant effect but also is very poorly absorbed systemically.

4.2.3.3 Red Squill It is the red variety of **European squill** *Urginea maritima* (L.) Baker belonging to family *Liliaceae*. In fact, the red colour is attributed due to the presence of anthocyanin pigments present in the mesophyll cells of scales.

The glycoside present in the **red squill** is known as **scilliroside** having the following structure.



It is slightly soluble in water, but is soluble freely in lower alcohol, ethylene glycol, dioxane and glacial acetic acid. It is mostly used as a rat poison.

4.2.3.4 *Strophanthus*

Synonyms Semino stropanthi.

Biological Source These are the dried and ripe seeds of *Strophanthus hispidus* De*, or of *Strophanthus kombe* Oliver, belonging to the family *Apocynaceae*, deprived of the awns.

* De = De Candolle

Geographical Source *Strophanthus* plants are elimbers, which being perennial large & woody, found to be indigenous in the vicinity of Shire river, Nyanza and Tanganyika lakes of Eastern tropical Africa.


Preparation The ripe *strophanthus* fruit comprises of two fully developed follicles each about 30cm broad with tapering at both ends and consisting of a number of seeds. The ripe fruits are collected from the wild plants, the seeds are subsequently separated and freed from their awns.

Description

- Colour** : Greyish green to light yellowish brown
Odour : Slight unpleasant
Taste : Bitter
Size : Length 1- 2 cm; Breadth = 3-5 mm; Thickness = 2 mm
Shape : Lanceolate to linear-lanceolate, acute at the apex, rounded or blunt at the base
Weight : For 100 seeds 3-4 g
Specific feature : On treating with 80% H₂SO₄ the endosperm exhibits a deep Emerald green colour.

Chemical Constituents The seeds of *strophanthus* usually contain *three* vital glycosides, namely: **K-strophanthoside**, **K-strophanthride β** and **cymarín**. Interestingly, all these glycosides undergo hydrolysis to yield **strophanthidin**.

The structure of **strophanthidin** and its allied glycosides are given below:




K-Strophanthoside It is the main constituent of *S.kombe*, the aglycone is known as **strophanthidin** that has the following characteristic features, namely:

- (a) Three—OH moieties at positions C-3, C-5 and C-14.
- (b) An aldehydic (—CHO) function is present at C-10 which being an essential requirement.
- (c) At C-17 an unsaturated 5-membered lactone ring, and
- (d) At C-3 an 'O' atom forms a bridge to the sugar component(s) essentially comprising of cymarose, β -D-glucose and α -D-glucose.

Acidic hydrolysis of K-strophanthoside gives rise to the aglycone strophanthidin along with a triose sugar known as **strophanthotriose** that comprise of one mole of **cymarose** and two moles of **glucose**.

Enzymatic hydrolysis of **K-strophanthoside** using the enzymes-*glucosidase*, usually present in yeast, helps in cleaving off the terminal α -D glucose thereby yielding the secondary glycoside known as **K-strophanthidin β** . Further hydrolysis of the resulting product with *strophanthobiase*, the former yields the glycosides **cymarin** which comprises of the aglycone **strophanthidin** along with one mole each of **cymarin** and **β -D-glucose**.



Strophanthobiase

However, it is worth noting that the acidic hydrolysis of **K-strophanthoside** gives rise to the aglycone **strophanthidin** and **strophanthobiase** which being a disaccharide (or **biose**) It may be observed that the terminal *glucose* possesses an *alpha linkage*, while the one attached to *cymarose* bears a *beta linkage*.

Chemical Tests

1. Generally, the **strophanthus glycosides** exhibit an emerald green colouration on the addition of sulphuric acid.
2. Dissolve about 0.1 g of **strophanthin** in 5 ml of water and add to it a few drops of ferric chloride solution followed by a 1-2 ml of concentrated sulphuric acid; the appearance of an initial red precipitate that finally turns green within a period of 1-2 hours.
3. To 50 mg of **strophanthin** add 5 ml of water, shake and add 2 ml of 2% tannic acid solution, the appearance of a distinct precipitate affirms its presence.
4. It shows positive **Baljet Test**, **Legal Test** and **Keller Killiani Test** (see Section 4.2.3.1).

Uses

1. It is used intravenously for treating emergency cardiac conditions. However, orally strophanthin is not so active.

2. These glycosides have been found to exert less cumulative effect unlike the digitalis glycosides.
3. Overall their therapeutic actions are very much similar to those of *Digitalis*.

4.2.3.4.1 Allied Drugs


G-Strophanthin

Synonyms Ouabain; Gratus strophanthin; Acocantherin; Gratibain; Astrobain; Purostrophan; Stropherim; Strodiral.

Biological Source **G-Strophanthin** is obtained from the seeds of *Strophanthus gratus* (Wall & Hock)Baill. It also occurs in the wood of *Acokanthera ouabio* Cathel and *A. schumperi* belonging to the family *Apocynaceae*.

Geographical Source The wood is grown in Ethiopia and Somaliland.

Chemical Constitutents The seeds contain the glycoside **G-strophanthin** (or **Ouabain**) as given below:



Ouabain on hydrolysis gives rise to an aglycone termed as G-strophanthidin (or Ouabagenin) and L-rhamnose as the residual sugar moiety.

Description **G-Strophanthin** mostly occurs as colourless small shining crystals, which are odourless and have an extremely bitter taste. It is readily affected by light, but is quite stable in air. It is practically insoluble in ether, chloroform and ethyl acetate, whereas it is sparingly soluble in cold water (1:75); freely soluble in hot water and alcohol (1:100).

Chemical Tests

1. Mix a few crystals of ouabain with a mixture of conc. H_2SO_4 and water (4:1), the appearance of a brownish red colour, which deepens slowly and ultimately shows a green fluorescence.
2. **Froehde's Test:** Mix a few crystals with a drop of Froehde's reagent, evaporate to dryness, cool and add a drop of H_2SO_4 —the development of a blue colour takes place.


3. **Mandalin's Test:** Moisten a few crystals with Mandalin's reagent, evaporate to dryness, cool and then add one drop of conc. H_2SO_4 —the appearance of a green colour occurs.

Uses

1. It is an important cardiotonic, which is usually administered intravenously in acute cardiac failure, due to its inherent rapid onset of action.
2. It is invariably employed as a '*reference standard*' for comparison of **cardiac glycosides**.
3. It also exerts diuretic action.

4.2.4 Flavonoid Glycosides


Flavonoid constitute one of the largest class of naturally occurring plant products mostly phenols either in the free state or as their respective glycosides. As the very name suggests they are usually yellow-coloured compounds (*flavous* is a *latin* word yellow colour). Interestingly, more than 2000 different chemical compounds have been isolated, identified and reported from plant sources. In fact, their chemical structures are solely based upon a C_6 — C_3 — C_6 carbon skeleton having a **pyran** or **chroman** ring bearing a second benzene (aromatic) ring strategically positioned at C—2, C—3 or C—4 as shown below:




In nature they are invariably available as: **flavones, flavanones, flavonols, isoflavones,** and **anthocyanidins***. In certain specific instances either the 6-membered heterocyclic ring (**pyrones**) is replaced by a 5 membered heterocyclic ring (**aurones**) or exists in an open-chain isomeric form (**chalcones**). Besides, the normally existing glycosylated derivatives found in nature, other types of derivatives, such as methylated, acetylated, prenylated, or sulphated ones also exist. Nevertheless, it has been established that a large variety of **flavonoids** exert a wide range of activities in nature, namely: antimicrobial agents, signaling molecules, or stress metabolites.

The structures of a few typical **flavonoids** are represented here as follows:


* **Anthocyanidins** are the colored aglycones found as a large number of pigments from flowerd and fruits (**Gr. Antho** flower + **Gr. Kyanos**, blue). Investigations of these pigments were initiated by Willstatter in 1914 and later on extended by Karner R Robinson, GM Robinson and others.




Flavones







Flavanones



Flavonols






Isoflavones

Anthocyanidins
[Oxonium Form]Anthocyanidins
[Oxonium Form]Anthocyanidins
[Carbonium Form]Anthocyanidins
[Carbonium Form]

The structure (a) is more stable, because it has a **naphthalenoid system** of linkages, whereas (b) contains a **quinonoid system**.

The structures of only the key portions of **pyrones**, **aurones** and **chalcones** are as follows:

Pyrones
(6-membered)Aurones
(5-membered)Chalcones
(Open-Chain isomeric form)

The **flavonoid glycosides** mostly occur as **O-glycosides** or **C-glycosides** contained in the cell sap of relatively younger tissues of higher plants belonging to several families, such as: *Compositae*, *Leguminosae*, *Polygonaceae*, *Rutaceae* and *Umbelliferae*. It has been observed that a host of natural plant products containing **flavonoid glycosides** exert a variety of therapeutic effects, namely: antiasthmatic, antispasmodic, diuretic, fungicidal and oestrogenic activities.

A few typical **flavonoid glycosides** shall be discussed in the sections that follows, namely:

- (a) Flavone Glycosides,
- (b) Flavonol Glycosides,
- (c) Flavanone Glycosides,
- (d) Chalcone Glycosides,
- (e) Isoflavonoid Glycosides, and
- (f) Anthocyanidin Glycosides.

4.2.4.1 Flavone Glycosides

The two important members of this class of glycosides are **Apiin** and **Diosmin** which are described here under:

4.2.4.1.1 Apiin


Synonyms Apioside, Apigenin 7-apiosyglucoside.

Biological Source **Apiin** usually occurs in the seeds and leaves of *Petroselinum sativum* Hoffm, and *Apium petroselinum* Linn., known as **parsley**, and also in *Apium graveolens* L., and *Anthemis nobilis* L, called as **celery**, family *Umbelliferae*. It has also been found to be present in the ray florets of *Marticaria chamomilla* Linn., and some other ray florets belonging to the family *Compositae*.


Geographical Source The fruits are grown in Persia, Greece, Northwest Himalaya and Europe.

Description It is a very small fruit, almost globular in shape. The taste is at first like anise, but afterwards bitter. The colour is like anise, but generally faint. It occurs as colourless needles.


Chemical Constituents The structure of **apiin** is given below:



Apiin is neither hydrolysed by the enzyme emulsin, nor by yeast or yeast extract. However, it undergoes hydrolysis in an acidic medium to yield *glucose*, *apiose* (pentose sugar) and *apigenin* (aglycone) which is 4, 5, 7-trihydroxyflavone.



Apiin when treated with dilute sulphuric acid (0.5 to 1%) undergoes cleavage only at *apiose moiety* thereby giving rise to **7-glucoapigenin** as shown below:



Chemical Tests

1. It gives a yellow precipitate with basic lead acetate.
2. It produces a reddish brown colour with FeCl_3 .
3. It gives an intense yellow colour with NH_4OH solution
4. It yields a pale yellow colour with NaOH solution.


4.2.4.1.2 Diosmin

Synonym Barosmin.

Biological Source It occurs in the dried *Buchu* leaves *i.e.*; various species of *Barosma crenulata*, *Barosma serratifolia* and *Barosma betulina* belonging to family *Rutaceae*. It has also been isolated from *Serophularia nodosa*, *Hyssopus officinalis*, *Mentha crispera*, *Mentha pulegium* and from species of *Conium*, *Dahlia* etc.

Description It occurs as pale yellow needles.

Chemical Structure The chemical structure of **diosmin** (5, 7, 3'-trihydroxy-4'-methoxyflavone-7-rhamnoglucoside) is given below:



Diosmin upon hydrolysis yields rhamnose, glucose and the aglycone diosmetin.

Chemical Test A solution of **diosmin** in concentrated sulphuric acid exhibits a slight fluorescence.

4.2.4.2 Flavonol Glycosides

The two well known glycosides belonging to this class are namely: **Rutin** and **Quercetin**, whereas the less important ones are—**galangin**, **gossypin**, **hibiscitrin**, **kaempferin** and **avecularin**.

4.2.4.2.1 Rutin

Synonyms Melin; Phytomelin; Eldrin; Ilixanthin; Sophorin; Globularicitrin; Paliuroside; Osyritrin; Osyritin; Myrticolorin; Violaquercitrin; Birutan; Rutabion; Rutozyd; Tanrutin.

Biological Source **Rutin** is found in many plants, especially the **buckwheat plant** (*Fagopyrum esculentum* Moench; family: *Polygonaceae*); in **forsythia** [*Forsythia suspensa* (Thunb.) Vahl ver. *Fortunei* (Lindl.) Rehd., family: *Oleaceae*]; in **hydrangea** (*Hydrangea paniculata* Sieb., family: *Saxifragaceae*); in **pansies** (*Viola* sp. *Violaceae*); from leaves of *Eucalyptus macroryncha* F.v. Muell., family: *Myrtaceae*); in *Fagopyrum tartaricum* Gaertn: family: *Polygonaceae*); in *Ruta graveolens* L., (family: *Rutaceae*); in buds of *Sophora japonica* L., (family: *Leguminosae*); in fresh leaves of **tobacco plants**, *Nicotiana tabacum* L., (family: *Solanaceae*); in **cotton seed** *Gossypium hirsutum*; (family: *Malvaceae*); in *Viola tricolor*, (family: *Violaceae*).


Geographical Source The various plants like eucalyptus, tobacco, and cotton grow abundantly in tropical countries like India, Africa, Ceylon and United States, Australia and China.

Preparation Based on experimental evidences it has been observed that the glycoside content in plants declines sharply as they mature; however, the highest yields are usually obtained from the **buckwheat leaves and flowers** as soon as the plants attain the blossom-stage.

The dried and ground plant material (say 150 g) is extracted with two successive quantities (2 × 200 ml) of ethanol (80% v/v). The resulting filtered hydro-alcoholic extract is carefully evaporated under vacuum in a rotary evaporator till it reaches upto 50-60 ml. The content of the flask is mixed with an equal volume of ether and the ethereal layer is separated. The aqueous layer is once again extracted with the same volume of ether and the ethereal layer separated. Both the ethereal layers are discarded and the aqueous layer is evaporated under reduced pressure to 10 ml. Keep the concentrated residual liquid in a refrigerator (0-5°C) overnight when a solid crystalline substance appears. Separate it from the mother liquor. The crude rutin, thus obtained may be further purified by *Column Chromatography*, using magnesium silicate as an adsorbent and ethanol as an eluant.

Description It has a pale yellow crystalline needle like appearance. It is practically insoluble in water, ether, petroleum ether and chloroform. It is fairly soluble in ethanol and acetone.

Chemical Structure The structure of **rutin** (or 5, 7, 3', 4' tetrahydroxy flavonol -3-rhamnoglucoside) is given below:



Rutin on refluxing with dilute mineral acid (200 ml of 0.1N H_2SO_4 + 1 g rutin) for 90 minutes gives rise to the aglycone known as **quercetin** plus the corresponding sugars.

Chemical Tests

1. It gives a distinct yellow precipitate with basic lead acetate.
2. It yields a greenish brown colour with ferric chloride.
3. It produces a silver mirror with ammonical silver nitrate solution (Tollen's Reagent)

Uses

1. **Rutin** is used to decrease the capillary fragility (*i.e.*, to enhance the tensile strength of capillary walls), reduce capillary permeability by tissue injury, and minimise the destruction of epinephrine in body tissues.
2. It has been mostly used in certain disease condition to reduce capillary bleeding promptly.
3. It is found to be useful in the treatment of retinal hemorrhages.

4.2.4.2.2 Quercetin


Synonyms Quercitroside; Quercimelin; Quercetin – 3-L-rhamnoside; Thujin; Quercitin; Quercetin.

Biological Sources Quercetin occurs in the bark of *Quercus tinctoria* and some other species of *Quercus*. It is also obtained from *Alsculus hippocastarum* L., **horse chest nut**, belonging to family *Hippocastanaceae*. Besides, it is also found in *Thuja occidentalis* L., *Morus alba* L., *Humulus lupulus* L., *Fraxinus excelsior* L., *Vitis vinifera* and a host of other plants.

Geographical Source *A. hippocastanum* is found in India and the African continent.

Description The crystals are yellow in colour when obtained from ethanol or methanol. It is practically insoluble in cold water and ether. *Flavine yellow shade* obtained from the **quercitron bark** by extraction under high pressure steam which is used exclusively in dyeing fabrics.

Chemical Structure The structure of **quercetin** is as shown below:



Quercitin on hydrolysis in an acidic medium gives rise to **rhamnose** and **quercetin** (*i.e.*, 5, 7, 3', 4'-tetrahydroxy flavonol).

It has been observed that the fully methylated *quercetin* upon hydrolysis yields 5, 7, 3', 4'-tetramethyl flavonol which amply suggests that the residue is duly attached at C-3 of the aglycone moiety and also **quercitin** is nothing but quercetin-3-rhamnoside.

Chemical Tests

1. It exhibits a brown fluorescence under the UV-light.
2. It gives a distinct yellow precipitate initially with a solution of basic lead acetate, but it gets dissolved on further addition of the reagent in excess.
3. It reduces Tollen's reagent to give a silver mirror.
4. It gives a negative test with Fehling's solution *i.e.*, not yielding brick red precipitate.

Uses It has been used as textile dye.

4.2.4.3 Flavanone Glycosides

Flavanone glycosides are most abundantly distributed amongst the citrus fruits. **Hesperidin** is the glycoside most commonly found in this particular class. However, the comparatively less important glycosides belonging to this category are, namely: **naringin**, **citronin** and **liquiritin**. **Hesperidin** shall be discussed in details here.

4.2.4.3.1 Hesperidin

Synonyms Cirantin; Hesperitin 7-rhamnoglucoside; Hesperetin-7-rutinoside.

Biological Source **Hesperidin** is the most predominant flavonoid in lemons and sweet orange *Citrus sinensis* (Linn.) Osbeck. It is also found in the rind or peel or unripe, green citrus fruits, for instance: **Bitter Orange** (*Citrus aurantium* Linn.); Lemons (*Citrus limon* Linn); **Citron** (*Citrus medica* Linn.).


Geographical Source Citrus fruits are abundantly grown in the tropical African countries *e.g.*; Togo, Nigeria, Ghana, besides the Mediterranean regions.

Preparation The glucoside may be isolated by adopting the laid down detailed procedures as in Section 4.2.4.2.1. It is also present in the dried orange peel upto 8% and it occurs in the highest concentration in the white portion of the peel usually termed as *albedo*.

Description **Hesperidin** is a colourless needle like crystals. It seems to be closely related to Vitamin P (*Citrin*). It is readily soluble in hot water, sparingly soluble in alcohol and cold water, and practically insoluble in ether, benzene and chloroform.

Chemical Structure The Structure of **hesperidin** is given as under:

The hesperidin chalone, comprising of an embeded phloroglucinol ring is promptly converted to flavones (*e.g.*, **hesperidin**) in an acidic medium, when heated or allowing it to stand in the dry state for a long duration. However, **methylation of the hesperidin chalone** helps in the ultimate methylation of one of the phenolic moieties present in the phloroglucinol portion of the chalone not only stabilizes the corresponding **methyl chalone** but also prevents closure of the ring to produce the **flavones** [see structure on next page].



Uses

1. It is normally used in conjunction with ascorbic acid to minimise capillary fragility.
2. It is also indicated in the prevention and management of capillary fragility or permeability in hypertension, cardiovascular and cerebrovascular disease, and also in habitual and threatened abortion.

4.2.4.4 Chalcone Glycosides


Chalcones are characterised usually by the presence of the following essential structure:



Evidently, the two aromatic rings (A and B) are linked by the *three carbon -aliphatic chain* which does not actively take part in the formation of a *hetero-ring* as is normally found in other types of flavonoid compounds. Likewise, the **dihydrochalcones** shall have the following general structure:





Also, the **chalone** yields a flavanone on treating with 10% H_2SO_4 . These *two* reactions may be represented as follows:



Interestingly, the **chalcones** and **flavanones** are closely related to each other and they are also fully *interconvertible*.

Examples:



A few typical examples are described below:

4.2.4.4.1 Carthamin


Synonyms Safflor carmine; Safflor red; Carthamic acid.

Biological Source **Carthamin** is the coloring principle obtained from *Carthamus tinctorius* L., belonging to family *Compositae* (Safflower).


Geographical Source It is grown throughout a large portion of India. It is one of the most ancient crops cultivated in Egypt as a dye-yielding herb. It is also grown in Russia, Mexico, United States, Ethiopia and Australia.

Description It is a dark red granular powder having a green luster. It is slightly soluble in water and practically insoluble in ether. It is found to be soluble in alcohol and in dilute alkali carbonates.

Chemical Structure The structure of **carthamin** is as given below:



Carthamin on being treated with dilute hydrochloric acid gets converted to its isomeric yellow compound known as **isocarthamin**.






Uses

1. It is used as a dye.

4.2.4.5 Isoflavonoid Glycosides


Unlike most other flavonoids, the isoflavonoid glycosides possess a rather limited taxonomic distribution, chiefly confined to the family *Leguminosae*. On the contrary, their chemical structures display an exceptionally broad spectrum of modifications. Wong (1975)* carried out an extensive review of the chemistry as well as taxonomic distribution of isoflavonoids. A few major classes of isoflavonoids along with their enormous prevailing structural variations are exemplified in Table 4.1.

Table 4.1 Major Classes of Isoflavonoids


S.No	Name	Biological Source	Chemical Structure	Uses
1	Tephrosin (Toxicarol; Hydroxydequelin)	From leaves of <i>Tephrosia vogelli</i> Hook.f., (<i>Leguminosae</i>) In derris root, cube root;		Toxic to fish, insects and crustaceae. Not toxic to human
2.	Sophoricoside (Genistein-4'- glucoside)	From the green pods of <i>Sophora</i> <i>japonica</i> L., (<i>Leguminosae</i>)		Not reported
3.	Prunetrin (Glucoside prunetin)	From the bark of <i>Prunus avium</i> L., var Bigarreau Napoleon (<i>Rodaceae</i>)		Not reported

However, there exists a close relationship with the structure of **isoflavones** to the skeleton of the **rotenoids** (or **rotenone**), whereas both of them may be regarded as being derived from *3-phenyl chroman* as shown below:


* Wong, E 'The Flavonoids' Chapman & Hall, London, p. 743-800, 1975.



Isoflavone
[Genistein]



Rotenoid
[Rotenone]



3—Phenyl Chroman

4.2.4.6 Anthocyanidin Glycosides

In addition to chlorophyll, **anthocyanins** represent the most important class of natural plant pigments visible to the naked eye. **Anthrocyanidine** (aglycones) are structurally closely related to each other. It has been established that there exist *six* prominent **anthrocyanidines** whose chemical structures are entirely based on the structure of **pelargonidin**. These six aglycones differ from **pelargonidin** by having one or two additional hydroxyl or methoxyl moieties strategically positioned at C-3' and C-5' in the latter, as shown in Table 4.2.




Table 4.2 Six Major Anthocyanidins and Biological Source


S.No.	Anthocyanidins	R	R'	Biological Source(s)
1.	Pelargonidin	H	H	Flowers of <i>Pelargonium graveolens</i> L'Herit., and <i>P. roseum</i> (R. Br.in) Ait. Family: <i>Geraniaceae</i> .
2.	Cyanidin	OH	H	Flowers of <i>Althaea roaea</i> cav., Family: <i>Malvaceae</i> .
3.	Peonidin	H ₃ CO	H	Tubers from <i>Peony officinalis</i> Linn., Family: <i>Ranunculaceae</i> .
4.	Delphinidin	OH	OH	From the whortleberry <i>Vaccinium myrtillus</i> Linn., family: <i>Ericaceae</i> .
5.	Petunidin	OH	OCH ₃	Flowers of <i>Petunia hybrida</i> , family: <i>Solanaceae</i> .
6.	Malvidin	H ₃ CO	OCH ₃	Flowers of <i>Malva sylvestris</i> Linn., family: <i>Malvaceae</i> .

It is, however, pertinent to mention here that the sugar components are invariably attached to C-3' position, and more rarely to C-5' position. It is worth mentioning here that in the flavone glycosides the attachment is normally at C-7 position. These sugar moieties may be:

- Monosaccharides** : e.g.: arabinose, galactose, glucose and rhamnose
Disaccharides : e.g.: rhamnoglucosides (present in *Antirrhinum* sp.)
Trisaccharides : e.g.: 5-glucosides-3-rutinoside (present in some *Solanaceae* species such as: *Atropa* and *Solanum*.)

Leucoanthocyanidins (or Proanthocyanidins) **Procyanthocyanidins** represent a group of highly water soluble naturally occurring plant pigments that are closely related to anthocyanidins. However, they may be easily differentiated from other flavonoids by virtue of the fact that the **procyanthocyanidins** are easily converted into their corresponding anthocyanidins on being boiled either with alcoholic or aqueous hydrochloric acid.

In general, they are considered equivalent to *flavan 3, 4-diols* chemically, which may be present either in their monomeric/ polymeric forms or as their corresponding derivatives. Examples: **(-)-Melacacidin** obtained from *Acacia melanoxylan* and other related species.



(-) – Melacacidin (2,3 – *cis* & 3, 4 – *cis*)

The **isomelacacidin** has the 2,3-*cis* and 3,4 –*trans*. configuration

4.2.5 Coumarin and Furanocoumarin Glycosides

Generally, **couramin** and its derivatives **furanocoumarin** are found to be present in a plethora of naturally occurring plants. Nevertheless, the coumarin is present either in the free state or its corresponding glycosides form in nature, but it has been observed that the former being most common.

The *two* different types of glycosides of this group shall be discussed separately under the following heads:

4.2.5.1 Coumarin Glycosides

These are reported to be present in about 150 different species spreading over to nearly 30 different families, of which a few important ones are, namely: *Caprifoliaceae*, *Leguminosae*, *Oleaceae*, *Rubiaceae*, *Solanaceae*, and *Umbelliferae*.

This basic nucleus of coumarin is considered to be derived from *o*-hydroxy cinnamic acid (or *o*-coumarin acid) by its dehydration to yield the fused lactone ring as shown below:





It has been observed that invariably most naturally occurring **coumarins** essentially bear an oxygen atom either as hydroxyl (OH) or alkoxy (—OCH₃ or —OC₂H₅) at C-7 position.

A few important naturally occurring **coumarin glycosides** along with their respective biological sources have been summarised in Table 4.3.

2.5.1.1 Coumarin **Coumarin** is abundantly found in a variety of natural products which are used profusely as a flavouring agent in pharmaceutical preparations.

Synonyms Tonka bean camphor; Coumarin; Coumarimic anhydride.

Table 4.3 Coumarin Glycosides and Biological Sources

S.No	Name	Biological Source	Chemical Structure	Uses
1.	Acsulin	Horse chest nut tree; Fruit and bark of <i>Aesculus hippocastanum</i> Linn., family <i>Hippocastanaceae</i>		Fruit and bark in diarrhoea
2.	Cichorin	Flowers of the chicory plant: <i>Cichorium intybus</i> Linn., family: <i>Compositae</i>		Used as tonic, febrifuge, and in diarrhoea
3.	Daphnin	Bark of various species of <i>Daphne</i> eg: <i>Daphne mezereum</i> family: <i>Thymelaceae</i>		As a febrifuge
4.	Fraxin	From bark of the common European ash: <i>Fraxinus excelsior</i> Linn., family: <i>Oleaceae</i> .		Bark used as a bitter astringent, tonic and febrifuge.

Biological Source It is found in the *tonka seed*, also known as **tonquin beans** or **tonco seed** *i.e.*, *Dipteryx odorata* Wild, and *Dipteryx oppositifolia* L., belonging to family *Leguminosae*. It is also obtained from *Woodruff* (*Asperula species*) and in **sweet clover** *i.e.*; *Melilotus alba* Dess., family: *Leguminosae*.

Geographical Source These plants are found in Europe, India and in the African continent.

Description


Colour : Colourless

Odour : Pleasant and fragrant odour resembling that of vanilla beans.

Taste : Burning Taste

Shape : Orthorhombic, rectangular plates.

Chemical Structures **Coumarin** has the following structure:



Coumarin

Uses It is mostly employed as a pharmaceutical aid.

4.2.5.2 Furanocoumarin Glycosides

In general, the **furanocoumarins** are obtained by the fusion of the 'furan ring' to the coumarin nucleus either at C-6 and C-7 positions or at C-7 and C-8 positions. A few typical examples belonging

to this class of glycosides are discussed in the sections that follows, namely: **Khellol glucoside**; **Psoralea**; and **Cantharides**.

4.2.5.2.1 Khellol Glucoside


Synonyms Khellinin; 2-Hydroxy-methyl-5-methoxyfuranochrome glucoside.

Biological Source It is obtained from the seeds of *Eranthis hyemalis* Linn., family *Ranunculaceae*, *Ammi visnaga* Lam family *Umbelliferae*.


Geographical Source The drug is indigenous to Egypt specially the Nile Delta and also to the Mediterranean region. It is cultivated in India.

Preparation The annual herb plant usually bears flowers from March to April. The harvesting is carried out when the ripening of first fertilized flowers takes place. The plants are cut and preserved in stacks, preferably in a dry place, whereby all the fruits are ripened.

Chemical Structure The **furanocoumarin** derivative **khellol glucoside** has the following structure.



Khellol Glucoside: $R_1 = H$;



Khellin: $R_1 = OCH_3$; $R_2 = H$;

Visnagin: $R_1 = R_2 = H$;

The drug also contains *two* well known aglycones **khellin** and **visnagin** as shown above.

Description The characteristic features of **khellol glucoside**, **khellin** and **visnagin** are summarised in Table 4.4.

Table 4.4 Characteristic Features of Khellol Glucoside, Khellin and Visnagin

S.No	Characteristic Features	Khellol Glucoside	Khellin	Visnagin
1.	Colour	Colourless crystals	Colourless crystals	Colourless thread like needles.
2.	Odour	No specific odour	Odourless	Odourless
3.	Taste	—	Bitter	—
4.	Solubility	Soluble in acetic acid, hot ethanol; slightly soluble in hot methanol; practically insoluble in acetone, ethyl acetate, ether, chloroform, cold alkali	Sol. In water 0.025 g/100 ml at 25°C, in acetone 3.0 g; in methanol 2.6 g; in isopropanol 1.25 g; in ether 0.5 g; in skelly solve B 0.15 g; More soluble in hot water and methanol	Very slightly soluble in water, sparingly soluble in alcohol; and freely soluble in chloroform.

Uses

1. It is mostly used as a coronary vasodilator
2. Khellin has proved to be a potent smooth muscle relaxant
3. Khellin is employed as coronary vasodilator, in angina pectoris, renal and uterine colic pains, bronchial asthma and whooping cough.

Note Khellin is found to be fairly stable when mixed with the usual tableting excipients.

4.2.5.2.2 Psoralen

Synonyms *Lata-kasturi* (Bengali); *Bahuchi* (Sanskrit).


Biological Sources They are the dried ripe fruits of *Psoralea corylifolia* Linn., belonging to the family *Leguminosae*. **Psoralen** is also found naturally in more than two dozen plant sources, namely: Bergernot, Limes, Cloves: family *Rutaceae*; Figs.: family *Moraceae*.

Geographical Sources It is grown almost throughout India as a weed in abandoned locations. It is also found in Ceylon. Several species of *Psoralea* have been used medicinally in America.


Description

- Colour** : Dark chocolate to black
Odour : Pungent and characteristic after crushing the fruits
Taste : Unpleasant, bitter and acrid
Size : 3 to 5 × 2 to 3 mm
Shape : Pods are ovoid, oblong beam shaped

Chemical Constituents The fruits of *P. corylifolia* invariably contain **fluorocoumarin** compounds known as **psoralen** and **isopsoralen** as shown below:




Psoralen



Isopsoralen


The seed kernel of *P. corylifolia* is found to contain **psolaridin** as given below:





Psolaridin

The plant also contains substituent components of the *linear* molecule, such as: **8-methoxypsoralen** and **5-methoxypsoralen** (or **bergapten**); besides *angular* molecules, such as: **anglicin** and **isobergapten**.*


* It has been observed that the naturally occurring **psoralens** lower the phototoxic potential whereas the **anglicins** may enhance it.



8-Methoxypsoralen

5-Methoxypsoralen
or
Bergapten

Angelicin



Isobergapten

However, in the recent past *two* more compounds, namely: **psoralenol** and **bavachromanol** have been reported.

Besides the fruit contains a variety of other chemical constituents, for instance: fixed oil (10%); resin (8.9%); essential oil (0.05%) and small amounts of raffinose and a pigment.

Chemical Tests (For Psoralen)

1. To a small amount of drug add a minimum quantity of alcohol for complete dissolution. Add to this 3 volumes of propylene glycol, 5 volumes of acetic acid and 43 volumes of water and shake well. The appearance of a blue fluorescence under UV-light indicates its presence.
2. The drug is dissolved in minimum amount of alcohol and on addition of a little sodium hydroxide solution exhibits a yellow fluorescence in UV-light.

Uses

1. The seeds are recommended in leprosy, leucoderma and other skin manifestations. They are also used for snake bite and scorpion sting.
2. The oleroesin extracts of seeds are employed to cure leucoderma patches.
3. The seeds also find their use as stomachic, anthelmintic, diuretic and diaphoretic.
4. It is used orally as a laxative.

4.2.5.2.3 Cantharides Beetles

Synonyms Spanish fly; Blistering fly; Blistering beetles.

Biological Sources *Cantharides* comprises of the dead and dried insects of *Cantharis vesicatoria* Linn., (*Lyatta vesicatoria*) belonging to the family *Meloidae*. **Cantharides** contains the **furanocoumarin** derivatives **cantharidin** ranging from 0.6 to 1%.

Geographical Sources These beetles are invariably found in the Southern and Central Europe residing on the plants belonging to the family *Oleaceae* and *Caprifoliaceae*. The various countries that are commercially engaged in the collection of **cantharides** are namely: Russia (now known as CIS countries), Rumania, Italy, Spain, Sicily and India.

Preparation The fully developed insects, that are brilliant green in appearance with a distinct metallic lustre, are invariably collected in the early morning on a large spread cloth by vigorously shaking the branches of the shrubs. The beetles are sacrificed either by exposing them to the vapours of chloroform, sulphur dioxide and ammonia in a closed chamber or dipping them into vinegar. The dead beetles are dried artificially at a controlled temperature not exceeding 40°C.

Description


Colour : Brilliant green or Bronze green

Odour : Characteristic odour

Size : Length = 10-20 mm: Width = 3-6 mm.

Chemical Constituents **Cantharides** essentially contains an important **vesicating*** principle termed as **cantharidin**, which is nothing but the anhydride of cantharidic acid, located in the soft portions of beetles.

Besides, **cantharides** also contains resin, formic, acetic, uric acids and fat (12-15%)



Cantharidin

Substituents/Adulterants **Cantharides** beetles are mostly substituted by *Mylabris species*, the well known Chinese cantharis having a close resemblance to the former ones. Mylabris essentially comprises of the dried beetles of *Mylabris cichorii* or *Mylabris pustulata* abundantly found in China and India, and contains **cantharidin** ranging between 1 to 1.2%.

Uses

1. **Cantharidin** has proved to a hair growth stimulant and hence used in hair oil.
2. **Cantharides** beetles, in general, is a vesicant, rubefacient and counter irritant.

4.2.6 Cyanogenetic Glycosides

The **cyanogenetic glycosides** are named so because they yield either hydrocyanic acid upon hydrolysis or they essentially possess a hydrocyanic acid in the aglycone moiety. They are also designated as 'cyanophore glycosides'. Interestingly, about 110 families belonging to the plant kingdom have been reported to contain the **cyanogenetic glycosides**; however, *Rosaceae* being the most prominent one amongst them.

It is pertinent to mention here that **cyanogenetic glycoside** containing drug substances, as such do not exert any specific therapeutic activity, but they are invariably employed as viable pharmaceutical aids, such as: flavouring agents.

A large number of **cyanogenetic glycosides** were isolated and identified from various plant sources, namely: Linamarin, Linustatin, Lotaustralin and Lucumin as shown in Table 4.5.





A few important examples of naturally occurring drug substances containing **cyanogenetic glycosides** shall be discussed here, namely; Bitter almond, Wild cherry bark and Linseed. These drugs shall be discussed in the pages that follows:

4.2.6.1 Bitter Almond

Synonym Amygdala amara.

* **Vesicating** = Causing blisters.

Table 4.5 Cyanogenetic Glycosides and Biological Source

S.No	Name	Biological Sources	Chemical Structure
1.	Linamarin (Manihotoxine; Phaseolunatin)	Clover (<i>Trifolium repens</i>); Birds foot trefoil (<i>Lotus corniculatus</i>); and other legume pasture plant. <i>Acacia</i> spp. (<i>Leguminosae</i>); seedlings of flax (<i>Linum usitatissimum</i> , fam <i>Linaceae</i>);cassava (<i>Mahihot esculentum</i>) fam: <i>Euphorbiaceae</i>); <i>Passiflora</i> spp. (<i>Passifloraceae</i>) and several other families	
2.	Linustatin	Seed meal of flax (<i>Linum usitassimum</i> , fam: <i>Linaceae</i>); also present in certain <i>Passiflora</i> spp. (fam: <i>Passifloraceae</i>)	
3.	Lotaustralin	<i>Lotus corniculatus</i> Linn., <i>australis</i> , <i>Trifolium repens</i> and other legume herbs; <i>Haloragis ereta</i> (<i>Haloragidaceae</i>); <i>Linum usitatissimum</i> (<i>Linaceae</i>); <i>Passiflora</i> spp. (<i>Passifloraceae</i>) and certain <i>Tricum</i> spp. Including <i>T. monococcum</i> (<i>Graminaceae</i>)	
4.	Locumin (Lucuminoside; Prunasin xyloside)	Seeds of <i>Calocarpum sapota</i> (<i>Sapotaceae</i> ; (S) – Epilucumin i.e.; the epimer is present in <i>Anthemis altissima</i> (<i>Compositae</i>), alongwith two cyanogenic glycoside, <i>Anthemis</i> glycosides A and B	

Biological Source **Bitter almond** comprises of the dried ripe kernels of *Prunus amygdalus* Batsch. *Var amara* (DC) Focke; *Prunus communis* Arcang., *P. amygdalus* Bail; and *Amygdalus communis* Linn., belonging to family *Rosaceae*.

Geographical Source **Bitter almond** trees are mostly native of Persia and Asia Minor. They are also cultivated in the cooler parts of Panjab and Kashmir, Italy, Sicily, Portugal, Spain, Southern France and Morocco.


Description

- Colour** : Brown
Odour : No specific odour
Taste : Bitter


- Size** : Length = 1.5 to 2 cm; Breadth = 12.5 mm;
Thickness = 8 mm
- Shape** : Oblong, ellipsoidal, rounded at one end and pointed at the other.
- Solubility** : Insoluble in ether, but soluble in water and boiling alcohol.

Preparation The cyanogenetic glycoside **amygdalin** is usually obtained from either the cake of bitter almond or other prunaceous seeds after the expression of the fixed oil. The cake is subjected to extraction with ethanol (95%, v/v), and the resulting alcoholic extract is concentrated to a small volume preferably under vacuum and mixed with a large volume of ether, when the desired glycoside will separate out as a crystalline product.

Chemical Constituent **Bitter almond** contains a colourless crystalline cyanogenetic bitter glycoside commonly termed as **amygdalin** present to the extent of 1-3% as given below:



Amygdalin upon enzymatic hydrolysis with **emulsin** gives rise to one mole each of benzaldehyde and hydrocyanic acid plus two moles of glucose as follows:




Besides, **bitter almond** contains fixed oil (40-50%), proteins (20%), volatile oil (0.5%) and an enzyme emulsin.

The enzymatic hydrolysis of **amygdalin** takes place in the following *three* steps, namely:

- The enzyme **amygdalase** helps to cleave the glycoside amygdalin first into one mole each of glucose and **prunasin** (or mandelonitrile glucoside),
- The enzyme **prunase** helps to liberate the second molecule of glucose with the formation of the aglycone *mandelonitrile* (or benzaldehyde cyanohydrin), and
- The enzyme **hydroxynitrilase** helps to break down the mandelonitrile into one mole each of benzaldehyde and hydrocyanic acid.

All these *three* steps may be summarised as given below:



It is, however, assumed that the enzyme **emulsin**, isolated from the kernels of bitter almonds, comprises of several enzymes, such as: **amygdalase**, **prunase**, **hydroxynitrilase** etc.

Chemical Tests The general tests of the **cyanogenetic glycosides** by means of microchemical reactions in naturally occurring crude drugs are based on their hydrolysis to yield hydrocyanic acid. In fact, there are *four* specific and characteristic reactions to detect the presence of liberated HCN, namely:

- 1. Ferriferrocyanide Test:** Macerate 1 g of the powdered drug with 5 ml of alcoholic KOH (5% w/v) for five minutes. Transfer it to an aqueous solution containing FeSO_4 (2.5 %w/v) and FeCl_3 (1% w/v), and maintain at 60-70°C for 10 minutes. Now, transfer the contents to HCl (20%) when the appearance of a distinct prussian blue colour confirms the presence of HCN.
- 2. Precipitation of Hg from HgNO_3 :** The reduction of aqueous mercurous nitrate solution (3% w/v) to metallic Hg by HCN being observed by an instant formation of black metallic Hg in the cells.
- 3. Grignard Reaction Test:** First of all, dip a strip of white filter paper into a solution of picric acid (1 % w/v in water) drain and then dip into a solution of sodium carbonate (10% w/v in water) and drain. Now, place the crushed and moistened drug material in a small Erlenmeyer flask, and subsequently suspend the strip of the prepared sodium picrate paper above the material and stopper the flask with an air tight cork. Maintain the flask in a warm place for 1 hour when the liberated HCN would turn the sodium picrate paper from its original yellow colour to brick red colour due to the formation of sodium isopurpurate (**Grignard's Reaction**).
- 4. Cuprocyanate Test:** First of all, saturate pieces of filter paper in a freshly prepared solution of guaic resin dissolved in absolute ethanol and allow them to dry completely in air. Now, carefully moisten a piece of the above paper with a very dilute solution of CuSO_4 and place it into contact with a freshly exposed surface of the drug. In case, HCN is generated, it will give rise to a distinct stain on the paper.

Uses

1. Bitter almonds are employed as sedative due to HCN content.
2. The fixed oil of bitter almond finds its use as demulscent in skin-lotion.
3. It is also employed in the preparation of amygdalin and bitter almond water.

Note

1. The misleading term *Vitamin B₁₇*, has sometimes been applied to **amygdalin**.
2. Bitter almond oil must not be used for flavouring of foods and confectionaries.

4.2.6.2 Wild Cherry Bark

Synonyms Viginian Prune Bark; Wild Black Cherry Cortex; Pruni.

Biological Source It is the dried bark of *Prunus serotina*, Ehrh., and *Prunus macrophylla* Sieb et Zucc, belonging to family *Rosaceae*.


Geographical Source **Wild Cherry bark** is found to be indigenous to the Eastern States of USA and certain parts of Canada. However, in the United States it is found abundantly in Dakota, Florida, Missisipi, North Carolina and Virginia.

Preparation It has been established that the **wild cherry bark** possesses the highest potency only during the autumn. Therefore, the bark is mostly collected during this period. As the inner layer of the bark contains a substantial amount of HCN, hence soonafter collection it is necessary to get rid of the inner layer of cork. Consequently, after the removal of cork as well as a portion of the cortex, the exposed surface of the bark exhibiting phloem more or less give rise to an uniform dark brown coloured product, which is commercially known as **Rossed Bark**. The resulting rossed bark is dried in the shade and stored carefully in a dry place for onward trasmission to several countries as a valued export material.

Description



- Colour** : Dark-brown colour
- Odour** : Mostly very faint; but when slightly moisten it has an odour resembling to that of benzaldehyde (bitter almond like)
- Taste** : Bitter and astringent
- Size** : Length = 10 cm; Width = 4 cm; Thickness = 3-4 mm
- Shap** : Mostly curved or channelled
- Fracture** : Short and granular
- Inner Surface** : Reddish brown and longitudinally striated
- Outer Surface** : '*Rossed Bark*' - Rough with pale buff coloured lenticel scars; '*Unrossed Bark*'—Reddish brown to brownish black, smooth, glassy and exfoliating cork having prominent whitish lenticels.

Chemical Constituents **Wild cherry bark** essentially contains a **cyanogenetic glycoside** termed as **prunasin** (or **mandelonitrile glucoside**) as shown below:


*d*-Prunasin

***d*-Prunasin** undergoes hydrolysis in the presence of the enzyme *prunase*, usually present in the bark itself, to yield one mole each of benzaldehyde, glucose and hydrocyanic acid.


Besides, the drug also contains *p*-coumaric acid, scopoletin *i.e.*, β -methylesculetin, benzoic acid and trimethyl gallic acid.

*p*-Coumaric Acid

Scopoletin



Benzoic Acid

Trimethyl
Gallic Acid

Chemical Tests The chemical tests are same as described under Section 4.2.6.1.

Uses

1. The syrup of wild cherry is mostly employed as a flavoured vehicle in cough syrup.
2. It is also used as a sedative expectorant.

4.2.6.3 Linseed

Synonym Flax seed.

Biological Source It consists of the dried fully ripe seeds of *Linum usitatissimum* Linn. belonging to family *Liliaceae*.

Geographical Sources It is cultivated extensively as a source of fibres in Algeria, Egypt, Greece, Italy and Spain; as a source of oil in Afghanistan, India and Turkey; and in Russia (now CIS – countries) for both oil and fibre. It is also found in several temperate and tropical zones.


Preparation The cyanogenetic glycoside *linamarin* is prepared from the defatted oil meal, seed-skins or embryos of flax by standard methods available for glycosides.

Description

- Colour** : Reddish brown
Odour : Characteristic odour
Shape : Oval and strongly flattened
Size : Length = 4-6 mm; Width = 2-3 mm.

Chemical Constituents The ripe seeds of linseed contain small quantities of a **cyanogenetic glycosides** known as *linamarin* (or *phaseolunatin*) as given below:

Interestingly, **linamarin** evolved HCN with linseed meal only but not with emulsin. However, pure linamarin is a bitter needle like crystalline substance. It is freely soluble in water, cold alcohol, hot acetone, slightly in hot ethyl acetate, ether, benzene, chloroform and practically insoluble in petroleum ether.



Linamarin

Besides, linseed seeds comprise of fixed oil (33-43%) mucilage present in testa (6%), proteins (25%) and an enzyme called *linase*.

Linamarin upon enzymatic hydrolysis yields HCN which actually renders the seeds highly poisonous.

Chemical Test The mucilage of linseed seed gives a distinct red colour on being treated with Ruthenium Red Solution.


Uses

1. Therapeutically, the linseed oil is mostly recommended for the external applications only; ointments and lotions.
2. It is employed in the treatment of scabies and other skin disease in combination with pure flowers of sulphur.
3. As the linseed oil has an inherent very high '**iodine value**' it is used mostly in the preparation of non staining '**Iodine Ointment**' and several other products such as: '**Cresol with Soap**'.
4. Commercially, it is one of the most important '**drying oil**'; and, therefore, substantially huge amounts are exclusively used for varnishes and paints.
5. Linseed oil finds its extensive application in the manufacture of soap, grease, polymer, plasticizer, polish and linoleum.


4.2.7 Thioglycosides

This specific group of glycosides is also referred to as '**Thiocyanate Glycosides**' or '**Sulphurated Glycosides**' or '**Glucosinolate Compounds**' or '**Isothiocyanate Glycosides**' in various literatures. The aglycone portion of such glycoside essentially contains isothiocyanate residue having sulphur plus nitrogen atoms. The general structure originally assigned to these aglycones (Formula A) has now been replaced by a more favourable one (Formula B).

In general, the **thioglycosides** are specifically abundant in several families, such as: *Cruciferae*, *Capparidaceae* and *Rosaceae*. More than forty **thiocyanate glycosides**, having a variety of



(Formula-A)



(Formula-B)

configurations in the side chain, have been isolated and identified. In fact, most glycosides belonging to this category invariably comprise of a sulphuric acid residue which on hydrolysis gives rise to a potassium salt respectively.

The *three* principal **thioglycosides** commonly known are as follows: **Sinigrin**—in **Black Mustards**; **Sinalbin**—in **White Mustards**; and **Gluconapin**—in rape seeds. These naturally occurring plant drugs shall be discussed individually in the sections that follows.

4.2.7.1 Black Mustard

Synonym Brown Mustard.

Biological Sources These are dried ripe seeds of *Brassica nigra* Linn., Koch or *Brassica juncea* Linn, Czern & Coss, belonging to family *Cruciferae*.

Geographical Sources *B. nigra* is extensively cultivated in various parts of Europe and United States. *B. juncea* is widely grown in different parts of India and the CIS-countries (*i.e.*, Russia).

Preparation The **thioglycoside** *sinigrin* is obtained from the defatted black- mustard seed by employing standard methods. It is usually present in the seeds to the extent of 4%.

Black mustard seeds are powdered and defatted with petroleum ether. The defatted meal is boiled with ethanol to destroy the enzyme. The resulting marc is squeezed while hot, dried at 100°C and maceraed in cold water for 3-4 hours with constant stirring, since **sinigrin** is fairly soluble in cold water. The liquid content is decanted and maceration is repeated a number of times to ensure complete extraction of the **thioglycoside**. The combined aqueous extract is collected and treated with mild alkalis, such as : BaCO₃, so as to neutralize any free acidity. The liquid is now concentrated under vaccum to a syrupy consistency. The resulting syrup is boiled with ethanol (95% w/v) for about 2-3 hours to allow sinigrin to dissolve and at the same time to precipitate the mucilageous components. The alcholoic extracts are filtered and allowed to cool slowly when **sinigrin** crystallizes out (approximately 4%).

Description

Colour : Black, dark brown or reddish brown

Odour : Whole seed-none; Crushed seed-pungent characteristic odour.


Taste : Bitter

Size : Approx. 0.9-1.0 mm in diameter


Shape : Mostly spherical in shape

Special Features Seeds are normally covered with a brittle testa and the kernel is oily and greenish yellow in colour. The approx. weight of 100 seeds ranges between 150 to 170 mg.

Chemical Constituents The **black mustard** seed contains a thioglycoside *i.e.*, a **β-glucopyranoside** termed as **sinigrin**. It is also known as myronate potassium or allyl glucosinolate as shown under:



However, **sinigrin** undergoes complete hydrolysis in the presence of an enzyme *myrosin* to yield one mole each of allyl iso-thiocyanate, glucose and potassium hydrogen sulphate as given below:



The **allyl iso-thiocyanate** *i.e.*, the volatile oil component of mustard is solely responsible for the characteristic pungent dour of mustard oils.

Besides, it contains fixed oil (30%), proteins (20%), and volatile oil (0.7-1.3%).

Chemical Tests

1. The powdered **black mustard** seeds on being treated with sodium hydroxide solution yields bright yellow colouration
2. Chromatographic Evaluation
 - (a) Paper chromatography of the mustard oil in a solvent system consisting of butanol-acetic acid water; and subsequently spraying the chromatogram with 0.02 N silver nitrate solution, drying at 100°C and finally spraying with 0.02 N potassium dichromate produces yellow spots against a red background of silver chromate thereby confirming the presence of *sinigrin*.
 - (b) The thiourea derivatives are used as reference compounds along with the mustard oil spots in paper chromatography using the solvent system consisting of water saturated chloroform or butanol-ethanol-water. The chromatogram is sprayed with *Grote's Reagent* (*i.e.*, a mixture of sodium nitroprusside, hydroxylamine and bromine) which distinctly yields blue spots with thiourea derivatives as well as *sinigrin*.

Uses

1. A paste of **black mustard** seed is mostly employed in the form of plaster or poultice as a rubefacient and counter irritant.
2. In higher doses, when administered internally, it acts as an emetic.
3. **Black mustard** seeds are invariably used as a widely accepted condiment in the preparation of pickles, curries and vegetables.
4. The fixed oil is widely employed as a popular edible oil.

4.2.7.2 White Mustard


Synonyms *Brassia alba* Hook f. & Th., *Sinapis alba* Linn.

Biological Source These are the dried ripe seeds of *Brassia alba* H.f. & T., belonging to family *Cruciferae*.


Geographical Source The plant is grown in India as a garden crop. It is a weed usually arising from cultivation in Panjab.

Preparation The powdered **white mustard** seeds are defatted with a suitable solvent (*e.g.*, petroleum ether, n-hexane) and the dried marc is extracted with boiling ethanol (95% v/v). The thioglycoside is purified by dissolving in warm water, decolourised with activated charcoal, filtered and the resulting filtrate is crystallied out.

Chemical Constituents The main constituent of **white mustard** is the **thioglycoside** known as **sinalbin** (or *sinapine glucosinalbate*) having the following structure:



The enzymatic hydrolysis of **sinalbin** by the enzyme *myrosin* gives rise to one mole each of glucose, acrylyl isothiocyanate and sinapine acid sulphate as shown (see page 214).



Chemical Tests

1. The hydrolysed product of sinalbin *e.g.*; sinapine acid sulphate and other salts are crystalline and give rise to a distinct bright yellow colouration in an alkaline medium .

Uses

The paste of **white mustard** seed is frequently employed in the form of a plaster or poultice as counter irritants and rubefacients.

4.2.8 Saponin Glycosides

In general a group of plant glycosides commonly referred to as **saponin glycosides**, usually share in different extents, the following *two* specific characteristics namely:

- (a) They produce foam in aqueous solution, and
- (b) They cause haemolysis of Red Blood Corpuscles (RBC).

The **saponin glycosides** are broadly regarded as haemotoxic in nature by virtue of the fact that they afford the haemolysis of erythrocytes, which render most of them as '*fish poisons*'. Invariably, they possess a bitter and acrid taste, besides causing irritation to mucous membranes. They are mostly amorphous in nature, soluble in alcohol and water, but insoluble in non-polar organic solvents like benzene, n-hexane etc.

Interestingly, the naturally occurring plant materials consisting of **saponin glycosides** have been extensively employed in various parts of the globe for their *exclusive detergent characteristics*, for instance: In South Africa the bark of *Quillaia saponaria* belonging to family *Rosaceae* and in Europe the root of *Saponaria officinalis* belonging to family *Caryophyllaceae*.

Sapogenins—The *aglycone* of the **saponin glycosides** are collectively known as *sapogenins*. *Sapotoxins*—the harmful and poisonous sapogenine/ saponins are often referred to as *sapotoxins*.

Based on the nature of the '**aglycone**' residue present in the **saponin glycosides**, they are broadly classified into the following *two* categories, namely:

- (i) Tetracyclic triterpenoid saponins (or Steroidal saponins), and
- (ii) Pentacyclic triterpenoid saponins.

These *two* categories of **saponin glycosides** will be discussed with suitable examples from plant sources in the sections that follows:

4.2.8.1 Tetracyclic Triterpenoid Saponins (or Steroidal Saponins)

Due to the enormous pharmaceutical importance a plethora of plants have been screened thoroughly for the detection of **steroidal saponins**. They are not only confined to *monocot plants* but also extended to *dicot plants*, such as:

Monocot Plants : Family—*Amaryllidaceae*, *Dioscoreaceae* and *Liliaceae*

Dicot Plant : Family—*Apocynaceae*, *Leguminosae* and *Solanceae*

However, from a commercial angle the steroidal saponins occupy a very important position in the therapeutic armamentarium which is evidenced by the following glaring examples, such as: used as raw material for the synthesis of a number of medicinally potent **steroids** e.g., vitamin D, **sex hormones**—like testosterone, progesterone, oestradiol etc., **cardiac glycosides** e.g., digoxin, digitoxin; **corticosteroids** e.g., cortisone acetate, corticosterone, aldosterone; **oral contraceptives** e.g., mestranol, norethisterone; and **diuretic steroid** e.g., spironolactone.

A few typical examples of naturally occurring medicinal plants containing **tetracyclic triterpenoid saponins** shall be described in the sections that follow, namely: *dioscorea*, *solanum khasianum* and *shatvari*.

4.2.8.1.1 Dioscorea

Synonyms Rheumatism root; Yam.

Biological Source It essentially comprises of the dried tubers of *Dioscorea deltoidea* Wall., *Dioscorea tokora* Makino, and *Dioscorea composita* and other species of *Dioscorea* belonging to the family *Dioscoreaceae*.

Geographical Source *D. deltoidea* is grown in United States and Mexico. It is cultivated from Nepal to China at an altitude ranging from 3,000 to 10,000 feet. It is also found growing abundantly in North Western Himalayas from Kashmir to Panjab in India.

Preparation Generally, the saponins do have high molecular weight and hence their isolation in the purest form poses some practical difficulties.

The tubers are washed, sliced and extracted with hot water or ethanol (95% v/v) for several hours. The resulting extract is filtered, concentrated under vacuo and the desired glycosides is precipitated with ether.

However, there are various ways and means to obtain the respective steroidal saponins from the aqueous/ alcoholic extracts as stated below:

For Acid Saponins : Lead Acetate is employed to precipitate the steroidal saponins from aqueous extract.

For Neutral Saponin : Basic lead acetate is used to precipitate the neutral steroidal saponins from aqueous extract.

Nevertheless, a few saponins are also precipitated from their aqueous solutions either by the addition of $\text{Ba}(\text{OH})_2$ solution or by the addition of $(\text{NH}_4)_2\text{SO}_4$ solution.

Note The Barayata-Saponin Complex is sometimes employed for the estimation of Saponins.

Description


Colour : Slightly brown

Odour : Odourless

Taste : Bitter and acrid

Size : Varies dependig on the actual age of the rhizomes (tubers)

Chemical Constituents The major active constituent of **dioscorea** is *diosgenin* usually present in the range of 4-6%. **Diosgenin** is the *aglycone* of saponoin **dioscin**



Diosgenin

Besides, the rhizomes contain starch to the extent of 75% but it has no edible utility because of its bitter taste. They also contain phenolic compounds and an enzyme *sapogenase*.

Uses

1. **Dioscorea** is mostly employed in the treatment of rheumatic arthritis.
2. **Dioscorea** has a tremendous potential as a commercial product because of its high content of diosgenin, which in turn is invariably employed as a starting material for the synthesis of a host of important therapeutic drugs, for instance: *sex-hormones*, *oral contraceptives* and several *corticosteroids*.

4.2.8.1.2 Solanum khasianum

Biological Source It consist of the dried and full grown berries of *Solanum khasianum* CR belonging to the family *Solanaceae*.


Geographical Source The plant grows indigenously on the *Khasia Mountains* in Assam (India).

Preparation The plant usually grows in various climatic and agricultural conditions. Almost after a duration of six months the plants are normally harvested for the collection of berries. They are dried immediately either in an artificial environment at low temperature (50-60°C) or dried preferably in shade so as to bring down the initial large moisture content to enable its prolonged storage life.

The dried berries are powdered by mechanical grinders and the oil is removed by solvent extraction. The defatted material (marc) is then extracted in a soxhlet assembly with ethanol (95% v/v) The resulting alcoholic extract is filtered, concentrated under vacuo, treated with HCl (12N) and refluxed for at least six hours. The alcoholic extract thus obtained is made alkaline by the addition of ammonia and the contents are again refluxed for a duration of 1 hour. The contents of the flask is filtered and the residue is washed, dried and taken up in chloroform. The resulting mixture is filtered and the steroidal alkaloid solasodine is obtained as a solid residue soon after evaporating the solvent.

Description The berries have a yellowish to greenish colouration with flattened smooth brown seeds.

Chemical Constituents The berries mostly comprise of the steroidal saponin **Solasonine** as shown below:



Solasonine also occurs in various *Solanum species* namely : *Solanum aviculare* Forst F., *Solanum sodomeum* L., *Solanum xanthocarpum* Schrad & Wendl, m *Solanum nigrum* Linn., *Solanum torvum* Sw., and *Solanum verbascifolium* Linn.

Besides, the berries contain a **steroidal glycoalkaloid** known as **solasodine** (Approx. 3%) and a greenish yellow fixed oil (8-10%).

Uses **Solasodine** is the hydrolysed product of **solasonine** which is mostly used as a starting material for the synthesis of steroidal drugs, such as: **19-NOR steroids, pregnane** etc.

4.2.8.1.3 Shatavari


Synonyms Shatamuli.

Biological Source The **shatavari** mostly comprises of the dried roots and the leaves of the naturally occurring plant known as *Asparagus racemosus* Will, belonging to the family Liliaceae.

Geographical Source It is widely distributed throughout the tropical regions of Africa, Australia, Asia and India. It is also found in the Himalayan range up to an altitude of 4000-4500 feet. It occurs as a widely grown plant in the dry and deciduous forests of Maharashtra State in India.

Preparation The roots usually occur in the form of a cluster or fascicle at the base of the stem. The leaves are mostly linear green and needlelike. The **steroidal saponin** is extracted by the standard methods.

Chemical Constitutents The **shatavari** contains *four steroidal saponins* usually designated as **shatavarin I-IV** present collectively to the extent of 0.2%; however, **shatavarin I** is the major glycoside present.




Uses

1. The roots are employed mostly as galactagogue to promote the flow of milk.
2. The roots are used invariably as tonic and diuretic.
3. The **steroidal saponin Shatavari-I** is reported to exert antioxytotic activity.
4. The roots are extensively employed as a medicinal oil for the control and management of nervine disorders and rheumatism.
5. In the Ayurvedic System of Medicine it is widely used both in threatened abortion and safe delivery because of its distinct uterine blocking activity.

4.2.8.2 Pentacyclic Triterpenoid Saponins

This particular class of saponin essentially contains the sapogenin component with **pentacyclic triterpenoid** nucleus, that is eventually linked with either sugars or uronic acids. It is pertinent to mention here that the sapogenin may be further classified into *three* major categories namely: **α -Amyrin**, **β -Amyrin** and **Lupeol**.




- Odour** : None
Shape : Tuberos and corpulent
Appearance : Translucent and bears the stem scars.

Chemicals Constituents **Ginseng** chiefly comprises of a complex mixture of **triterpenoid saponins** which may be either a **steroidal triterpene** or a pentacyclic related to oleanolic acid. However, these glycosides have been classified into *three* major heads, namely:


- (a) Ginsenosides,
- (b) Panaxosides, and
- (c) Chikusetsu Saponins.

Ginsenoside Rg₁, is one of the major saponins that has been isolated and identified in ginseng, with a **steroidal triterpene aglycone** known as (20S)-**protopanaxatriol** as shown below:



Ginsenoside Rg₁

In all, about 13 **ginsenosides** have been isolated and identified. Interestingly, **panaxasides** undergo decomposition yielding *oleanolic acid*, *panaxadiol* and *panaxatriol* as given below:



Uses

1. In the Chinese system of medicine ginseng is the most favourite remedy for a variety of ailments *e.g.*, as a general tonic, stimulant, carminative and diuretic activities.
2. It also possesses adaptogenic (antistress) properties and is found to exert positive action on the metabolism, the endocrine system and the central nervous system.
3. In the **orient ginseng** is used abundantly in the treatment of anaemia, diabetes, insomnia, gastritis, neurasthenia and specifically to cure sexual impotence.
4. It is found to enhance the natural resistance (*i.e.*, non-specific resistance) and increases the ability to overcome both exhaustion or illness to a great extent.
5. It prolongs the life of elderly persons and cures giddiness.

4.2.8.2.2 Liquorice

Synonyms Glycyrrhiza; Liquorice root; Glycyrrhizae radix.

Biological Sources **Liquorice** is the dried, peeled or unpeeled, roots, rhizome or stolon of *Glycyrrhiza glabra* Linn., invariably known in commerce as Spanish liquorice, or of *Glycyrrhiza glabra* Linne. var *Glandulifera* Waldstein et Kitaibel, mostly known in commerce as Russian liquorice, or of other varieties of *Glycyrrhiza glabra* Linne., which produce a sweet and yellow wood, belonging to family *Leguminosae*.

The word **Glycyrrhiza** has been derived from the Greek origin that means *sweet root*; and **glabra** means *smooth* and usually refers to the smooth, pod-like fruit of this particular species. Nevertheless, the fruits of the *glandulifera* variety has a distinct gland like swellings.

Geographical Sources **Liquorice** is grown in the sub-Himalayan tracts and Baluchistan. It is cultivated on a large scale in Spain, Sicily and Yorkshire (England) *G. glabra* var *violaceae* is found in Iran; whereas *G. glabra* var *glandulifera* exclusively grows in Russia (the '*Russian Liquorice*').

The following are the *three* commonly grown varieties of *Glycyrrhiza glabra*, namely:


- (a) *G. glabra* var. *violaceae* (or *Persian Liquorice*): This specific species bears violet flowers,
- (b) *G. glabra* var *gladulifera* (or *Russian Liquorice*): It has a distinct big stock together with a number of elongated roots, but it has not got any stolon, and
- (c) *G. glabra* var. *typica* (or *Spanish Liquorice*): This specific plant bears only purplish-blue coloured papilionaceous flowers. It possesses a large number of stolons.

Preparation The roots are usually harvested after 3 to 4 years from its plantation when they mostly display enough growth. The rhizomes and roots are normally harvested in the month of October, particularly from all such plants that have not yet borne the fruits. thereby ascertaining maximum sweetness of the sap. The rootlets and buds are removed manually and the drug is washed with running water. The drug is first dried under the sun and subsequently under the shade till it loses almost 50% of its initial weight. The large thick roots of the Russian Liquorice are usually peeled before drying. It is an usual practice in Turkey, Spain and Israel to extract a substantial quantity of the drug with water, the resulting liquid is filtered and evaporated under vacuo and the concentrated extract is molded either into sticks or other suitable forms.

Description

- Colour** : *Unpeeled Liquorice*-Externally, yellowish brown or dark brown; and internally, yellowish colour
- Odour** : Faint and characteristic
- Taste** : Sweet
- Size** : Length = 20 to 50 cm; Diameter = 2 cm
- Shape** : *Unpeeled drug*—Straight and nearly cylindrical
Peeled drug—Mostly angular
- Fracture** : Fibrous in bark; and splintery in the wood.


Chemical Constituents **Glycyrrhiza (liquorice)** comprises of a saponin like glycoside known as **glycyrrhizin** (or **glycyrrhizic acid**) as shown below:




Glycyrrhizin
[Glycyrrhizic Acid]

Glycyrrhizin is found to be 50 times as sweet as sugar. **Glycyrrhizin** upon hydrolysis loses its sweet taste and gives rise to the aglycone **glycyrrhetic acid (glycyrrhetic acid)** together with two moles of glucuronic acid. The former is a pentacyclic triterpene derivative of the β amyryn type. A host of other chemical constituents essentially include are namely: **coumarin derivatives** e.g., *umbelliferone* and *herniarin*; **flavonoid glycoside** e.g., *liquiritoside*; *isoliquiritoside*, *liquiritin*; *isoliquiritin*, *rhanoliquiritin*, and *rhamnoisoliquiritin*; **asparagine**; **22-33-dihyrostigmasterol**; **glucose**; **mannitol** and **about 20% of starch**.

Interestingly, **carbenoxolone**, which is an *oleanane* derivative is prepared from **glycyrrhiza** and possesses considerable **mineralocorticoid activity**. It is used as an anti-ulcer drug.



Glycyrrhetic Acid



Carbenoxolone

Chemical Tests

1. When sulphuric acid (80%) is added to a thick section of the drug or powder, it instantly produced a deep yellow colour.

Substituents/Adulterants *Glycyrrhiza uralensis*, also known as **Manchurian Liquorice**, which is pale chocolate brown in appearance having wavy medullary rays and exfoliated cork is mostly used as an adulterant for *G. glabra*. This particular species is from sugar, but contains **glycyrrhizin**. Sometimes, the **Russian Liquorice** is also used as an adulterant, because the drug is purplish in appearance, has long roots but having no stolons.

Uses

1. **Glycyrrhiza** has demulscent and expectorant properties
2. It is used as a masking agent for bitter drugs in pharmaceutical formulations, such as: quinine, aloe, ammonium chloride etc.
3. **Ammoniated glycyrrhiza** is employed as a flavouring agent in beverages, pharmaceuticals and confectionary.
4. The inherent surfactant activity due to the presence of saponins helps to facilitate the absorption of poorly absorbed drugs, for instance: anthraquinone glycosides.
5. It is invariably added to beer to form stable and enhanced foaminess.
6. It imparts a distinct and characteristic bitter taste to a number of beverages, such as: stout, root beer and porter.
7. The presence of **glycyrrhetic acid** exert mineralocorticoid activity and hence it is used in the treatment of inflammations, rheumatoid arthritis and Addison's disease.
8. **Liquorice** is an important ingredient in '**Liquorice compound powder**' wherein it augments the action of senna.
9. **Liquorice liquid extract** is employed as a foam stabilizer in the foam type-fire-extinguisher.
10. **Liquorice liquid extract** is used in the treatment of peptic ulcer.
11. In Europe the **glycyrrhetic acid** is employed exclusively in dermatological formulations for its remarkable antiinflammatory properties.

Caution As glycyrrhizin appreciably enhances sodium and fluid retention and promotes potassium depletion. Therefore, patients with history of either cardiac problems or hypertension must avoid consumption of significant amount of liquorice.

4.2.8.2.3 Senega

Synonyms Senega snakeroot; Seneca snakeroot; Rattlesnake root; Radix senegae; Senega root.

Biological Source **Senega** is the dried root and root stock of *Polygala senega* L., or *Polygala senega* var *latifolia* Torret Gray or *Polygala alba* Nutt. belonging to family *Polygalaceae*.

Geographical Source The plant is grown in North America and Eastern Canada. Presently, the drug is chiefly sourced from the cultivated species in Japan. However, the species grown in North West United States is known as *Northern Senega*, whereas the one found in Canada, Minnesota and Manitoba is called as *Western Senega*.


Preparation The root is collected from the wild plants normally in summer. The stems are promptly cut off and the roots are sorted out, washed thoroughly and dried either in the shade or artificial environment between 50-60°C.

Description

- Colour** : Brownish grey
Odour : Characteristic odour of methyl salicylate
Taste : First sweet and then acrid taste
Size : Length = 5 to 20 cm; Diameter = 3 to 10 cm
Appearance : A large knotty crown with a long tapering root normally curved, twisted having two or more large branches
Fracture : Short in the bark and splintery in the wood.

Chemical Constituents **Senega** essentially contains two **saponin glycosides** that are *triterpenoid in character*, namely: **senegin** (4%) and **polygallic acid** (5.5%).

Hydrolysis of **senegin** gives rise to one mole each of **senegenin**, **senegenic acid** and **presenegenin**. It has been established that **senega** contains certain other derived forms of **presenegenin** known as **Senegin II** as shown below:



The sweet taste of the drug is owing to the presence of **1, 5-anhydrosorbitol** (or **polygalitol**). Besides, the **senega root** contains fixed oil, resin, sucrose, proteins, sterol and methyl salicylate (which is formed by the enzymatic hydrolysis of the glycosides called as **primveroside**).

Substituents and Adulterants The roots obtained from *Polygala chinensis* Linn., grown almost throughout India at an altitude of 5000 feet is mostly used as an adulterant in **Senega root**.

Uses

1. The **senega root** is used extensively as an expectorant and in chronic bronchitis to relieve the spasms.
2. It is also employed as an emetic.

4.2.8.2.4 Bacopa

Synonyms Herpestis; Brahmi.

Biological Sources It comprises of the fresh stems and the fresh leaves of *Bacopa monnieri* Linn., Pennell or *Bacopa monniera* Wettst., or *Herpestis monniera* Linn., H.B. & L., belonging to family *Serophulariaceae*.

Geographical Sources The plant is grown extensively throughout the marshy places in India, Ceylon and Singapore. The plant is glabrous, succulent and creeping herb.

Preparation The leaves along with stems are collected from the fully grown plant and are dried preferably in shade. The leaves are separated from the stems and packed separately in polybags.

Description

Colour : Green


Odour : None

Taste : Bitter

Size : Length = 1.2-1.8 cm; Breadth = 2.5-10 mm

Shape : Leaves sessile, broad, entire, ovate-oblong or spatulate with black spots.

Chemical Constituents The leaves contain **saponin glycosides** known as **bacoside A** and **bacoside B** which on acid hydrolysis give rise to **triterpenoid aglycone** termed as **bacogenin A** and **bacogenin B** respectively. It also contains **asiatic acid** and **brahmic acid** as depicted below:



Uses

1. It is used in the treatment of insanity and epilepsy

2. It is also employed as a potent nervetonic, cardiogenic and diuretic
3. It is mostly used in the treatment of asthma and as an aperient ie; acts as a mild laxative.

4.2.8.2.5 Quillaja

Synonyms Soap bark; Quillay bark; Panama bark; China bark; Murillo bark; Quillaia bark.

Biological Source **Quillaja bark** consists of the inner dried bark of *Quillaja saponaria* Molina belonging to family *Rosaceae*.

Geographical Source The plant is grown in South America *i.e.*, Peru and Chile. It is also cultivated in Northern India.

Preparation The bark is collected from the trunk of wild plants. Careful incisions are made on the trunk and the bark is stripped off. The bark is freed from the outer dark coloured cork, cut into small pieces, dried, graded and packed in polybags.

Description

Colour : *Outer surface*: Pale yellowish brown, smooth with occasional reddish or blackish patches

Inner surface: Yellowish white, smooth and hard

Odour : Odourless


Taste : Astringent and acrid

Size : Length = 100 cm; Breadth = 10-20 cm; Thickness = 5-10 mm

Shape : Hard, tough and flat

Fracture : Splintery.

Chemical Constituents **Quillaja (quillaia)** contains 9-10% of colourless amorphous **triterpenoid saponin glycosides**. The glycosides on hydrolysis give rise to **quillaic acid** and **quillaia saponotoxin**. The acrid and astringent taste of the bark is due to the presence of **quillaia saponotoxin**.



Quillaic Acid

Besides, the drug also contains tannin, starch, sucrose and calcium oxalate.

Uses

1. It is used in mineral water industry.
2. It is employed for making shampoo liquid.

3. It is mostly used as a foam producer.
4. It is generally employed as an emulsifying agent.

4.2.9 Aldehyde Glycosides

Vanilla pod is the most glaring example of a naturally occurring plant that contains an **aldehyde glycoside** e.g.; *glucovanillin*; and **cinnamon bark** is another important example which contains *cinnamic aldehyde*.

4.2.9.1 Vanilla

Synonyms Vanilla beans; Vanilla pods; Fructus vanillae; Baunilha.

Biological Sources **Vanilla** consists of the cured, full grown, unripe fruit of *Vanilla planifolia* Andrews, commonly known as Bourbon, Madagascar or Mexican vanilla; or *Vanilla tahitensis* J.W. Moore, frequently termed as Tahiti vanilla, belonging to family *Orchidaceae*.


Geographical Sources The plant is indigenous to the Eastern coast of Mexico. Presently, it is widely cultivated in West Indies, Reunion, Mauritius, Seychelles, Tahiti, Java, Madagascar, Ceylon, France, Polynesia, Indonesia, Uganda, Hawaii and India.

Preparation The full grown and unripe fruits are hand picked at that particular stage when their colour changes from green to yellow. These fruits are allowed to undergo fermentation whereby a characteristic flavour and aroma develops gradually. However, the very important and critical process of fermentation essentially consists of slow drying in shades at a controlled temperature. Fermentation, in fact, helps in the conversion of **vanilloside** to **vanillin** and glucose respectively.

Description

- Colour** : Green to yellowish-green
- Odour** : Characteristic pleasant odour
- Taste** : Acrid and astringent.
- Size** : Length = 15-25 cm; Diameter = 8-10 mm
- Surface** : Longitudinally wrinkled.

Chemical Constituents The fruits of **vanilla** pod contains between 2 to 2.75% **vanillin**, which is essentially present in the form of **glucovanillin** as given below:



Glucovanillin (Avenein)

It also contains another glycoside known as **glucovanilline alcohol**, which upon hydrolysis yields **vanillic alcohol** and glucose. The vanillic alcohol on oxidation gives **vanillin**.

Uses

1. It is mostly used as a pharmaceutical aid for flavouring various liquid preparations.
2. Interestingly, the pleasant odour and flavour of **vanilla** are not only confined to **vanillin**, but more or less collectively on account of **vanillin** along with other fragrant chemical constituents.
3. **Vanillin** enhances the chocolate flavour of cocoa based malted milk foods.

4.2.10 Bitter Glycosides

In general, bitters are the edible natural products mostly consumed before any normal meals to stimulate as well as enhance the appetite. However, the **bitter glycosides** as a class do possess almost similar activities like the bitters such as: digestive, stomachic and febrifuge.

Therapeutically, the bitters have been found to exert their stimulant effects on the gustatory (*i.e.*; related to the sense of taste) nerves located in the mouth and ultimately give rise to an improved gastric juice secretion in the stomach.

The **bitter glycosides** have been found not confined to the same chemical class, but the most important ones amongst them essentially possess the *pyran cyclopentane ring*.

A number of **bitter glycosides** isolated from natural plants have been put into actual therapeutic practice, namely: **Picrorhiza**, **Gentian**, and **Chirata**, which shall be discussed in the sections that follow.

4.2.10.1 Picrorhiza

Synonyms Indian Gentian; Katki.

Biological Source It is the dried rhizome of *Picrorhiza kurroa* Royle ex Benth., belonging to family *Serophulariaceae*.

Geographical Sources It is a perennial herb grown abundantly and distributed on the Alpine Himalayas extended from Kashmir to Sikkim at an altitude between 3000 to 5000 meters. It is also found in China.

Preparation The plant is either propagated by seeds or by rhizomes. The rhizomes are collected from the cultivated and naturally growing plants washed dried and packed.

Description

Colour : *Externally*—Dark Greyish brown
Internally—White blackish

Odour : Slight and unpleasant

Taste : Bitter




Size : Length = 3-5 cm; Diameter = 0.5-1 cm

Shape : Mostly cylindrical small fragments having longitudinal wrinkled and annulations at the tip.

Special

Features : The stems and conical buds along with the drugs usually form a part of the drug itself. The roots are invariably wrinkled in the longitudinal fashion having transverse cracks. They are greyish to brown in appearance, while the fracture is tough.

Chemical Constituents The therapeutically potent constituents of the drug essentially comprises of *three vital bitter glycosides*, namely: **Picroside I**, **Picroside II** and **Kutkoside** as given below:



S.No	Glycoside	R ₁	R ₂	R ₃
1.	Picroside I	H	H	<i>Trans-Cinnamoyl</i>
2.	Picroside II	Vanilloyl	H	H
3.	Kutkoside	H	Vanilloyl	H

In fact, chemically both **Picroside** and **Kutkoside** are **C-9 monoterpene iridoid glycosides** having an epoxy moiety present in the cyclopentane ring.

Besides, it also contains organic acids, resin, sugar and tannins.

Uses

1. It is mostly employed as a vital bitter tonic
2. It is also used as a stomachic and febrifuge.
3. In large doses it exerts its action as a laxative
4. It also finds its usefulness in the treatment of jaundice.
5. Its alcoholic extract exhibits remarkable antibacterial effect.

4.2.10.2 Gentian

Synonyms Yellow Gentian; Pale Genetian; Bitter Root; Gentian Root; Radix; Radix Gentianae; Gentiana.

Biological Source **Gentian** is the dried rhizomes and roots of *Gentiana lutea* L., belonging to family *Gentianaceae*.


Geographical Source It is perennial herbaceous tree which is found to be native to the hilly zones in Central and Southern Europe. It is also grown on Vosges mountains, Yugoslavia (now known as Serbia and Croatia) and Jura.

Preparation The long rhizomes and fully grown fleshy roots of 2 to 5 year aged plants are dug up carefully and collected preferably in autumn. The roots and rhizomes are washed thoroughly to get rid of the adhered soil and then sliced into a longitudinal fashion. The freshly sliced pieces of roots and rhizomes generally appear white in colour and do not have any odour. However, during the process of gradual drying in small heaps at a controlled temperature of 50-60°C fermentation commences which eventually turns them into dark or yellow coloured product that have a characteristic odour.

Description The description of root and rhizome of **gentian** are summarised below:

Description	Root	Rhizome
Colour	Yellowish brown to dark brown	Yellowish brown to dark brown
Odour	Characteristic and agreeable	Characteristic and pleasant
Taste	Initially sweet in taste but rapidly turns very bitter	Initially sweet in taste but rapidly becomes externally bitter
Size	Length = 15 to 20 cm; Diameter = 2.5 cm;	Length = 15 cm; Diameter = 6 cm
Shape	Cylindrical and branched	Cylindrical, but usually have one or more conical buds at the apex
Outer surface	Longitudinally striated	Transversely wrinkled having marks of leaf scars and root scars


Chemical Constituents The principal *bitter glycosides* of common **gentians**, which was isolated in 1862 from *G. lutea*, is gentiopicroin, also known as **gentiopicroside**. It is present upto 2%.



It is a highly water soluble crystalline substance having a bitter value to the extent of 12,000.

The process of drying and fermentation helps in the cleavage of the above glycoside to **gentiogenin** and glucose.

The drug also contains **amarogenin**, which is considered to be a strongly **bitter glucoside** that even imparts a distinct bitter taste at 580 thousand time dilutions.



Besides, it contains **amaroswerin** and **gentioside**; and **gentinin** which is a mixture of **gentiopicrin** and **gentinin**. It also contains a flavonoid alkaloid commonly known as **gentisin** (*Syn: Gentianic acid; Gentianin; Gentin*) and **gentisic acid**.

Uses

1. It is invariably used as a bitter tonic in anorexia and dyspepsia.
2. It appreciably improves the relatively dull appetite.

4.2.10.3 *Chirata*

Synonyms Bitter stick; Chiretta; Chirayita; East Indian Balmony.

Biological Source It is the dried plant of *Swertia* (*Ophelia*) *chirata* (Roxb) Buch-Ham; belonging to family *Gentianaceae*.

Geographical Source It is found in India from Himalaya to the mountainous regions in Kashmir, Bhutan, Meghalaya and Khasi Hills at an altitude ranging between 1200-1500 meters. It is also grown in Nepal.

Preparation The plant usually flowers from July to October. It is collected for medicinal utilities as and when the capsules are fully formed. The dried plants are tied into bundles weighting approximately 1-1.25 kg.

Description

Colour : Leaves, flowers and fruits-yellowish shade; stem-yellowish brown to purple

Odour : Odourless

Taste : Extremely bitter

Size : Stems = Length: 1.0 meter; Breadth = 6 mm

Shape : Stems are mostly cylindrical, glabrous and quadrangular at the apex having a large pith.

Chemical Constituents It invariably contains **bitter principles**, namely: **ophelic acid; bitter glycosides: amarogentin and chiratin; alkaloids; gentianine and gentiocrucine**.

Substituents/Adulterants In fact, there are *three* widely known substituents for *chirata*, namely:

(a) *Swertia paniculata* Wall: The plant grows in the temperate climate of Western Himalaya, from Kashmir to Nepal at an altitude of 2000-3000 meters;

(b) *Swertia angustifolia* Buch-Ham: The plant is found in the subtropical region of Himalaya at an altitude ranging between 300 to 2000 meters from the Chenab to Bhutan; and

(c) *Swertia densifolia*: It grows in the Konkan region and usually attains a height of about 30 to 90 cm.

Uses

1. It is invariably used as a bitter tonic.
2. It also finds its use as a febrifuge.
3. It is employed in dyspepsia.
4. It has been recommended as a diuretic and in epilepsy.
5. Industrially, it is extensively used in dyeing cotton cloth.


4.2.11 Miscellaneous Glycosides

There are a number of glycosides which do not fall into the various classifications discussed under Sections 4.2.1 to 4.2.10 ; therefore, they have been grouped together under the present head *i.e.*, ‘**Miscellaneous Glycosides**’ A few important members of this group shall be described here briefly.


4.2.11.1 Steroidal Alkaloidal Glycosides

They are specifically abundant in *two* families, namely: *Liliaceae* and *Solanaceae*. Just like saponins, the **steroidal alkaloidal glycosides** do possess significant haemolytic activities, such as:


S.No.	Name	Biological Source (Family)	Steroidal Alkaloidal Glycosides	Uses
1.	Bitter Sweet	<i>Solanum dulcamara</i> Linn., (<i>Solanaceae</i>)	Solanidine	Skin disease, psoriasis
2.	Tomato	<i>Solanum lycopersicum</i> Linn., (<i>Solanaceae</i>)	Rubijervine	Antifungal
3.	Potato or White Flower Potato	<i>Solanum tuberosum</i> Linn., (<i>Solanaceae</i>)	Solanine	—



Solanidine



Rubijervine




Solanine

The sugar components are usually attached at C-3 in **solanidine** and **rubijervine** which may be galactose, glucose, rhamnose or xylose and their quantity may vary from one to four.

4.2.11.2 Antibiotic Glycosides

Streptomycin is the glaring example of an **antibiotic glycosides** produced by the soil Actinomycete, *Streptomyces griseus* (Krainsky) Waksman et Henrici belonging to family *Actinomycetaceae*. It is usually formed by the combination of the genin **Streptidine** a nitrogen containing cyclohexane derivative and **Streptobiosamine** a disaccharide representing two-thirds of the streptomycin molecule, through a glycosidic linkage as shown below:



4.2.11.3 Glycosidal Resins


The *Jalap* and *Scammony* are the two well known examples of the **glycosidal resins** occurring in the natural products, namely: *Jalap*—from the dried tuberous root of *Exogonium purga* (Hayne) Lindl. (*E. jalap* Baill., *Ipomea purga* Hayne), belonging to family *Convolvulaceae*; *Scammony*—from the dried roots of *Convolvulus scammonia* L., belonging to family *Convolvulaceae*. The glycosidal nature of these resins are evidenced by the presence of sugars, such as glucose, rhamnose and fucose on hydrolysis.

4.2.11.4 Nucleosides (Nucleic Acids)

These naturally occurring substances are of prime biological importance and essentially possess *three* vital components namely: *first*, a **sugar moiety** e.g.; *ribose* or *2-deoxyribose*; *secondly*, a **purine or pyrimidine base** e.g.; *adenine*, *guanine* and *cytosine*; and *thirdly*, a **phosphoric acid**.

A **base-sugar unit** is known as a **nucleoside**, whereas a **base-sugar phosphoric acid unit** is known as **nucleotide**.

An example of a **nucleotide** and a **nucleoside** is given here under:



Nucleotide : An adenylic acid unit of RNA

Nucleoside : Adenosine with the adenine as the heterocyclic base.

4.3 BIOSYNTHESIS OF GLYCOSIDES

Generally, the naturally occurring living plant could be regarded as the most sophisticated and meticulously designed biosynthetic laboratory not only confined to the **primary metabolites** such as: Amino acids, carbohydrates, terpenes, fatty acids which are mostly consumed as a source of edible food material by human beings, but also for a plethora of **secondary metabolites** of enormous pharmaceutical significance, for instance: glycosides, flavonoids, alkaloids, essential oils and the like. Interestingly, such naturally found chemical substances which specifically attribute plant drugs their marked and pronounced therapeutic activities are collectively termed as '**phytopharmaceuticals**'. Therefore, a higher plant is nothing but an intricate **solar energised biochemical reactor** that is exclusively responsible for the mass production of primary as well as secondary metabolites from air, water, minerals and sunlight - a source of UV radiations.

However, the *primary metabolites* are more or less widely distributed in nature practically in all organisms that are essentially required for the overall growth as well as physiological development by virtue of their basic cell metabolism. Nevertheless, the **secondary metabolites** are biosynthetically engineered products solely derived from the primary ones and are confined in their distribution strategically i.e.; being restricted to a particular taxonomic group. These products may be regarded either as various chemical adaptations to environmental stresses or they may be considered as nature's protective, defensive or offensive chemical entities against the host of microorganism, fungi, insects and higher herbivorous predators.

Thus, with regard to cellular economic cognizance the secondary products are mostly tedious to form and subsequently accumulate, and hence, invariably show up in the plant kingdom in relatively much lesser amounts in comparison to the **primary metabolites**. The **secondary metabolites** are also regarded to be as the waste products of the plant metabolic processes.

The different biosynthetic reactions taking place in the plant cells are based on certain enzymes. In fact, it is the control of enzymatic activity on the plant metabolism which ultimately governs a specific biosynthetic pathway. In general, the enzymatic reactions in plants are reversible. Under the influence of specific enzymes the **secondary metabolites** are either synthesized or hydrolysed in plants.

The biosynthetic pathways in plants may be duly elucidated and extensively studied by the aid of isotopically labelled precursors. Nowadays, with the advent of '**tracer technology**', it is a lot


easier to introduce isotopes into the anticipated precursors of plant metabolites and employed as specific ‘**markers**’ in the elaborated biogenetic experiments. It is now quite possible to unfold the mysteries of biosynthetic pathways with the use of radioactive carbon (^{14}C), hydrogen (^3H), sulphur (^{35}S) and phosphorus (^{32}P).

The biosynthesis of different categories of glycosides shall be discussed briefly in the sections that follow.

4.3.1 Biosynthesis of Anthracene Glycosides


The **biosynthesis of anthracene glycosides** may be considered under the following *two* heads, namely:

(a) *Emodin and Other Related Derivatives*: The indepth knowledge with regard to the biosynthesis of **anthracene aglycones** has been duly established from an elaborated study with microorganisms, specifically *Penicillium islandicum* as shown below:



In this particular instance, an intermediate **poly β -ketomethylene acid** is assumed to have formed from 8 acetate units which on being subjected to intramolecular condensation gives rise to **anthraquinones** *i.e.*; **emodin** and other related derivatives.


(b) *Alizarin*: Another metabolic pathway for the formation of **anthraquinone** is established and recognised through the **shikimic acid—mevalonic acid mediators** as could be seen functional in certain plants belonging to the family *Rubiaceae* as given below:



The **biosynthesis of alizarin** reveals that the *ring A* in alizarin molecule has been derived from the **shikimic acid**, whereas the *ring C* in alizarin has been incorporated by the **mevalonic acid** component.

4.3.2 Biosynthesis of Phenol Glycosides

The **biosynthesis of Phenol Glycosides** *arbutin* takes place from the shikimic acid *via* phenylalanine (an amino acid)—cinnamic acid—hydroquinone and finally to the desired glycoside as indicated below:



4.3.3 Biosynthesis of Steroid Glycosides

Biotransformation of **steroids and cardiac glycosides** (*e.g.*, **gitoxin, digitoxin**) by plant cell cultures have been studied extensively and have been reviewed by Reinhard* (1974), Stohs and Rosenberg** (1975), Stohs*** (1977), and Furuya**** (1978).

However, in general the **steroidal aglycones of cardioactive glycosides** may be assumed to have formed as a broad based overall mechanism of **steroid biogenesis** as shown below:



The steroidal molecule is considered to have generated with the head to tail linkage of several acetate units.

4.3.4 Biosynthesis of Flavonoid Glycosides

Recently, both extensive and intensive research at the enzymatic level has more or less confirmed the original hypothetical steps postulated for the incorporation of acetate and phenylalanine into flavonoids. In fact, studies related to enzymology and regulation of flavone and flavonol glycoside biosynthesis has unfolded many further details of the individual reactions. It has been reported that

* Reinhard, E., In **Tissue Culture and Plant Science**—1974, (H.E. Street Ed.), Academic Press, New York, pp. 433-459, 1974.

** Stohs S.J., and H. Rosenburg., *Lloydia*, 38, 181-194, 1975.

*** Stohs, S.J., 'In plant **Tissue Culture and its Biotechnological Applications**', (W. Barz, E. Reinhard and M.H. Zenk, eds), Springer-Verlag, New York, pp. 142-150, 1977.

**** Fuarya, T., In '**Frontiers of Plant Tissue Culture**', (T.a. Thorpe-ed.) The Boostore, University of Calgary, Alberta, Canada, pp. 191-200, 1978.

more than 20 different **flavonoid glycosides** occurring in irradiated parsley cells are based on only three **flavone** and three **flavonol** aglycones all of which have essentially very similar substitution modes (Kreuzaler and Hahlbrock*, 1973). The chemical structures of the aglycones and their probable reactions are as shown here under.

It may be observed that except for the characteristic C-3 hydroxyl moiety of **flavonols** the six aglycones essentially differ only with respect to substitution at C-3' position.

Another school of thought suggests that the flavonoid glycoside aglycones may be obtained from the major pathways ultimately leading to the synthesis of aromatic compounds in the biological systems, namely:

- (a) Acetate Pathway, and
- (b) Shikimic Acid Pathway

It has been observed that one 6-carbon fragment of the C₆-C₃-C₆ compounds derived from the **acetate pathways** gets combined with the remaining 9 carbon fragment obtained from the **shikimic (phenyl propanoid) pathway** as stated below:

Explanation

1. The C₆-C₃ segment, perhaps in the oxidation form of a cinnamic acid molecule, gets combined with three molecules of acetate to yield first a C₁₅ **chalcone** moiety an intermediate and subsequently the **flavanone** residue.
2. The simultaneous introduction or removal of OH moieties from the aromatic rings B and A gives rise to the production of a good number of derivatives,
3. Flavonoids are first formed by the introduction of the hydroxy group at position 3, whereas dehydrogenation at positions 2 and 3 results in the formation of flavonols, and
4. Evidently, the simultaneous occurrence of a variety of glycosides having the same aglycone in a specific plant species strongly supports the well established hypothesis that **glycosylation** usually takes place at a late stage of **flavonoid biosynthesis**.

4.3.5 Biosynthesis of Coumarin and Furanocoumarin Glycosides


It has been established experimentally that by using grafts of *Melilotus alba* on *Trigonella foenum graecum*, practically no **coumarin** is formed in the shoots of *M. alba*; therefore, it may be inferred that the roots are absolutely essential for coumarin synthesis, perhaps since they provided an important precursor (Reppel and Wagenbreth, 1958)**. However, more recently it has been shown with the aid of reciprocal grafting experiments involving parsnip (*Pastinacia sativa*) that **furanocoumarins** in this species are usually generated in the fruits where they accumulate, and of course, no evidence for translocation could be observed (Beyrsh, 1967)***.

There are two experimentally demonstrated pathways whereby natural products incorporating the bezopyran nucleus are usually formed, namely:


* Kreuzaler, F., and K., Hahlbrock, *Phytochemistry*, **12**, 1149-1152, 1973.

** Reppel L., and D. Wagenbreth, *Flora* (Jena), **146**, 212-227, 1958.

*** Beyrsh, T., *Planta Med.* **15**, 306-310, 1967.



- (a) In 3- and 4-phenylcoumarins, the aromatic components of this nucleus is derived from polyketide, wherein the 3 aliphatic carbon atoms and the phenyl substitute is found to originate from shikimic acid *via* a **phenylpropanoid** intermediate, and
- (b) The coumarin originates *via* the shikimate-chorismate pathway leading to **phenylpyruvic acid**, from which arise L-phenylalanine by transamination and *trans*-cinnamic acid in turn by the action of phenylalanine ammonia lyase. Generally, two types of **furanocoumarins** are recognised, namely: *Linear furanocoumarin* and *Angular furanocoumarin* as shown below:



In (A), the furan ring is fused at C-6 & C-7 positions of the benzopyran nucleus (eg., *psoralens*), whereas in (B) the fusion is between C-7 and C-8. However, the latter is less widely distributed than the former.

4.3.6 Biosynthesis of Cyanogenetic Glycosides

Cyanogenesis is the ability of living organisms that exist freely in the higher plants but instead it is released from the **cyanogenetic precursors** as a result of the enzymatic action. Now, it has been well established that these precursors are normally glycosides of **hydroxynitriles** (or **cyanohydrins**). Once the cellular integrity of a cyanophoric plant tissue is disrupted, the **cyanogenetic glycosides** in turn are brought in contact with the respective catabolic enzymes that helps to hydrolyze the glycosides and ultimately give rise to the formation of hydroxynitriles.

There are two important **cyanogenetic glycosides**, namely: **dhurrin** nad **prunasin** which are present in a variety of plant families and genera as stated below*:


S.No.	Glycoside	Family	Genera
1.	Dhurrin	<i>Gramineae</i> <i>Proteaceae</i> <i>Trochodendraceae</i>	<i>Sorghum</i> spp. <i>Macadamia</i> , <i>Stenocarpus</i> <i>Trochodendron</i>
2.	Prunasin	<i>Caprifoliaceae</i> <i>Compositae</i> <i>Leguminosae</i> <i>Myoporaceae</i> <i>Myrtaceae</i> <i>Oliniaceae</i> <i>Polypodiaceae</i> <i>Rosaceae</i> <i>Saxifragaceae</i> <i>Scrophulariaceae</i>	<i>Sambucus</i> <i>Achillea</i> , <i>Centaurea</i> , <i>Chaptalia</i> <i>Acacia</i> Spp. (Australia), <i>Holocalys</i> <i>Eremoophila</i> <i>Eucalyptus</i> <i>Olinia</i> <i>Cystopteris</i> , <i>Pteridium</i> <i>Cotoneaster</i> Spp. <i>Cydonia</i> <i>Jamesia</i> <i>Linarria</i> Spp.

* Seigler, D.S., *Prog. Phytochem.*, **4**, 83-120, 1977.


Seigler, D.S., *Rev Latinoam Quim*, **12**, 39-48, 1981.

The two amino acids *Tyrosine* and *Phenylalanine* are considered to have derived from the **Shikimic Acid Pathway** and **Phenylalanine Pathway** as depicted below:


(a) **Shikimic acid Pathway:** It is based on studies carried out with *E. coli* as follows:




(b) **Phenylalanine Pathway:** It starts with **Shikimic acid** as the starting material as shown under:




(c) **Tyrosine from Phenylalanine:** It is obtained by the oxidation of phenylalanine.



(d) **Dhurrin from Tyrosine:** It is obtained from tyrosine as shown below:



(e) **Prunasin (or Prulaurasin) from Phenylalanine:** It is obtained from phenylalanine as given below:




It has been established that labelled **shikimic acid** and labelled **tyrosine** were equally effective precursors of the hydroxylated aglycone of *Dhurrin*, a **cyanogenetic glycoside** produced by *Sorghum vulgare* Linn. belonging to family *Gramineae*.

Likewise, introducing labelled phenylalanine to young cherry laurel plants ie; *Prunus laurocerasus*, has proved that the amino acid acts as a precursor of *Prunasin* in young peach seedlings.

4.3.7 Biosynthesis of Thioglycosides

The seeds of a number of plants belonging to the mustard family comprise of glycosides, the aglycones of which are invariably isothiocyanates. Hence, these glycosides are also known either as **thioglycosides** or as **glucosinolates** and mostly form a group of bound toxins.


It has been observed that the aglycone portions of **thioglycosides** may largely consist of either aliphatic or aromatic derivatives. It is established experimentally that the carboxy-labelled acetate is being incorporated in the allyl moiety of *Sinigrin* usually present in *Brassica juncea*.



4.3.8 Biosynthesis of Saponin Glycosides


Saponins are usually of *two* types, namely: *first*, the **steroidal saponins** which essentially have a spiroketal side chain, and *secondly*, the **triterpenoid saponins**. It has been proved that labelled

acetate and mevalonate are duly incorporated into **spiroketal steroids** as well as **pentacyclic triterpenoids**. It is, however, pertinent to mention here that the major pathway adopted by both types of sapogenins is more or less identical and that it involves head to tail coupling of various acetate units. It is assumed that a branching takes place most probably after the formation of the triterpenoid hydrocarbon **squalene**, that ultimately leads to the **spiroketal steroids** in one direction and to the **pentacyclic triterpenoids** in the other as shown below:



4.3.9 Biosynthesis of Aldehyde Glycosides

It has been established that the aromatic nuclei of **aldehyde glycosides** are usually from the $\text{C}_6\text{-C}_3$ precursors formed *via* the *Shikimic Acid Pathway*. However, the conversion of **cinnamic acid** to **vanillin** is considered to be the most propable route as shown under:



4.4 PROFILE OF GLYCOSIDES IN NATURAL PLANT SOURCES

The variety of glycosides occurring in natural plant sources are so numerous that it is not quite possible to include all of them in one chapter in the present context. Hence, it is thought worthwhile to summarize them in the following Table 4.6:

Table 4.6 Summary of Glycosides in Natural Plant Sources

S. No.	Class	Name	Biological Source (Family)	Major Glycosides	Uses
I	Anthracene Glycosides	Aloe	<i>Aloe barbadensis</i> ; A. Ferox; A. perryi; A. africana; A. spicata (Liliaceae)	Aloin, Aloe-emodin,	Purgative
		Rhubarb	Rheum officinale; R. palmatum; R. emodi; R. webbianum; (Polygonaceae)	Rhein; Aloe emodin;	Bitter Stomachic; Purgative
		Cascara sagrada	Rhamnus purshiana (Rhamnaceae)	Barbaloin; Deoxybarbaloin	Cathartic, Tonic and Stomachic
		Frangula	Rhamnus frangula (Rhamnaceae)	Carcaroside-A, B, C & D Frangulin A & B; Glucofrangulin-A & B	Cathartic
		Senna	Cassia senna; C., angustifolia; (Leguminosea)	Sennoside A, B, C & D	Purgative
		Chrysarobin	<i>Andira araroba</i> (Leguminoseae)	Chrysophano anthranol	Ringworm
II	Phenol Glycosides	Bearberry (Uvs ursi)	<i>Bergenia crassifolia</i> (Saxifragaceae)	Arbutin	Diuretic
		Canadian Wintergreen Plant	Gaultheria procumbens and Monotropa hypopitys (Ericaceae)	Gaultherin	—
		Poplar and Willow	Salix fragilis and Salix purpurea (Salicaceae)	Salicin Populin	Analgesic Bitter Stomachic
		Khatta	Citrus aurantium (Rutaceae)	Hesperidin	—
		Irridis rhizoma (Orris root)	Iris kumaonensis (Iridaceae)	Iridin	Cathartic, stimulant, Diuretic
		Apple	<i>Malus sylvestris</i> (or <i>Pyrus malus</i>) (Rosaceae)	Phlorizin	—
III	Steroid Glycosides	Digitalis (Fox Glove)	<i>Digitalis purpurea</i> (Serophulariaceae)	Purpurea Glycosides A, B, & C; Digitoxin	Cardiotonic
		Digitalis	<i>Digitalis lanata</i> (Scrophulariaceae)	Lantosides A, B & C; Digoxin	Cardiotonic
		European Squill (Sea Onion)	<i>Urginea maritima</i> (Scilla maritima) (Liliaceae)	Glucoscillaren Scillaren -A; Scillarenin;	Cardiotonic
		Indian Squill (Scilla)	<i>Urginea indica</i> (Liliaceae)	Proscillaridin Scillaren-A;	Rat poison
		Red Squill (Red variety of European Squill)	<i>Urginea maritima</i> (Liliaceae);	Scillaren-B Scilliroside	Cardiotonic
		Strophanthus	<i>Strophanthus hispidus</i> (or <i>S. kombe</i>) (Apocynaceae)	Cymarín; K- Strophanthidin-β; K-Strophanthoside	Cardiotonic
		Ouabain	<i>Strophanthus gratus</i> (Apocynaceae)	G-Strophanthin (Ouapain)	Cardiotonic
		Oleander	<i>Nerium oleander</i> (Apocynaceae)	Oleandrin	Cardiac Stimulant
		Black Hellebore	<i>Helleborus niger</i> (Ranunculaceae)	Hellebrin	Cardiotonic

(Contd.)

(Table 4.6 contd.)

	Adonis	Adonis vernalis (Renunculaceae)	K-Strophanthin; Adoniotoxin	Cardiotonic
	Thevetia	Thevetia nerifolia (Apocynaceae)	Thevetin- A	Cardiotonic
IV Flavonoid Glycosides				
A Flavone Glycosides	Parsley	Petroselinum sativum and Apium petroselinum (Umbelliferae)	Apin	Stimulant; Diuretic;
	Buchu	Barsoma crenulata B. serratifolia; B. betulina; (Rutaceae)	Diosmin	—
B Flavonol Glycosides	Buck Wheat	Fagopyrum esculatum (Polygonaceae)	Rutin	Decrease capillary fragility
	Ring	Crategus oxycantha (Rosaceae)	Quereitrin	Textile dye
C Flacanone Glycosides	Lemons, Sweet Oranges; Bitter Oranges;	Citrus sinensis; Citrus limon; C. aurantium, C. medica, (Rutaceae)	Hesperidin	Minimise capillary fragility
D Chalone Glycosides	Safflor Red	Carthamus tinctorius	Carthamin	Dye
E Isoflavonoid Glycoside	Sharapunkha	Tephrosia purpurea (Leguminosae)	Tephrosin	Fish poison
	Gilas Green Pods	Prunus avium (Rosaceae) Sopora japonica (Leguminosae)	Prunetrin Sophoricoside	— —
F Anthocyanidin Glycosides	Pelargonium Flower	Pelargonium gravecolens and P. roseum (Geraniaceae)	Plargonidin	Diuretic
	Althaea Flower	Althaea rosea (Malvaceae)	Cyanidin	—
	Peony Tuber	Peony officinalis (Ranunculaceae)	Peonidin	—
	Whortle Berry	Vacinium myrtillus (Ericaceae)	Delphinidin	—
	Petunia Flower	Petunia hybrida (Solanaceae)	Petunidin	—
	Malva Flower	Malva sylvestris (Malvaceae)	Malvidin	—
V Coumarin and Furanocoumarin Glycosides				
A Coumarin Glycosides	Horse-chestnut Tree;	Alsculus hippoeastanm (Hippocastanacea)	Aesculin	Fruit and bark in diarrhoea Tonic
	Chicory Flower	Chicorium intybus (Compositae)	Chicorin	Febrifuge Diarrhoea
	Daphne Bark	Daphne mezerium (Thymelaceae)	Daphnin	Febrifuge
	European Ash	Fracinus exclsior (Oleaceae)	Fraxin	Tonic; Febrifuge; Bitter Astringent;
	Tonka Seed	Dipteryx odorata D. oppositifolia (Leguminoseae)	Coumarin	Pharmaceut ical aid
B Furanocourmarin	Honey Plant (Visnaga)	Ammi Visnaga (Umbelliferae)	Khellin; Khellol; Glucoside;	Coronary vasodiltor
	Psoralea Fruit	Psoralea Corylifolia (Leguminosae)	Psoralen, Isopsoralen	In leprosy and leucoderma
	Cantharides	Cantharis	Cantharidin	Vesicant

(Contd.)

(Table 4.6 contd.)

	Beetles (Spanish Fly)	vesicatoria (Mesloidae)			rubefacient and hair growth stimulant		
	Mylabris (Chinese cantharides)	Mylabris sidae (Coleoptera)	Cantharidin		Rubefacient, Counter irritant		
VI	Cyanogenetic Glycosides	White Clover	<i>Trifolium repens</i> (Leguminosae)	Linamarin	—		
		Flax	<i>Linum usitassimum</i> (Linaceae)	Linustatin	Demulscent as poultices		
		Lotus	<i>Lotus corniculatus</i> (Leguminosae)	Lotaustralin	—		
		Legumes					
		Calcocarpum seeds	<i>Calcocarpum</i> <i>sapota</i> (Sapotaceae)	Lucumin	—		
		Anthemis Seeds	<i>Anthemis altissima</i> (Compositae)	Anthemis Glycosides A and B	—		
		Bitter Almond	<i>Prunus amygdalus</i> <i>P. communis</i> ; <i>Amygdalus</i> <i>communis</i> ; (Roseceae)	Amygdalin	Sedative; Demulscent in skin lotion		
Wild cherry Bark	<i>Prunus serotina</i> <i>P. macrophylla</i> (<i>Rosaceae</i>)	d-Prunasin	Sedative expectorant				
VII	Thioglyco- sides	Black Mustard	<i>Brassica nigra</i> <i>Brassica juncea</i> (Cruciferae)	Sinigrin	Rubefacient Counter irritation		
		White Mustard	<i>Brassica alba</i> (<i>Cruciferae</i>)	Sinalbin	Rubefacients, Counter Irritation;		
VIII	Saponin Glycosides	A	Tetracyclic Triterpenoid Saponins	Rheumatism Root	<i>Dioscorea</i> <i>delitoidea</i> ; <i>C. tokora</i> ; <i>D. composita</i>	Diosgenin	As a starting material for some potent steroidal drugs
				Solanum Berries	<i>Solanum</i> <i>khasianum</i> (Solanaceae)	Solasonine	Starting material for synthesis of steroidal drugs
				Asparagus Roots	<i>Asparagus</i> <i>recemosus</i>	Sarsapogenn; Shatavarin-I, Shatavarin-IV	Tonic Diuretic
		B	Pentacyclic Triterpenoid Saponins	Ginseng	<i>Panax ginseng</i> , <i>P. japonica</i> ; <i>P. notoginseng</i> ; <i>P. quinquefolium</i> ; <i>P. pseudo ginseng</i> ; (Araliaceae)	Ginsenoside- Rg ₁ Panaxosides	General tonic; Stimulant; Carminative, Diuretic
				Liqorice	<i>Glycyrrhiza glabra</i> ; (Leguminosae)	Glycyrrhizin	Demulscent Expectorant
				Senega	<i>Polygala senega</i> <i>Polygala alba</i> ; (Polygalaceae)	Senegin-II	Expectorant; Emetic
				Bacopa	<i>Bacopa monnieri</i> (Scrophulariaceae)	Bacogenin A, Bacogenin B	Insanity; Epilepsy
				Soap Bark	<i>Quillaja saponaria</i> (Rosaceae)	Quillaia Sapotoxin	Foam producer
				Sarsaparilla	<i>Smilax aristolochiaefolia</i> ; <i>S. regelii</i> (Liliaceae)	Sarsapogenin; Smilagenin	Rheumatism, Skin ailments, Syphilis

(Contd.)

(Table 4.6 contd.)

IX	Aldehyde Glycosides	Vanilla Beans (Vanilla Pods)	<i>Vanilla planifolia</i> <i>Vanilla tahitensis</i> (<i>Orchidaceae</i>)	Glucovanillin	Pharmaceutical aid;
X	Bitter Glycosides	Indian Gentian Bitter Root	<i>Picrorhiza kurroa</i> (<i>Scrophulariaceae</i>) <i>Gentiana lutea</i> (<i>Gentianaceae</i>)	Picoside I, Picoside II, Kutposide Gentiopeirin, Amarogentin	Tonic, Stomachic, Febrifuge, Laxative Bitter tonic in dyspepsia and anorexja
		Bitter stick (East Indian Balmomy)	<i>Swertia</i> (<i>Ophelia</i>) <i>Chirata</i> (<i>Gentianaceae</i>)	Chiratin, Amarogenin	Bitter tonic Febrifuge Diuretic Epilepsy
XI	Miscellaneous Glycosides				
A	Steroidal Alkaloidal Glycosides	Bitter Sweet Tomato	<i>Solanum dulcamara</i> (<i>Solanaceae</i>) <i>Solanum lycopersicum</i> (<i>Solanaceae</i>)	Solanidine Rubijervine	Psoriasis skin diseases Antifungal
B	Antibiotic Glycosides	Streptomycin	<i>Streptomyces griseus</i> (<i>Streptomyce-taceae</i>)	Streptomycin	Antibiotic
C	Nucleosides (Nucleic Acids)	Adenosine	Base-sugar unit as a nucleoside	Adenosine	RNA, DNA formation

FURTHER READING REFERENCES

1. Baker, W., and W.D., Ollis., In: **Recent Developments in the Chemistry of Natural Phenolic Compounds**, Ed by W.D. Ollis, Pergamon Press, Oxford, 152, 1961.
2. Bonner, J., and J.E. Varner., '**Plant Biochemistry**', Academic Press, London, 1965.
3. Britton, G., '**The Biochemistry of Natural Pigments**', Cambridge University Press, Cambridge, 1983.
4. Conn, E.E., '**The Biochemistry of Plants**', Vol. 7, *Secondary Plant Products*, Academic Press, London, 1981.
5. Ciba Foundation Symposium: 140, **Cyanide Compounds in Biology**, John Wiley & Sons Ltd., Chichester (U.K.), 1988
6. Griffith, J.Q., Krewson, C.S. and J. Naghski, Eds., '**Rutin and Related Flavonoids**', Mack Publishing Company, Easton Pennsylvania, 1959.
7. Glasby, J.S. '**Encyclopedia of the Terpenoids**', Vols I & II, Jphn Wiley & Sons Ltd. Chichester (U.K.), 1982.
8. Harborne, J.B., Ed. '**Biochemistry of Phenolic Compounds**', Academic Press Inc., New York, 1964.
9. Mann J., Davidson R.S., Hobbs J.B., Banthrope D.V., and Hardborne J.B.: **Natural Products: Their Chemistry and Biological Significance**, Longman, Harlow, 1994.
10. Poulton, J.E., **Cyanogenic Compounds in Plants and Their Toxic Effects. In Handbook of Natural Toxins** Vol I, Keeler, R.F., Tu A. T. Eds., Marcel Dekker Inc., New York, 1983.
11. Shibata, S., Tanaka, O., Shoji, J., and H. Saito., **Chemistry and Pharmacology of Panax, In Economic and Medicinal Plant Research**, Vol I, Wagner, H., Hikino, H., Farnsworth, N.R., Eds., Academic Press, Inc., Orlando, 1985.
12. Thompson, R.H., '**Naturally Occurring Quinones**', Butterworth, London, 1957.
13. Torsell K.B.G.: **Natural Product Chemistry. A Mechanistic, Biosynthetic and Ecological Approach**, Apotekarsocieteten, Stockholm, 1997.
14. Wright, S.E., '**The Metabolism of Cardiac Glycosides**', Springer Verlay, Berlin, 1960.


5

Terpenoids

- Introduction
- Classification
- Further Reading References

5.1 INTRODUCTION

A plethora of naturally occurring plant products have been found to be related wherein they are comprised of one or more units of **isoprene** (C₅H₈)-a hydrocarbon:




In general, **terpenoids**, may be defined as natural products whose structures are considered to be divided into several isoprene units; therefore, these compounds are invariably termed as **isoprenoids**. Besides, this particular group of compounds is sometimes collectively referred to as the **terpenes** in relatively older texts. Logically, the *-oid* suffix seems to be more acceptable and convincing, as it is in the same vein for steroids, alkaloids, flavonoids, etc., However, the *-ene* suffix must be solely confined to the unsaturated hydrocarbon belonging to this specific class of compounds.

It has now been established experimentally that the **isoprene units** come into being through the biogenetic means starting from acetate *via* mevalonic acid. Each such unit essentially consists of five-carbons having two unsaturated bonds and possesses a branched chain. The **terpenoids** usually have a number of such **isoprene units** joined together *in a head to tail manner*, as exemplified below:

Terpenoids are broadly classified on the basis of the number of **isoprene units** incorporated into a **specific unsaturated hydrocarbon terpenoid molecule**, such as:

- (a) **Monoterpenoids:** These are built up of *two* isoprene units and have the **molecular formula C₁₀H₁₆**;
- (b) **Sesquiterpenoids:** These are composed of *three* isoprene units and have the **molecular formula C₁₅H₂₄**;
- (c) **Diterpenoids:** These are comprised of *four* isoprene units and have the **molecular formula C₂₀H₃₂**;







(d) **Triterpenoids:** These contain *six* isoprene units and have the **molecular formula $\text{C}_{30}\text{H}_{48}$** ; and

(e) **Tetraterpenoids (or Carotenoids):** These are made up of *eight* isoprene units and have the **molecular formula $\text{C}_{40}\text{H}_{64}$** .

Biogenetic Isoprene Rule The very idea and basic concept that **terpenoids** are essentially built up of several **isoprene units** is commonly termed as the **biogenetic isoprene rule** as could be observed from the various typical examples cited earlier.

Meroterpenoids It has been observed that a good number of other natural products do exist which essentially belong to mixed biosynthetic origin and are mostly made up from **isoprene** as well as **nonisoprenoid** entities.



It has been observed that when the substance is **monomeric*** the corresponding nitrosochloride provides a distinct blue colouration, which also ascertains the presence of tetrasubstituted ethylenes.

5.2 CLASSIFICATION


Based on the extensive distribution of **terpenoids** in the vast plant kingdom they are classified broadly as follows, namely:

- (i) Monoterpenoids
- (ii) Sesquiterpenoids
- (iii) Diterpenoids
- (iv) Triterpenoids
- (v) Tetraterpenoids and Carotenoids
- (vi) Volatile Oils (or Essential Oils)
- (vii) Resins and Resin Combinations
- (viii) Oleoresins
- (ix) Oleo-Gum-Resins
- (x) Balsams

These different classes of naturally occurring substances shall be discussed individually in the sections that follows:

5.2.1 Monoterpenoids

In general, **monoterpenoids** represent a structurally diverse class of compounds may be categorised into nearly 35 varying structural analogues. However, the most commonly occurring structural variations are of the following types, namely:

Name	Chemical Structure	Type
Myrcene		Acyclic
p-Menthane		Monocyclic
α-Pinene		Bicyclic

* **Monomeric**: An entity or a unit from which polymer is formed.

It has been found that a large number of **monoterpenoid** derivatives belonging to these categories invariably occur naturally in the purest optically active form; however, certain plant species do have both enantiomers, such as: *Pinus* species contain both (+)- and (-)- α – pinene.

A few typical examples of monopenoids found in naturally occurring plant species are described under: *camphor*, *eucalyptol*, *menthol* and *thymol*.

5.2.1.1 Camphor

Synonyms Gum camphor; Japan camphor; Formosa camphor. Laurel camphor.

Biological Source It occurs in all parts of the camphor tree, *Cinnamomum camphora*. T. Nees & Ebermeier, belonging to family *Lauraceae*.

Geographical Source The word *camphora* is derived from the Arabic *Kafur*, meaning chalk. The **camphor** tree, which is a huge evergreen plant, is found to be indigenous to Japan, China and Taiwan. It has also been naturalized specifically in the Mediterranean region eg; Algeria, Tunisia, Libya, Egypt, Italy and Greece. Besides it is grown in South Africa, Ceylon, Brazil, Jamaica, Florida and California. History reveals that Borneo camphor (from Borneol) arrived in Arabia in the sixth century and in Europe in the twelfth century. Earlier, the world's 80% supply of natural camphor was provided by Taiwan (Formosa) alone and the rest 20% by Japan and Southern China. Soon after the second World War (1945) the commercial production of **synthetic camphor** has more or less catered for the ever increasing demand of **camphor** in the world market.


Preparation It is prepared from the chipped wood by subjecting it to steam distillation and subsequently collecting the distillate in specifically designed chambers where camphor will solidify on its inner walls upon cooling and may be collected later on from the bottom of the chamber. The crude solidified camphor is purified by mixing it with a suitable proportion of soda lime, sand and charcoal; and subjecting the mass to sublimation at controlled temperature when pure crystals of camphor would be collected as a sublimate. It is finally compressed into either small cubes or thin plates, wrapped and exported.

Camphor from Volatile Oils It may be prepared from volatile oils by *two* simple methods, namely:

Methods-I In case, the oil contains a substantially large proportion of **camphor**, it may be separated by deep freezing or sudden chilling; and if the camphor content in oil is not so much it is mostly fractionated and the camphor containing fraction is chilled to recover camphor.

Method-II **Camphor** may be recovered from volatile oils by the instant production of insoluble complexes with strong mineral acids eg; sulphuric acid 80% (30N).

Synthetic Camphor (or Borneol Camphor) The **camphor** is obtained commercially from α -pinene present in the turpentine oil through several steps sequentially *e.g.*, treatment with HCl, isomerization, treatment with KOH and finally oxidation with HNO_3 as given below:



Description


Colour : Translucent mass with crystalline fraction

Odour : Characteristic odour

Optical Activity : Natural camphor = Dextro rotatory (+ 41° to 43°) Synthetic camphor = Racemic mixture;



Solubility : Soluble in water (1:600)

Chemical Structure Camphor is a **bicyclic terpenoid ketone** as given below:




Camphor

In the presence of platinum black it undergoes hydrogen at ambient temperature giving rise to **isborneol** as the major product and traces of **borneol**.

 α -Borneol

1-Isborneol

Its prolonged hydrogenation often yields **camphene**.



Camphene

Chemical Tests

1. Its semicarbazone derivative using semicarbazine hydrochloride has a m.p. 247-248°C for d-camphor.
2. Its 4-dinitrophenylhydrazone derivative has different mp *e.g.*, *d*- and *l*- = 175°C and *dl* = 164°C.

Camphor in presence of Borneol

1. **Borneol** is first esterified with stearic acid to yield a high-boiling ester. The resulting mixture on steam distillation removes camphor as the product while the ester remains in the flask.
2. **Borneol** forms an adduct with either succinic anhydride or phthalic anhydride upon heating to yield borneol acid succinate and borneol acid phthalate respectively, which are soluble in

NaOH solution. Camphor (unreacted) is subsequently extracted with ether from the alkaline medium.

3. **Camphor** forms an oxime ($>C=N-OH$) with hydroxylamine which is then dissolved in sulphuric acid; the unreacted borneol is removed by extraction with ether.

Distinction between Natural Camphor and Synthetic Camphor A drop of freshly prepared vanillin solution (1: 100 in dilute HCl) and sulphuric acid when added to powdered natural camphor, it gives rise to an instant yellow colouration changing to red, violet and finally blue. The **synthetic camphor** fails to respond to this test and gives a distinct bright smoky flame.

Uses

1. It is used as a topical antipruritic in concentrations ranging between 0.1 to 0.3%
2. It is mostly used as a counterirritant (11%) particularly for fibrositis and neuralgia associated with inflamed joints, sprains and other inflammatory manifestations.
3. It is also employed as antipyretic, antiseptic, antifungal and carminative agent.
4. It is employed as a safe and effective measure for reducing cough when applied externally, in the form of an ointment, on the chest and throat of children.
5. It exerts its stimulant, rubefacient, antispasmodic and analgesic activities.
6. It stimulates the nerve endings in the skin and causes substantial relief of pain due to the masking of deeper visceral pain with a milder pain arising from the skin at the same level of innervation.

5.2.1.2 Eucalyptol

Synonyms Cineole; Cajeputol.

Biological Source It is obtained from the leaves of *Eucalyptus globulus* Labill, belonging to family *Myrtaceae*.

Geographical Source The eucalyptus tree is a native of Australia and Tasmania. It is largely cultivated in California, Spain, Portugal and India. In India it is abundantly found in the Himalayan region, Nilgiri district, Kumaon Hills and Assam.

Preparation A number of volatile oils from certain *Eucalyptus species* invariably contain eucalyptol as high as 30 to 70%. It also occurs in **cajuput oil** (40%) and in **laurel leaf oil** (50%). However, **eucalyptol** may be isolated from these oils by adopting one of the following methods:


Method 1 By subjecting the volatile oil to fractional distillation and collecting the fractions between 170-180°C to obtain crystals of **cineole** at -10°C (m.p. + 1.5°C)

Method-2 **Cineole** forms addition compounds with halogen acids, e.g., $C_{10}H_{18}O \cdot HCl$ and $C_{10}H_{18}O \cdot HBr$; with phosphoric acid as $C_{10}H_{18}O \cdot H_3PO_4$ which also serve as a means of its purification, and

Method 3 **Eucalyptol** yields an addition product with a 50% (w/v) alcoholic solution eg; $C_{10}H_{18}O \cdot C_6H_6O_2$ (mp 82-85°C), from which the former may be generated.

Note: This method is mostly applicable to such volatile oils that have a higher cineole content.

Synthetic Method **Eucalyptol** may be prepared synthetically by the dehydration of **terpin hydrate** as given below:



Description


Colour : Colourless or pale yellow liquid.

Odour : Camphoraceous and aromatic.

Taste : Pungent and leaves a cold sensation.

Solubility : Water insoluble; soluble in paraffin, fixed oils and ethanol 90%.

Chemical Structure



Eucalyptol is an epoxy or oxido derivative of p-menthane. and is also known as 1,8-epoxy-p-methane or 1,8-oxido-p-menthane. It is found to be quite stable and hence may be distilled over metallic sodium safely without undergoing any change whatsoever. It is not affected by the action of reducing agents.

Chemical Tests

1. When a drop of **eucalyptol** is carefully treated with a drop of 5% (w/v) solution of hydroquinone in alcohol on a slide, it forms either colourless prisms or rhomboids; but with a 50% (w/v) solution of resorcinol in alcohol leaf-like crystals are obtained.
2. It forms characteristic addition compounds with HCl, HBr and H₃PO₄ with well defined melting points.

Uses

1. It is used internally as a stimulating expectorant to relieve severe cough and in bronchitis in the form of inhalations.
2. It is abundantly employed externally as a mild anaesthetic and antiseptic for the treatment of various inflammatory conditions.

3. It also finds its use as a decongestant nasal drops.
4. It is profusely used in room sprays, lotions and all types of cosmetic preparations.
5. It is also used as a flavouring agent in pharmaceutical preparations *e.g.*; mouth washes, and gargles.

5.2.1.3 Menthol


Synonyms 1-Menthol; 3-Menthanol; Menthan-3-ol; Peppermint camphor, Hexahydrothymol.

Biological Sources It is found in the **peppermint oil** obtained from the fresh flowering tops of the plants commonly known as *Mentha piperita* Linn., or other allied species of *Mentha*, belonging to family *Labiatae*.

Geographical Source Various mentha species are duly cultivated in various parts of the world. It grows both abundantly and widely in Europe, while it is cultivated in Japan, Great Britain, Italy, France, United States, CIS countries, Bulgaria and India.

Preparation It is normally prepared from *Japanese Peppermint Oil*, from the flowering tops of *Mentha avensis* Linne' var *piperascens*, by subjecting it to refrigeration below -22°C whereby the menthol crystallizes out distinctly. The crystals of **menthol** are separated by filtration and squeezed between layers of filter papers to remove the adhering oil and finally purified by recrystallization.

Synthetic racemic menthol is prepared by the hydrogenation of either **pulegone** or **thymol** as shown below:



It may also be prepared from **pinene**.

Description


Colour : Colourless

Odour : Pleasant peppermint like odour

Taste : Characteristic, aromatic and cooling taste


Shape : Hexagonal crystals usually needle like, prisms; crystalline powder; fused masses.

Chemical Structure **Menthol** has three chiral centres (*), hence it would give rise to eight (2^3) optically active isomers and four racemic forms. **Menthol** on oxidation gives menthone (a ketone), by the sacrifice of one chiral centre; therefore, the resulting menthone must exist in four (2^2) optically active isomers and two racemic forms and all, these have been actually prepared.




Special Features following are the special features of **menthol**, namely:

- (a) **Dehydrogenation:** **Menthol** first on dehydration yields two isomeric forms of *p*-menthane, which on subsequent dehydrogenation gives rise to ***p*-cymene** as follows:



- (b) **Reduction:** **Menthol** on reduction with hydroiodic acid yields ***p*-menthane** as under:



Chemical Tests

- When 10 mg crystals menthol are first dissolved in 4 drops of concentrated sulphuric acid and then a few drops of vanillin sulphuric acid reagent are added it shows an orange yellow colouration that ultimately changes to violet on the addition of a few drops of water.

2. A few crystals of **menthol** are dissolved in glacial acetic acid and to this solution a mixture of 3 drops of H_2SO_4 and 1 drop of HNO_3 are added. It fails to produce either green or bluish green colouration (**Thymol** gives a green colouration).
3. **Menthol** provides a plethora of compounds of diagnostic value for differential identification, for instance: **menthoxy acetate; p-nitrobenzoate; d-camphor sulphonate; acid phthalate; phosphoric acid-complex; and 3,5-dinitrobenzoate.**

Uses

1. It is used profusely in various types of mouth washes, toothpastes and similar oral formulations.
2. It finds its enormous use as a flavouring agent for chewing gums, candies, throat lozenges and also certain mentholated cigarettes.
3. It is mostly used on the mucous membranes or on the skin to serve as a counterirritant, antiseptic and as a mild stimulant at a concentration ranging between 1 to 16%.
4. It is employed in conjunction with other allied substances *e.g.*, **camphor, eucalyptus oil (eucalyptol)** in various pharmaceutical preparations, such as: expectorants, nasal sprays, and inhalants to cause immediate relief from symptoms of nasal congestion, sinusitis and above all bronchitis.
5. **Menthol** at a lower concentration ranging from 0.1 to 1%, when applied to the skin helps in the dilatation of blood vessels affording a cold feeling followed usually by a depression of the sensory cutaneous receptors thereby exhibiting an antipruritic action. Perhaps this could be the reason for its logical inclusion in formulations meant for treating sunburns, minor burns, douche powders, athlete's foot and poison ivy rash.

5.2.1.4 Thymol

Synonyms Thyme camphor; m-Thymol; 3-*p*-Cymenol; 3-Hydroxy-*p*-cymene;

Biological Sources It is obtained from the essential oil of *Thymus vulgaris* L., (Thyme oil); *Monarda punctata* L., (**Horsemint oil**), and *Monarda didyma* L., (**Oswego tea oil**), belonging to family *Labiatae*. It may also be derived from *Carum capticum* Bentham er Hooker, (**Ajowan oil**), belonging to family *Umbelliferae*, and several species of *Ocimum*, for instance: *Ocimum gratissimum* L. (**Tulsi oil**), belonging to family *Labiatae*.


Geographical Source *T. vulgaris* is grown and cultivated abundantly in many parts of Europe, Australia and North Asia.

Preparation **Thymol** may be extracted from **thyme oil** by agitation with dilute aqueous alkali solution (= 5% w/v in water). The aqueous layer is first separated and subsequently made acidic with dilute acid, when **thymol** gets separated as an oily layer floating on the surface that may be recovered either by extraction with ether or by steam distillation.


Another means of obtaining **thymol from thyme oil** is to subject the latter to very low temperature (-25°C) when thymol separates as crystals.

Synthetic Thymol The **thymol** of commerce may be prepared synthetically by anyone of the following routes, namely:


- (a) **From Menthone:** Menthone is first treated with bromine. and then quinoline to produce **thymol**:



- (b) **From m-Cresol:** *m*-Cresol on being treated with isopropanol in the presence of a suitable catalyst yields **thymol**.



- (c) **From Piperitone:** When pipertone, usually obtained from the Australian Eucalyptus oils, is treated with ferric chloride it gives rise to **thymol**.



Description


Colour : Transparent, colourless

Odour : Aromatic thyme—like odour


Taste : Pungent taste

Solubility : In water (1: 1200); in alcohol (1:1), in glycerol (1: 1000); Freely soluble in ether, chloroform, carbon disulphide, benzene and glacial acetic acid; soluble in fixed oil and volatile oil.

Chemical Structure The phenolic OH moiety present in **thymol** enables it to form salts of acetate and carbonate easily which are used as antiseptic and anthelmintic respectively.



Thymol when dissolved in NaOH solution and treated with an I₂-KI solution it forms thymol iodide that finds its use as an anti-infective and antifungal agent.



Chemical Tests

1. **Thymol** when fused with phthalic anhydride develops bright violet red to intense red colouration, and on adding dilute alkali it gives an intense blue colouration due to the formation of thymolphthalein.
2. **Thymol** on being dissolved in concentrated sulphuric acid yields the corresponding thyme-sulphuric acid [C₆H₂(SO₃H)(CH₃).(C₃H₇).OH], which produces a distinct violet colour with ferric chloride solution.
3. An alcoholic solution of **thymol** on being treated with FeCl₃ solution does not produce any colouration.

Note: Carvacrol on identical treatment gives a green colouration.

4. A small crystal of **thymol** is dissolved in 1 ml of glacial acetic acid and to this is added one drop of HNO₃ and six drops of sulphuric acid, when it exhibits a deep bluish green colour.
5. Dissolve 0.1 g of **thymol** in 2 ml of NaOH solution (10% w/v) and heat in a water bath to produce either a clear colourless solution or a pale red solution, that ultimately turns darker in shade on keeping without the separation of oily drops. If the resulting solution is shaken with a few drops of chloroform it gives a violet colouration.
6. **Thymol** forms definite derivatives with various reagents *e.g.*, naphthylurethane derivative (m.p. 160°C); phenylurethane derivative (106-107°C).

Uses




1. It is invariably employed as an antifungal and antibacterial agent.

- It is a vital component in several analgesic and topical antiseptic formulations in low concentrations ranging between 0.1 to 1% in personal health care products.
- It is widely employed in preparation exclusively intended for mouthwashes, gargles, oral preparations and as a local anaesthetic in toothache.

5.2.2 Sesquiterpenoids

The **sesquiterpenoids** are found to be extensively distributed in nature and by all means represent the most abundantly prevailing class of **terpenoids**. A few typical examples are cited below:






S. No.	Name	Biological Source (Family)	Geographical Source	Uses
1.	β-Cadinene	<i>Juniperus communis</i> Linn., (<i>Cupressaceae</i>); Oil of Cade	United States, Europe, Canada, India in Western Himalayas (12,500-14,000 ft)	For scenting soaps
2.	β-Caryophyllene	<i>Syzygium aromaticum</i> (l. Merrill & Perry) (<i>Jambrosia caryophyllus</i> Niedenzu; <i>Eugenia caryophyllata</i> Thumb) (<i>Myrtaceae</i>) Clove oil	Moluccas Islands (volcanic island in Eastern Indonesia), Zanzibar, Tanzania (Pemba Islands), Madagascar, Ceylon, Malaysia, Haiti and Southern India	Antitoothache; Antiseptic in tooth-pastes, mouth washes; As a spice; stimulant aromatic carminative; Perfumery; preparing vanilline (from Eugenol)
3.	Abscisic Acid	Found in sycamore, birch, rose leaves, cabbage, potato, lemon, avocado	Mostly in tropical countries	An essential plant growth and development hormone



Sesquiterpenoids may be classified into *four* major categories namely: **acyclic**, **monocyclic**, **bicyclic** and **tricyclic sesquiterpenoids**, as summarized in Table 5.1.


Sesquiterpenoid Lactones Interestingly, another class of compounds essentially bearing such characteristic features as an α -methylene γ -lactone system; α , β -unsaturated carbonyls, and **epoxides** and obviously chemically distinct from the **sesquiterpenoids** are collectively termed as **sesquiterpenoid lactones**. The specific and vital **biological nucleophilic** e.g.; thiol and amino moieties present in the enzymes, help in the augmentation of faster and reactive approach to receptor sites by these **sesquiterpenoid lactones**. Thus, the overall effect is evidenced by a marked and pronounced biological activities, for instance: modified antimicrobial activity, enhanced antitumour properties.

Table 5.1 Classification of Sesquiterpenoids with Summarized Details


S. No.	Type of Sesquiterpenoids	Name	Biological Source (Family)	Geographic Source	Nos of Double Bonds	Specific Gravity	Structure	Characteristic Features	Uses
I	Acyclic	α -Farnesene, β -Farnesene,	Citronella oil from leaves of <i>Cymbopogon winterianus</i> and <i>C. flexuosus</i> (Steud.) Melabar Lemongrass Wats (<i>Graminaceae</i>)	Ceylon, India (Malabar Coast)	4	0.84	 <p>β-Farnesene α-Farnesene</p>	Ozonolysis cleaves- β farnesene into acetone, formalin, succinic and levulinic acid components	Perfume in toiletaries;
II	Monocyclic	Zingiberene	From rhizomes of <i>Zingiber officinale</i> Roseoe, (<i>Zingiberaceae</i>); Jamaica Ginger	South East Asia, West Indies, Australia, Africa, Jamaica, Taiwan, Mauritius, India	3	0.87-0.98	 <p>Zingiberene</p>	Has 3 double bonds and 2 of which are conjugated Dehydrogenation with S yields cadalene-a naphthalene hydrocarbon	Stomachic Carminative
III A	Bicyclic Naphthalene Derivatives	β -Cadinene	From essential oils of juniper species and cadars (oil of cade) <i>Juniperus communis</i> Linn., (Cuperessaceae)	Canada, USA, Europe, India	2	0.90-0.92	 <p>β-Cadinene</p>	Heating with S it gets transformed into cadalene-a naphthalene hydrocarbon	Scenting soaps
B	Azulene Derivatives	Guaiazulene	From chamomile oil, <i>Achillea millefolium</i> Linn., (<i>Compositae</i>): [Syn: <i>A lanulosa</i> Nutt]	Europe, Africa	2	0.90-0.92	 <p>Guaiazulene</p>	Unsaturated hydrocarbons Catalytic hydrogenation yields deahydroazulenes	Anti inflammatory
IV	Tricyclic	α -Santalene	Sandalwood oil from the wood of <i>Santalum album</i> Linn., (<i>Santalaceae</i>)	India, Malaysia	1	0.91-0.935	 <p>α-Santalene</p>	A tricyclic saturated hydrocarbon with a 6-carbon side chain	Scenting soaps

In general, the **sesquiterpenoid lactones** are classified into *three* major group as summarized below:


S. No.	Class	Name	Biological Source	Geographical Source	Special Features	Uses
1.	Germacranolides	Germacranolide	Leaf of Labrador Tea <i>Geranium macrorrhizum</i> (<i>Geraniaceae</i>)	Europe,	Has a ten membered carbon skeleton ring	—
2.	Eudesmanolides	Eudesmanolide	<i>Magnolia obovata</i> (<i>Magnoliaceae</i>)	Europe North America	Has two fused six membered carbon skeleton ring	Neurotrophic activity
3.	Guaianolides	Guaianolide	<i>Guaicum officinale</i> Linn., (<i>Zygophyllaceae</i>) <i>Packwood Tree</i> ; <i>Brazil Wood</i>	Tropical America	Has a five membered ring fused to a seven membered ring	Antioxidant



Germacranolide



Eudesmanolide



Guaianolide


It would be worthwhile to mention some typical examples of natural products that have the **sesquiterpenoid lactone** functions, namely: **artemisinin**.

5.2.2.1 Artemisinin

Biological Source It is obtained from the leaves and the closed, unexpanded flower heads of *Artemisia annua* Linn., belonging to family *Asteraceae*. This particular herb has been used in the Chinese system of medicine exclusively for the treatment of malaria since more than one thousand years.

Geographical Source The plant grows abundantly in China.

Chemical Structure Though the herb was used for its wonderful proven therapeutic efficacy for more than a decade centuries, but its active principal **artemisinin** was isolated and identified in 1972.



Artemisinin

It has been established experimentally that the presence of an internal peroxide linkage strategically located in the seven membered ring is an absolute necessity for it to exert the unique antimalarial property.

Modifications in Structure On account of the poor water solubility of **artemisinin** an attempt was made to improve either its water solubility or its lipid solubility. In the former instance, *Sodium artesunate i.e.*, the sodium salt of its hemisuccinate ester was developed; while in the latter instance, *Artemether i.e.*, its corresponding methyl ether analogue was produced. Evidently, **sodium artesunate** is employed for intravenous injections and artemether is used as a potent long acting drug.

Uses

1. The drug and its derivatives are used as fast acting blood schizontocides in the control and management of malarial fever caused by *plasmodium vivax* strain.
2. These drugs are found to be active against both chloroquine resistant and chloroquine sensitive strains of *Plasmodium falciparum*.
3. These drugs are found to show extremely encouraging therapeutic effects specifically in the treatment of *Cerebral malaria* by virtue of their significant rapid clearance of the prevailing parasites when compared to either chloroquine or quinine (synthetic antimalarials)

Note: The chances of recurrence is quite substantial by the treatment of artemisinin and its derivatives; therefore, it is always necessary to adopt the course of a combination therapy employing other antimalarials.


5.2.2.2 Parthenolide

Biological Source It is obtained from the leaves of *Tanacetum parthenium* (L.) Schultz-Bip, belonging to family *Asteraceae*. It is commonly known as *feverfew* and has been employed for centuries as an effective *febrifuge* (antipyretic) which perhaps suggested the original nomenclature.

It is also obtained from *Chrysanthemum parthenium* (L.) Bernh. Family *Compositae*; and *Magnolia grandiflora* (L.) family *Magnoliaceae*.

Geographical Source The plant *M. grandiflora* is a native of North America and also cultivated in Indian gardens.

Chemical Structure Parthenolide is a **sesquiterpenoid lactone** having the following structure with the chemical name 4, 5 α -epoxy-6 β -hydroxy germacra-1 (10), 11(13)-dien-12-oic acid γ -lactone.



It has an additional epoxide bridge between 4-and 5 α -positions.

Uses

1. It is found to act as a **serotonin antagonist** thereby causing an inhibition of the release of serotonin from blood platelets.


- Based on the findings conducted by an elaborated double blind placebo-controlled clinical trials have established that the drug is significantly effective in the prophylaxis of migraine by reducing considerably the severity as well as the frequency of the pain due to headache.
- A normal dose of 125 mg per day of good quality dried leaves either in the form of tablets or capsules are used in the therapeutic practice as an antipyretic or febrifuge.

5.2.2.3 *Matricarin*

Biological Source It is obtained from the dried flower heads of *Matricaria chamomilla* L., and *Artemisia tilesii* Ledeb, belonging to family *Compositae*. It is also found in *Matricaria recutita* Linne., family *Asteraceae* which represent the drug usually termed as **German Chamomile**. Besides, an allied plant source *Chamaemelum nobilc* Linne, normally known as **Roman Chamomile** also comprises of identical components and used alike.


Geographical Source In general, the above two **chamomiles** are cultivated abundantly in various parts of Europe.

Chemical Structure Its chemical name is 8 α -acetoxy-4 α -hydroxyguaia-1(10), 2-dien-12, 6 α -olide.




Uses

- Chamomile** has acclaimed to be the most popular 'herbal tea' in the United States because of its definite anti-inflammatory and antispasmodic therapeutic properties.
- The volatile oil of *M. chamomilla* contains the *sesquiterpenoid* (-)- α -Bisabolol (**Bisabolane**) which exerts anti-inflammatory activity.
- An infusin (tea) when consumed over a long span results into a cumulative overall positive effect which certainly justifies its age-old usage as an unique home-remedy and healthy beverage not only in Europe but also in the United States.



5.2.3 Diterpenoids

Generally, **diterpenoid** represent a broad class of non-volatile C₂₀ compounds that have been essentially obtained from **geranyl pyrophosphate**.



It has been observed that they mostly originate from the plant or fungal sources, but they are invariably formed by certain insects as well as **marine organisms**.

Characteristic Features Following are some of the characteristic features of **diterpenoids**, namely:

- (a) Most of them are carboxylic compounds having upto five aromatic rings.
- (b) Certain members of this class are acyclic compounds,
- (c) Mostly occur as hydrocarbons or highly oxygenated compounds based on their degree of oxidation,
- (d) Invariably isolated as optically active solids which may occur as the antipodal stereochemical configurations and the normal configurations; the former ones are called the *ent* series, while the latter ones possess and A/B ring fusion and are related to the steroid stereochemically.

A few typical examples of **diterpenoids** shall be discussed in the sections that follow.

5.2.3.1 Colforsin


Synonyms Forskolin; Beforsin (obsolete)

Biological Source It is obtained from the root of *Coleus forskohlii*, Briq., belonging to family *Labiatae*. The word *coleus* has been derived from the Greek *coleus* equivalent to sheath *i.e.*; the natural formation of the fused filaments of the flower which form a staminal sheath around the style; and the word **forskohlii** is due to the honour bestowed upon the Finnish botanist Forskal.

Geographical Source The plant is extensively distributed within the subtropical to temperate climatic zones on the hilly regions of Burma, Africa, Nepal, Ceylon and Thailand. It is also found on the dry, sunny slopes of hills at an attitude ranging between 300 to 1800 meters. It is cultivated in India.

Chemical Structure It was first discovered in India during a general screening studies of **potential medicinal plants***. Its chemical name is 7 β -acetoxy-8, 13-1 α , 6 β , 9 α -trihydroxylate-14 en-11-one.

* Bhat, S.C. *et al.*, *Tetrahedron Letters*, 1669, 1977



Uses


1. It is used in the purification of adenylate cyclase; and as a result it serves as a vital research tool in cyclic AMP-related investigations.
2. It also finds enormous use in glaucoma and hypertension.
3. It possesses significant therapeutic potential in diseases like: congestive cardiomyopathy and bronchial asthma wherein the excessive long term usage of β -adrenergic agonist drugs (*e.g.*; **propranolol, labetalol**) ultimately results into the desensitization of the receptors and a loss of drug efficacy.

5.2.3.2 Ginkgolide-B

Biological Source It is obtained from the root bark and leaves of *Ginkgo biloba* L., belonging to family *Ginkgoaceae*.

Geographical Source It is cultivated in the south eastern United States. The priests in China and Japan have confined this specimen to their temple grounds. It is a dioecious tree attaining a maximum height of 30 meters and has been cited in literatures as a living fossil that still survived unchanged in the region of eastern Asia since 200 million years.

Chemical Structure **Ginkgolide-B** is the most active member of the family significant therapeutic efficacy in the treatment of severe sepsis, whereas the corresponding A and C analogues are devoid of such activities.



Uses

1. The standardized dehydrated acetone–water extract of the dried leaves equivalent to 6% **terpenoids** and 24% **flavone glycosides** is sold commercially in Europe as an approved drug to enhance blood fluidity and circulation.

2. In the United States only the tablets containing 40mg of the Ginkgo is allowed to be sold as a dietary supplement.
3. **Ginkgolides A,B, C and M** have been shown to check the platelet activating factor (PAF) thereby preventing the bronchoconstriction, hypotension, cutaneous vasodilatation and finally the release of inflammatory compounds.

5.2.3.3 *Taxol*

Synonym Paclitaxel; Taxol A; NSC – 125973.

Biological Source It is obtained from the bark of the **Pacific Yew tree**, *Taxus brevifolia* Nutt belonging to the family *Taxaceae*.

Geographical Source The plant is a native to the northwest United States. It is a small, not so growing evergreen tree.

Preparation Keeping in view the paucity of the drug it look quite sometime to isolate **taxol** and establish its chemical structure. The very complexity of its chemistry has more or less turned its total synthesis into a not so viable and feasible economic exercise. However, an attempt is being made to enhance its availability through the semisynthetic route whereby the **taxol precursors** are usually obtained by extraction from the needles of largely available species of *Taxus*.

Example The chemical component, 10-descetylbaaccatin III, isolated from the needles of *Taxus baccata* Linn., may be conveniently converted to **taxol** via simple synthetic route.

Note: **The needles, in comparison to the bark, may be harvested without causing any injury to the plant whatsoever, and thus provides a rather more easily renewable plant source for the drug.**

Chemical Structure The chemical structure is provided in Chapter 1 (Table 1.1).

Characteristic Features **Taxol** has the following characteristic features, namely:

- (a) It has a taxane ring system,
- (b) It has a four membered octane ring
- (c) An ester side chain at C-13 of the taxane ring is a prime requirement for taxol's cytotoxic activity, and
- (d) The presence of an accessible hydroxyl moiety at C-2 of the ester side chain renders an appreciable enhancement of the cytotoxic activity.


Uses

1. **Taxol** is primarily employed in the treatment and management of *metastatic carcinoma of the ovarian glands* after the failure of follow-up chemotherapy.
2. It is also used in the treatment of breast cancer usually after the observed failure of combination chemotherapy for metastatic disease.
3. Because of its hydrophobic nature the injectable concentrate of taxol formulation meant for intravenous infusion is normally solubilized duly in polyoxyethylated castor oil. However, before injection it should be appropriately diluted in normal saline or dextrose solution or combination thereof.


5.2.4 Triterpenoids

Triterpenoids, generally are obtained by biogenesis from six isoprene units, They are found to share commonly the acyclic precursor **squalene** (C₃₀). Based on the various possible modes, whereby ring closure in squalene takes place may ultimately give rise to a large number of triterpenoids having a variety of skeleton structures. In actual practice, more than 4000 naturally occurring triterpenoids have been isolated and identified, and over 40 varying skeleton types have been established.

The **triterpenoids** may be categorized into two major groups, namely: the tetracyclic and the pentacyclic compounds: the former ones of the steroidal types with C-27 carbon atoms present in the skeleton while the latter are of the triterpenoid types with C-30 carbon atoms as shown below:



(Steroidal Triterpenoid)
[C-27]



(Pentacyclic Triterpenoid)
[C-30]

However, the steroidal types (C-27) and the triterpenoid types (C-30) may be distinguished by virtue of the fact that the former yields Diel's hydrocarbon on dehydrogenation with Se at 360°C, while the latter gives either naphthalene or pinene end products.

Interestingly, both these types of compounds may easily combine with sugar moieties at C-3 position to yield the corresponding **glycosides**. Nevertheless, the **free triterpenoids** are invariably associated with natural latex, resins, or cuticle of plants.

A few typical examples shall be discussed in the sections that follows:

5.2.4.1 Cucurbitacin-B


A group of **tetracyclic triterpenoids**, usually termed as "**bitter principles of cucurbits**" that essentially possess distinct antineoplastic and anti-gibberellin activity.

Biological Sources **Cucurbitacins** are obtained from a number of species belonging to cucurbitaceous plants known since antiquity due to their useful as well as toxic properties.

It is obtained primarily from most plants belonging to the *Cucurbitaceae* family, namely: *Luffa acutangula* (Linn.) Roxb. (**Ridged or Ribbed gourd**); *Luffa cylindrica* (Linn.) M. Roem (*Luffa aegyptiaca* Mill ex Hook f. (**Dish-cloth gourd, Vegetables sponge, Spongogourd**)), *Luffa echinata* var *longystyla* Clarke (supposed to be a hybrid of *L. graveolens* Roxb. And *L. aegyptiaca* Mill); and *Luffa graveolens* Roxb. It is also found in various species belonging to family viz., *Begoniaceae*, *Cruciferae*, *Datisceae*, *Euphorbiaceae*, and *Scrophulariaceae*.

Geographical Source It is found in Eastern Himalayas, Sikkim, Bihar and Bengal abundantly.

Chemical Structure It has been observed that **cucurbitacin B** and E are the most commonly identified principles of these plant sources.



Cucurbitacin-B

Uses

1. It has antineoplastic and anti-gibberellin activity.
2. The plants have been employed as vermifuges, narcotics, emetics and antimalarials.
3. They have also been implicated in sporadic livestock poisoning in South Africa.


5.2.4.2 Quassin

Synonyms Nigakilactone D; Nortriterpenoid, Quassane.

Biological Source It is one of the bitter constituents of the wood of *Quassia amara* L., belonging to family *Simaroubaceae*, usually known in commerce as *Surinam quassia*. It is also obtained from the dried wood of *Pierasma quassioides* Benn., *Pierasma excelsa* (*Picroena excelsa* or *Aeschrion excelsa*) belonging to family *Simaroubaceae*.

Geographical Source It is found abundantly in Surinam and Jamaica.

Chemical Structure It has a **triterpenoid** structure as given below:



Quassin

Uses

1. Its bitterness threshold is found to be 1: 60,000; and hence used as bitter tonic.
2. It also finds its application as an insecticides and an anthelmintic for the expulsion of threadworms.
3. It is also used as a febrifuge.
4. **Quassin** possesses antifertility activity, thereby inhibiting testosterone secretion of rat Leydig cells.


5.2.4.3 Azadirachtin

It is a **tetranontriterpenoid** obtained from the seeds of the **Neem tree** and the **Chinaberry tree**.

Biological Sources It is obtained from the seeds of *Azadirachta indica* A. Juss., (*Melia azadirachta* L.) and *Melia azedarach* L., belonging to family *Meliaceae*.

Geographical Sources The plant is found abundantly in tropical countries like India, Africa and Burma.

Chemical Structure Its chemical structure was first reported by *Thornton** in 1993, as shown below:



Uses

1. It is a highly active feeding deterrent and growth regulator.
2. It is used experimentally as insect control agent.
3. It helps in insect ecdysis and growth inhibition**.

It is, however, pertinent to mention here that the **triterpenoids** which are exclusively used as drugs are described in the chapter on '**Glycosides**'.

5.2.5 Tetraterpenoids and Carotenoids

A plethora of natures yellow, orange, red and purple colours are mostly by virtue of the presence of **carotenoids**. The essentially consist of an important group of C_{40} **tetraterpenoids**. Invariably, there are two specific regions in a living plant wherein the **biogenesis of carotenoids** usually occur, namely: chloroplasts and chromatophores of bacteria and fungi.

There are two characteristic features that have been observed in such types of naturally occurring compounds, namely:

- (a) Additional isopentenyl moieties ($H_3C-CH=CH-CH_2CH_3$) could be embedded onto the tetraterpenoid backbone to result into the formation of either C_{45} or C_{50} carotenoids, as seen in certain microbes.

Example **Homocarotenoids**, and

* Thornton, M.D., '*Phytochemistry*', **12**, 391, 1973.

** Reimhoed, H, and K.P. Sieber, '*Z. Naturfoesch.*', **36C**, 466, 1981.

(b) Oxidation of C_{40} carotenoids often yields such carotenoids that do possess less than 40 carbon atoms,

Example Apocarotenoids

So far nearly 600 **carotenoids** have been duly isolated and identified from naturally occurring sources, such as: plants, bacteria, fungi and marine organisms. The ones obtained from the marine sources are found most abundantly and usually contain acetylenic moieties ($HC\equiv CH$).

Characteristic Features of Carotenoids Some characteristic features of **carotenoids** are enumerated below:

- (i) Most widely known carotenoids are either simple unsaturated hydrocarbons having the basic lycopene structure or their corresponding oxygenated analogues, usually termed as **Xanthophylls**,
- (ii) Eight isoprene units are found to be joined head to tail in **lycopene** to give it a conjugated system that eventually is responsible for attributing the chromophoric character to the molecule *i.e.*; producing colour, and
- (iii) Cyclization of **lycopene** at both terminals of the molecule yields a bicyclic hydrocarbon commonly known as **β -carotenes**, which occur most abundantly in the higher plants.


Interestingly, both in plants and micro-organisms the **carotenoids** have been observed to serve *three* major roles, namely: *first*, as **photosynthetic pigments**; *secondly*, as **photoprotective agents**; and *thirdly*, as **membrane stabilization substances**. In contrast, **carotenoids** in animals serve as a precursor of vitamin A and other retinoids. Besides, they also act not only as cancer preventive agents but also as **photoprotective agents**. Perhaps the protective characteristic features of carotenoids, in general, may be due to the easy accessibility to various singly oxygen atoms and ample free radicals, collectively checking the oxidation damage to cells and catering as **antioxidants**.

With the advent of various innovative aspects of biotechnology a quantum jump in the availability of carotenoid production is very much on the cards.

5.2.5.1 Vitamin A

Synonyms Retinol, All-trans-retinol; Oleovitamin A; Bioosterol; Lard factor; Vitamin A alcohol; Acon; Afaxin; Agiolan; Atav; Avibon; Avitol; Axerol; Epiteliol; Testanol; Vaflo; Vogan;

It mostly occurs in animals (not in plants) such as: milk fat, fish liver oil. However, the naturally occurring carotenoids are duly converted into **Vitamin A** by the liver. It is mainly extracted from fish liver oils where it invariably occurs in the esterified form.



Vitamin-A

However, **total synthesis of Vitamin A** has been accomplished from **β -ionone** and a propargyl halide.

It gets easily absorbed from the normal intestinal tract to the extent of 80-90% and is subsequently stored in body tissues, mostly in the liver. It has been observed that approximately one third of the Vitamin A consumed is usually stored in the body.

The most abundantly found dietary sources of Vitamin A are namely: fish- liver oils (*e.g.*, cod-liver oil); dairy products (*e.g.*, butter, cream, whole milk powder, cheese), animal organs (*e.g.*, liver, kidney, heart).

Biochemically, **carotene** and **provitamin A** substances *i.e.*, allied **β -carotenoids** undergo cleavage by the presence of **β -carotene oxygenase** in the mucosal cells of the intestine to give rise to **retinal**; and a substantial portion of which is readily gets reduced by NADH to **retinol**.

Uses

1. It is particularly useful in the proper maintenance of vision, growth and tissue differentiation.
2. The vitamin is employed mainly as a prophylactic when there exists an insufficient normal dietary intake
3. It helps in the synthesis of specific **glycoproteins**.


Vitamin A Derivatives There are *two* important derivatives of **vitamin A** which find their usage in therapeutic domain, namely:

(a) **13-*cis* Retinoic Acid** (Synonym Isotretinoin; Isotrex; Accutane; Roaccutane).

Uses

1. It is used in acute recalcitrant cystic acne.
2. It is invariably employed in keratinization disorders of the skin, that are mostly preneoplastic.

(b) **All-*trans* Retinoic Acid** (Synonyms Tretinoin; Vitamin A acid)



All-*trans*-Retinoic Acid

Uses

1. It is generally used for the treatment of acne vulgaris by virtue of the fact that it enhances epidermal cell mitosis and epidermal cell turnover.
2. It is used in several formulations like cream, solution and gel meant for tropical applications.

5.2.6 Volatile Oils (or Essential Oils)

Volatile oils are the odorous and volatile products of various plant and animal species. As they have a tendency to undergo evaporation on being exposed to the air even at an ambient temperature, they are invariably termed as **volatile oils**, **essential oil** or **ethereal oils**. They mostly contribute to the odorous constituents or '*essences*' of the aromatic plants that are used abundantly in enhancing the aroma by seasoning of eatables.

The nature has so meticulously provided specialized secretary structures within the plants which are primarily responsible for the generation of volatile constituents, as shown in the following Table 5.2.

Table 5.2 Specialized Secretary Structures Vs Plant Sources

S.No	Specialized Secretary Structures	Biological Source	Family
1.	Glansular hairs	<i>Lavandula angustifolia</i> (Lavender oil)	<i>Lamiaceae</i>
2.	Oil tubes (or vittae)	<i>Foeniculum vulgare</i> (Fennel) <i>Pimpinella anisum</i> (Aniseed)	<i>Apiaceae</i>
3.	Modified parenchymal cells	<i>Piper nigrum</i> (Black pepper)	<i>Piperaceae</i>
4.	Schizogenous (or lysigenous) passages	<i>Pinus palustris</i> (Pine oil) Citrus limon (Lemon oil)	<i>Rutaceae</i>

These volatile oils are usually formed by two modes namely; *first*, by hydrolysis of some glycosides; and *secondly*, by the protoplasm directly. It has been observed that the volatile oils are present in different parts of a plant as given in Table 5.3.

Table 5.3 Plant Organs Containing Volatile Oils

S.No	Plant Organs	Biological Source (S)	Family
1.	Petals (Rose oil)	<i>Rosa gallica</i> ; <i>R. alba</i> ; <i>R. damascena</i> and <i>R. centifolia</i>	<i>Rosaceae</i>
2.	Flowering Tops (Lavender oil)	<i>Lavandula angustifolia</i>	<i>Lamiaceae</i>
3.	Leaves (Citronella oil)	<i>Cymbopogon winterianus</i> and <i>C. nardus</i> ;	<i>Poaceae</i>
4.	Bark (Cinnamon oil)	<i>Cinnamomum camphora</i>	<i>Lauraceae</i>
5.	Fruit (Caraway oil)	<i>Carum carvi</i>	<i>Apiaceae</i>
6.	Wood (Pine oil)	<i>Pinus palustris</i>	<i>Pinaceae</i>
7.	Buds (Clove oil & Chamomile oil)	<i>Syzygium aromaticum</i> and <i>Matricaria chamomilla</i>	<i>Compositae</i>
8.	Rhizomes (Ginger oil & Calamus oil)	<i>Zingiber officinale</i> , <i>Acorus calamus</i> ;	<i>Zingiberaceae</i> <i>Araceae</i>
9.	Seeds (Gramis of Paradise & Cardamom)	<i>Aframomum melegueta</i> <i>Elletaria cardamomum</i>	<i>Zingiberaceae</i>

Volatile oils differ from the fixed oils in many respects which may be enumerated below:

S.No	Characteristic Features	Volatile oil	Fixed oil
1.	Chemical Constituents	Mostly consist of terpenoids;	Mostly consist of glyceryl esters of fatty acids;
2.	Spot Test	Does not leave any spot on filter paper	Leaves a permanent grease spot on paper
3.	Saponification Test	Not applicable	Saponifies with alkalies.
4.	Rancidity	Not applicable	Becomes rancid on storage
5.	Exposure to air and light	Easily oxidised and undergo resinification	Not applicable
6.	Fragrance	Distinctly marked and specific	Not applicable

In general, it has been observed that a single volatile oil invariably comprises even more than 200 different chemical components, and mostly the trace constituents are solely responsible for attributing its characteristic flavour and odour.

5.2.6.1 Preparation of Volatile Oils

There are in all *four* established methods whereby the preparation of **volatile oils** from various plant sources may be accomplished, namely:

- (i) Direct Steam Distillation,
- (ii) Expression,
- (iii) Extraction and
- (iv) Enzymatic Hydrolysis.

These methods are described in the sections that follows:

5.2.6.1.1 Direct Steam Distillation In case of direct steam distillation, the freshly cut drug is introduced into the distillation flask. The generated steam is made to pass through the drug material as shown in Fig. 5.1, and the volatile oil content along with the steam on being passed through the water condenser is collected in **Florentine Flasks** of the type FLW or FHW depending on whether the resulting oil is lighter than water or heavier than water.




Fig. 5.1 Assembly for Preparation of Volatile Oils by Steam Distillation

The various parts of the assembly for the preparation of **volatile** oils by steam distillation are as follows:

- A = Steam generator (Copper),
- B = Distillation Flask,
- C = Sand bath,
- D = Water condenser,
- E1 = Florentine Flask for **oils lighter than water** (OLW),

E2 = Florentine Flask for **oils heavier than water** (OHW), and
F = Beaker.

Caution The distillation flask is heated initially to start the process of steam distillation. Once the distillation commences the heat of the steam entering the flask not only maintains the high-temperature required but also in removing the volatile components to the water condenser for ultimate collection in the respective Florentine Flask.

In actual practice, however, there are *three different modes of distillation* depending exclusively on the condition of the plant substance, namely:

(a) **Water Distillation:** It is mostly applicable to such plant material which is dried initially in air and the constituents are not degraded by boiling upto 100°C.

Example: Turpentine Oil—In this instance, the crude turpentine oleoresin is added directly into the distillation flask and subsequently subjected to distillation.

(b) **Water and Steam Distillation:** It is often suitable for such plant material, whether fresh or dried, the constituents of which undergo degradation by direct boiling.

Example: Clove Oil, Cinnamon Oil—In this case, the crude drug is first macerated with water for several hours, prior to steam distillation.

(c) **Direct Steam Distillation:** It is invariably applicable to fresh drugs that is loaded with sufficient natural moisture and hence no maceration is required.

Example: Peppermint Oil, Spearmint Oil—In this instance the freshly cut drug is added directly into the distillation flask prior to steam distillation.

5.2.6.1.2 Expression A number of volatile oils mostly undergo decomposition on being subjected to distillation. Likewise, volatile oils found in the rind of the fruit, such as: **orange, lemon and bergamot peel**, are best obtained by extrusion *i.e.*, by the application of pressure. Even on a commercial scale these oils are produced by extrusion so as to preserve the natural fragrance that otherwise get deteriorated by distillation process.

In actual practice, however, the expression may be accomplished by any one of the *four* following processes, namely:

(a) **Sponge Method:** The citrous fruit (*e.g.*, **orange, lemon, grape fruit, bergamot**) is first washed to remove the dirt, and then cut into halves to remove the juice completely. The rind is turned inside out by hand and squeezed when the secretory glands rupture. The oozed volatile oil is collected by means of the sponge and subsequently squeezed in a vessel. The oil floating on the surface is separated.

(b) **Scarification Process (Ecuelle a Piquer): Ecuelle a piquer** is a specially designed apparatus (Fig. 5.2) first introduced on the Revieras in France, which is nothing but a large bowl meant for pricking the outer surface of citrous fruits. It is more or less a large funnel made of copper having its inner layer tinned properly. The inner layer has numerous pointed metal needles just long enough to penetrate the epidermis. The lower stem of the apparatus serve two purposes; *first*, as a receiver for the oil; and *secondly*, as a handle. Now, the freshly washed lemons are placed in the bowl and rotated repeatedly when the oil glands are punctured (scarified) thereby discharging the oil right into the handle. The liquid, thus collected, is transferred to another vessel, where on keeping the clear oil may be decanted and filtered.




Fig. 5.2 Ecuelle a Piquer

- (c) **Raspings Process:** In this process the outer surface of the peel of citrus fruits containing the oil gland is skillfully removed by a grater. The '**raspings**' are now placed in horsehair bags and pressed strongly so as to ooze out the oil stored in the oil glands. Initially, the liquid has a turbid appearance but on allowing it to stand the oil separates out which may be decanted and filtered subsequently.
- (d) **Mechanical Process:** A substantial quantum of volatile oil across the globe is now prepared by various mechanical means solely based on the above principles. However, the use of heavy duty centrifugal devices may also be incorporated so as to ease the separation of oil/water emulsions invariably formed. It is pertinent to mention here that with the advent of modern mechanical devices the oil out put has increased appreciably and the older methods have only remained for the sake of history.

5.2.6.1.3 Extraction The extraction process is particularly useful for such plant sources which either contain very small amount of volatile oils or the oil contents are extremely susceptible to decomposition by the exposure to steam. In such cases the recovery of volatile oils is not commercially feasible.

Examples: Volatile oils obtained from various flowers like **Jasmine** (*Jasminum officinale* Linn. Ver *grandiflorum* Bailey: family – *Oleaceae*); **Sweat violet** (*Violaodorata* Linn, family – *Violaceae*); **Gardenia** (*Gardenia lucida* Roxb., family – *Rubiaceae*); **Acacia** (*Acacia farnesiana* Willd., family – *Leguminosae*); **Narcissus** (*Narcissus tazetta* Linn., family – *Amaryllidaceae*); and **Mimosa** (*Mimosa pudia* Linn., family – *Leguminosae*).

In general, the **extraction of volatile oil** from natural sources is carried out by *two* different methods, namely:

- (i) Extraction with volatile solvents *e.g.*, Hexane, Benzene and
- (ii) Extraction with non-volatile solvents *e.g.*, Tallow, Lard, Olive oil

These two extraction processes shall be discussed briefly in the sections that follows:

5.2.6.1.3.1 Extraction with Volatile Solvents The plant material containing the volatile oil is usually extracted with a low boiling volatile solvent, such as n-hexane, benzene, petroleum ether etc., either by adopting the method of hot continuous extraction (Soxhlet extraction) or by percolation. The resulting volatile oil containing solvent is removed under reduced pressure when the volatile oil will remain in the flask.

Advantages There are several advantages of this process, namely:

1. It is possible to maintain a uniform temperature (usually 50°C) during most of these extractions which ultimately ensures the retention of a more intense and natural fragrance which otherwise cannot be achieved by distillation (perhaps due to chemical degradation of constituents).
2. **Floral Concretes:** The ultimate concentrated and purified volatile oils are collectively designated as '**floral concretes**'. In actual practice, these floral concretes represent an admixture of natural odoriferous components of flowers, plant waxes, colour pigments and certain albuminous material. Hence, most of them are solid in consistency and partly soluble in 95% alcohol.

5.2.6.1.3.2 Extraction with Non-Volatile Solvents This process is usually employed for the preparation of the finest brands of perfume oil *i.e.*, the natural flower oils. In this instance, the volatile oil content usually present in the fresh plant sources *eg.*, flower petals, is so scanty that oil removal is not commercially viable by any other methods. *Grassein* Southern France, is the well-known centre for the extraction of **flower volatile oil** in the world.

There are *three* methods that are used for the **extraction of volatile oils** from flowers with non-volatile solvents, namely:

- (i) Enfleurage Method,
- (ii) Pneumatic Method, and
- (iii) Mecceration Method.

These methods would be described briefly as under.

- (a) **Enfleurage Method:** A thick layer of molten lard and tallow (beef fat) is applied on either surfaces of pre-cleaned glass plates that are securedly placed in a covered wooden frame (or the chasis) Each glass plate is liberally sprinkled with fresh flower petals to cover its top surface only. These plates are now stacked one over the other and enclosed in the wooden frame, whereby each layer of the flower shall be enclosed between two layers of the fat. Such batteries of loaded plates are allowed to remain for 24 hours, after which the flowers are removed and recharged with fresh lots. This very process is repeated religiously for several weeks till the fatty layers appear to be fully saturated with the essential oils of the flowers or until a certain desired concentration of it is accomplished.

Example: Jasmine flowers—The whole process lasts nearly seventy days.

The flowers are subsequently removed (**defleurage**) and the fat is separated carefully and stirred with absolute alcohol. The latter will dissolve the volatile oil portion thereby leaving the former undissolved in alcohol. The alcoholic extract is reasonably chilled and filtered to get rid off any traces of residual fat. Three successive extraction procedures are repeated so as to affect the complete recovery of volatile oil and the resulting solution is employed as such in the *perfume industry* and is commonly termed as the '**Tripple Extract**'.

The volatile oil may be recovered from the '**Triple Extract**' by anyone of the following methods, namely: *first*, fractional distillation under vacuum at 0°C; *secondly*, evaporation under vacuum at 0°C; *thirdly*, the alcoholic extract is diluted with water and saturated with NaCl, when the oil will separate with the retention of fresh natural odour.


- (b) **Pneumatic Method:** The basic principle of this method is very much like the '**enfleurage method**'. In this particular instance, the current of warm-air is made to pass through the

flowers, and the subsequent air loaded with suspended volatile oil particles is then routed through a fine spray of molten fat in a closed chamber wherein the volatile oil gets absorbed promptly, and


- (c) **Maceration Method:** The fresh flower petals are gently and carefully heated in molten fat (lard, tallow, or fixed oil), stirred frequently until complete exhaustion takes place. The flowers are then strained, squeezed and the exuded fat is returned to the main bulk of the fat, unless and until a desired concentration is achieved. The volatile oil containing fat is allowed to cool and is recovered by three successive extractions with absolute alcohol.

5.2.6.1.4 Enzymatic Hydrolysis It has been observed that the volatile oil is normally found in plant substances in the form of odourless glycosidal combinations. However, the odoriferous components are liberated free only by hydrolysis of such aforesaid **glycosides**. A few typical examples of such volatile components are given below:


5.2.6.1.4.1 Volatile Oil of Bitter Almond (Benzaldehyde) It is found to be present in the kernels of bitter almond in the form of the glycoside **amygdalin**:




5.2.6.1.4.2 Volatile Oil of Black Mustard The volatile oil component is present is *allyl isothiocyanate* in the form of the naturally occurring glycosides **sinigrin**:



5.2.6.1.4.3 Eugenol It occurs invariably in the form of glycosides combination as **gein** present in the roots of *Geum urbanum* Linn. belonging to family *Rosaceae*.



5.2.6.1.4.4 *Methyl Salicylate* It is found to occur in the form of glycosidal combination as **gaultherin** (*Synonym: monotropin or monotropitiside*) in the leaves of *Gaultheria procumbens* Linn. belonging to family *Ericaceae*. The glycosides **gaultherin** undergoes hydrolysis in the presence of the enzyme *gaultherase* to yield the aglycone methyl salicylate and the corresponding sugars glucose and xylose.



5.2.6.2 Quantitative Determination of Volatile Oil in a Plant Material

The necessity to determine the volatile oil quantitatively from a plant material is mostly accomplished by a specially designed apparatus, which helps in ascertaining the raw material to be employed for commercial production. Such a determination is also extremely helpful in establishing and appraising the quality of spices and **oleoresins**.

Clavenger devised an apparatus to determine the volatile oils which essentially has several advantages, namely: (a) **Compactness in size**, (b) **Cohobation of distillation waters**, and (c) **Reasonably accurate estimation of volatile oil content by employing relatively smaller quantum of the raw material**. However, the apparatus also possesses an additional merit for *steam-rectification* of small quantities of essential oils.

Apparatus It consists of a round bottomed flask of varying capacity from 1 L to 2 L which is provided with a hanging type heating mantle and regulator. The mouth of the round bottomed flask is connected with a specifically designed trap to collect the volatile oil which could be either heavier than water or lighter than water as shown in Figure 5.3.

The various vital components of the apparatus for the **quantitative estimation of volatile oils** are as follows:

- A = RB-flask,
- B = Heating Mantle (Hanging type),
- C = Drug and water
- D = Bend insulated with asbestos/ cotton pad
- E = Water condenser
- F = Inlet – for water
- G = Outlet – for water
- H = Volatile oil collected in a graduated stem
- I = Excess water reintroduced in round bottomed flask

Methodology A known weight of the drug either as such powdered or cut into small pieces is introduced into the round bottomed corning flask (1 L or 2 L capacity) along with a distillation




Fig. 5.3 Apparatus for Quantitative Estimation of Volatile Oils

medium which often is fresh water or a mixture of water and glycerin. The quantity of this medium is usually 3 to 6 times the weight of the drug substance. The distillation is done for about 5-6 hours. The distillate is collected in a specially designed trap (or receiver). The stem of which is graduated upto 5 ml with each ml mark is subdivided into 1/10 the ml. In case, the volatile oil is heavier than water the trap on the left hand side of Fig. 5.3 is used; and when the volatile oil is lighter than water the trap on the right hand side is employed.

5.2.6.3 Physical Characteristics of Volatile Oils

It is a well known fact that the volatile oils usually differ from each other with regard to their chemical constitutions. However, they invariably possess a number of physical characteristics as stated below, namely:

- (a) **Odour:** Most volatile oils do possess very pleasant and characteristic odour which vary considerably from one specimen to another.

Detection When a drop of the volatile oil is soaked on a filter paper, an expert may judge its quality and genuinity and may also differentiate between the authentic pure sample from the adulterated one by their individual odours.

- (b) **Nature:** In general, the volatile oils are mobile liquids at ordinary temperatures. However, there are a few exceptions, for instance:
- (i) **Anise Oil:** It solidifies at 15°C and melts at 17°C,
 - (ii) **Rose Oil:** It solidifies at 17°C and melts at 19°C, and

(iii) **Oil of Mentha and Oil of Thyme:** They distinctly deposit a solid substance upon chilling *i.e.*, menthol and thymol respectively, and leaving behind a liquid portion as a 'mother liquor'. The former is termed as "**Stearoptene**" and the latter is known as "**Oleoptene**".

(c) **Volatility:** The essential oils are mostly volatile completely, with the exception of a few of them *e.g.*, '**oil of lemon**', '**oil of orange**', that contain also an additional non-volatile substance of gummy nature. Both the volatile and their pure components do possess high vapour pressures, and hence evaporate completely and rapidly when exposed to atmosphere.

Detection Volatile oils do not leave a stain when soaked on a piece of filter paper, whereas a fixed oil does. Thus, it also checks its adulteration.

(d) **Colour:** Invariably, the colour of freshly obtained volatile oils are more or less colourless, but on prolonged storage they usually undergo both oxidation and resinification thereby rendering it dark in colour. The darkened volatile oil can be redistilled to obtain once again the colourless sample.

Prevention The volatile oils must be stored in a cool, dry place and preferably filled upto the brim in amber glass bottle having an airtight stopper.

(e) **Refractive Index:** The refractive index of volatile oils vary from 1.42 to 1.61. They are mostly characterized by high refractive indices.

Detection The pure authentic volatile oils have definite refractive index as specified in official compendia, whereas the adulterated oils will show different values.

(f) **Optical Rotation:** A large number of volatile oils exhibit optical activity by virtue of the chemical constitution of the oil(s) or its constitution. It gives some vital informations with regard to the source and authenticity of the oil sample, namely:

(i) Both optical rotation and specific rotation offer a fairly dependable and reliable clue whether the volatile oil is either genuine or adulterated,

(ii) It also establishes the source and variety of the volatile oil, for instance: **American oil of Turpentine is dextro-**, whereas the French oil of **Turpentine is levo-**, and

(iii) It ascertains whether the chemical constituent is either isolated from the volatile oil or obtained synthetically, for example: **Menthol** isolated from **peppermint oil** is exclusively *levo*-rotatory, whereas the **synthetic menthol** could be either *racemic* or *levo*. Likewise, the natural camphor is *dextro*, whereas the synthetic one could be either *racemic* or *levo*.

(g) **Specific Gravity:** The specific gravity of volatile oils ranges between 0.8 to 1.17. Interestingly, the volatile oil listed in various official compendia are lighter than water (*i.e.*, specific gravity less than 1), such as: **Oil of Anise** d_{25}^{25} 0.978-0.988; **Oil of Balm** d_{15}^{15} 0.89-0.925; **Oil of Basil** d_{20}^{20} 0.905-0.930; **Oil of Bergamot** d_{25}^{25} 0.875-0.880 etc. In contrast, there are certain volatile oils whose specific gravity is more than 1 *i.e.*, these are heavier than water, for instance: **Oil of Cherry Laurel** d_{20}^{20} 1.054-1.066; **Oil of Cinnamon** d_{25}^{25} 1.045-1.063; **Oil of Clove** d_{25}^{25} 1.038-1.060; **Oil of Garlic** d_{15}^{15} 1.055-1.098; **Oil of Parsley** d_{15}^{15} 1.040-1.100; **Oil of Sassafras** d_{25}^{25} 1.065-1.077 etc.

- (h) **Solubility:** The majority of volatile oils are immiscible with water, but are soluble in absolute alcohol and several other organic solvents *e.g.*, ether, chloroform, carbon disulphide, acetone, hexane, ethyl acetate etc.

Exceptions

- (i) **Oil of Rose** gives a turbid solution due to the very presence of paraffin hydrocarbons that are very sparingly soluble in alcohol, and
 (ii) Many a times certain volatile oils on being dissolved in organic solvents render them turbid due to the presence of traces of moisture which may be eliminated by treating the volatile oil with a small amount of powdered anhydrous sodium sulphate crystals.

In addition to the above stated physical parameters there are certain other equally valuable and important characteristic data for the identification as well as detection of adulteration in a sample of volatile oil, namely: **boiling range, flash point, evaporation residue, molecular refraction** and the like.

5.2.6.4 Chemical Characteristics of Volatile Oils


It has been observed that plethora of volatile oils are found to be more or less '*complex mixtures*' essentially comprising of different class of chemical constituents. Therefore, they are found to vary widely in the chemical composition and *vis-à-vis* their therapeutic applications.

However, there are a few exceptions to the above observation wherein only one chemical entity is solely present in the naturally occurring volatile oil namely:

- (a) **Oil of Bitter Almond**—contains *benzaldehyde* exclusively, and
 (b) **Oil of Winter Green**—contains *methyl salicylate* exclusively

In fact, there are more than 500 different chemical compounds that have been duly isolated, purified and identified in volatile oils over the years with the advent of most sophisticated physico-chemical methods of analysis, such as: **UV-visible spectroscopy, IR-Spectroscopy, NMR – spectrometry, GC – analysis, HPLC–analysis, Mass spectrometry, X-ray diffraction analysis, optical rotary dispersion (ORD) analysis, HPTLC** and the like.


The chemical constituents of volatile oils are recognized as '**terpenes**' that may contain one or several **isoprene units** as shown below:



A few examples of some **terpene hydrocarbons** are summarized below:

Group	Empirical Formula	Isoprene Units	Example
Hemiterpene	C_5H_8	01	Isoprene
Monoterpene			
Acyclic	$C_{10}H_{16}$	02	Myrcene
Monocyclic	$C_{10}H_{16}$	02	Limonene
Bicyclic	$C_{10}H_{16}$	02	α -Pinene
Sesquiterpene	$C_{15}H_{24}$	03	Santalene
Diterpene	$C_{20}H_{32}$	04	Resin of Turpentine
Triterpene	$C_{30}H_{48}$	06	Saponins
Polyterpene	$(C_5H_8)_n$	∞	Rubber

Phenylpropanoids There is another major class of volatile oil constituents that invariably contains a C_6 phenyl ring and an attached C_3 -propane side chain.



5.2.6.5 Classification of Volatile Oils

The most acceptable classification whereby volatile oils and volatile-oil containing drugs may be grouped together are as follows, namely:


- (i) Hydrocarbon volatile oils,
- (ii) Alcohol volatile oils,
- (iii) Aldehyde volatile oils,
- (iv) Ketone volatile oils,
- (v) Phenol volatile oils,
- (vi) Phenolic ether volatile oils,
- (vii) Oxide volatile oils, and
- (viii) Ester volatile oils.

These volatile oils shall be discussed in the sections that follows.

5.2.6.5.1 Hydrocarbon Volatile Oils It has been observed that **terpene hydrocarbons** usually occur in most of the volatile oils obtained from natural sources. They may be further classified into *three* categories, namely:

- (a) Unsaturated acyclic hydrocarbons,
- (b) Aromatic hydrocarbons, and
- (c) Alicyclic hydrocarbons.

5.2.6.5.1A Unsaturated Acyclic Hydrocarbons Two typical examples of chemical constituents belonging to the category of **unsaturated acyclic hydrocarbons** are given below:

(i) β -Myrcene

Chemical Structure It is 7-methyl-3-methylene-1, 6-octadiene.

Occurance It is found in several essential oils, such as: **Oil of Bay** (or Myrcia oil) – *Myrcia acris* (Family : *Myricaceae*); **Oil of Hops** – *Humulus lupulus* Linn., (family: *Moraceae*); and **Oil of Turpentine** – *Pinus logifolia* Roxb., (family: *Pinaceae*).

Isolation The **oil of bay** is treated with sodium hydroxide solution and the remaining undissolved portion which mostly contains myrcene, is repeatedly subjected to fractional distillation under vacuo (it is also obtained by pyrolysis of **β -pinene**).

Characteristic Features It has a pleasant odour. It is lighter than water d_4^{20} 0.794, n_D^{20} 1.4709 and UV_{max} (ethanol): 226 nm (ϵ 16, 100). It is practically insoluble in water, but soluble in alcohol, chloroform, ether and glacial acetic acid.


Identification

- (a) **β -Myrcene** on reduction with sodium and alcohol (absolute) gives rise to **dihydromyrcene** ($C_{10}H_{18}$) which on subsequent bromination yields **tetrabromodihydromyrcene** (mp 88°C), and
- (b) It readily forms addition compounds with α -naphthoquinone (mp 80-81.5°C) and with maleic anhydride (mp 34-35°C).

Use It is used as an intermediate in the manufacture of perfumery chemicals.

(ii) Ocimene (or *trans*- β -Ocimene)

Chemical Structure It is 2,6 dimethyl 2,5,7 octatriene.



Occurrence *trans*- β Ocimene is found in the volatile oil obtained from the leaves of *Ocimum basilium* L., (*Labiatae*); *Baronia dentigeroides* Cheel (*Rutaceae*); *Litsea zeylanica* C & T Nees (*Lauraceae*) and *Homoranthus flavescens* A. Cunn., (*Myrtaceae*).

Isolation The volatile oil obtained from the fresh leaves of *O. basilicum* is treated first with NaOH solution to get rid of the phenolic constituents i.e; *eugenol* present in the range of 30-40% of the oil. The undissolved fraction of the oil is taken up in an appropriate solvent, solvent removed under vacuum and the resulting volatile oil is subjected to fractionation under vacuum so as to obtain the desired main constituent.

Characteristic Features

trans- β -Ocimene : d_4^{20} 0.799; n_D^{20} 1.4893; UV_{max} (ethanol)
232 nm (ϵ 27, 600)

cis- β -Ocimene : d_4^{20} 0.799; n_D^{20} 1.4877; UV_{max} (ethanol)
237.5nm (ϵ 21, 000)

The **Ocimene** (*trans*- or *cis*-) undergo oxidation most readily and with relatively shorter exposure to air to form a yellow resin. However, in an atmosphere free from oxygen ocimene may be preserved unaltered. Its bp ranges between 176-178°C (decomposes).

Identification

- Reduction with sodium and absolute alcohol yields **dihydromyrcene** which on bromination yields **tetrabromodihydromyrcene** (mp 88°C),
- It yields ocimenol – an alcohol on hydration with sulphuric acid (50%) in glacial acetic acid solution,
- Its phenylurethane derivative has mp 72°C, and
- Ocimene upon oxidation with $KMnO_4$ in alkaline solution affects complete degradation to form acids, the lead salts of which has a rhombic crystalline form, whereas the corresponding lead salts of myrcene treated in a similar fashion has a needle form thereby differentiating between **ocimene** and **myrcene** distinctly.

Uses It is used in perfumery.

5.2.6.5.1B *Aromatic Hydrocarbons* A typical example of aromatic hydrocarbon is that of **para-cymene** as detailed below:

(i) para-Cymene

Chemical Structure It is 1-methyl-4 (1-methyl ethyl) benzene (*Syn.*: *Dolcymene*)



para-Cymene

Occurrence It occurs in a number of essential oils, such as: oils of lemon, nutmeg, coriander, cinnamon, sage and thyme. **p-Cymene** has been reported in a number of volatile oils either due to conversion from cyclic terpenes *e.g.*, **pinene** or **terpinene** or from various terpene analogues *e.g.*, citral, carvone, sabinol etc.

Isolation The **p-cymene** fraction obtained by the fractional distillation of volatile oils may be freed from terpenes having identical boiling points by subjecting it to oxidation with cold $KMnO_4$

solution, whereby the former being resistant to the oxidising agent is recovered in its pure form. However, pure *p*-cymene may be prepared from **thymol**.

Characteristic Features *para*-Cymene is a colourless liquid and is found to be inactive optically. Its fragrance resembles to that of the aromatic hydrocarbons closely.

Identification

- Its boiling point is 177.10°C.
- It melts at -67.94°C.
- Its specific density d_4^{20} 0.8573 and d_4^{25} 0.8533.
- Its specific rotation n_D^{20} 1.4909 and n_D^{25} 1.4885.
- p*-Cymene** on oxidation with hot concentrated potassium permanganate solution gives rise to *p*-hydroxy- isopropylbenzoic acid having a melting point 155-156°C.

Uses

- It is employed in the formulation of certain imitation (artificial) essential oils.
- It is used profusely for the preparation of scented soaps and toileteries.
- It also finds its application in the masking of undesirable odours.


5.2.6.5.1 Alicyclic Hydrocarbon The alicyclic hydrocarbons are also termed as '**monoterpenes**' or '**true terpenes**' having the empirical formula $C_{10}H_{16}$. Generally, they may be classified into *two* categories, namely:

- Monocyclic Terpenes, and
- Bicyclic Monoterpenes

These two types of alicyclic hydrocarbons shall be discussed individually with some typical examples as under:


A. Monocyclic Terpenes

Basically, the cyclic terpenes are the extended structural homologues of *cyclohexane* usually derived by varying extent of dehydrogenation. The parent molecule is **methyl-isopropyl cyclohexane** (or ***para*-Menthane**)




The structure of the monocyclic terpenes is expressed with reference to the saturated parent substance '**menthane**' *i.e.*; **hexahydrocymene**. Consequently, the three **isomeric menthanes** viz; *ortho*-, *meta*- and *para*-, theoretically yield the **monocyclic terpenes** respectively.

A number of isomere that have been derived form various degree of dehydrogenation of ***p*-menthane** resulting into the formation of a series of *p*-menthenes are given on page 255:



Interestingly, all the six different species of menthenes have been systematically characterized and identified. However, the most important and abundantly found in various essential oils is Δ^3 menthene, which is observed as a natural constituent of thymol oil and is very closely related to menthol, the main constituent of **pippermint oil**.


Furthermore, the subsequent dehydrogenation of *para*-menthane yields correspondingly the **dihydro-*p*-cymenes**, also termed as *para*-menthadienes.



There are *five* important members belonging to this particular group, namely: **α -terpene**, **β -terpene**, **α -phellandrene**, **β -phellandrene** and **limonene** that are very frequently found in a variety of essential oils.

It is pertinent to mention here that the alicyclic (cyclic) hydrocarbons are invariably found to be more stable than the corresponding acyclic hydrocarbons. Nevertheless, the **monocyclic terpenes** usually undergo isomerization, oxidation and polymerisation very rapidly especially when these are subjected to distillation at atmospheric pressure.


Bearing in mind the diagnostic and therapeutic efficacies of the **monocyclic terpenes** one has to consider the possibility that certain structural configurations like: **geometrical isomerism, stereoisomerism, boat and chair form of isomers** do exist amongst them as depicted below:



A few typical examples of the ‘**monocyclic terpenes**’ are described here under:

(i) Limonene


Chemical Structure It is 1-methyl-4-(1-methyl ethynyl) cyclohexane (Synonym: Cinene, Cajeputene, Kautschin)



Occurrence It occurs in various ethereal oil, specially oils of lemon, orange, caraway, dill and bergamot. It is also found in grapefruit, bitter orange, mandarin, fennel, neroli and celery.

Isolation **d-Limonene** is isolated from the **mandarin peel oil*** (*Citrus reticulata* Blanco, Rutaceae). It may also be isolated from the ethereal oils of lemon, orange, caraway and bergamot either by careful *fractional* distillation under reduced pressure (vauum) or *via* the preparation of adducts, such as: tetrabromides (mp 104-105°C) and the desired hydrocarbon may be regenerated with the help of pure zinc powder and acetic acid.

Characteristic Features It is colourless liquid having a pleasant lemon-like odour. It is practically insoluble in water but miscible with alcohol. **Limonene** when protected from light and air is reasonably stable, otherwise it undergoes oxidation rapidly. When it is heated with mineral acids, the former gets converted to **terpentine** and to some extent **p-cymene**. On the contrary, the action of mineral acids on limonene in cold yields **terpin hydrate** and **terpineol** (alcohols) due to hydration. However, limonene could be regenerated from these alcohols upon heating. The racemic mixture *i.e.* **dl-limonene** is also termed as **dipentene (inactive limonene)**, which on being treated with HCl in the presence of moisture yields **dipentene dihydrochloride** (mp 50-51°C) from methanol. Dehydrogenation of dipentene or limonene with sulphur rapidly yields **p-cymene**. Autoxidation of limonene gives rise to **carveol** and **carvone** which may be observed in poorly stored orange oils by a distinct and marked caraway like odour.



Identification


- (a) **Limonene** on bromination yields tetrabromide derivative which is crystallized from ethyl acetate (mp 104-105°C).
- (b) It forms monohalides with dry HCl or HBr, and the corresponding dihalides with aqueous HCl or HBr.
- (c) Its nitrosochloride derivative** serves as an useful means of identification having mp ranging between 103-104°C.

Uses

- (i) It is used in the manufacture of resins.
- (ii) It is employed as a wetting and dispersing agent.
- (iii) It is widely employed for scenting cosmetics, soaps as well as for flavouring pharmaceutical preparations.


* Kugler Kovate, *Helv Chim Acta*, **46**, 1480, 1963

** Prepared by the action of amyl nitrite and hydrochloric acid

(ii) Sylvestrene**Chemical Structure**

Sylvestrene is generally found to be a mixture of two hydrocarbons (a) and (b) as shown above, wherein one of these forms predominates over the other. It is mostly available as its *d*- and *l*-isomers; whereas the racemic mixture is known as **carvestrene**.

Occurrence It is observed that **sylvestrene** does not occur as a *natural product*, but it is obtained from either of the two **bicyclic monoterpene hydrocarbons**, namely: **3-Carene** and **4-Carene**, during the course of its isolation from the respective dihydrochloride.



Isolation The **turpentine** obtained from *Pinus sylveris* L., may contain as much as **42% of 3-carene**, whereas turpentine from *Pinus longifolia* Roxb. (*Pinaceae*) about **30% of 3-carene**. **Sylvestrene** is isolated in a relatively pure form by preparing the corresponding dihydrochloride.

Characteristic Features It is a colourless oil with an agreeable **limolene** – like odour. It is considered to be one of the most stable **terpenes**. It is neither isomerized by heating nor by the interaction of alcoholic sulphuric acid. On being heated to 250°C it undergoes polymerization.

Identification

- (a) **Sylvestrene** yields the following '*dihalides*' by interaction with solutions of glacial acetic acid-hydrogen halides, for instance: dihydrochloride (mp 72°C); dihydrobromide (mp 72°C); and dihydroiodide (mp 66-67°C).
- (b) The nitrosochloride derivative prepared by the action of amyl nitrite and hydrochloric acid has a mp 107°C.
- (c) It is dextrorotatory.

Uses It does not find any substantial usage either in the perfume or flavour industries.

B. Bicyclic Monoterpenes

The **bicyclic monoterpenes**, as the name suggests essentially possess two cyclic rings which are condensed together. This class of compound is relatively more complex in nature in comparison to the monocyclic species. The second ring system usually contain 2, 3 or 4 C-atoms in common and the rings may be having 3, 4, 5 or 6 membered rings.

The **bicyclic monoterpenes** may be regarded as chemical entities derived from:


- (a) *para-Menthane* – by direct fusion of 2-C atoms and the formation of a simple bridge, and
 (b) *Methylated Cyclohexanes*– by having a bridge with either $-\text{CH}_2-$ or $\text{C}(\text{CH}_3)_2$ – moieties.

In general, the '**bicyclic monoterpenes**' are classified into *five* categories, namely:

- (i) Thujane; (ii) Pinane; (iii) Carane
 (iv) Camphane; and (v) Fenchane.

These *five* distinct categories shall be discussed briefly with typical examples as given below:

I. Thujane




4-Methyl-1-(1-methyl ethyl) bicyclo[3.1.0] hexane.

Eventually, **thujane** is derived from **p-menthane** with direct union between C-2 and C-4. It comprises of a 3-membered and a 6-membered ring. The '*bridge*' in this particular instance does not have the isopropyl group in it.

Example: *Sabinene*

A Sabinene

Chemical Structure



4-Isopropyl-p-methylene bicyclo-2, 4-hexane.


Occurrence It is the major constituent ($\approx 30\%$) in **oil of savin** obtained from young shoots of *Juniperus sabina* L., *Cupressaceae*. It is also present in oils of cardamom and majoram.

Isolation It is obtained by the fractional distillation of **oil of savin** under reduced pressure.


Characteristic Features It is a liquid, lighter than water. It is found to be isomeric with **α -thujane**.

Identification **Sabinene** either on boiling with dilute sulphuric acid or on shaking with cold dilute sulphuric acid yields:

- (i) different forms of terpinene, and
- (ii) 1, 4-terpin.



II. Pinane It is formed from ***p*-menthane** by forming a bridge between C-3 and C-6 positions, thereby resulting into the formation of a 4-membered ring system and a parent 6-membered ring system.



Example α -Pinene.

Chemical Structure 2,6,6-Trimethyl bicyclo[3,1,1] hept-2-ene;

Occurrence It is obtained from oil of turpentine which contains 58-65% **α -pinene** along with 30% **β -pinene**. It is also widely distributed in essential oils belonging to the family *Coniferae*. It has been reported to be present in oils of American peppermint, corriander, cumin and lemon.

Isolation

- (i) It is isolated from the essential oils stated above by the help of chromatographic techniques.
- (ii) Mostly isolated by the fractional distillation from essential oils, preferable under reduced pressure followed by further purification. The fraction collected between 155-165°C is converted to crystalline form of nitrosochloride (treated with amyl nitrite and hydrochloric acid) from which the desired product is liberated by treatment with aniline.

Characteristic Features It is a colourless oil which has a tendency to resinification on exposure to air. The various physical parameters of its isomers are given below:

dl-form : bp₇₆₀ 155-156°C; d₄²⁰ 0.8592; n_D²⁰ 1.4664

d-form : bp₇₆₀ 155-156°C; d₄²⁰ 0.8591; n_D²⁰ 1.4661;

l-form : bp₇₆₀ 155-156°C; d₄²⁰ 0.8590; n_D²⁰ 1.4662.


The **l-form** is usually found in the *French Turpentine Oil*, whereas the **d-form** is found in the *American, German and Swedish Turpentine Oils*.

Identification It may be characterized by—

- Preparation of its nitrosochloride derivative mp 115°C, which is devoid of optical activity,
- Preparation of its hydrochloride derivative mp 132°C, and $[\alpha]_D^{20} - 33.24^\circ \text{C}$ (in alcohol), and
- Preparation of its adduct with malic anhydride (crystalline) mp 169°C.


Uses

- It is abundantly used as a starting material for the large-scale preparation of **synthetic camphor** as given below:



- Turpentine oil** is cooled to -10°C first and then hydrogen chloride gas is passed through it to obtain the pinene hydrochloride. The latter undergoes isomerization to yield bornyl chloride which on treatment with alkali gives rise to borneol. This on oxidation with nitric acid yields pure synthetic camphor.
- It also finds its application in the production of insecticides, solvents, plasticizers, perfume bases and synthetic pine oil.



III. Carane *para*-Menthane with a bridge between C-3 and C-8 results into the formation of **carane**, which comprises of a 3-membered ring imbedded into the 6-membered parent ring as given below:



Example

A 3-Carene

Chemical Structure 3,7,7 Trimethylbicyclo [4,1,0] hept-3-ene (a); 4,7,7-Trimethyl-3 norcarene (b).



Occurrence It is a constituent of **turpentine**. The **turpentine** obtained from *Pinus sylvestris* L., contains upto 42%; turpentine from *Pinus longifolia* Roxb; *Pinaceae* about 30%.


Isolation It is isolated from the **turpentine oil** by the usage of chromatographic techniques.

Characteristic Features It is a sweet and pungent odour essential oil having a more agreeable odour than that of **turpentine**. It is practically insoluble in water but miscible with most fat solvents and oils. The **d-form** possess physical characteristics, e.g.; d_{15}^{15} 0.8668; d_{30}^{30} 0.8586; bp₇₀₅ 168-169°C; $[\alpha]_D^{20} + 17.69$; n_D^{30} 1.468.


Identification The **d-form** gives rise to the nitrosoate derivative (C₁₀H₁₆ N₂O₄), which may be prepared by treating **d-Carene** with amyl nitrile, acetic acid and nitric acid. Its prism decomposes at 147.5°C.

Uses It is used as an antiseptic, carminative, stimulant, stomachic and diuretic.

IV. Camphane It is formed with a direct bondage between C-1 and C-8 in the structure of **p-menthane**. It essentially comprise of *two* five-membered rings besides a six-membered ring.

**Example****A Camphene**

Chemical Structure 2,2, Dimethyl-3-methylenebicyclo-[2,2,1] heptane;



Occurrence It mostly occurs in a large variety of essential oils, for instance:

- (i) Turpentine oil (*levo* and *dextro* forms),
- (ii) Cypress oil (*dextro* form),
- (iii) Camphor oil (*dextro* form in species of *Lauraceae*),
- (iv) Bergamot oil, and
- (v) Oils of Citronella, Neroli, Ginger, and Valerian).

Camphene occurs in a number of species, namely: *Achillea*, *Milefolium*, *Acorus calamus*, *Anethum graveolens*, *Artemisia*, *Cinnamomum*, *Foeniculum vulgare*, *Juniperus*, *Kaempferia galanga*, *Myristica fragans*, *Peumus boldus*, *Pinus ellottii*, *Piper nigrum*, *Pistacia lentiscus*, *Rosamarins officinalis*, *Satureja*, *Schinus molle*, *Thymus*, *a* and *Valeriana officinalis*.

Isolation **Camphene** is isolated by the chromatographic techniques from rectified turpentine oil.

Characteristic Features **Camphene** obtained from alcohol found in cubic crystals (*dl*-form) having an insipid odour.

dl-form: mp 51 to 52°C; bp₇₆₀- 158.5 to 159.5°C; d₄⁵⁴ 0.8422; n_D⁵⁴ 1.45514.

Solubility Soluble in ether, dioxane, cyclohexane, cyclohexene and chloroform. Practically insoluble in water and moderately soluble in alcohol.

d-form : mp 52°C; $[\alpha]_D^{17} + 103.5^\circ$; (C=9.67 in ether); $d_4^{50} 0.8486$; $n_D^{50} 1.4605$;

l-form : mp 52°C; $[\alpha]_D^{21} - 119.11^\circ$; (C=2.33 in benzene); $d_4^{54} 0.8422$; $n_D^{40} 1.4620$.

Identification It forms large dodecahedra on being subjected to slow sublimation


Uses

1. As an important constituent of **eucalyptus oil** which is used as a counter-irritant, antiseptic and expectorant.


V. Fenchane It is a trimethyl cyclohexane with a methylene ($-\text{CH}_2-$) bridge. It consists of *two* five-membered and a six-membered ring.

Example *d*-Fenchone

Chemical Structure



Fenchane



d-Fenchone

(1S) – 1,3,3,-Trimethylbicyclo [2,2,1]-heptan-2-one.

Occurrence It occurs in fennel oil and in the essential oil of *Lavondula stochas* L., *Libitatae*.

Isolation It is isolated from the **fennel oil** by column chromatography which mostly contains this ketone to the extent of 20%.

Characteristic Features It is a colourless oily liquid having a camphor like odour. It attributes the bitter taste to the drug. It is very soluble in absolute alcohol and ether; but practically insoluble in water.

$D_4^{18} 0.948$; mp 6.1°C; bp₇₆₀ 193.5°C; $[\alpha]_D^{20} + 66.9^\circ$; $n_D^{18} 1.4636$.

Identification The pH of its saturated solution is 6.82.

Uses

1. It is employed extensively in foods and in perfumes.
2. It also finds its application as counterirritant.

5.2.6.5.1 Biosynthesis of Monoterpenoids The hypothetic mechanism for the biosynthetic formation of monoterpenoids *viz.*, **myrcane, carane, thuzane, bornane, menthane, pinane** and **fenchane** as individual class has been shown in Fig. 5.4.

5.2.6.5.2 Alcohol Volatile Oils A good number of alcohols occur abundantly in a plethora of volatile oils, which may be judiciously classified into the following heads, namely:

- (a) Acyclic (aliphatic) alcohols,
- (b) Monocyclic (aromatic) alcohols,




Fig. 5.4 Probable Mechanism for Biosynthesis of Various Monoterpenoids.
 (Adapted from 'Pharmacognosy and Pharmacobiotechnology' by Robbers J.E. *et al.*, 1996)

(c) Alicyclic (terpene and sesquiterpene) alcohols.

These *three* distinct categories of ‘**alcohol volatile oils**’ shall be discussed briefly along with certain typical examples from the plant kingdom.

5.2.6.5.2.1 Acyclic (Aliphatic) Alcohols In general, a number of acyclic alcohols, such as: methyl, ethyl, isobutyl, isoamyl, hexyl and other higher alcohols occur widely in volatile oils, but being water soluble they are usually eliminated during steam distillation.


They may be further sub divided into *two* important categories, namely:

- (a) Saturated aliphatic alcohols, and
- (b) Unsaturated aliphatic alcohols.


which shall be discussed along with suitable examples.

5.2.6.5.2.1A Saturated Aliphatic Alcohols Volatile oils normally contain a few saturated monohydroxy alcohols belonging to the paraffin series, most of which are found to be esterified with fatty acids. In the course of steam distillation these esters undergo hydrolysis to yield the lower members of saturated aliphatic alcohols together with the lower fatty acids rarely.

A variety of substances are duly formed on account of the degradation of complex plant constituents *e.g.*, methanol, ethanol (a by product of fermentation due to plant starches), furfural and butanedione (diacetyl), which ultimately are located in the **distillation waters of volatile oils**.



Furfural



Butanedione (Diacetyl)

Isolation of aliphatic alcohols may be accomplished from the volatile oils by fractional distillation, by forming their respective derivatives *e.g.*, *para*-hydroxybenzoates, acid phthalates and calcium chlorides.

The saturated aliphatic alcohols may be identified by the preparation of their respective crystalline derivatives, such as: *para*-nitrobenzoates, 3,5 dinitrobenzoates, phenylurethanes, and naphylurethanes.


The presence of ethanol as an ‘**adultrant**’ in volatile oil may be carried out by treating it with iodine, potassium iodide, sodium hydroxide solution (0.5N) and heating the resulting mixture to give rise to the yellow crystals of iodoform (mp 119°C).

5.2.6.5.2.1B Unsaturated Aliphatic Alcohols The unsaturated aliphatic alcohols frequently occurring in volatile oils are nothing but **terpene-derivatives** wherein the six membered carbon ring is found to be broken at one point only. A few important typical members of this category are as follows:

Examples

1. Geraniol

Chemical Structure 3,7-Dimethyl-2,6-octadien-8-ol; Lemonol.



Geraniol

Occurrence It is an **olefinic terpene alcohol** which constitute the major part of **oil of rose**, **oil of palmarose** (95%), **oil of geranium** (40-50%), **oil of citronella** (30-40%) and also in the essential **oil of lemon grass** etc.

Isolation

Method 1: **Geraniol** may be readily isolated in its pure form from volatile oil fractions by virtue of the fact that it readily forms a distinct crystalline derivative with anhydrous calcium chloride [$2C_{10}H_{18}O \cdot CaCl_2$]. The resulting compound is practically insoluble in organic solvents, such as: chloroform, ether, petroleum ether or benzene and hence can be readily decomposed with pure distilled water into geraniol and calcium chloride. The separated oil thus obtained is rapidly washed with luke-warm water and subjected to steam distillation finally.

Method 2: It may also be isolated and purified, of course, much less conveniently, by forming its solid acid phthalate (mp $47^\circ C$) which yields a crystalline silver salt.

Characteristic Features It is an oily liquid having a marked and pronounced agreeable rose-like odour. However, the odour of its **geometrical isomeride 'Nerol'** is definitely found to be more refreshing than that of geraniol.

Its physical characteristics are as under:


bp₇₅₇ 229-230°C; d_4^{20} 0.8894; n_D^{20} 1.4766; UV max: 190-195 nm (ϵ 18000).

It is practically insoluble in water, but soluble in ether, ethanol. The characteristic features of its corresponding acetate, butyrate and formate analogues are stated below:


Derivative	Mol. Formula	Odour	bp (°C)	d	Solubility
Acetate	$C_{12}H_{20}O_2$	Sweet, fragrant,	242 (decomposes)	$\sim D_{15}^{15}$ 0.9174	Insoluble in water. Very soluble in ethanol; miscible with ether
Butyrate	$C_{14}H_{24}O_2$	Fragrant odour	bp ₁₈ 152	D_4^{17} 0.901	Insoluble in water; soluble in ethanol and ether
Formate	$C_{11}H_{18}O_2$	Odour of roses and of green rose leaves	bp ₁₅ 113-114°	D_4^{20} 0.927	Insoluble in ether and ethanol

Identification


1. It is characterized conveniently by preparing its specific derivatives, for instance: 3-nitrophthalate (mp $109^\circ C$) diphenyl urethane (mp $82.2^\circ C$), and α -naphthylurethane (mp $47-48^\circ C$).
2. When treated with 5% sulphuric acid **geraniol** gives rise to mainly **terpin hydrate** as given below:



3. The interaction of **geraniol** with phosphoric acid and gaseous hydrogen chloride yields diterpene together with other terpenes as depicted below:



4. In the presence of mineral acids **geraniol** undergoes *cyclization* to give rise to **α -terpineol** as given under:




Uses

1. It finds its wide application in a plethora of formulations used as rose scents.
2. It is also employed as insect attractant.
3. It is employed extensively in perfumery *e.g.*; **butyrate** for compounding artificial attar of rose; **formate** as an important constituent of *artificial neroli oil* and of *artificial orange blossom oil*.
4. It is used in soap, cosmetic and flavour industries.

3. Nerol

Chemical Structure *cis*-2,6-Dimethyl-2,6-octadien-8-ol; It is the *cis*-isomer of **geraniol**



Occurrence **Nerol** is found in a number of essential oils, specifically **oil of Neroli** (usually obtained from the fresh and tender flower of orange), **oil petit grain** (normally prepared from not fully matured

fruits of bitter orange) and also in **oil of bergamot** (conventionally prepared from *Citrus auranti* var. *bergamia*).

Isolation Nerol may be isolated from admixture with geraniol in volatile oils by treatment with anhydrous calcium chloride, when the former that does not form any complex with CaCl_2 is separated conveniently by either centrifugation or filtration techniques.

Characteristic Features It is an oily liquid having the odour of sweet rose. It is optically inactive. It has the physical parameters as : bp_{745} 224-225°C; d^{15} 0.8813; UV max: 189-194 nm (ϵ 18000). It is soluble in absolute alcohol.


Identification

1. It is characterized by the preparation of its tetrabromide derivative $\text{C}_{10}\text{H}_{18}\text{Br}_4\text{O}$ (mp 116-118°C).
2. It also gives rise to the diphenylurethane analogue (mp 52-53°C).
3. It forms needles of allophanate ($\text{C}_{12}\text{H}_{20}\text{N}_2\text{O}_3$) (mp 84-86°C) from petroleum ether (40-60°C).

Uses It is used extensively as a base for the manufacture of perfumes.

2. Linalool

Chemical Structure 3,7-Dimethyl-1,6-octadien -3-ol;




Occurrence It is the major constituent of **linaloe oil**. It also occurs in a variety of essential oils, namely: Ceylon Cinnamon (*Cinnamomum verum*), *Artemisia balchanorum*, *Acorus calamus*, *Aloysia triphylla*, *Artemisia dracuncululus*, *Camellia sinensis*, *Cananga odorta*, *Glechoma hederaceae*, *Humulus lupulus*, *Lantana camara*, *Laurus nobilis*, *Lavanchula angustifolia*, *Myrica*, *Myristica fragrans*, *Narcissus tazetta*, *O. imum basilicum*, *Peunus boldus*, *Piper nigrum*, *Prunus armeniaca*, *Robinia pseudoacacia*, *Rosmarinus officinalis*, *Salvia*, *Satureja*, *Syzygium aromaticum*, *Thymus* and *Tilla europaea*.

Isolation It is conveniently isolated from the saponified volatile oil by subjecting it to careful fractional distillation.


Characteristic Features The various typical examples whereby **linalool** reacts with organic acids, anhydrides and inorganic acids are given below:

- (a) **Organic Acids:** It is very sensitive to organic acids and gets rapidly isomerized to **geraniol**. Hence, its esters cannot be obtained in the purest form by ordinary methods.




(b) Inorganic Acids


(i) With Chromic Acid: It undergoes oxidation to yield **Citral**.



(ii) With Formic Acid or Conc. Sulphuric Acid: It undergoes dehydration to yield **α -terpinene** and **dipentene**.



(iii) With 5% (w/w) Sulphuric Acid Solution



(iv) With Glacial Acetic Acid and Acetic Anhydride: Linalool on being heated with glacial acetic acid and acetic anhydride gives rise to a mixture of esters of **geraniol**, **α -terpineol** and **nerol** as follows:


- (c) Aromatic terpene aldehydes, and
 (d) Heterocyclic terpene aldehydes.

These *four* types of **terpene aldehydes** shall be discussed briefly with the help of some typical examples in the sections that follows:


5.2.6.5.3.1 **Aliphatic Terpene Aldehydes** The two important members of this particular class are namely, **citral** and **citronellal**.

A. Citral

Chemical Structure 3,7-Dimethyl-2,6-octadienal; ($C_{10}H_{16}O$); **Citral** from natural sources is a mixture of two geometric isomers **Geranial** and **Neral**.



Geranial (Citral-a)



Neral (Citral-b)

Occurrence It occurs abundantly in the **oil of lemon grass** (75 to 85%) [*Cymbopogon flexuosus* (Ness) stapf. And *Cymbopogon citratus* (DC) stapf. Family : *Graminae*]. It is also present to a limited extent in **oils of verbena, lemon, lime, orange and ginger root**. It is reported to be present in various other species, namely: *Ocimum pilosum* (35%), *Liptospermum citratum*, *Eucalyptus staigeriana* and in the leaf oils of several *Citrus species* etc.


Isolation The rich **citral** containing volatile oils *e.g.*, **lemon grass oil** is thoroughly shaken with 5%(w/v) sodium bisulphite solution for about 25-30 minutes. The resulting crystalline adduct is first separated on a Büchner funnel, and subsequently washed with solvent ether or ethanol to remove the impurities. The crude citral is usually regenerated by decomposing the sodium bisulphite adduct with dilute sodium hydroxide solution carefully. Finally, the pure citral is obtained by distilling the crude citral cautiously under reduced pressure (bp_{26} 92–93°C).

Separation of Geranial (Citral-a) and Neral (Citral-b) Tiemann* observed that **geranial** may be obtained free from **neral** during the process of regeneration from the bisulphate adduct, by taking the strategic advantage of the fact that the crystalline sodium bisulphite adduct of **geranial** is sparingly soluble, whereas the corresponding adduct of **neral** is readily soluble in water.

Tiemann** further observed that **neral** may be isolated from the regular citral (mixture) by shaking it for a short time with alkaline cyanoacetic acid solution ($NC.CH_2.COOH$), when **geranial** reacts with this acid much faster than **neral**. Thus, we may have:


* Tiemann, Semuler, *Ber.* **26**, 2708 (1893).

** Tiemann, Semular, *Ber.* **31**, 3310, 3317 (1898).




Identification

1. By virtue of the presence of two ethylenic and one aldehydic linkage **citral** is very sensitive to oxidizing agents (even exposure to air) to yield **linalool** having an intensified yellow colour.
2. Geraniol on treatment with ammoniacal silver nitrate (**Tollen's Reagent**) gives rise to **geranic acid** ($\text{C}_{19}\text{H}_{15}\text{COOH}$).
3. Hydrogenation of **geraniol** with sodium amalgam in faintly acidic solution yields **citronellal** and **citronellol**.
4. Treatment with potassium bisulphate or diluted sulphuric acid **geraniol** gets converted to **para-cymene** with the loss of a molecule of water.
5. Citral when digested with acetic anhydride and sodium acetate gives rise to its *enolic* form as given below:




6. **Syntheses of Pseudo- and α and β -Ionones:** Citral undergoes condensation with substances containing a *reactive methylene group* as depicted below in sections (a) and (b) respectively:

(a)



Interaction of acetone and **citral** gives rise to the formation of **pseudo-ionone** (or ψ -ionone) with the loss of water molecule.

(b)



In general, the aliphatic ketone **pseudo ionone** undergoes cyclization with the aid of a variety of reagents, namely: sodium acetate, conc. sulphuric acid, formic acid, dilute mineral acids, sodium bisulphate etc., as stated above.

7. Citral may also be identified by the preparation of derivatives such as:


- (i) the 2,4-dinitrophenyl hydrazones: Citral-a mp 108-110°C; and Citral-b mp 96°C;
- (ii) the semicarbazones: Citral-a mp 164°C and Citral-b mp 171°C.

Uses

1. It is used extensively in the synthesis of **vitamin A**, **ionone** and **methylionone**.
2. It is employed as a flavour for fortifying **lemon oil**.
3. It is used widely in perfumery for its distinct citrus effect in lemon and verbena scents, in cologne odours and in perfumes for coloured toilet soaps.

B. Citronellal


Chemical Structure 3,7-Dimethyl-6-octenal; (C₁₀H₁₈O);



Occurrence It is the chief constituent of **citronella oil** (*Cymbopogon winterianus* and *Cymbopogon nardus*, family: *Poaceae*). It is also found in a variety of volatile oils, for instance: lemon, lemon grass, melissa* (*Melissa officinalis*, fam: *Labiatae*) and rose. Due to the presence of one asymmetric C-atom in citronellal it can exist in racemic (*dl*-) form, *d*- and *l*-forms. However, the *d*-form occurs as the chief constituent in the **oil of citronella** obtained from *Eucalyptus citriodora* and other species of *Eucalyptus* (fam: *Myrtaceae*) whereas the *l*-form occurs exclusively in **Java Lemon Oil**.

* Spoon, *Chem. Weekbl.*, 54, 236 (1958).

Isolation It may be conveniently isolated from essential oils by the formation of its crystalline bisulphite adduct. Interestingly, **citronellal** essentially possesses an ethylenic and an aldehyde moiety by virtue of which three different bisulphite adducts are possible theoretically as given below:



Note These three structural analogues have been prepared actually under various experimental parameters. However, the complete decomposition and subsequent regeneration of the desired aldehyde (i.e. Citronellal) may be accomplished easily by treating the 'normal bisulphite adduct' either with dilute mineral acids or with alkali carbonates. Strong alkalies eg., NaOH and KOH must be avoided so as to cause resinification of the aldehyde.

Separation of Citronellal from Citral There are two separate procedures adopted for the separation of **citronellal** from **citral** as discussed here under:

- (a) **Tiemann's Method***: It is based on the fact that citronellal reacts exclusively with a concentrated solution of sodium sulphite and sodium bicarbonate, whereas citral reacts even with a dilute solution.
- (b) **Gildemeister Hoffmann's Method****: It is solely guided by the fact that with *neutral sodium sulphite*, citronellal yields hydrosulphonic derivatives from which the latter cannot be recovered. However, the reaction shall commence only if:
 - (i) right from the beginning a strong current of pure CO₂ is made to pass through the reaction mixture, or
 - (ii) another acid is added gradually to the reaction mixture in sufficient quantities.

The said reaction of **citronellal** with neutral sulphite may afford its separation from citral, which also reacts rapidly with neutral sodium sulphite.


Characteristic Features It is a colourless liquid having a pleasant melissa like odour. It has the following physical characteristic features. bp₁ 47°C; bp₇₆₀ 203-204°C; n_D²⁰ 1.4460; [α]_D²⁵ + 11.50°; d = 0.848-0.856; It is very slightly soluble in water but readily soluble in alcohols.

Under improper storage conditions it slowly undergoes decomposition, polymerization and resinification.

Under direct sun-light it yields a complex mixture which consists of acetone, **β-methyl adipic acid**, **isopulegol** and **menthone**.


* Tiemann, *Ber*, **32**, 834 (1899).

** Gildemeister Hoffmann *die Aetherischen Oele vol IV*, 307-356 (4th ed. 1956)



Strong alkalis, such as: NaOH and KOH, usually resinifies citronellal. Therefore, it is always preferred to make use of relatively weaker alkalis, for instance: Na_2CO_3 , K_2CO_3 , for its regeneration from its corresponding bisulphite adduct.

Reduction of citronellal with sodium amalgam yields **citronellol** *i.e.*, a terpene alcohol, due to catalytic hydrogenation as shown below:



Identification

1. Its semicarbazone derivative has mp 91-92°C.
2. Its dinitrophenyl hydrazone derivative has mp 76.5°C.

Uses

1. For the manufacture of **citronellal** used in perfumery.
2. It is used largely in soap perfumes and as insect repllant.
3. It is employed as artificial citrus flavour.


5.2.6.5.3.2 Cyclic Terpene Aldehydes The **cyclic terpene aldehyde** are of two types *viz.*, monocyclic and bicyclic. A few typical examples from each category shall be discussed in the sections to follow:

A. Monocyclic Terpene Aldehydes

Examples: Perillaldehyde, Safranal and Phellandral

1. Perillaldehyde

Chemical Structure 4-(1-Methylethenyl)-1-cyclohexene-1- carboxaldehyde; ($\text{C}_{10}\text{H}_{14}\text{O}$).



Occurrence It is found in essential oils of *Perilla arguta* Benth; *Labiatae*; *Sium latifolium* L., *Umbelliferae*; Mandarin peel oil *Citrus reticulata* Blanco; *Rutaceae* etc.

Isolation It may be isolated from the respective essential oil, *first* by forming its crystalline sodium bisulphite adduct; and *secondly*, regenerating the desired product with alkali very carefully.

Characteristic Features The physical characteristic of the *d*- and *l*-isomers are as follows:

***d*-form** : Liquid, bp₇₄₅ 273°C; bp₇ 98-100°C; d₄²⁰ 0.953; n_D²⁰ 1.5058; [α]_D²⁰ + 127° (C = 13.1 in CCl₄).

***l*-form** : Liquid, bp₁₀ 104-105°C; d₄²⁰ 0.9645; n_D²⁰ 1.5069; [α]_D²⁰ - 146°.


Identification It forms oxime (C₁₀H₁₅NO) known as: **1-perillaldehyde-α-syn-oxime; perillartine; “perilla sugar”**. Previously, it was commonly referred to as **1-perillaldehyde-α-anti-oxime**. The needles have mp 102°C, UV max (alcohol). 232 nm (ε 21800). It is about 2000 times as sweet as sucrose.

Uses


1. The oxime is used as a sweetening agent in Japan.

2. Safranal

Chemical Structure 2,6,6-Trimethyl-1,3-cyclohexadiene-1-carboxaldehyde; (C₁₀H₁₄O).



Occurrence It is the chemical constituent obtained from the dried stigmas and tops of the styles of *Crocus sativus* (Fam; *Iridaceae*); The fresh drug comprises of **protocrocine** which upon drying undergoes decomposition to yield one mole of **crocin** (a coloured glycoside) and two moles of **picrocrocin** (a colourless **bitter glycoside**). It is the latter that on hydrolysis gives rise to safranal which is solely responsible for attributing the characteristic odour of the drug (saffron). These transformations are as given below:



Isolation The stigmas and tops of the styles of *C. sativus* are first dried under shade and then subjected to hydrolysis in controlled conditions to yield safranal. The resulting product is treated with pure sodium bisulphite to obtain the safranal sodium bisulphite adduct from which the desired product is regenerated by treatment with dilute alkaline solution.


Characteristic Features It is a liquid having a pleasant characteristic odour. It has the following physical properties namely:

bp_{1.0} 70°C (bath temperature); d₄¹⁹ 0.9734; n_D¹⁹ 1.5281. It is freely soluble in organic solvents like: methanol, ethanol, petroleum ether and glacial acetic acid.

Uses It is employed as a flavouring agent in confectionery products.

3. Phellandral


Chemical Structure 4-Isopropyl-1-cyclohexen aldehyde; (C₁₀H₁₅O).



Occurrence It was first and foremost found in the essential oil of *Phellandrium aquaticum*. It also occurs in the essential oil obtained from the flowers of lavender (*Santolina chamaecyparissus* L., family: *Asteraceae*) and several *Eucalyptus species*.

Isolation It is usually isolated through its sparingly soluble crystalline sodium bisulphite compound, from which **phellandral** is generated by treatment with dilute alkaline solution carefully.

Characteristic Features It is an oil having an odour very much reminiscent of cuminaldehyde. It undergoes rapid oxidation either on exposure to air or with silver oxide to give rise to **phellandric acid** (mp 144-145°C)




Identification It may be identified by preparing its corresponding derivative, such as: oxime, semicarbazone and phenyl hydrazine etc.

Uses Due to its resemblance of odour of cuminaldehyde it finds application in perfumery.

B. Bicyclic Terpene Aldehydes

Example: Myrtenal.

Chemical Structure



It is a naturally occurring **oxygenated pinane derivative**.

Occurrence The leaves of boldo contains essential oil to the extent of 2%* (*Peumus boldus* Molina, family: *Monimiaceae*).


Isolation The aldehyde is successfully isolated by preparing its sodium bisulphite adduct first and then regenerating it by treatment with mild alkaline solution cautiously.

Uses

1. The aromatic leaves are frequently used as mild diuretic, especially in liver ailments like jaundice.
2. It is also recommended for urogenital inflammations *e.g.*; gonorrhoea in Latin America

* Bruns, K., and Kohler, M, *Über die Zusammensetzung des boldoblatteröls*, *parf kosm*, **55**, 225, 1975


5.2.6.5.3.3 **Aromatic Terpene Aldehyde** In comparison to aliphatic aldehydes, the aromatic aldehydes invariably play a vital role in essential oils. It has been observed that a major portion of certain volatile oils mainly comprise of aromatic terpene aldehydes, such as : **bitter almond** and **cassia**. A variety of such aromatic aldehydes commonly found in essential oils are, namely: anisaldehyde, benzaldehyde, cinnamaldehyde, cuminaldehyde and salicylaldehyde.



Examples: A few typical examples of aromatic terpene aldehyde are discussed below, for instance: **Cuminaldehyde, Vanillin** etc.

A. Cuminaldehyde

Chemical Structure 4(1-Methylethyl) benzaldehyde.



Occurrence It occurs as a constituent of essential oils present in eucalyptus, myrrh, cassia and cumin. Cumin mainly comprises of the dried ripe fruits of *Cuminum cyminum* Linn., (family: *Umbelliferae*).

Isolation The essential oil obtained from the dried ripe fruits of cumin ranges between 2-4%, the major constituents of which cuminaldehyde (35-60%). The aldehyde may be separated by forming its sodium bisulphite adduct and subsequently regenerating the desired product by treatment with alkaline solution carefully.

Characteristic Features It is a colourless to yellowish, oily liquid. It possesses a strong persistent odour, acrid and burning taste.


The physical characteristic of **cuminaldehyde** are as follows: d_{20}^{20} 0.978; bp_{760} 235-236°C; n_D^{20} 1.5301. It is practically insoluble in water, but freely soluble in ether and ethanol.

Identification It is identified by forming its **thiosemicarbazone** derivative (C₁₁H₁₅N₃S).

Uses It is extensively employed as an adjunct in perfumery.

B. Vanillin

Chemical Structure 4-Hydroxy-3-methoxybenzaldehyde (C₈H₈O₃).




Occurrence It occurs naturally in vanilla (**vanilla bean**) especially found in cured, fully-grown unripe fruit (pods) of *Vanilla planifolia* Andrews, known in commerce as Bourbon or Mexican vanilla. It is also found in the vanilla pods of *Vanilla tahitensis* J.W. More, usually recognised in commerce as *Tahiti vanilla*. Both these species belong to natural order *Orchidaceae*.

The term *vanilla* has been derived from the spanish word *vania*, meaning sheath like pod and *illa*, meaning small; *planifolia* is derived from the Latin word *planus*, meaning flat, and *folium*, meaning leaf; *tahitensis* refers to Tahiti its adopted home.

It also occurs in small quantities in a variety of essential oils e.g., **clove oil; gums and oleoresins** e.g., *benzoin*, *Peru balsam*. Interestingly, plants do not contain vanillin as such, but they exist in the form of *glycosides*, which upon hydrolysis in the presence of enzymes release **vanillin**.


Isolation It may be accomplished by any one of the following *five* methods, namely:

- (a) **From Vanilla Pods: Vanillin** is obtained by the extraction of powdered vanilla pods with ether, by at least three successive extractions, evaporating the combined ethereal fraction (crude vanillin). It is further purified by crystallization of the crude product from ethanol.
- (b) **From Lignin Waste:** More conveniently, **vanillin** is obtained from the lignin waste, a byproduct in the manufacturer of paper pulp. Lignin is a complex polymeric natural material of woody plants and essentially has the following fragment structural unit:




Thus, **lignin** when subjected to oxidation yields **vanillin** in a large extent which ultimately has rendered this process an economically feasible and viable one.

- (c) **From Eugenol:** It may also be prepared from **eugenol** which is a major constituent of **clove oil** as given below:



Eugenol on alkaline treatment undergoes intramolecular rearrangement to give rise to **isoeugenol** which on acetylation with acetic anhydride yields the corresponding acetate derivative. The resulting product on oxidation yields 4-acetoxy-3-methoxy benzaldehyde which upon treatment with HCl yields vanillin.

- (d) **From Guaiacol:** It may be prepared on an industrial scale by the help of **Reimer-Tiemann's reaction**, whereby **o-vanillin** is obtained from guaiacol *i.e.*, **catechol methyl ether** as stated below:



Guaiacol on treatment with sodium hydroxide in the presence of chloroform yields **vanillin** and **o-vanillin**.

- (e) **From Bisulphite Adduct:** **Vanillin** may also be isolated from its bisulphite adduct. The ethereal solution containing vanillin is extracted completely with saturated aqueous sodium bisulphite solution. Usually specialized techniques are adopted to ensure pure isolates from its tautomers and closely related isomers.

Characteristic Features It consists of fine, white to slightly yellow, needle-shaped crystals having an odour and taste quite resembling to that of vanilla fruits. It is usually affected by light. On prolonged heating at 105°C, it decomposes with the formation of non-volatile byproducts. It is soluble in hot water (1 g dissolves in 16 ml of water at 80°C), and in glycerol (~ 20 ml of glycerol per 1g of vanillin). It is freely soluble in ethanol, ether, chloroform, carbon disulphide, glacial acetic acid, pyridine, oils and aqueous solutions of alkali hydroxides.

It has the following physical parameters: d 1.056; mp 80-81°C; bp 285°C.

Identification It may be identified by preparing a number of derivatives, such as: semicarbazone (mp 230°C); dinitrohydrazone (mp 271°C); *para*-nitrophenylhydrazone (mp 227°C); benzoate (mp 75°C) and acetyl derivative (mp 77°C).


Uses

1. It is employed as a pharmaceutical aid (flavour).
2. It is extensively used as a flavouring agent in beverages, confectionery, foods and perfumery.
3. It is used in the manufacture of liqueurs.
4. It has been established that 1 part of vanillin equals 400 parts vanilla pods; and 2.5-3 parts equals 500 parts tincture vanilla.
5. It also finds its use as a reagent in analytical chemistry.

5.2.6.5.3.4 **Heterocyclic Terpene Aldehyde** **Heterocyclic terpene aldehyde** is relatively quite rare as compared to other class of compounds discussed in this context. Example: **Furfural**.

A. Furfural

Chemical Structure 2-Furfuraldehyde (C₅H₄O₂).



Furfural

Occurrence It occurs in the first fraction of a number of essential oils, belonging to the natural order *Pinaceae*. It is also found in **colophony** (*Pinus paulusteric*); **oil of orris rhizome** (*Iris florentina* Linn., Family: *Iridaceae*); **oil of lavender** (*Lavandula officinalis*; family: *Labiatae*); **oil of cinnamon** (*Cinnamomum cassia* Blums; Family: *Lauraceae*); and **clove oil** (*Eugenia Caryophylus*; Family: *Myrtaceae*).

Isolation It may be accomplished in *two* manners, namely:

- (i) *Extraction i.e.*, by washing the first fraction of the oil with water, extracting the aqueous layer with ether, and finally evaporating the ether under reduced pressure.
- (ii) *Addition Compound i.e.*, **furfural** forms an addition compound on being treated with a saturated aqueous solution of sodium bisulphite.

Characteristic Features It is a colourless oily liquid having a peculiar odour, somewhat resembling the odour of benzaldehyde. The physical characteristic features are: d_4^{20} 1.1563; bp_{760} 161.8°C; mp: -36.5°C; volatile in steam; n_D^{20} 1.5261. It is soluble in ethanol and ether; soluble in 11 parts of water.

Identification

1. It gives an intense red colour with aniline acetate.
2. It is also identified by forming its corresponding oxime and phenyl hydrazone derivatives.
3. It reduces Fehling's solution to give red precipitate of cupric oxide.
4. It also reduces **Tollen's Reagent** (*i.e.*, ammoniacal silver nitrate solution) to give silver mirror.



Uses

1. It is used extensively in the manufacture of furfural-phenol plastics, for instance: *Durite*.
2. It is employed in solvent-refining of petroleum oils.
3. It makes its use as a solvent for nitrated cotton, gums and cellulose acetate.
4. It finds its application for accelerating vulcanization.
5. It is used as insecticide, germicide and fungicide.
6. It is employed in the manufacture of varnishes.
7. It is commonly used as a reagent in analytical chemistry.

5.2.6.5.4 Ketone Volatile Oils The ketones that invariably occur in volatile oils may be classified in the following *two* categories, namely:

- (i) Aliphatic ketones, and
- (ii) Aromatic Ketones.

5.2.6.5.4.1 Aliphatic Ketones **Aliphatic ketones** do not occur abundantly in volatile oils. However, the relatively lower members of this group originate most probably by virtue of the decomposition of rather more complex compounds during the process of steam distillation. Two such species, for instance *acetone* and *diacetyl* are commonly found in the '**oils of cohobation**' (or the distillation waters) accomplished by redistillation (**cohobation**) of the distillation waters.



It has been observed that acetone and diacetyl are frequently accompanied by methanol and furfural.

5.2.6.5.4.2 Aromatic Ketones There are also termed as '**cyclic terpene ketones**'. Generally, the aromatic ketones are classified into two categories, namely:


- (i) Monocyclic terpene ketones, and
- (ii) Bicyclic terpene ketones.

These two distinct categories shall be discussed separately in the sections that follows:

A. Monocyclic Terpene Ketones A few typical examples of this specific class of ketones are: ***l*-menthone**; **carvone**.

A.1 *l*-Menthone

Chemical Structure (2*S*-*trans*)-5-Methyl-2-(1-methylethyl) cyclohexanone. As it has two asymmetric carbon atoms (*i.e.*, chiral centres) it can exist in two pairs of enantiomorphs or four optically active isomers (*d*-; *l*-; ***dl*-menthone** and **isomenthone**).



Occurrence It is found in a variety of volatile oils, such as: pennyroyal (*Mentha pulegium*, Family: *Lamiaceae*); peppermint (*Mentha piperita* Linn., Family: *Labiatae*); geranium (*Geranium maculatum* L., Family: *Geraniaceae*); and buchu (*Barsoma betulina* (Berg.) Bartl. and Wendel, Family: *Rutaceae*).

Isolation *l*-Menthone usually occurs in association with **isomenthone**. The former gets solidified at -6°C whereas the latter at -35°C . In normal practice, the **peppermint oil**, which contains upto 30% of **menthone**, is subjected to its oxime or semicarbazone formation and subsequently the ***l*-menthone** is regenerated by the aid of dilute sulphuric acid.

Characteristic Features It is a bitter liquid having peppermint-like odour. It is slightly soluble in water, whereas freely soluble in organic solvents. It has the following physical characteristics: bp 207°C ; mp -6°C ; d_4^{20} 0.895; n_D^{20} 1.4505; $[\alpha]_D^{20}$ -24.8° .


Identification The isomeric forms of menthones may be characterized by the preparation of specific derivatives, for instance: oximes, semicarbazones etc.

Uses

1. It is used extensively in perfume and flavour compositions.
2. It is also employed in the preparation of artificial essential oils.

A.2 Carvone

Chemical Structure 2-Methyl-5-(1-methylethenyl)-2-cyclohexene-1-one.



Occurrence It occurs in the **mandarin peel oil** (*Citrus reticulata* Blanco., Family: *Rutaceae*); **spearmint oil** upto 70% (*Mentha spicata* or *Mentha cardiaca*, Family: *Lamiaceae*); **gingergrass oil**

(*Zingiber officinale* Roscoe, Family: *Zingiberaceae*); **oil of caraway** upto 50-60% (*Carum carvi* Linn, Family: *Umbelliferae*).

Isolation It may be isolated by the following *two* methods:

Method 1: Formation of Sodium Sulphite Adduct. **Carvone** may be conveniently isolated from essential oils (e.g., spearmint oil, oil of caraway) by virtue of the fact that it (a lactone) readily forms the water soluble salt of a hydrosulphonic acid ($C_{10}H_{16}O_7S_2Na_2$) on treating with a neutral solution of sodium sulphite, whereby the corresponding addition takes place at the both ethylenic linkages. In order to achieve this, the fraction collected between the boiling range 220-235°C in the case of oil of caraway, is shaken with the requisite quantity of a concentrated aqueous solution of sodium sulphite and the sodium hydroxide thus liberated during the course of reaction is neutralized from time to time with a dilute mineral acid (e.g., HCl) very carefully. As soon as the above process is completed fully, the resulting fractions which have not involved in the above cited reaction may be eliminated by extracting the solution with ether successively (at least three times). At the end, the desired product *carvone* can be regenerated by the action of sodium hydroxide and finally distilled off with steam.

Method 2: Formation of Hydrogen Sulphide Adduct. Alternatively, **carvone** may be separated from the volatile oils by the formation of its hydrogen sulphide adduct [$(C_{10}H_{14}O)_2 \cdot H_2S$]. It is easily accomplished by the passage of a current of hydrogen sulphide (H_2S) gas into an ammoniated alcoholic solution of carvone. Ultimately, the pure ketone *i.e.*, **carvone** may be regenerated from the corresponding separated adduct by careful digestion with alkali.

Characteristic Feature It is a colourless liquid having a distinct odour typical of **caraway seed**. The various physical parameters of *d*-, *l*- and *dl*-forms are given below:

Form	bp(°C)	d	n_D^{20}	$[a]_D^{20}$
d-Carvone	230 (At 755 mm atmospheric pressure)	0.965 (d_4^{20})	1.4989	+61.2°
l-Carvone	230-231 (At 763 mm atmospheric pressure)	0.9652 (d_{15}^{15})	1.4988	-62.46°
dl-Carvone	230-231 (At 760 mm atmospheric pressure)	0.9645 (d_{15}^{15})	1.5003	—

It is miscible with ethanol but practically insoluble in water. It congeals at very low temperature.

Identification The different tests for the identification of **carvone** are stated below, namely;


(a) **Bromination** of carvone gives rise to a mixture of crystalline derivatives* having distinct melting points:

d- and *l*-form : mp 120 and 120-122°C

dl-form : mp 112-114°C

* These are probably the dibromo derivatives because the tetrabromoderivatives are liquids.

- (b) **Mono-hydrochloride salt** is formed when it is treated with HCl in acetic acid.
- (c) **Hydrobromide salt** is obtained by treating *d*-carvone with HBr (mp 32°C).
- (d) **Isomerization of Carvone to Carvacrol**. It undergoes isomerization to form carvacrol with a number of dehydrating agents, such as: H₂SO₄; H₃PO₄; NaOH; ZnCl₂.



- (e) It is also characterized by the preparation of several compounds, for instance: oxime, semicarbazone, H₂S-derivative, phenylhydrazone.

Uses

1. It is used as **oil of caraway**.
2. It is also used for flavouring liqueurs.
3. It is used extensively in perfumery and soaps.
4. It is employed for flavouring many types of food products and beverages.
5. It finds its enormous applications in *oral hygiene* preparations *e.g.*, toothpastes, gargles, mouth-washes.
6. It is also used in flavouring pharmaceuticals.

B. Bicyclic Terpene Ketones These class of compounds essentially contain two cyclic ring structures fused with each other along with a ketonic function. The two typical examples of pure chemical entities that belong to this group are **camphor** and **d-fenchone**, which have been duly dealt with earlier in this chapter under '**monoterpenoids**' and '**bicyclic monoterpenes**' respectively.

5.2.6.5.5 Phenol Volatile Oils The important drugs containing phenol volatile oils are, namely: **Clove oil, Myrcia oil (Bay oil), Organum oil, Pinetar, Thyme** etc., In fact, they essentially owe their value in the pharmaceutical domain almost exclusively by virtue of their antiseptic and germicidal properties of their *phenolic constituents*. A good many of them are employed as popular flavouring agents, for instance: **oil of anise, clove** and **sassafras**.

5.2.6.5.5.1 General Methods of Isolation Mostly phenols are weak acids. Hence, they react with dilute alkali solutions (3-5% w/v) to result into the formation of corresponding water-soluble salts known as '*phenolates*'. This specific characteristic property usually offers, a convenient method for carrying out the separation of *phenolic components* from the *non-phenolic ones*. To affect the separation, therefore, the volatile oils or fractions are subjected to treatment with dilute alkaline solutions with vigorous shaking. Once the two layers get separated, the water-soluble salts are decomposed by acidification carefully and the phenols thus generated (or liberated) are isolated either by means of steam-distillation or by extraction with ether.

Note

1. **Thymol** and **carvacrol** may be steam-distilled from alkaline solution without previous acidification.
2. Several phenols may be isolated by chilling the oil as such or its fraction to a very low temperature (-20 to -30°C) whereby these compounds normally separate in crystalline form.

5.2.6.5.5.2 **General Properties of Terpene Phenols** There are several characteristic features of **terpene phenols** which not only help them in their separation but also aid in their identification as stated below:

1. **Bromine Reaction:** Phenols react with bromine evolving the corresponding HBr. The resulting bromides are usually *crystalline* in nature and sparingly water-soluble. Hence, they may be separated easily and identified accordingly.
2. **Reaction with Ferric Chloride:** Phenols react with dilute aqueous solutions of ferric chloride (FeCl_3) (0.1-0.2% w/v) to give rise to intense coloured reactions, which attributes to the *specific colour-tests*.
3. **Formation of Phthaleins:** Several phenols react with phthalic anhydride to form their corresponding phthaleins.
4. Most phenols react with specific reagents, such as: acetic anhydride, benzoyl chloride, phenyl isocyanate and *p*-nitrobenzoyl chloride to give characteristic reaction products that also help in their identifications.

5.2.6.5.5.3 **Classification** The **terpene phenols** are classified into the following categories, namely:


- (i) Monohydric phenols and
- (ii) Dihydric phenols.

The above categories of phenols shall be discussed briefly as under:

5.2.6.5.5.3.1 **Monohydric Phenols** The typical examples of monohydric phenols are: **carvacrol**, **eugenol**, **thymol**.

A. Carvacrol

Chemical Structure 2-Methyl-5-(1-methylethyl)-phenol.



Occurrence It occurs in the essential **oil origanum** (*Origanum vulgare* Linn., Family: *Labiatae*); **oil of thyme** (*Thymus serpyllum* Linn., and *T. vulgaris* Linn., Family: *Labiatae*); **oil of marjoram** and **oil of summer savory**.

Characteristic Features It is a liquid oil having strong thymol odour. Its physical properties are as follows: d_4^{20} 0.976; bp_{760} 237-238°C; $mp \sim ^\circ C$; n_D^{20} 1.52295.

It is practically insoluble in water and freely soluble in alcohol and ether.

Uses

1. It is mostly employed as a disinfectant.
2. It is also used as an anthelmintic (Nematodes).


B. Eugenol Please see section 5.2.6.1.4.3 in this chapter.

C. Thymol Please see section 5.2.1.4 in this chapter.

5.2.6.5.5.3.2 **Dihydric Phenols** The various dihydric phenols found in natural products are namely: **Catechin (catechol)**. A few of these constituents shall be dealt with here under:

A. Catechin (Catechol)

Chemical Structure (2R-trans)-2-(3, 4-Dihydroxyphenyl)-3, 4-dihydro-4H-1-benzopyran-3, 5, 7-triol.



Catechin

Occurrence It is a flavonoid found primarily in higher woody plants as (–)-**catechin** along with (–)-**epicatechin** (*cis*-form). It is also found in catechu (gambir and acacia), mahogany wood etc. Besides, it occurs in a variety of medicinal-plants, such as: *Argimonia eupatoria* L., (*Rosaceae*)-agrimony; *Areca catechu* L., (*Arecaceae*)-areca-nut, betel-nut palm; *Camellia sinensis* (L.) Kuntze (*Theaceae*)-tea; *Catha edulis* Vahl (*Celastaceae*)-khat; *Cola acuminata* (Beauv.) Schott & Endl. (*Sterculaceae*)-kola nuts, cola, guru; *Caratadegus oxyacantha* L. (*Rosaceae*)-hawthorn; *Ephedra gerardiana* Wall. ex Stapf (*Ephedraceae*)-Paskistani ephedra; *Eucalyptus globulus* Labill. (*Myrtales*)-eucalypt, tasmanium bluegum; *Leonurus cardiaca* L., (*Lamiaceae*)-motherwort; *Malus sylvestris* Mill., (*Rosaceae*)-apple; *Paullinia cupana* Kunth. ex H.B.K. (*Sapindaceae*)-guarana, ubano, Brazilian cocoa; *Polygonum aviculare* L., (*Polygonaceae*)-prostrate knotweed; *Rheum officinale* Bail. (*Polygonaceae*)-Chinese rhubarb, Canton rhubarb, Shensi rhubarb; *Santolina chamaecyparissus* L. (*Asteraceae*)-Lavendar-cotton; *Solidago virgaureae* L. (*Asteraceae*)-European goldenrod, woundwort; *Uncaria gambir* (Hunter) Roxo. (*Rubiaceae*)-gombir, pale catechu; *Vanilla planifolia* Andr. (*Orchidaceae*)-vanilla.

Isolation The areca nut or kola nut is cut into small chips mechanically and filled into the extractors. The steam is passed through the drug profusely to affect maximum extraction. The crude extract is filtered and concentrated under vacuum. The concentrated extract is chilled in deep-freezer when catechin separates as its hydrated product (mp : 93-96°C) and its anhydrous product (mp : 175-177°C).

Characteristic Features Its needles obtained from water and acetic acid give rise to its hydrate form having m.p. 93-96°C, whereas its anhydrous form registers mp 175–177°C, and $[\alpha]_D^{18} + 16^\circ$ to $+ 18.4^\circ$.

The physical parameters of *l*-form and *dl*-form are stated below.

S. No.	<i>l</i> -Catechin	<i>dl</i> -Catechin
1.	Needles from water and acetic acid mp 93-96°C (hydrated form); mp 175-177°C (anhydrous form)	Needles from water and acetic acid mp 212-216°C
2.	$[\alpha]_D - 16.8^\circ$	–
3.	Solubility in aqueous medium.	Practically insoluble in benzene, chloroform, pet, ether; soluble in hot water, alcohol, acetons, glacial acetic acid; and slightly soluble in cold water and ether.

Note Catechin is called catechol (flaran) to distinguish it from catechol (pyrocatechol q.v).

Identification


1. **Catechin** on being treated with HCl yields phluroglucinol, that burns along with lignin to produce purple or magnata colour. The *tannin extract* is taken on the tip of a match-stick, dipped in HCl and burnt in the blue-flame of the Bunsen-burner.
2. It reacts with vanillin and HCl to produce a pink or red colour.

Uses

1. It is used as an antidiarrheal agent.
2. It is also employed for dyeing and tanning.

B. Protocatechuic Acid

Chemical Structure 3,4-Dihydroxybenzoic acid.



Protocatechuic Acid

Occurrence It is found in the dried fruit of *Illicium verum* Hook. f. (*Magnoliaceae*)-**star anise, Chinese anise**; in the leaves and seeds of *Perilla frutescens* (L.) Britt. (*Lamiaceae*)-**beef-steak plant, perilla, wild coleus**; and in the timbers of *Tabebuia Spp.* (*Bignoniaceae*)-Pao D' Arco. However, minute amounts are found in wheat grains and also in wheat seedlings.

Characteristic Features It is white to brownish crystalline powder. It undergoes discolouration on exposure to air. Its mp $\sim 200^\circ\text{C}$ and d 1.54. It is soluble in 50 parts of water and freely soluble in ether and alcohol.


5.2.6.5.6 Phenolic Ether Volatile Oils A good number of volatile oils essentially contain phenolic ethers which attribute powerful aromatic odour and flavour. Because of their distinct characteristic aroma they are used extensively as pharmaceutical aids, perfumery and confectionery. A few typical

examples of phenolic ether volatile oils are, namely: **Anethole; Safrole; Myristicin; Apiole; Cineole** and **Ascaridole**.


General Properties of Phenolic Ether Volatile Oils There are certain characteristic general properties of phenolic ether volatile oils that help in their identifications:

1. They are very stable *neutral compounds* which are sparingly water-soluble. They do not react with alkalis.
2. Phenolic ethers, in general, yield the corresponding phenols on treatment with HBr or HCl.
3. They form crystalline derivatives on account of various reactions, such as: bromination, nitration and oxidation.
4. Phenolic ethers give rise to the formation of **sulphonamides** in a two-step reaction depicted below:

Step 1: It reacts with chloro sulphonic acid to yield the corresponding sulphonyl chloride together with a molecule each of hydrochloric acid and sulphuric acid.




Step 2: The resulting sulphonyl chloride (**Step 1**) on reaction with ammonium carbonate gives rise to the desired **sulphonamide** and a mole each of ammonium chloride, carbon dioxide and water.




These chemical constituents shall be discussed individually in the sections that follows:

A. Anethole (**Synonym** Anise camphor, Monasirup)

Chemical Structure 1-Methoxy-4-(1-propenyl) benzene.




It has a monohydric phenolic ether function.




Identification

- Anethole** undergoes oxidation with $K_2Cr_2O_7$ in two steps; *first step*-yields anisaldehyde (*para*-methoxy benzaldehyde), and *second step*-gives rise to *para*-methoxy benzoic acid (mp $184^\circ C$) as depicted below:



- It gets condensed with maleic anhydride to yield a condensation product having mp $310^\circ C$ as shown below:




- It gives rise to the formation of nitroso derivative having mp $126^\circ C$.

Uses

- It is used as a flavouring agent in perfumery particularly for soap and dentifrices.
- It is also employed as a pharmaceutical and (flavour).
- It finds its application as an imbedding material in microscopy.
- It is employed as a flavouring agent in alcoholic, non-alcoholic beverages and confectionaries.
- It is used as a sensitizer in bleaching colours in colour photography.

B. Safrole

Chemical Structure 5-(2-Propenyl)-1,3-benzodioxole; 4-allyl-1,2-methylenedioxybenzene.



Safrole


Occurrence It is the constituent of a number of volatile oils, notably of *sassafras i.e.*, the dried bark of the roots of *Sassafras albidum* Nees, belonging to the family *Lauraceae*, in which it is present to the extent of 75%.

It is extensively found in a variety of other plant sources, namely: *Acorus calamus* L., *Araceae* (**sweet flag, flagroot, calamus**); *Angelica polymorpha* Max., *Apiaceae* (**dong quai**); *Cananga odorata* (Lam.) Hook. f. & Thoms., *Annonaceae* (**cananga, ylang-ylang**); *Cinnamomum comphora* (L.) J.S. Presl., *Lauraceae* (**camphor, hon-sho**); *Illicium verum* Hook. f. *Magnoliaceae* (**Star-anise, Chinese anise**); *Myristica fragrans* Houtt. *Myristicaceae* (**mace, nutmeg**); *Ocimum basilicum* L. *Lamiaceae* (**sweet basil, garden basil**); *Piper nigrum* L. *Piperaceae* (**black pepper**); *Theobroma cacao* L. *Sterculiaceae* (chocolate, cocoa, cacao); *Umbellularia californica* (Hook. and Arn.) Nutt. (**California bay, California sassafras, (California laurel)**).

Isolation Safrole may be isolated from the oil of sassafras, camphor oil and oil of star-anise and also the safrole-rich fraction of the oil to about -10 to -15°C . It may also be isolated by subjecting the above safrole containing oils to fractional distillation under reduced pressure, chilling the fraction and finally crystallization.


Characteristic Features It is colourless or slightly yellow liquid having a specific sassafras odour. Its physical properties are: d_{20}^{20} 1.096, mp $\sim 11^{\circ}\text{C}$, bp $232-234^{\circ}\text{C}$ and n_{D}^{20} 1.5383. It is insoluble in water, very soluble in alcohol and freely miscible with ether and chloroform.

It undergoes isomerization on being heated with alkalis to yield **isosafrole** as shown below:



Identification

1. **Bromination:** Safrole on bromination yields the corresponding pentabromosafrole (mp $169-170^{\circ}\text{C}$).
2. **Oxidation:** Safrole on oxidation with $\text{K}_2\text{Cr}_2\text{O}_7$ and dilute H_2SO_4 (6 N) gives rise to the aldehyde derivative piperonal as shown below:




3. *Colour test:* Both **safrole** and **isosafrole** on treatment with concentrated sulphuric acid instantly produces an intense red colouration.

Uses

1. It is widely used as a flavouring agent for a variety of products, such as: beverages, pharmaceuticals chewing gums, toothpastes, in perfumery and scenting soaps.
2. It is also used in denaturing fats in soap manufacturing process.
3. It is mostly employed for the conversion to isosafrole and the manufacture of **heliotropin**.

C. Myristicin

Chemical Structure 4-(Methoxy)-6-(2-propenyl)-1, 3-benzodioxole.




Occurrence The aromatic ether is extracted from nutmeg, mace, French parsley, carrots and **dill oils**.

The botanical sources of **myristicin** are as follows: *Anethum graveolens* L. (*Apiaceae*) (**Dil, Dill Seed, Garden Dill**); *Daucus Carota* subsp. Sativus (Hoffm.) Arcang [*Apiaceae*] (**Cultivated carrot, Queen Anne's Lace** (Wild)); *Myristica fragrans* Houtt. [*Myristaceae*] (**Mace, Nutmeg**); *Petroselinum crispum* (Mill) Nym. [*Apiaceae*] (**Parsley**); *Piper nigrum* L. [*Piperaceae*] (**Black Pepper**); *Sassafras albidum* (Nutt.) Nees [*Lauraceae*] (**Sassafras**).

Isolation The rich source of volatile oil containing **myristicin** is subjected to fractional distillation under reduced pressure when the latter is collected as a colourless oily liquid.

Characteristic Features It is an oily liquid having a characteristic aromatic odour. It does not congeal at low-temperature.

Myristicin on being treated with either metallic sodium or boiled with alcoholic KOH undergoes isomerism to yield **isomyristicin** as given below:



i.e., the allyl group in the former gets converted to the propenyl group in the latter.

It has the following physical parameters:

$$\text{bp}_{40} 173^{\circ}\text{C}; n_{\text{D}}^{20} 1.54032; d_{20}^{20} 1.1437.$$

Identification

1. On oxidation with KMnO_4 it gives rise to *two* products, namely:

(a) Myristicin aldehyde (mp 130°C); and

(b) Myristinic acid (mp 208-210°C).


2. On interaction with bromine it yields the corresponding dibromoderivative having mp 130°C.

Uses It is used as a flavouring agent in food products and confectioneries.

D. Apiole

Synonym Dill; Dill apiole; Parsley comphor.

Chemical Structure 4,5-Dimethoxy-6-(2-propenyl)-1,3-benzodioxole.




Occurrence It occurs abundantly in **dill oil** *Anethum graveolus* L., belonging to the natural order *Umbelliferae*. It is also found in the **Parsley seed oil** *Petroselinum crispum* (Mill.) Nym. (Family: *Apiaceae*). The volatile oil of *Sassafras albidum* (Nutt.) Nees (Family: *Lauraceae*) contains **apiole**.


Isolation It is obtained by chilling the volatile oil to a very low temperature in a deep-freezer and finally recrystallizing it either from ethanol or petroleum ether (mp 29.5°C).

Characteristic Features **Apiole** crystallises usually in the shape of long colourless needles with a faint specific odour of *Parsley*. Its physical parameters are: mp 29.5°C, bp 285°C; n_D^{17} 1.5305; d_{15}^{15} 1.1598. It is practically insoluble in water, but soluble in ethanol, ether and in fatty oils.

Apiole on boiling with alcoholic KOH undergoes isomerisation to yield isoapiole (mp 55-56°C) whereby the allyl group in the former gets isomerized to the propenyl function in the latter as given below:



Apiole on treatment with bromine yields a monobromide (mp 51°C), a dibromide (mp 75°C) and also a tribromide (mp 120°C) as depicted below:



On oxidation with KMnO_4 both **apiole** and **isoapiole** yield the corresponding **apioaldehyde** and **apiolic acid**.

Identification

1. It may be identified by forming its bromoderivatives as stated above having a specific melting point.
2. It may also be identified by preparing its oxidative products with KMnO_4 , such as: opioaldehyde (mp 102°C) and apiolic acid (mp 173°C).

Uses

1. It exerts a synergistic activity with insecticides.
2. Dill is frequently employed as an aromatic stimulant, carminative and flavouring agent.
3. **Dill oil** is an important ingredient of '**Gripe Water**' which is given to infants to relieve them from flatulence.

E. Cineole

Synonyms Eucalyptol; Cajeputol.

Chemical Structure 1, 8-Epoxy-p-menthane.



Occurrence It is the chief constituent of **oil of eucalyptus** obtained from the leaves of *Eucalyptus globulus* Labill (Family: *Myrtaceae*) and other species of *Eucalyptus*. It also occurs largely in a variety of plants, namely: *Acorus calamus* L., (*Araceae*); *Aloysia triphylla* Britton (Family: *Verbenaceae*)-**Lemon Verbena**; *Artemisia vulgaris* L., (Family: *Asteraceae*)-**Mugwort, Carlina Thistle**; *Chamaemelum nobile* (L.) All (Family: *Asteraceae*)-**Roman Camomile, English Camomile, Camomile**, *Cinnamomum verum* J.S. Presl (Family: *Lauraceae*)-**Ceylon Cinnamon**; *Crocus sativus* L., (Family: *Iridaceae*)-**Saffron, Saffron crocus**; *Croton eleutheria* Sw. (Family: *Euphorbiaceae*)-**Cascarilla**; *Illicium verum* Hook. f. (Family: *Magnoliaceae*)-**Star-Anise, Chinese Anise**; *Juniperus communis* L. (Family: *Cupressaceae*)-**Common Juniper**; *Juniperus sabina* L. (Family *Cupressaceae*)-**Sabine, Savin**; *Laurus nobilis* L., (Family: *Lauraceae*)-**Bay, Grecian Laurel, Green Bay**; *Melaleuca leucadenron* L. (Family: *Myrtaceae*)-**Cajeput**; *Pimenta diocia* (L.) Merr. (Family: *Myrtaceae*)-**Allspice, Jamaica Pepper, Clove Pepper**; *Rosmarinus officinalis* L. (Family: *Lamiaceae*)-**Rosemary**; *Salvia sclarea* L., Family: *Lamiaceae*)-**Clary, Muscatel Sage**; *Tanacetum vulgare* L., (Family: *Asteraceae*)-**Tansy**; *Umbellularia californica* (Hook and Arn.) Nutt.-**California Bay, California Laurel, California Sassafras**.

Isolation Cineole may be isolated from **Eucalyptus oil**, which contains this ingredient to the extent of 80% by any one of the following *four* methods, namely:

Method I: Fractional Distillation. It may be obtained from fractional distillation under vacuo and the colourless liquid is collected over powdered anhydrous sodium sulphate. The clear oily substance is obtained finally in the pure crystalline form by chilling it (mp + 1.5°C).

Method II: Addition Products with Halogen Acids (HCl, HBr). It forms addition compounds with HCl and HBr as: $C_{10}H_{18}O \cdot HCl$ and $C_{10}H_{18}O \cdot HBr$, from which the pure cineole may be regenerated conveniently.

Method III: Addition Product with Resorcinol. It forms an addition compound with 50% (w/v) solution of pure resorcinol as $[(C_{10}H_{18}O)_2 \cdot C_6H_6O_2]$ having mp 80-85°C, from which cineole may be regenerated easily.

Note This reaction may cater for the separation of cineole from essential oils having a high cineole content (more than 50-60%), otherwise the volatile oil must first be fractionated.

Method IV: Addition Product with Phosphoric Acid. Cineole readily forms addition product with phosphoric acid as: $[C_{10}H_{18}O \cdot H_3PO_4]$ having mp 84°C, that may be decomposed by hot water.

Note This method is also utilized for the estimation of cineole in volatile oils in v/v percentage.

Characteristic Features It is a colourless liquid having a camphor-like odour. It possesses a spicy and cooling taste. Its physical characteristics are: d_{25}^{25} 0.921-0.923, bp 176-177°C, mp + 1.5°C, n_D^{20} 1.455-1.460, flash point (closed-up) 48°C. It is almost insoluble in water but miscible with alcohol, chloroform, ether, glacial acetic acid and oils.

Cineole forms addition compounds with resorcinol and phosphoric acid, that are found to be fairly stable, having mp 80-85°C and 80°C respectively.

It is not attacked by ordinary reducing agents, such as: glucose etc.

Identification

1. **Cineole** may be characterized by a host of derivatives/addition compounds obtained from pure chemical substances, for instance: halogen acids, resorcinol, phosphoric acid, *orthocresol* etc.
2. **Microchemical Test of Cineole:** A drop of pure cineole or a drop of Eucalyptus oil or a few drops of an alcoholic extract of eucalyptus leaf, is made to react with a drop of 5% (w/v) solution of *hydroquinone* on a microscopic slide and subsequently examined under a low-power microscope one may observe either colourless prisms or rhomboid crystals.


However, an identical treatment with a 50% (w/v) solution of *resorcinol* gives rise to beautiful leaf-like crystals.

Uses

1. It is used quite extensively in pharmaceutical preparations both meant for internal and external utilities, such as:
Internal usage—as a stimulating expectorant in cases of chronic bronchitis
External usage—as a mild antiseptic, anaesthetic in cases of inflammatory conditions.
2. It is also employed in room-sprays, hand lotions and all types of cosmetic formulations.
3. It is invariably used as a pharmaceutical aid *i.e.*, flavouring agent.

F. Ascaridole (Synonym Ascarisin)

Chemical Structure 1, 4-Peroxido-p-menthene-2. It is an organic peroxide which constitutes 60-80% of **oil of chenopodium**. It is the only naturally occurring **terpenoid peroxide**.




Occurrence **Ascaridole** is the major constituent (65-70%) in the **chenopodium oil**, *i.e.*, a volatile oil, obtained by the steam distillation from the fresh flowering and fruiting plants (except roots) of the botanical species *Chenopodium ambrosioides* var *anthelminticum* Linn., belonging to the family *Chenopodiaceae*.

Isolation It is isolated by the repeated fractional distillation of the volatile oil of chenopodium (**American wormseed oil**) *under vacuo* and collecting the fraction boiling at 95-98°C.

Characteristic Features **Ascaridole** is a viscid yellow oily liquid having a very peculiar and most disagreeable odour and flavour. It is highly unstable and is prone to explode when either subjected to heat or when treated with organic acids, *e.g.*, acetic acid; and with inorganic acids, *e.g.*, sulphuric, nitric; hydrochloric and phosphoric acids. It being a peroxide liberates I₂ from KI in acetic acid solution.

It is soluble in hexane, pentane, ethanol, toluene, benzene and castor oil. Its physical characteristics are: mp + 3.3°C; bp_{0.2} 39-40°C; [α]_D²⁰ 0.00; d₄²⁰ 1.0103;


Prepared Synthetically **Ascaridole** may be synthesized from **α-terpinene** by treatment with oxygen, chlorophyll and light as given below:



Identification As **ascaridole** does not produce any crystalline derivative, therefore, it is usually characterized by the help of the following **two specific reactions**:

1. **Formation of cis-1:4-Terpin [C₁₀ H₁₈ (OH)₂]:** Ascaridole on reduction with H₂ and Pd as a catalyst gives rise to the formation of **cis-1, 4-terpin** as follows:

The resulting *cis*-1, 4-terpin is *optically inactive* and also it is not identical with 1, 8-terpin, although the two compounds have similar melting points *i.e.*, 116–117°C.



2. **Formation of Ascaridole Glycol:** Ascaridole upon oxidation with FeSO₄ yield chiefly a ‘*glycol*’ which is not steam-volatile *viz.*, **ascaridole glycol**. Consequently, this glycol may be further characterized by the formation of its monobenzoate (mp 136-137°C) and dibenzoate derivatives (mp 116.5°C).

Uses

1. It has been used as an anthelmintic (Nematodes).
2. It is also employed for eliminating hookworms and roundworms.
3. It is used most frequently in large number of medical and veterinary formulations.

Note Estimation of Ascaridole* Ascaridole may be determined quantitatively by the method described in the Extra Pharmacopoea, which is developed by Cocking and Hymas and based upon the oxidizing property of the ‘peroxide function’ exclusively present in it on the strongly acidified solution of KI (with HCl and glacial acetic acid). Thus, the liberated I₂ is titrated with sodium thiosulphate (Na₂S₂O₃) using freshly prepared starch solution as an indicator (colour change from blue to colourless) under the specified experimental parameters.

Precautions

1. Addition of oxygenated constituents must be avoided in the assay procedure that may give rise to erroneous results.
2. As the liberated iodine is capable of being absorbed by unsaturated components present in the volatile oil, it is absolutely necessary to carry out the assay at low temperature so as to maintain such secondary reactions at the lowest level.

5.2.6.5.7 Oxide Volatile Oils The various chemical constituents containing oxide function present in naturally occurring volatile oils are namely: **Safrole, Myristicin, Apiole, Cineole and Ascaridole**. These compounds have already been discussed under section *Phenolic Ether Volatile Oils* (5.2.6.5.6) in this chapter.

* *Analyst*, 55, 180, 1930.

5.2.6.5.8 Ester Volatile Oils The 'Ester Volatile Oils' essentially attribute their flavouring characteristics, odour, aroma and specific perfume by virtue of the presence of a good number of naturally occurring esters, most common among which are the acetates of borneol, geraniol and terpineol. However, it is an age-old practice to allow the maturation or ageing of such ester-containing perfumes in order to enhance the process of esterification *in situ* thereby ultimately improving the overall aroma and bouquet of the volatile oil. Incidentally, there are certain exceptions, such as: the 'Oil of Wintergreen' which contains upto 99% of **methyl salicylate** (an ester).

Classification The ester volatile oils may be classified conveniently into *three* categories as follows:


- (i) Esters of Aliphatic Acids,
- (ii) Esters of Aromatic Acids, and
- (iii) Esters containing Nitrogen.

These different categories of ester volatile oils shall be discussed along with their typical examples as under:

5.2.6.5.8.1 Esters of Aliphatic Acids The typical examples of esters of the aliphatic acids are, namely: **Geranyl acetate, Linalyl acetate.**

A. Geranyl Acetate

Chemical Structure 3, 7-Dimethyl-2, 6-octadien-8-acetate; (C₁₂ H₂₀ O₂). It is an olefenic terpene acetate mostly found in a number of essential oils.



Geranyl Acetate

Occurrence The ester is very widely distributed in a variety of essential oils, such as: **oil of citronella, petit grain, lemon-grass, coriander, lavender** etc. It is also found in *Satureja montana* L. (*Lamiaceae*)-**Winter Savory, White Thyme, Spanish Savory**; and *Tilia europaea* L. (*Tiliaceae*)-**Lime Tree** (Europe), **Linden Tree** (America).

Isolation It may be obtained from the rich source of volatile oil containing **greanyl acetate** by fractional distillation under vacuum.

Characteristic Features It is a colourless liquid having a pleasant aromatic fragrance resembling to that of rose. It boils at 242-245°C with decomposition at atmospheric pressure.

Identification The ester on saponification with alcoholic KOH yields geraniol and acetic acid as the products of reaction. The former may be identified by examining its physical parameters, for instance: bp₇₅₇ 229-230°C; bp₁₂ 114-115°C; d₄²⁰ 0.8894; n_D²⁰ 1.4766; uv_{max} 190-195 nm (ε 18000).


Uses

1. It is used abundantly in perfumery.
2. It is also employed in making cosmetics and various types of toilet soaps.

B. Linalyl Acetate

Synonym Bergamot.

Chemical Structure 3, 7-Dimethyl-1, 6-octadien-3-yl acetate; (C₁₂ H₂₀ O₂). It is also an olefinic terpene acetate and regarded as the most valuable constituent of bergamot and lavender oils.



Linalyl Acetate

Occurrence It is found in a number of volatile oils, namely: *Lavandula angustifolia* Mill. (*Lamiaceae*)-**Lavender, True or Common Lavender**; *Salvia selarea* L. (*Lamiaceae*)-**Clary, Cleareye, Muscatel Sage**; *Satureja montana* L. (*Lamiaceae*)-**Winter Savory, White Thyme, Spanish Savory**; *Thymus vulgaris* L. (*Lamiaceae*)-**Common Thyme**; *Tilia europaea* L. (*Tiliaceae*)-**Linden Tree** (America), **Lime Tree** (Europe).

Isolation It may be obtained from **lavender oil** or **bergamot oil** by subjecting it to distillation under very high vacuum, because on distillation at atmospheric pressure or with steam distillation linalyl acetate gets hydrolysed rapidly and decomposed eventually.

Characteristic Features It is a colourless oily liquid having a very pleasant fruity odour of **bergamot oil**. Its physical properties are: d_4^{20} 0.885; bp 220°C; n_D^{20} 1.4460. It is almost insoluble in water but miscible freely with ether and alcohol.

Identification Linalyl acetate upon saponification with alcoholic KOH yields **linalool (linalol)** and acetic acid. The *dl*-form of **linalool** has a bp_{720} 194-197°C.


Uses It is used extensively in perfumery.

5.2.6.5.8.2 **Esters of Aromatic Acids** The various esters that are associated with aromatic acids and found in volatile oils are: **Benzyl benzoate, Cinnamyl Cinnamate; Methyl Salicylate.**

A. Benzyl Benzoate

Synonyms Ascabin; Venzonate; Ascabiol.

Chemical Structure Benzoic acid phenyl methyl ester; (C₁₄H₁₂O₂).



Benzyl Benzoate

Occurrence It is contained in **Peru and Tolu balsams**. It is also found in a variety of volatile oils, such as: *Cananga odorata* (Lam.) Hook. f. & Thomas (*Annonaceae*)-**Ylang-Ylang, Cananga**; *Cinnamomum verum* J S Presl (*Lauraceae*)-**Ceylon Cinnamon**; *Myroxylon balsamum* Var. *Pereirae*

(Royle) Harms. (*Fabaceae*)-**Balsam of Peru**; *Peumus boldus* Molina (*Monimiaceae*)-**Baldo**; *Vanilla planifolia* Andr. (*Orchidaceae*)-**Vanilla**.

Isolation Benzyl benzoate may be isolated by cooling the corresponding fraction to a very low temperature when it gets separated as a solid (mp 21°C). It may also be further recrystallized from chloroform or ether.

Characteristic Features It is an oily liquid or leaflets, having a faint, pleasant aromatic odour. It possesses a sharp burning taste. Its physical characteristics are: mp 21°C; d_4^{20} 1.118; bp 323-324°C; bp₁₆ 189-191°C; bp_{4.5} 156°C, and n_D^{21} 1.5681. It is sparingly volatile with steam. It is insoluble in water or glycerol, but miscible with alcohol, chloroform, ether and oils.

Identification The benzyl benzoate on saponification yields the two products of reaction *i.e.*, benzoic acid and benzyl alcohol that may be identified by carrying out specific tests for these compounds.


Uses

1. It is used extensively as a diluent and solvent of solid aromatics *e.g.*, artificial musk.
2. By virtue of its low volatility benzyl benzoate is employed as a fixative in perfume composition.
3. It is used as a solvent for cellulose acetate and nitrocellulose.
4. It serves as a substitute for camphor in celluloid and plastic pyroxylin compounds.
5. It is also employed in confectionery and chewing gum flavours.

B. Cinnamyl Cinnamate

Synonyms Cinnyl cinnamate; Styracin.

Chemical Structure 3-Phenyl-2-propenoic acid 3-phenyl-2-propenyl ester.



Cinnamyl Cinnamate

Occurrence It occurs in the buds of *Populus balsamifer* L., (Family: *Salicaceae*)-Goris; *Lavanga scandens* Buch-Ham., *Lavangalata*-Baslas; and *Styrox Benzoin* Dryander (Family: *Styracaceae*)-**Benzoin, Sumatra Benzoin, Styrax**.

Isolation It may be obtained from the volatile oil fraction by chilling it to low temperature and collecting the solids having mp 44°C.

Characteristic Features The characteristic features of the *trans-trans-cinnamyl cinnamate* are: mp 44°C; uv_{max} (95% ethanol): 216, 223 nm (log ϵ 3.45, 3.25). It is practically insoluble in water but sparingly soluble in cold ethanol and soluble in ether (1 g in 3 ml).

Identification On saponification with alcoholic KOH this ester gives rise to cinnamic acid and cinnamyl alcohol which may be further identified by performing their specific tests.


Uses

1. It is used in perfumery.
2. It is also employed in making toilet soaps etc.

C. Methyl Salicylate

Synonyms Winter green oil; Betula oil; Sweet birch oil; Teabery oil.


Chemical Structure 2-Hydroxybenzoic acid methyl ester; (C₈ H₈ O₃)



Methyl Salicylate

Occurrence It is largely found in a variety of medicinal plants, namely: Flowers of *Acacia farnesiana* (L.) Willd. (Family: *Fabaceae*)-**Cassie, Huisache**; *Cananga odorata* (Lam.) Hook. f. & Thoms (Family: *Annonaceae*)-**Cananga, Ylang-Ylang**; Leaves of *Chenopodium ambrosioides* L. (Family: *Chenopodiaceae*)-**Wormseed**; *Erythroxylum coca* Lam. (Family: *Erythroxylaceae*)-**Coca**; Flowerbuds of *Filipendula ulmaria* (L.) Maxim (Family: *Rosaceae*)-**Meadowsweet, Queen of the Meadow**; Twigs of *Gaultheria procumbens* L. (Family: *Ericaceae*)-**Wintergreen, Teaberry, Boxberry**; Bark of *Betula lenta* L. (Family: *Betulaceae*)-**Sweet Birch**. However, oil of wintergreen contains upto 99% **methyl salicylate**.

It is pertinent to mention here that in several aromatic medicinal plants, for instance: *Wintergreen*, the active chemical constituent *i.e.*, **methyl salicylate** does not occur as such, but is present in the form of a *glucoside* known as **Gaultherin** which upon enzymatic hydrolysis gives **methyl salicylate** and **primeverose (glucoxylose)** as shown under:



Isolation **Methyl salicylate** may be obtained from gaultherin by enzymatic hydrolysis and then subjecting the resulting products of reaction to very low temperature when the former gets solidified at -8.6°C and hence may be separated easily.

Characteristic Features It is a colourless, yellowish or reddish, oily liquid. Its odour and taste resembles to that of gaultheria. Its physical parameters are: mp -8.6°C ; bp $220-224^{\circ}\text{C}$; d_{25}^{25} 1.184; d of the natural ester is ~ 1.180 ; and n_D^{20} 1.535-1.538. It is very sparingly soluble in water (1 g in 1500 ml), but freely soluble in chloroform and ether. It is, however, miscible with alcohol and glacial acetic acid.

Identifications

1. It develops a red-violet colouration on being treated with cold saturated aqueous solution of FeCl_3 , that lasts for about 15 minutes.
2. **Methyl salicylate** readily forms a soluble ester-salt with a moderately concentrated aqueous solution of KOH as *potassium methyl salicylate*.
3. Upon saponification the ester yields salicylic acid (mp 158°C) and methanol respectively.
4. **Methyl salicylate** may also be identified by the formation of several derivatives as stated below:
 - (i) Methyl-o-acetoxy benzoate (mp $52\text{--}52.5^\circ\text{C}$)—with Acetic Anhydride,
 - (ii) Methyl-o-benzyloxy benzoate (mp 92°C)—with Benzoyl Chloride,
 - (iii) o-Cabomethoxyphenyl-N-phenyl urethane—with Phenylisocyanate.

Uses


1. **Methyl salicylate** has local irritant, antirheumatic and antiseptic properties. It is an important ingredient of **Iodex^(R)** ointment for relief of pain in several conditions like, pulled muscle, muscular pain, pain in joints, etc.
2. It also finds its extensive usage in a variety of products, such as: flavouring of food products, beverages, candies, confectionery, toothpastes, mouth washes, gargles, and pharmaceutical preparations.
3. It is also used in perfumery.

5.2.6.5.8.3 **Esters Containing Nitrogen** The specific example of an ester containing *nitrogen* is **Methyl Anthranilate** which is present in several volatile oils. It is described below:

A. Methyl Anthranilate

Synonyms Neroli Oil (Artificial).

Chemical Structure 2-Aminobenzoic acid methyl ester, ($\text{C}_8\text{H}_9\text{NO}_2$).



Methyl Anthranilate

Occurrence It occurs in a good number of medicinal herbs, for instance: Flowers of *Robinia pseudoacacia* L. (Family: *Fabaceae*)-**Black Locust, False Acacia**; *Citrus sinensis* (Linn.) Osbeck. (Family: *Rutaceae*)-**Sweet Orange**; *Cananga odorata* (Lam.) Hook. f. & Thoms. (Family: *Annonaceae*)-**Ylang-Ylang, Cananga**; *Jasminum officinale* Linn. var *grandiflorum* Bailey., (Family: *Oleaceae*)-**Jasmine**. It is also found in bergamot, other essential oils and in grape juice.

It is also obtained synthetically by carrying out the esterification of anthranilic acid with methanol in the presence of HCl.

Isolation **Methyl anthranilate** may be isolated from the essential oils very conveniently by shaking the volatile oil with cold dilute sulphuric acid (2 N) *i.e.*, 1 ml of conc. H_2SO_4 dissolved slowly in

17 ml of distilled water. The resulting methyl anthranilate sulphate thus obtained gets crystallized in the cold, which may be further purified by recrystallization from alcohol. Finally the pure desired ester is regenerated by treatment with dilute NaOH solution (2 N) carefully.

Characteristic Features It is a crystalline mass having a powerful pleasant taste. It has a peculiar odour that mostly resembles to orange blossoms and certain varieties of grape. It gives an inherent blue-violet fluorescence which is distinctly visible in any volatile oil containing it. **Methyl anthranilate** has the following physical parameters, namely:

d 1.168; mp 24-25°C; bp₁₅ 135.5°C.

It is slightly soluble in water, but freely soluble in ethanol and ether.

Identification It may be identified by preparing its derivatives, such as: Picrate (mp 104°C); Benzoate (mp 100°C).

Uses

1. It is used frequently as a perfume for ointments.
2. It is also employed for the manufacture of synthetic perfumes.

5.2.7 Resins and Resin Combinations

Resins, in general, are amorphous solid or semisolid substances that are invariably water insoluble but mostly soluble in alcohol or other organic solvents. However, physically they are found to be hard, translucent or transparent and fusible *i.e.*, upon heating they first get softened and ultimately melt. But chemically, they are complex mixtures of allied substances, such as: **resin acids, resin alcohols (or resinols), resinotannols, resin esters, glucoresins** and the like.

Another school of thought considers **Resins** as amorphous products having an inherent complex chemical entity. These are normally produced *either* in schizogenous or in schizolysigenous ducts or in canals and are regarded as the end products of metabolism. The physical general characteristic features of resins are namely: hard, transparent, or translucent and, when heated they yield usually complex mixtures that comprise of resin acids, resin alcohols, resinotannols, esters and resenes. Some researchers do believe that the resins are nothing but the oxidation products of the **terpenes**. They are found to be mostly insoluble in water, but soluble in ethanol and organic solvents. They are electrically non-conductive and combustible in nature.

Resins shall now be discussed at length in their various aspects as enumerated here under:

- (a) Distribution of Resins in Plants
- (b) Occurrence in Plants
- (c) Physical Properties of Resins
- (d) Chemical Properties of Resins
- (e) Solubility
- (f) Preparation of Resins
- (g) Chemical Composition of Resins
- (h) Classification of Resins.

5.2.7.1 Distribution of Resins in Plants

Interestingly, the **resins** and resinous substances are more or less extensively distributed throughout the entire plant kingdom, specifically the *Spermatophyta i.e.*, the seed plants. Notably, their presence

is almost rare and practically negligible in the *Pteridophyta i.e.*, the ferns and their allies. However, the resins have not been reported in the *Thallophyta i.e.*, the sea-weeds, fungi etc.

Therefore, all these findings and observations lead one to the fact the *resins* are the overall and net result of metabolism in the *higher plants*, since the majority of them belong to the phylum *Angiospermum i.e.*, seed-enclosed flowering plants, and *Gymnosperm i.e.*, naked-seed non-flowering plants.

In general, the most important and extensively studied resin-containing families are, namely: *Pinaceae* (*Colophory* or *Rosin*); *Leguminosae* (*Tolu Balsam* and *Balsam of Peru*); *Dipterocarpaceae* ('*Garijan*'—a Balsam substitute for copaiba); *Burseraceae* (Myrrh) and *Umbelliferae* (Asafoetida).

5.2.7.2 Occurrence in Plants

In the plants resins usually occur in different secretory zones or structures. A few typical examples of such plant sources along with their specific secretory structures are given below:

- (i) **Resin Cells** : *Ginger–Zingiber officinale* Roscoe (Family: *Zingiberaceae*);
- (ii) **Schizogenous Ducts** : *Pine Wood–Pinus polustris* Miller.
or Schizolysoygenous Ducts (Family: *Pinaceae*);
- (iii) **Glandular Hairs** : *Cannabis–Cannabis sativa* Linne'.
(Family: *Moraceae*)

The formation of **resins** in the plant is by virtue of its normal physiological functions. However, its yield may be enhanced in certain exceptional instances by inflicting injury to the living plant, for instance: *Pinus*. Furthermore, many resinous products are not formed by the plant itself unless and until purposeful and methodical injuries in the shape of incisions are made on them and the secretions or plant exudates are tapped carefully, such as: **Balsam of Talu** and **Benzoin**. In other words, these resins are of pathological origin. One school of thought has categorically termed the secretion exclusively obtained from the naturally occurring secretory structure as the **Primary Flow**, whereas the one collected through man-made-incisions on the plant *i.e.*, abnormally formed secretory structures, as the **Secondary Flow**.

In normal practice, it has been observed evidently that resins are invariably produced in ducts as well as cavities; sometimes they do not occur in the so called specialized-secretory structures, but tend to get impregnated in all the elements of a tissue, for example: **Guaiacum Resin**—is obtained from the heartwood of *Guaiacum officinale* Linn. and *G. sanctum* Linn., (Family: *Zygophyllaceae*) *i.e.*, it is found in the vessels, fibres, medullary ray cells and wood parenchyma. In this particular instance, the resins occur as **tyloses**, achieved by chopping off the conduction in these areas so as to enhance the effective usage of root pressure and the capillaries in forcing both the nutritive contents and forcing water to reach the top end of these tall trees.

It is pertinent to mention here that in some exceptionally rare instances the resin occurs as a result of sucking the juice of the plant by scale insects and converting the sucked-juice into a resinous substance that ultimately covers the insect itself and twigs of the plant as well, for instance: *Laccifer lacca* (Family: *Coccidae*)-**Shellac**.

5.2.7.3 Physical Properties of Resins

The various physical properties of **resins** can be generalized as detailed below:

1. **Resins**, as a class, are hard, transparent or translucent brittle materials.
2. They are invariably heavier than water having the specific gravity ranging from 0.9-1.25.
3. **Resins** are more or less amorphous materials but rarely crystallisable in nature.
4. On being heated at a relatively low temperature **resins** first get softened and ultimately melt down thereby forming either an adhesive or a sticky massive fluid, without undergoing any sort of decomposition or volatilization.
5. On being heated in the air *i.e.*, in the presence of oxygen, resins usually burn readily with a smoky flame by virtue of the presence of a large number of C-atoms in their structure.
6. On being heated in a closed container *i.e.*, in the absence of oxygen, they undergo decomposition and very often give rise to **empyreumatic products** *i.e.*, products chiefly comprising of hydrocarbons.
7. Resins are bad conductors of electricity, but when rubbed usually become negatively charged.
8. They are practically insoluble in water, but frequently soluble in ethanol, volatile oils, fixed oils, chloral hydrate and non-polar organic solvents *e.g.*, benzene, n-hexane and petroleum ether.

5.2.7.4 Chemical Properties of Resins

The various chemical properties of **resins** may be summarized as stated below:

1. Resins, in general, are enriched with carbon, deprived of nitrogen and contain a few oxygen in their respective molecules.
2. Majority of them undergo slow atmospheric oxidation whereby their colour get darkened with impaired solubility.
3. Resins are found to be a mixture of numerous compounds rather than a single pure chemical entity.
4. Their chemical properties are exclusively based upon the functional groups present in these substances.
5. Consequently, the resins are broadly divided into **resin alcohols, resin acids, resin esters, glycosidal resins** and **resenes** (*i.e.*, inert neutral compounds).
6. Resins are regarded as complex mixtures of a variety of substances, such as: **resinotannols, resin acids, resin esters, resin alcohols and resenes**.
7. One school of thought believes that resins are nothing but **oxidative products of terpenes**.
8. They may also be regarded as the end-products of *destructive metabolism*.
9. The acidic resins when treated with alkaline solutions they yield soaps (or **resin-soaps**).

Note The solutions of resins in alkalis distinctly differ from ordinary soap solutions by virtue of the fact that the former cannot be easily 'salted-out' by the addition of NaCl, unless it is used in large excess quantity.

5.2.7.5 Solubility

The solubility of various types of **resins** are as follows:

1. Majority of **resins** are water-insoluble and hence they have practically little taste.

2. They are usually insoluble in petroleum ether (a non-polar solvent) but with a few exceptions, such as: colophony (freshly powdered) and mastic.
3. Resins mostly got completely dissolved in a number of polar organic solvents, for instance: ethanol, ether and chloroform, thereby forming their respective solutions which on evaporation, leaves behind a thin-varnish-like film deposit.
4. They are also freely soluble in many other organic solvents, namely: acetone, carbon disulphide, as well as in fixed oils and volatile oils.
5. Resins dissolve in chloral hydrate solution, normally employed for clarification of certain sections of plant organs.

5.2.7.6 Preparation of Resins

So far, no general method has either been suggested or proposed for the preparation of resins. In fact, there are *two* categories of resinous products, namely: (a) **Natural Resins**; and (b) **Prepared Resins**, have been duly accepted and recognized. Therefore, this classification forms the basis of the methods employed in the preparation of the *two* aforesaid resins.

A. Natural Resins: These resins usually formed as the exudates from various plants obtained either normally or as a result of pathogenic conditions (*i.e.*, by causing artificial punctures), such as: mastic, sandarac. These are also obtained by causing deep incisions or cuts in the trunk of the plant, for instance: **turpentine**. They may also be procured by hammering and scorching, such as: **balsam of Peru**.

B. Prepared Resins: The resins obtained here are by different methods as described below:

- (i) The crude drug containing resins is powdered and extracted with ethanol several times till complete exhaustion takes place. The combined alcoholic extract is either, evaporated on a electric water-bath slowly in a fuming cup-board or poured slowly into cold distilled water. The precipitated resin is collected, washed with cold water and dried carefully under shade or in a vacuum desiccator,

Examples: Podophyllum; Scammony and Jalap.

- (ii) In the case of *alco-resins*, organic solvents with lower boiling points are normally employed *e.g.*, solvent ether (bp 37°C); acetone (bp 56.5°C), for their extraction. However, the volatile oil fraction can be removed conveniently through distillation under vacuo.
- (iii) In the instance of **gum-resins**, the resin is aptly extracted with 95% (v/v) ethanol while leaving the insoluble gum residue in the flask (or soxhlet thimble).

5.2.7.7 Chemical Composition of Resins

The copious volume of information with regard to the '**chemistry of resins**' is mainly attributed by the meaningful research carried out by Tschirch and Stock, who advocated that the proximate constituents of resins may be classified under the following heads, namely:

- (i) *Resin Acids*
- (ii) *Resin Esters and their Decomposition Products i.e., Resin Alcohols (Resinols) and Resin Phenols (Resinotannols).*

* Tschirch, A, and L. Stock: *Die Harze, Borntraeger*, Berlin, Vols. 1 & 2, 1933-36.

(iii) *Resenes i.e.*, the chemical inert compounds.

However, it has been observed that in majority of the known **resins** these *three* aforesaid categories evidently predominates and thus the resulting product consequently falls into one of these groups. It is worth mentioning here that representatives of all the three said groups are rarely present in the same product.

Given below are some typical examples of resin substances that predominates the *three* classes suggested by Tschirch and Stock, namely:

- A. **Resin-Esters** : *Examples:* Ammoniacum; Asafoetida; Benzoin; Balsam of Peru and Tolu; Galbanum; Storax;
 B. **Resin-Acids** : *Examples:* Colophony; Copaiba; and
 C. **Resenes** : *Examples:* Bdellium; Dammar; Mastic; Myrrh; Olibanum.




A few important and typical chemical constituents that have been duly isolated and characterized from various **naturally occurring resins** are discussed below:

1. Resin Acids

Synonyms Resinolic Acid.

The **resin acids** essentially contain a large portion of carboxylic acids and phenols. However, they occur both in the *free state* and as their *respective esters*. They are usually found to be soluble in aqueous solutions of the alkalis, thereby forming either soap like solutions or colloidal suspensions. **Resinates**, *i.e.*, the metallic salts of these acids find their extensive usage in the manufacture of inferior varieties of soaps and varnishes.


A few typical examples of resin acids are enumerated below:

S. No.	Name	Chemical Structure	Source(s)
1	Abietic Acid		Colophony, Rosin
2	Copaivic Acid and Oxycopaivic Acid	—	Copaiba
3	Guaiaconic Acid		Guaic
4	Pimaric Acid		Burgandy Pitch, Fankicense
5	Sandracolic Acid	—	Sandarac
6	Commiphoric Acid	—	Myrrh

Out of all the six commonly found resin acids *Abietic Acid* shall be discussed here under:

Abietic Acid (Synonym Sylvic Acid)

Chemical Structure 13-Isopropylpodocarpa-7, 13-dien-15-oic acid; (C₂₀ H₃₀ O₂).



Abietic Acid

It is a **tricyclic diterpene** embedded with four isoprene units. It is studded with four methyl moieties and a carboxylic acid function. Besides, it also has two double bonds one each in ring-B and ring-C of the phenanthrene nucleus.

Preparation It is a widely available organic acid, prepared by the isomerization of **rosin**.* It may also be synthesized from **dehydroabietic acid****.

The commercial grade of abietic acid is normally obtained by heating either rosin alone or with mineral acids. The product thus achieved may be glassy or partly crystalline in nature. It is usually of yellow colour and has a mp 85°C *i.e.*, much lower than the pure product (mp 172-175°C).

Characteristic Features It is obtained as monoclinic plates from alcohol and water. Its physical parameters are: mp 172-175°C; $[\alpha]_D^{24} -106^\circ$ ($c = 1$ in absolute alcohol); $uv_{\max} 235, 241.5, 250$ nm (ϵ 19500, 22000, 14300). It is practically insoluble in water, but freely soluble in ethanol, benzene, chloroform, ether, acetone, carbon disulphide and also in dilute NaOH solution.

Identification It readily forms the corresponding methyl ester as methyl abietate (C₂₁ H₃₂ O₂), which is colourless to yellow thick liquid bp 360-365°C, $d_{20}^{20} 1.040$, and $n_D^{20} 1.530$.

Uses

1. It is used for manufacture of esters (ester gums), such as: methyl, vinyl and glyceryl esters for use in lacquers and varnishes.
2. It is also employed extensively in the manufacture of 'metal resinates' *e.g.*, soaps, plastics and paper sizes.
3. It also assists in the growth of butyric and lactic acid bacteria.

2. Resin Alcohols In general, resin alcohols are complex alcohols having higher molecular weight. These are of *two* types, namely:

* Harris, Sanderson, *Org. Syn. Coll. Vol. IV*, 1 (1963); and Fieser and Fieser, *The chemistry of Natural Products Related to Phenanthrene* (New York, 3rd. edn., 1949).

** A.W. Burgastahler, and L.W. Worden., *J. Am. Chem. Soc.*, **83**, 2587, (1961) E. Wenkert *et al.*, *ibid*, **86**, 2038, (1964).

(a) **Resinotannols:** The resin alcohols which give a specific tannin reaction with iron salts are termed as **resinotannols**.

A number of **resinotannols** have been isolated from the plant kingdom. It is an usual practice to name them according to the resins in which they are found, such as:

- Alocresinotannol** – From *Aloe* species viz., *Aloe barbedensis* Miller, (Curacao Aloes); *Aloe perryi* Baker, (Socotrine Aloes); *Aloe ferrox* Miller, *Aloe africana* Miller, *Aloe spicata* Baper. All these belong to the natural order *Liaceae*.
- Ammoresinotannol** – From *Ammoniacum i.e.*, the oleo-gum-resin from *Dorema ammoniacum* D. Don. (Family: *Umbelliferae*).
- Galbaresinotannol** – From *Galbanm i.e.*, the oleo-gum-resin from *Ferula galbaniflua* Boiss et Bubse (Family: *Unbelliferae*).
- Peruresinotannol** – From Balsam of Peru *i.e.*, the balsam obtained from *Myroxylon balsamum* var *Pereirae* (Royle) Harms (Family: *Fabaceae*);
- Siaresinotannol** – From Sumatra Benzoin (Benzoin, *Styrax*) *i.e.*, the gum exuded from *Styrax benzoin* Dryander (Family: *Styracaceae*).
- Toluresinotallol** – From Balsam of Tolu *i.e.*, the Balsam obtained from *Myroxylon balsamum* (Linn.) Harms. (belonging to the family. *Leguminosae*).


(b) **Resinols:** The resin alcohols that fail to give a positive reaction with tannin and iron salts are known as **resinols**. *The following* are some typical examples of **resinols**, for instance:

- Benzoresinol** – From Benzoin which is purely a pathological product obtained either from *Styrax benzoin* Dryander and *Styrax paralleloneurus* Brans. (*Sumatra Benzoin*) or from *Styrax tonkinensis* Craib. (*Siam Benzoin*) belonging to family *Styraceae*.
- Storesinol** – From storax which is the balsamic resin usually obtained from the trunk of *Liquidamber orientalis* Mill. family *Hamamelidaceae*.
- Gurjuresinol** – From Gurjun Balsam that is the aleo-resin obtained from *Dipterocarpus turbinatus* Gaertn. F. belonging to family: *Dipterocarpaceae*.
- Guaiaresinol** – From Guaiacum Resin obtained from the heartwood of *Guaiacum officinale* Linn. and *Guaiacum sanctum* Linn. belonging to family: *Zygophyllaceae*.

3. Resenes These are oxygenated compounds, but are not affected either by alkalies or acids. In fact, they are more or less neutral substances being devoid of characteristic functional groups, and, therefore, do not exhibit any characteristic chemical properties. Interestingly, they are immune to oxidizing agents and variant climatic conditions, a fact which essentially attributes the resins containing them one of their major plus points for the manufacture of **varnishes**. A few important examples of **resenes** are as follows:

- Dracoresene** – Derived from the scales of the fruit of Dragon's Blood *i.e.*, *Daemonorops draco* Bl. (and other species) belonging to the natural order (*Areceae*).
- Masticoresene** – Derived from Mastic-an oleo-resin obtained from *Pistacia lentiscus* Linn belonging to family: *Anacardiaceae*.
- Fluavil** – Obtained from **Gutta-percha** and also from the bark of various trees. Gutta-percha is hard and has a very low elasticity. X-ray diffraction studies have

revealed that rubber is composed of long chains built up of isoprene units arranged in the *cis*-form, whereas gutta-percha is the *trans*-form as shown below:



5.2.7.8 Classification of Resins

The *resins* are broadly classified under *three* major categories, namely:

A. Taxonomical Classification The resins are grouped together according to their botanical origin exclusively, such as:

- Coniferous Resins** – *e.g.*, Colophony; Sandarac;
- Berberidaceae Resins** – *e.g.*, Podophyllum; and
- Zygophyllaceae Resins** – *e.g.*, Guaiacum.

In this particular instance, it has been observed that the resins that usually occur in plants of the same natural order (*i.e.*, family), may exhibit more or less related characteristic features.

B. Chemical Classification The resins may also be classified as per the presence of the predominating chemical constituents for instance:

- Acid Resins** – *e.g.*, Colophony (Abietic acid); Sandarac (Sandracolic acid); Shellac (Alleuritic acid); Myrrh (Commiphoric acid);
- Ester Resins** – *e.g.*, Benzoin (Benzyl benzoate); Storax (Cinnamyl cinnamate);
- Resin Alcohols** – *e.g.*, Balsam of Peru (Peruresinotannol) Guaiacum resin (Guaicresinol); Gurjun balsam (Gurjuresinol);
- Resene Resins** – *e.g.*, Dragon's Blood (Dracoresene); Gutta-percha (Fluavil);

Glycoresins – e.g., Jalap Resin from Jalap *i.e.*, *Ipomea purga* Hayne; (Family: *Convolvulaceae*) Podophylloresin from the dried roots and rhizomes of *Podophyllum hexandrum* (*P. emodi*) Royle. (Family *Berberidaceae*).

C. Constituents of Resin Invariably, to maintain the simplicity, resins may also be classified according to the major constituents present either in the *resin* or *resin combinations*.

Examples: Resins; Oleo-resins; Oleo-gum resins; Balsams.

After having been exposed to the various aspects of *resins* with regard to their physical and chemical, properties, occurrence and distribution, preparation, chemical composition and classification, it would be worthwhile to gain some in-depth knowledge about certain typical examples belonging to Resins; **Oleo-resins; Oleo-gum-resins; Balsams;** and **Glycoresins.**

5.2.7.8.A Resins

The various **resins** that will be discussed in the section that follows are, namely:

1. Colophony (Synonym Rosin)

Biological Source It is a yellow resin, and abietic anhydride. It is the residue left after distilling off the volatile oil from the **oleoresin** obtained from *Pinus palustris* and other species of *Pinus* belonging to family *Pinaceae*. Generally, it is offered as **wood rosin** obtained from southern pine stumps, **gum rosin** collected as the exudate from incisions in the living tree *viz.*, *P. palustris* and *P. caribaea*, and finally from **tall oil rosin**. It is chiefly produced in the USA.

Characteristic Features **Colophony** fuses gradually at 100°C and at a higher temperature it burns with a smoky flame, while leaving not more than 0.1% of ash as a residue. The alcoholic solution of colophony turns into milky-white on addition of water. When fragments of rosin are heated with water, they first melt then flow together and ultimately forms a sticky-mass.

It is a pale yellow to amber, translucent fragments, brittle fracture at ordinary temperature. It has a slight turpentine-like odour and taste.

Its acid number is not less than 150 and d 1.07-1.09. It is almost insoluble in water, but freely soluble in ethanol, benzene, ether, glacial acetic acid, oils, carbon disulphide and also soluble in dilute solutions of fixed alkali hydroxides.

Chemical Constituents

1. **Colophony** contains 90% resin acids known as abietic acid (see Section 5.2.7.7). The remaining 10% as resene-an inert substance and esters of fatty acids.
2. It also contains a mixture of dihydroabietic acid ($C_{20}H_{30}O_2$) and dehydroabietic acid ($C_{20}H_{28}O_2$).
3. On being heated at 300°C, abietic acid undergoes further molecular rearrangement to produce neo-abietic acid.

Chemical Tests

1. Dissolve 0.1 g of powdered **colophony resin** in 10 ml of acetic anhydride, add one drop of sulphuric acid and shake well. The appearance of a purple colour which rapidly changes to violet colour.
2. The alcoholic solution of **colophony** is acidic to litmus paper *i.e.*, it turns blue litmus paper to red.

3. Dissolve 0.2 g of **colophony** with 5 ml of petroleum ether (60-80°C) and filter to discard the undissolved resin, if any. Shake the resulting clear solution with twice its volume of 0.1% (w/v) cupric acetate solution. The petroleum ether layer attains an *emerald-green colouration* due to the formation of the copper salt of abietic acid.

Uses

1. **Colophony** is used in pharmacy for the preparation of zinc oxide plasters, ointments and other adhesive plasters.
2. It is widely used in the manufacture of printing inks, rubber, dark varnishes, sealing wax, linoleum and thermoplastic floor tiles.
3. It also finds its application as varnish and paint driers, cements, soaps, wood polishes, paper, plastics, fireworks, tree wax, sizes, rosin oil
4. It is used for waterproofing cardboard, walls etc.

2. Eriodictyon

Synonyms Yerba Santa; Consumptive's weed; Bear's weed; Mountain balm; Gum plant.

Biological Source It is obtained from the dried leaves of *Eriodictyon californicum* (H. & A.) Greene belonging to family: *Hydrophyllaceae*.

The plant is an evergreen shrub indigenous to the mountains of California (USA) and northern Mexico. The Indians of California have been using this as a drug since many years.

Chemical Constituents It contains a volatile oil, **eviodictyol** *i.e.*, the aglycone of **eriodictin**, **homoeriodictyol**, **chrysoeriodictyol**, **xanthoeriodictyol**, **eriodonol**, **eriodictyonic acid**, **ericolin**, formic acid, butyric acid, tannin and a *resin*.

Uses

1. It is used as a pharmaceutic flavouring aid to disguise the bitter taste of quinine formulations.
2. It has also been employed as a stimulating expectorant.

3. Guaiac

Synonyms Guaiacum resin; Gum guaiac; Resin guaiac.

Biological Source **Guaiac** is obtained from the heartwood of *Guaiacum officinale* L. or *G. sanctum* L., belonging to family: *Zygophyllaceae*.

Preparation The **resin** is obtained by cutting the tree and the log is suspended horizontally. The either ends of the log are set on fire and the resin which oozes out is collected carefully in earthen or metallic cups and allowed to harden in shade.

Characteristic Features The **guaiac resin** is brown or greenish-brown irregular lumps. It has an aromatic odour. The fracture is brittle and splintery and the exposed surface is usually glossy. It melts at 85-90°C. It is insoluble in water, but freely soluble in ethanol, ether, chloroform, creosote, solution of chloral hydrate and alkalies. It is, however, slightly soluble in carbon disulphide and benzene.

It is found to be incompatible in liquid preparations containing **acacia**, mineral acids, ferric chloride, gold chloride, water, spirit nitrous ether and permanganates.

Chemical Constituents A few of the major resinous constituents belong to the group of 'lignans'. These are essentially phenolic compounds with a C₁₈ structure and made up from two C₆-C₃ units. About 10% of the **guaiac** is *guaiaretic acid* which is nothing but diaryl butane. It also contains both **α- and β-guaiaconic acids** (70%) and **guaiacic acid**. Besides, it contains traces of vanillin, saponin and volatile oil.

Chemical Tests When a small quantity of the resin is oxidised it gives rise to a distinct blue colouration (*guaiac blue*) due to the oxidation of **α-guaiacic acid**.

Uses

1. It is mostly employed as a diaphoretic and an expectorant.
2. It is used as a clinical reagent for the testing of blood and haemoglobin.
3. An ethanolic solution of guaiac is used for the detection of oxidase enzymes and cyanogenetic glycosides.


4. Cannabis

Synonyms Indian Hemp; Indian cannabis; Marihuana; Marijuana; Pot; Grass; Weed; Bhang; Ganja; Charas, Hashish.

Biological Source **Cannabis** consists of the dried flowering tops of pistillate plants of *Cannabis sativa* L., (*C. sativa* var. *indica* Auth.), belonging to family: *Moraceae*.

Preparation After years of intensive and extensive research carried out on the selective cultivation of *Cannabis*, two of its *genetic types* have been evolved, namely: (i) Drug Type, and (ii) Hemp Type. These *two* distinctly separate genetic types of **Cannabis** shall be described briefly as stated below:


A. Drug type (Cannabis): It is, in fact, the rich (upto 15%) in the psychoactive constituent (–)-Δ⁹-*trans*-tetra-hydrocannabinol (Δ⁹-THC) as shown below:



(–)-Δ⁹-*trans*-Tetrahydrocannabinol (Δ⁹-THC)

The Δ⁹-THC is usually concentrated into a **resin** that is secreted right into the *trichomes* located on the small leaves (bracts) and *bracteoles* (*i.e.*, the leaf-like structure which encloses the ovary) of the flowering tops of the *female plant*. Interestingly, for the specific drug usages either the resin (*hashish*) is employed or the flowering tops of the female plant (*marijuana*). Nevertheless, the male plant also generates an equivalent quantity of the active constituents; however, it is not concentrated into a resin but found throughout the entire plant.


B. Hemp Type (Cannabis): It contains surprisingly very little active principal. **Cannabidiol** is the predominant **cannabinoid** present in it as given below:




Cannabidiol

The hemp-type cannabis also possesses the elongated **bast*** fibers which is very much desired in the manufacture of ropes.

Chemical Constituents The chemical constituents Δ^9 THC and cannabidiol present in the drug-type and hemp-type cannabis have already been discussed above. Besides, the resin contains several active constituents, such as: **cannabinol**, **cannin**, **cannabol**, **tetrahydrocannabinol**, **cannabigerol**, **cannabichromene** and **Δ^8 -tetrahydrocannabinol**.



Cannabinol



Tetrahydrocannabinol

It also contains choline, volatile oil and **trigonelline**. However, the Indian Hemp seeds contain 20% of fixed oil.

Chemical Tests

1. Shake 0.1g of resin with 5 ml petroleum ether (60-80°C) and filter. To 1 ml of the filtrate, add 2 ml of 15% solution of HCl gas in ethanol, when a red colouration appears at the junction of the two layers. However, after shaking, the upper layer becomes colourless while the lower one attains a distinct orange pink colour, which finally vanishes upon addition of water.
2. Extract 1g of resin with methanol, filter and evaporate to complete dryness. Again extract the resulting residue with petroleum ether (60-80°C), filter directly into a separating funnel and extract the ethereal layer successively with 5% (w/v) Na_2CO_3 and 5% (w/w) H_2SO_4 . Wash the ethereal layer with distilled water, decolourise with powdered activated carbon, if necessary, and evaporate the filtrate. Add to the residue a few drops of N/10 alcoholic KOH solution, when a purple colouration is obtained.

Uses It has been used as a sedative in equine colic.

* **Bast:** Fibrous material obtained from the pith of jute, flax etc., used for making rope, matting etc.,

5. Mastic

Synonyms Mastiche; Mastich; Balsam Tree; Pistachia Galls; Mastix; Lentisk; Mastisol;

Biological Source **Mastic** is the concrete resinous exudate obtained from *Pistacia lentiscus* Linne' belonging to the natural order *Anacardiaceae*.

Preparation The resinous juice gets collected in cavities present in the inner bark. Fairly long incisions are made in the trunk and also in the larger branches, through which the resin exudes. The resin finally gets collected in the form of small tears on the outside. These are hand-picked and stored in dry place.

Characteristic Features It is a pale yellow or greenish yellow, globular, elongated or pear-shaped tears. It has a slightly balsamic odour and a **terebene*** taste. It is practically insoluble in water, but completely soluble in ethanol, chloroform (1g/0.5 ml), ether (0.1g/0.5 ml), partially soluble in **oil of turpentine**.

Chemical Composition **Mastic** contains 90% of a resin, comprising of **mastichic acid (α -resin)**, which is soluble in ethanol and **masticin (β -resin)**, which is insoluble in ethanol, and a volatile oil, 1 to 2.5%, that has the specific balsamic odour of the drug and largely contains (+)-**pinene**. A **bitter principle** is also present.

Uses

1. It is employed as an 'enteric coating' material in the formulation of tablets.
2. It is also used as a microscopical mountant.
3. It is widely used in the manufacture of varnishes.
4. Mastic is used in the form of a 'dental varnish' in dentistry to seal off cavities.
5. It is also used in tooth cements, plasters, lacquers, chewing gums and incense.
6. It is employed for retouching negatives.

6. Podophyllum

Synonyms Podophyllum resin; May apple; Mandrakes Root; Indian apple; Vegetable calomel.


Biological Sources **Podophyllum** is the dried rhizomes and roots of *Podophyllum peltatum* L., family: *Berberidaceae*, known as *American Podophyllum*; and from *Podophyllum hexandrum* Royle (*Syn. P. emodi* Wall. ex Hook. f. & Th.) usually called *Indian Podophyllum*.

Preparation Extract powdered **podophyllum** (1 killo) by means of slow percolation until it is almost exhausted of its resin content, using ethanol as the menstruum. Carefully concentrate the percolate by evaporation until the residue attains the consistency of a thin syrup. Pour the resulting syrupy liquid with constant stirring into 1 L of distilled water containing 10 ml of concentrated HCl and previously cooled to a temperature less than 10°C. Allow the precipitate to settle down completely, decant the clean supernatant liquid and wash the precipitate with two 1000 ml portions of cold distilled water slowly, dry the resin and powder it.

* **Terebene**: A mixture of hydrocarbons prepared from oil of turpentine and sulphuric acid, used to make paints and varnishes and medicinally as an expectorant and antiseptic.

Characteristic Features It is a light brown to greenish-yellow powder, or small, yellowish, bulky, fragile lumps usually becoming darker in shade on exposure to either heat ($> 25^{\circ}\text{C}$) or light (uv-rays). It has a characteristic faint odour and a bitter acrid taste. It is freely soluble in ethanol, usually with a slight opalescence. It is also soluble in dilute alkaline solution. It is found to be not less than 65% soluble in chloroform and 75% soluble in ether.

Chemical Constituents **Podophyllum** contains 3.5 to 6% of a resin whose active principles are **lignans**, which are essentially C_{18} -compounds related biosynthetically to the **flavonoids**, and are derived by dimerisation of two $\text{C}_6\text{-C}_3$ units. The most important ones present in the **podophyllum resin**, are **podophyllotoxin** (20% in *American Podophyllum*) and in much higher quantum almost upto 40% in *Indian Podophyllum*. Besides, it also contains **α -peltatin** (10%) and **β -peltatin** (5%). It is pertinent to mention here that a host of **lignan glycosides** are also present in the plant, but by virtue of their water-soluble properties, they are almost eliminated during the normal preparation of the resin.



Interestingly, all the three above mentioned chemical constituents are present both in free state and as their respective glycosides. The Indian *Podophyllum* is devoid of **α - and β -peltatins**. The resin also comprise of the closely related **dimethylpodophyllotoxin** and its glycoside; and **dehydropodophyllotoxin**, as well as **quercetin**-a tetra-hydroxy flavonol.

Chemical Tests

1. **Podophyllotoxin** (*active lactone*) present in the resin when dissolved in alkali, cooled to 0°C and subsequently treated with an acid it yields an unstable gelatinous *podophyllic acid*.
2. The resulting **podophyllic acid** when treated with dehydrating agents easily loses a molecule of water and gives rise to **picropodophyllin** (*inactive lactone*), which being an isomer of **podophyllotoxin**.

The resins obtained from the American and Indian *podophyllum* are not quite identical and these two drugs of the trade may be distinguished chemically as given below:

- (a) Prepare an alcoholic extract from *each* resin and filter. Add a few drops of strong solution of cupric acetate 5% (w/v) to each of the above two filtrates. The **American podophyllum** containing α - and β -peltatin produces an instant bright green colouration, while the **Indian podophyllum** (devoid of peltatin) fails this test.
- (b) An alcoholic solution of **Indian podophyllum** resin readily gelatinizes on being treated with alkali hydroxide, while the American resin does not gelatinize. This is due to the fact

that the former contains **podophyllic acid** and it gives the alkali salt of this acid which is gelatinous in nature.

Uses

1. It is used as a drastic but slow-acting purgative.
2. Podophyllotoxin possesses anti-tumour (antineoplastic) properties and may be used in the treatment of cancer.
3. It is invariably prescribed with other purgatives, henbane or belladonna to prevent gripping in infants.

7. Shellac

Synonyms Lacca; Lac.

Biological Sources **Shellac** is the resinous excretion of the *insect Laccifer* (Tachardia) *lacca* Kerr, order *Homoptera* belonging to family: *Coccidae*. The insects usually suck the juice of the tree and excrete 'stick-lac' more or less continuously. The various host trees are, namely: *Butea frondosa* Koen. ex. Roxb. (Family: *Leguminosae*) and *Butea monosperma* (Lam.) Kuntze; *Aleurites moluccana* (L.) Willd. (Family: *Euphorbiaceae*)-**Varnish Tree**; *Ficus benjamina* Linn., (Family: *Moraceae*); *Zizyphus jujuba* (Lam.) (Family: *Rhamnaceae*). However, the *whitest shellac* is produced while the **Kusum tree** is the host *i.e.*, *Schleichera trijuga* (Willd.) (Family: *Sapindaceae*).


Preparation The resin which is stuck on the smaller twigs and branches is normally serapped by means of knives. The resulting resin is subsequently powdered and extracted either with water or with alkaline solution so as to remove the colouring matter. The residual product is dried, melted in narrow bags suspended over a fire. The contents of the bags *i.e.*, the molten shellac, are squeezed out mechanically so as to force the liquid shallac through the cloth on to a previously cleaned surface of tiles to obtain the product as flat cakes. The product may also be obtained as thin sheets by stretching the semi-cooled product on the tiles with the help of a scrapper (or spreader). The thin sheets thus obtained get hardened after cooling and are subsequently broken up to obtain the flakes of shellac for the commercial market.

Characteristic Features **Shellac** is a brittle, yellowish, transparent/translucent sheets or crushed pieces or powder. It does not has any specific odour and taste. Its mp is 115-120°C and d 1.035-1.140. Its solubility in alcohol is 85-95% (w/w) (very slowly soluble); in ether 13-15%; in benzene 10-20% and in petroleum ether 2-6%. It is sparingly soluble in **oil of turpentine**. It is practically insoluble in water, but soluble in alkaline solutions, in aqueous solution of ethanalamines and in borax solutions with slightly purple colouration.


Chemical Constituents The major component of **shellac** is a resin that on being subjected to mild hydrolysis yields a complex mixture of aliphatic and alicyclic hydroxy acids and their polyesters respectively. Interestingly, the composite of the resultant hydrolysate solely depends on the source of shellac and the time of collection.

The major component of the aliphatic fraction is **aleuritic acid**, while the major component of the alicyclic fraction is **shellolic acid**.*

* *Notes Field Tetrahedron*, **26**, 3135 (1970)




Aleuritic Acid
(9, 10, 16-Trihydroxy palmitic Acid)




Shellolic Acid
(10 β , 13-Dihydroxycedr-8-ene-12, 15-dioic acid)

However, it also contains the isomers of **shellolic acid** along with small amounts of **kerrolic acid** and **butolic acid**. The colouring matter is due to the presence of **laccaic acid**, which is water-soluble, as given below:



Laccaic Acid -A: R = CH₂CH₂NHCOCH₃
Laccaic Acid -A: R = CH₂CH₂OH
Laccaic Acid -A: R = CH₂CH(NH₂)COOH



Laccaic Acid-D

Note Laccaic acid-A is the major component while the rest are present in relatively smaller quantities.

Uses

1. It is used chiefly in laquers and varnishes.
2. It is also employed in the manufacture of buttons, sealing wax, cements, inks, grinding wheels, photograph records, paper.
3. It also finds its use in electrical machines and for stiffening hats.
4. It is also used for finishing leather.
5. It is extensively used for coating tablets and confections.
6. It has also been used for preparing *sustained release medicament* formulations.

8. Tar

Synonyms Chair Tar; Pine Tar; Stockholm Tar.

Biological Source Tar is a bituminous liquid obtained from the wood of various species of the natural order *Pinaceae*, such as: *Pinus longifolia* Roxb. [*Pinus roxburghii* Sargent. It is also present in *Pinus elliottii* (Engelm.) belonging to family *Abietaceae* (Slash Pine).

Preparation It is usually obtained by the destructive distillation of the wood cuttings from the various species of *Pinus* as stated above.

Characteristic Features It is dark brown or sometimes black viscous liquid, but its very thin layer on a clean glass plate is almost transparent. It possesses a very strong to moderate specific naphthalene-like odour and has a bitter and pungent taste. It is practically insoluble in water, partially soluble in ethanol, whereas completely soluble in ether, chloroform, volatile oils and fixed oils. It has been observed that when tar is stored for a longer duration, it separates into a layer which is granular in nature by virtue of the fact that minute and critical crystallization of resin acids and catechol take place. **Tar** is found to be *acidic* in reaction.

Chemical Constituents **Tar** contains a good number of chemical constituents in various proportions depending upon the particular species of *Pinus* and its geographic location, such as: hydrocarbons, resin acid, resinous matter, and includes phenols, phenolic ethers, cresols, catechol, methyl cresols, guaiacol, benzene, toluene, xylene and styrene.

Chemical Tests

1. Shake 1g of drug in 20 ml of water and filter:
 - (a) To a portion of the filtrate dip a blue-litmus paper which turns red showing acidic reaction.
 - (b) To another portion of the filtrate add 2-3 drops ferric chloride solutions (0.1%) a red colouration is obtained.

Uses

1. It serves as an expectorant when used in the form of a syrup.
2. Pine tar is frequently employed as antipruritic and antibacterial.
3. It is used largely in ointments externally for the treatment of chronic skin diseases and eczema.

Note This 'Pine Tar' distinctly differs from the 'Coal Tar' in the following aspects, namely:




- (a) **Coal tar** is obtained by the destructive distillation of bituminous coal at a temperature less than 1000°C.
- (b) **Coal tar** mostly contains, benzene, naphthalene, phenols and pyridine.
- (c) It is alkaline to litmus paper.
- (d) It has more disinfectant and irritating properties to pine tar.
- (e) It becomes more viscous on exposure to air.

9. Kava

Synonyms Kava-Kava; Ava-ava; Kawa.

Biological Source It is the dried rhizome and roots of *Piper methysticum* Forst. belonging to the natural order *Piperaceae*.

Chemical Constituents Besides, an appreciable quantity of starch present in it, the drug also comprises of about 5 to 10% of a resin from which six different and closely related *styrilpyrones* have been duly isolated and characterized, namely: **Yangonin, Desmethoxy yangonin, Kawain, Dihydrokawain, Methysticin** and **Dihydromethysticin**.



mp	105-106°C
uv_{max}	In methanol 210, 245, 282 nm (log ε 4.38, 4.44, 2.81)
Soluble in	Acetone, ether, methanol

mp	155-157°C
uv_{max}	In ethanol 360 nm (log ε 4.33) Ethanol (hot), acetone, Ethyl acetate, glacial acetic acid.

mp	132-134°C
uv_{max}	In ethanol 226, 267, 306n (log ε 4.40, 4.14, 3.93) Ethanol (hot), acetone, Ethanol, ether, acetone

Uses

1. The drug (**kava pyrones**) acts as potent centrally acting skeletal muscle relaxants.
2. It also possesses antipyretic and local anaesthetic properties.
3. Its underground parts have been used extensively by the natives (Oceania Islands) in the preparation of an intoxication drink prepared from the roots of this plant.

5.2.8 Oleoresins

Oleoresins are homogenous mixtures of resins and volatile oils. These are, in fact, the vegetative secretions obtained as natural products and composed of resin(s) dissolved in essential oils.

Nevertheless, based on the presence of the relative quantum of volatile oil in the naturally occurring mixture, the **oleoresins** may be either liquid, or semisolid, or solid. In normal practice, there exists a small amount of “natural” exudate from **oleoresin** containing trees attributed to insect damage, traces of broken twigs, and other similar injuries, but the commercial supplies are invariably accomplished by deliberate methodical and artificial incisions made on the bark and even into the wood.

A few important **oleoresins** shall be discussed in the Sections that follow, namely: **Capsaicin, Capsaiba, Male Fern, Ginger, Turpentine.**

1. Capsaicin


Synonyms Axsain; Mioton; Zostrix.

Biological Sources It is the pungent principle present in fruit of various species of *Capsicum*, namely: *Capsicum annuum* L. (Family: *Solanaceae*)-**Paprika, Chili, Sweet Peppers**; *Capsicum frutescens* Linn., [*Synonyms* *C. minimum* Roxb.]-**Bridchilli.**

Preparation **Capsaicin**, the **oleoresin** from capsicum is prepared by extracting the crushed fruit with either hot acetone or hot ethanol by the method of percolation. The solvent *i.e.*, hot ethanol or acetone is evaporated on an electric-water bath in a fume-cupboard. The resulting residue is once again subjected to successive extraction with cold acetone or ethanol until the residue is free from the pungent odour. The solvent is removed and the **capsaicin** collected is not less than 8%.

Characteristic Features It has a monoclinic and rectangular plates, or scales from petroleum ether. Its mp is 65°C, bp_{0.01} 210-220°C (air-bath temperature), uv_{max} 227, 281 nm (ϵ 7000, 2500). It has a burning taste, one part in 100,000 can be detected easily by tasting. It is practically insoluble in cold water. It is freely soluble ethanol, ether, benzene, chloroform and slightly soluble in CS₂.

Chemical Constituents The capsicum contains 8-12% of an oleoresin **capsaicin** and a red colouring principle known as **capsanthin** as given below:



However, the pungency of **capsaicin** is not affected by dilute alkali, but is destroyed almost completely by subjecting it to oxidation with either KMnO₄ or K₂Cr₂O₇.

Uses

1. It is used as a tool in neurobiological research.
2. Pretreatment with capsaicin induces long-lasting desensitization of airway mucosa to various mechanical and chemical irritants.


2. Copaiba

Synonyms Balsam copaiba; Balsam capivi; Jesuit's Balgar.


Biological Source **Copaiba** is the oleoresin obtained from the South American species of *Copaifera* (*Copaiba*) belonging to family: *Leguminosae*.

Preparation The **oleoresin** is collected by incisions made on the trunk of various species of *Copaifera* Linn., (a method similar to **colophony** described under Section 5.2.7).

Characteristic Features It is a transparent, viscid to pale-yellow to brownish-yellow liquid. It has a peculiar odour and bears a nauseating, bitter and acrid taste. Its acid number is 28-95 and d 0.930-0.995. It is practically insoluble in water, but soluble in benzene, chloroform, ether, oils, CS₂, absolute ethanol, petroleum ether and partly soluble in 95% ethanol. It is incompatible with mineral acids, magnesia and water. **Copaiba** is found to contain a volatile oil, **resin acids** (e.g., **capaivic acid** and **illurinic acid**), besides a small quantity of a **bitter principle** and a fluorescent substance. The major constituents of the *volatile oil* are **cryophyllene**, **isocaryophyllene** and that of the *resin acid* is **β -metacapaivic acid** as given below:



Caryophyllene



Isocaryophyllene

Uses

1. It is used in varnishes.
2. It is also employed for removing old oil varnish from oil paintings.
3. It is used in the manufacture of photographic paper.

3. Male Fern

Synonyms Aspidium; Filix mas (B.P.); Male shield-fern; Male fern rhizome.


Biological Source **Male fern** comprises of rhizomes and stipes of *Dryopteris filix-mas* (L.) Schott.; *D. marginata* (Wall.) Christ; *D. odontolma* (Hochst.) C. Chr., and other species of *Dryopteris* belonging to family: *Polypodiaceae*.

Preparation The **male ferns** are prepared by first collecting the rhizomes in the autumn, washed, roots and the stipes except their bases are removed. Finally, the trimmed rhizomes are dried by applying a moderate heat very carefully.


Characteristic Features The rhizomes are dark brown or reddish brown externally and surrounded by **stipes bases**. The stipes bases are covered with membranous scales (ramenta). It has a slight and characteristic odour. It gives initially a sweetish taste, followed by bitter, astringent and nauseous taste. The rhizomes are cylindrical to conical in shape.

Chemical Constituents The main active constituents of male fern are derivatives of **phloroglucinol** and **butyric acid**.


It has been observed that two or more molecules of simple *monocyclic derivatives*, such as: **aspidinol**, **filicinic acid** and **acylfilicinic acid** may get condensed to give rise to *bicyclic derivatives*, for instance; **albaspidin**, **flavaspidic acid** and **filicic acid** as given here under:




Aspidinol




Filicinic Acid




Acylfilicinic Acid



Albaspidin



Flavaspidic Acid



Filicic Acid

BBB : $R_1 = R_2 = C_3 H_7$; mp 172-174°C;

PBB : $R_1 = C_2 H_5$; $R_2 = C_3 H_7$; mp 184-186°C;

PBP : $R_1 = R_2 = C_2 H_5$; mp 192-194°C;

Crystals from
ethylacetate

Note: Filicic acid is a mixture of six homologues, the three main components (i.e., BBB, PBB, PBP) which are obtained by recrystallization from ethyl acetate.

Filicin is the lactone of **filicic acid**, which occurs as granular sediment in all male fern extracts and may be obtained by collecting it and subsequently washing with ether-ethanol mixture (1 : 1). The insoluble portion is dissolved in ethyl acetate or methanol and allowed to crystallize slowly when it yields yellow flakes.

Uses

1. Male fern **oleoresin** is an anthelmintic, specifically a taeniafuge.
2. It is also used as its extract for the expulsion of tapeworms.



4. Ginger

Synonyms Gingerin.



Biological Source It is the **oleoresin** obtained by the method of percolation of the powdered rhizomes of *Zingiber officinale* Roscoe, belonging to the Family: *Zingiberaceae*.

Preparation The rhizomes are sliced, dried and powdered. The powdered ginger is extracted either with acetone or ether or ethylene dichloride by the method of cold percolation repeatedly till the gingerin is no longer present in the marc. The solvent is removed by distillation under reduced pressure. Ethanol gives the max yield of the oleoresin. The average yield of the **oleoresin** is 6.5% but it may range between 3.5 to 9.0% based solely upon the source of the plant product and to a great extent on the technique adopted in the course of preparation.

Characteristic Features It is a dark brown, aromatic and pungent viscous liquid.



Chemical Constituents Ginger contains volatile oil (1-3%), which comprises of **zingiberene**, **α -curcumene**, **β -sesquiphellandrene** and **β -bisabolene**. **Zingiberene** (a) has two chiral centres. The *acyclic chiral centre* has been stereochemically related to that in (+)-**citronellal**, and the *cyclic chiral centre* to that in (-)-**phellandrene**. Hence, (-)-**zingiberene** has the absolute configuration (b). The **oleoresin** contains the pungent **gingerols** and **shogaols**.



Uses

1. It is used as a flavouring agent, carminative, aromatic and stimulant to gastrointestinal tract (GIT).
2. **Ginger** finds its wide applications in soft drinks, beverages, ginger beer and wine.
3. It is extensively used for culinary purposes in ginger-bread, biscuits, puddings, cakes, soups and pickles.

5. Turpentine

Synonyms Gum turpentine; Gum thus.

Biological Source **Turpentine** is the **oleoresin** obtained from *pinus palustris* Miller and from other species of *Pinus*, belonging to the natural order *Pinaceae*.


Preparation **Turpentine** is usually collected from the slash pine *i.e.*, *Pinus elliottii* Engelmann var. *elliotti*, and *Pinus palustris* Miller, which grow in abundance in the Northern Florida, Georgia, and North and South Carolina. However, the yield of **turpentine** exclusively depends on the treatment

and the size of the tree. If proper skill and expertise are practiced the pine trees may yield turpentine for 15 to 20 years at a stretch.

The **oleoresin** is normally secreted in the ducts that are situated almost beneath the cambium in the sapwood. In spring the bark is neatly cut from the tree with the help of a long-handled cutting knife known as the “*bark-hack*”. After the removal of the chipped bark, the freshly exposed surface is quickly sprayed with a solution of 50% (w/w) sulphuric acid.* The flowing **oleoresin** is guided by galvanized metal gutters right into the various containers tied close to the tree-trunk. The thick-liquid thus collected is removed as **turpentine** by pot-still distillation periodically.

Characteristic Features The **gum turpentine** is an yellowish, opaque, sticky mass having a characteristic odour and taste. It is almost insoluble in water, but soluble in ether, ethanol, chloroform and glacial acetic acid.

Chemical Constituents The **gum-turpentine** when subjected to steam-distillation yields 15 to 30% of a volatile oil known in the trade as “**turpentine oil**”. It contains mainly the terpenes, such as: **dextro- and laevo- α -pinene, β -pinene and camphene**.



Uses

1. It is employed externally as a counterirritant.
2. It is also used as a rubefacient.
3. It is used as a constituent of stimulating ointments.
4. It is employed industrially as an insecticide.
5. It is used as a solvent for waxes.
6. It is utilized extensively in the production of synthetic camphor.
7. It is used in making various types of polishes, such as: shoe polish, furniture polish and stove polish.

5.2.9 Oleo-Gum-Resins

The **oleo-gum-resins** are the naturally occurring mixture of resin, gum, volatile oil, and mostly small quantities of other substances.

* Acid treatment collapses the thin-walled parenchymal cells which line the resin ducts. Thus, the duct channels get enlarged thereby allowing a faster uninterrupted flow of oleoresin and minimising the chances of hardened secretions blocking the outlets.

There are some potent **oleo-gum-resins** which exhibit remarkable medicinal values. A few such drugs shall be discussed briefly here under: **Asafoetida**; **Ammoniacum**; **Turmeric**; **Myrrh**; **Indian Bdellium** etc.

1. Asafoetida

Synonyms Asafetida; Asant; Devil's dung; Food of the Gods; Gum Asafoetida.


Biological Sources **Asafoetida** the oleo-gum-resin is obtained as an exudation of the decapitated rhizome on roots of *Ferula assafoetida* L.; *Ferula foetida* Regel, and some other species of *Ferula*, belonging to the nature order *Umbelliferae*.

Preparation **Asafoetida** is generally present as a milky liquid in the large schizogenous ducts and lysigenous cavities. However, these ducts and cavities are located more intensively in the cortex region of the stem and root. The drug is obtaining chiefly from the stem.


The fully grown plants are usually cut down to the crown region during the spring. The exposed surface is protected by a dome-like covering made up of twigs and leaves. After about a month, the hardened resinous substance is collected by scrapping. Likewise, the stems are also cut off and thereby additional collections of **asafoetida** are made frequently at an interval of 10 days unless and until the exudation ceases to ooze. Furthermore, it is also collected from the root by exposing its crown and excising the stem. The oleo-gum-resin exudes from the cut surface of the root and the former is collected soon after it gets dried. Thus, the entire collection of **asafoetida** from the various portions of the plant are mixed together and dried in the sun.

Characteristic Features The drug occurs normally as soft mass or irregular lumps or 'tears' or agglomeration of tears. The tears are brittle and tough. **Asafoetida** has a strong, alliaceous, persistent garlic-like odour and having a bitter acrid taste. This **oleo-gum-resin** when triturated with water it gives a milky emulsion.


Chemical Constituents **Asafoetida** contains volatile oil (8-16°C) gum (25%) and resin (40-60%). The volatile oil essentially consists of some organic sulphides solely responsible for attributing the characteristic garlic-like odour. The resin consists of **notannol**, **asaresinotannol** *i.e.*, the **resin alcohols**, which are present partially in the free state and partially in the combined form with **ferulic acid**. It also contains **umbellic acid** and **umbelliferone**; the latter is found combined with ferulic acid, but it gets generated on being treated with dilute HCl.



trans-Ferulic Acid



Umbellic Acid



Umbelliferone

There are *three* sulphur-compounds that have been isolated from the **asafoetida resin**, namely:

- (a) 1-Methylpropyl-1-propenyl disulphide,
- (b) 1-(Methylthio) propyl-1-propenyl disulphide, and
- (c) 1-Methylpropyl-3-(methylthio)-2-propenyl disulphide.

Interestingly, the latter two (*i.e.*, 'b' and 'c') have pesticidal properties.

Chemical Tests

1. It forms an instant milky-white emulsion when triturated with water owing to the presence of gum.
2. The freshly fractured surface when treated with a drop of sulphuric acid (conc.), it gives rise to a reddish-brown colour which on being washed with water changes to violet colouration.
3. Likewise, when the freshly fractured surface is treated with nitric acid (50%), it produces a green colour readily.
4. Boil 1 g asafoetida powder with HCl (50%), filter and make the filtrate strongly alkaline with NH₄OH (conc.), it gives a blue fluorescence. It is also known as the **Umbelliferone Test**.

Uses

1. It is abundantly used in India and Iran as a common condiment and flavouring agent in food products.
2. It is also an important ingredient in *Worcestershire Sauce*.
3. It is used as a repellent [2% (w/v) suspension] against dogs, cats, deer, rabbits etc.
4. It is used seldomly as an antispasmodic, carminative, expedorant and laxative.
5. It is still employed in veterinary externally to prevent bandage chewing by dogs.
6. It is also used as a powerful nerving stimulant especially in nervous disorders related to hysteria.

2. Ammoniacum

Synonym Gum ammoniac.

Biological Source It is a **oleo-gum-resin** exuded from the flowering and fruiting stem of *Dorema ammoniacum*, D. Don. and probably other species belonging to family: *Umbelliferae*.

Preparation The exudation of the milky-secretions obtained in the form droplets is usually caused by the *beetles* that puncture the fruiting stem of *D. ammoniacum*. While quite a few of these milky-droplets get hardened on the stem itself, and the rest falls on the ground. The solidified **oleo-gum-resins** are scrapped from the stem with a plant-knife and also collected from the droppings on the ground.

Characteristic Features The drug has an irregular, rounded tears, that are yellowish or browish outside and whitish from within; these are generally brittle when cold, but get softened on warming. It is also found, in the form of mass *i.e.*, agglomeration of small droplets. The mass is found to be darker in colour and less homogeneous.

It has a peculiar odour, slightly sweetish, bitter and somewhat acrid taste. The physical characteristics are: mp 45-55°C and d 1.207. Its acid number varies between 60-80, whereas the saponification number ranges between 97-114. It is partly soluble in water, ethanol, ether, vinegar, or alkaline solution. It readily forms an emulsion with water.

Chemical Composition **Ammoniacum**-the **oleo-gum-resin** consists of volatile oil (0.1-1.0%), resin (65-70%), gum (20%), moisture (2-12%), insoluble residue (3.5%) and ash (1%). **Ammoniacinol**, a phenolic substance is the main constituent of the resin, which is a colourless crystal, mp 110°C. It also contains traces of salicylic acid.

Chemical Tests

1. **Ammoniacum** when triturated with water, it forms a white emulsion.
2. A portion of the above emulsion when treated with a solution of chlorinated soda gives a deep orange-red colouration.
3. A portion of the emulsion on being treated with a potash solution yields a yellow colour.
4. A portion of the emulsion when treated with a 0.1% (w/v) solution of FeCl_3 , it gives an instant violet colouration due to the presence of traces of salicylic acid.

Uses

1. It is an important ingredient of porcelain cements.
2. It is a stimulant, and secreted by the bronchial mucous surface, thereby disinfecting the secretions.
3. It is used in plaster-of-paris (POP) plasters as a stimulant to the skin.
4. It is also used as a disinfectant expectorant in chronic bronchitis amalgamated with excessive discharge.

3. Turmeric

Synonyms Curcuma; Indian Saffron; Tumeric.


Biological Source **Turmeric** is obtained from the rhizome of *Curcuma longa* Linn. (*Curcuma domestica* Valetton) belonging to the natural order *Zingiberaceae*.

Preparation The plant is normally harvested after 9-10 months when the lower leaves start becoming yellow. The rhizome is carefully dug out from the soil with a blunt knife without damaging it. The fibrous roots are discarded. The raw green **turmeric** is cured and processed by boiling the rhizomes with water for a duration ranging between 12-14 hours. Subsequently, the cooked rhizomes are dried in the sun for 5-7 days. Cooking process helps in achieving *two* objects, namely:

- (a) Gelatinization of starch, and
- (b) Yellow colouration, due to *curcumin*, spreads over the entire rhizome.

Characteristic Features **Turmeric** has an aromatic pepper-like but somewhat bitter taste. It gives curry dishes their characteristic yellowish colouration.

Chemical Constituents It contains volatile oil (5-6%), resin and substantial quantity of **zingiberaceous starch grains**. The marked and pronounced yellow colour in **turmeric** is due to the presence of **curcuminoids** which essentially contains **curcumin** as given below:




Curcumin

(Orange-yellow crystalline powder, mp 183°C)

The **curcuma oil*** obtained from turmeric contains (\pm)-**ar-turmerone** as given below:

* H. Rupe *et. al. Helv. Chim. Acta*, **17**, 372 (1934).



The volatile oil contains a host of chemical substances, such as: ***d*- α -phellandene**, ***d*-sabinene**, **cineol**, **borneol**, **zingiberene**, and **sesquiterpenes**.

Turmeric also contains some other chemical constituents, namely: ***p,p*-dihydroxy dicinnamoylmethane**; ***p*- α -dimethyl benzyl alcohol**; ***p*-hydroxy-cinnamoylferuloylmethane**; **1-methyl-4-acetyl-1-cyclohexene**; and **caprylic acid**.

Chemical Tests

1. **Turmeric** powder when triturated with alcohol it imparts a deep yellow colour to the resulting solution.
2. The powdered drug when treated with sulphuric acid it imparts a crimson colour.
3. The aqueous solution of turmeric with boric acid gives rise to a reddish-brown colouration which on subsequent addition of dilute alkali changes instantly to greenish-blue.
4. **Turmeric** powder when reacted with acetic anhydride and a few drops of concentrated sulphuric acid (36 N), it readily gives a violet colouration. Interestingly, the resulting solution when observed under the ultraviolet light (preferably in a **uv-chamber**), it exhibits an intense red fluorescence, which is due to the presence of **Curcumin**.

Uses

1. It is extensively used across the globe as a condiment as curry powder.
2. It is employed as a colouring agent for ointments.
3. It is used medicinally as a tonic, as a blood purifier, as an anthelmintic and finally as an aid to digestion.
4. It is used extennally in the form of a facial cream to improve complexion and get rid of pimples.
5. A small quantity of turmeric when boiled with milk and sugar; it helps to cure common cold and cough symptoms.

4. Myrrh


Synonyms Gum Myrrh, Myrrha.

Biological Source **Myrrh** is an **oleo-gum-oresin** obtained from the stem and branches of *Commiphora obyssinica* (Berg) Engler or from other species of *Commiphora* belonging to family *Burseraceae*.

Preparation The plants usually exude yellow coloured resin after proper incisions are made in the bark of a tree. It gradually hardens and becomes dark or reddish-brown in appearance. The mass is collected by the native tribals of Somalia for trading.

Characteristic Features **Myrrh** normally occurs either in the form of isolated irregular, rounded tears of 2.5 cm in diameter or as masses duly formed by the agglomeration of these tears. The tears are dull, rough and reddish-brown in appearance. It has a strong aromatic odour and possesses an acrid, bitter taste.

Chemical Constituents Myrrh contains volatile oil (7-17%), resin (20-25%), gum (57-61%), and bitter principle (3 to 4%). The volatile oil consists of **eugenol**, **m-cresol** and **cuminaldehyde**. The resin is found to consist of a mixture of **α -**, **β -**, and **γ -commiphoric acids (resin acids)**. Besides, it also contains two phenolic resins **α -** and **β -heerabomyrrholic acids** which are ether insoluble. The oleo-gum-resin yields alcohol-soluble extract not less than 30%. It also contains phenolic compound such as: **pyrocatechin** and **protocatechuic acid**. The crude alcohol-insoluble fraction *i.e.*, 'gum', comprises of protein (18%) and carbohydrate (64%) made up of **arabinose**, **galactose** and **glucuronic acid**. However, the gum is found to be associated with an oxidase enzyme.



Chemical Tests

1. **Myrrh** when triturated with water produces a yellow-emulsion.
2. When **myrrh** (0.1 g) is triturated with 0.5 g of pure washed sand (SiO_2) in the presence of ether, filtered and evaporated on an electric water-bath, it forms a thin film of violet colour on being exposed to bromine vapours in a closed desiccator.

Uses

1. It is used chiefly in perfumes and incense.
2. It is frequently employed as an antiseptic and stimulant.
3. **Myrrh** acts as an astringent to the mucous membrane and hence it finds its application in oral hygiene formulations, such as: gargles, mouth-washes.
4. It is also used as a carminative.

5. Indian Bdellium

Synonym Scented bdellium.


Biological Sources **Indian bdellium** is the oleo-gum-resin obtained from the bark of the naturally occurring plant *Commiphora mukul* Engler., *Balsamodendron mukul* Hook. ex. Stocks., and *Commiphora weightii* (Arn) Bhand, belonging to family: *Burseraceae*.

Preparation The **oleo-gum-resin Indian bdellium** is obtained by the incision made on the bark and the exudates are collected. Each fully grown plant produces about 0.5 to 1 kg of the product which is normally collected from January through March every year.

Characteristic Features The **oleo-gum-resin** from **Indian bdellium** has a brown to pale yellow or sometimes dull green colour. It has an agreeable balsamic and aromatic odour with a typical bitter taste. The drug is usually obtained as irregular mass, rounded or agglomerated cluster of tears. The tears are found to be transparent, having a waxy surface and quite brittle in nature. It is sticky in touch and has a fractured surface. It is partially alcohol soluble; but when triturated with water it usually gives rise to a white emulsion.

Chemical Constituents This **oleo-gum-resin** mostly comprises of resin (60%), gum (30%), volatile oil (1-1.5%) moisture (5%) and foreign organic substances (3-4%). The volatile oil fraction contains

various terpenes, such as: **β -murcene**, **dimyrcene**, **polymyrcene**, **caryophyllene** and **isocaryophyllene** (Section 5.2.9.2).



5.2.10 Balsams

Balsams are the resinous mixtures that essentially contain large quantum of benoic acid, cinnamic acid or both, or esters of these organic aromatic acids.

A galaxy of typical examples of naturally occurring *balsams* will be discussed in the sections that follow, namely: **Storax**; **Peruvain Balsam**; **Tolu Balsam**; and **Benzoin**.

1. Storax

Synonyms Styrax; Sweet oriental gum; Levant Storax; Purified or prepared Storax; American Storax; Liquid Storax;

Biological Source **Storax** is the *balsam* obtained from the trunk of *Liquidamber orientalis* Mill., termed as **Levant Storax**, or of *L. styraciflua* L., known as **American Storax** belonging to the natural order *Hamamelidaceae*.

Preparation The natural balsam *storax* is a pathological product formed as a result of injury caused to the plant. It generally, exudes into the natural pockets between the bark and the wood and may be located by exerscences on the outside of the bark. These pockets, that may contain upto 4 kg of the balsam, *are* conveniently tapped with the help of strategically positioned gutters, and the product is ultimately allowed to fill into containers. The crude storax, thus collected, is further purified by dissolving in ethanol, filtration and subsequent evaporation of the solvent to obtain the pure storax.

Characteristic Features The **balsam storax** is a semiliquid grayish, sticky, opaque mass (*Levant Storax*), or a semisolid sometimes solid mass softened by gentle warming (*American Storax*). In general, **storax** is transparent in thin layers, possesses a characteristic agreeable balsamic taste and odour. It is, however, denser than water. It is almost insoluble in water, but completely soluble in 1 part of warm ethanol, ether, acetone and CS₂.

Chemical Constituents **Storax** contains the following chemical compounds, namely: **α -and β -storesin** and its **cinnamic ester** (30-50%), **styracin** (5-10%); **phenylpropyl cinnamate** (10%); **free-cinnamic acid** (5-15%); levorotatory oil (0.4%); small amounts of **ethyl cinnamate**, **benzyl cinnamate**, traces of **vanillin** and **styrene** (C₆H₅CH=CH₂).

Besides, *Levant storax* contains free **storesinol**, **isocinnamic acid**, **ethylvanillin**, **styrogenin**, and **styrocamphene**.

In addition to these, **American Storax** contains **styaresin** (*i.e.*, -cinnamic acid ester of the alcohol **styresinol**, an isomer of **storesinol**) and **styresinolic acid**. It also yields upto 7% of a dextrorotatory volatile oil, styrol and traces of vanillin.

Chemical Tests

1. **Benzaldehyde Test**—Treat 1 g of **storax** with 5 ml of $K_2Cr_2O_7$ solution (10% w/v) followed by a few drops of concentrated sulphuric acid (36 N) it produces benzaldehyde, which may be detected easily as the odour of bitter almonds.
2. Mix 1 g of **storax** with 3 g of pure sand (SiO_2) and 5 ml of $KMnO_4$ solution (5% w/v) and heat it gently. It gives a distinct odour of benzaldehyde.

Uses

1. It is used in fumigating pastilles and powders.
2. It finds its application in perfumery.
3. It is employed as an imbedding material in microscopy.
4. It is used as an expectorant, antiseptic and stimulant.
5. It is employed as a preservative for fatty substances *e.g.*, lard and tallow.
6. It is also used as a flavouring agent for tobacco.
7. It is a vital ingredient of “**Compound Benzoin Tincture**”.

2. Peruvian Balsam

Synonyms Balsam Peru; Indian balsam; Black balsam; China oil; Honduras balsam; Surinam balsam.

Biological Source **Balsam Peru** is obtained from *Toluiifer pereiare* (Klotzsch) Baill. (*Myroxylon pereiare* Klotzsch) belonging to family: *Leguminosae*.

Preparation **Peruvian Balsam** is a pathological product and is obtained usually by inflicting injury to the trees. Most of the world's commercial supply comes from El Salvador, although some is also produced in Honduras.


It is prepared by beating the stems of the trees with mallet. After a week the injured areas of the stem are scorched so as to separate the bark from the stem and after a similar duration the bark is peeled off completely. The desired balsam starts exuding freely from all the exposed surfaces, which are then covered carefully with cloth or rags to absorb the exuding balsam. The cloth or rags that are completely soaked with the balsam is then removed and boiled with water in a large vessel slowly. Thus, the balsam gets separated and settles at the bottom of the vessel. The supernatant layer of water is removed by decantation and the residual balsam is dried and packed in the containers.

Characteristic Features It is a dark brown, viscid liquid having a pleasant aromatic odour. It has a peculiar warm bitter taste and persistent aftertaste which resembles like vanilla. The **Balsam Peru** is transparent in thin films. It does not harden on being exposed to atmosphere. It is brittle when cold. It is almost insoluble in water and petroleum ether but soluble in ethanol, chloroform and glacial acetic acid.

Chemical Constituents Peruvian balsam contains free benzoic and cinnamic acids (12-15%); benzyl (40%); esters of these acids (5.2-13.4% **cinnamein**); and volatile oil (1.5-3%).


The fragrant volatile oil contains toluene, styrol, benzoic and cinnamic acids.

It also contains total **balsamic acids**, which is calculated on the basis of dry alcohol-soluble matter ranging between (35-50%).



Cinnamein
(Benzyl Cinnamate)

The **resins esters** (30-38%) are chiefly composed of **peruresinotannol cinnamate** and benzoate, vanillin, free cinnamic acid and **peruvicol** (or **nerolidol**).



Peruvicol (Nerolidol)

Uses

1. **Peru balsam** is a local protectant and rubefacient.
2. It also serves as a parasiticide in certain skin disorder.
3. It is used as an antiseptic and **vulnerary*** and is applied externally either as ointment or alone or in alcoholic solution.
4. It acts as an astringent to treat hemorrhoids.

3. Tolu Balsam

Synonyms Thomas balsam; Opobalsam; Resin Tolu; Balsam of Tolu.

Biological Source **Tolu Balsam** is a balsam obtained from *Toluidra balsamum* L., (*Myroxylon toluiferum* H.B.K.), belonging to family: *Leguminosae*. It is also obtained from *Myroxylon balsamum* (Linne') Harms. Family: *Fabaceae*.

Preparation **Tolu Balsam** is considered to be a pathological product produced in the new wood formed as a result of inflicted injury. For its preparation, it is an usual practice to make 'V' shaped incisions deep into the body of the main trunk. The exudate thus produced is collected either in cups or gourds held strategically just at the base of each incisions. **Balsam of Tolu** is collected from these cups, mixed and packed in air-tight sealed tins.

Characteristic Features It is a yellowish-brown or brown semifluid or nearly solid resinous mass. It has a characteristic aromatic vanilla-like odour and slightly pungent taste. It is usually brittle when cold. It is found to be transparent in thin layers, and shows numerous crystals of cinnamic

* **Vulnerary**: A folk remedy or herb to promote wound healing.

acid. It is almost insoluble in water and petroleum ether, but freely soluble in ethanol, benzene chloroform, ether, glacial acetic acid and partially in CS₂ or NaOH solution.

Chemical Constituents The drug contains **resin esters** (75-80%) *viz.*, **toluresinotannol cinnamate** along with a small proportion of the benzoate; volatile oil (7-8%)-containing chiefly **benzyl benzoate**; free cinnamic acid (12-15%); free **benzoic acid** (2-8%); **vanillin** and other constituents in small quantities. It also contains **cinnamein** (5-13%).

Chemical Tests

1. An alcoholic solution of **Tolu Balsam** (0.2% w/v) where treated with a FeCl₃ solution (0.5% w/v), the appearances of a green colour takes place.
2. Treatment of 1 g of the drug with 5 ml of 10% w/v KMnO₄ solution when subjected to gentle heating yields benzaldehyde.

Uses

1. It is used extensively in perfumery, confectionery and chewing gums.
2. It is used widely as an expectorant in cough mixture.
3. It also finds its application as an antiseptic in the form of its tincture.

4. Benzoin

Synonyms Bitter-almond-oil camphor.

Biological Source **Benzoin** in the **balsamic resin** obtained from *Styrax benzoin* Dryander and *Styrax paralleloneurus* Perkins, generally known in trade as **Sumatra Benzoin**; whereas, *Styrax tonkinensis* (Pierre) Craib ex Hartwich, or other species of the section Anthostyrax of the genus *Styrax*, known commonly in the trade as **Siam Benzoin** both belong to the family: *Styraceae*.

Preparation **Benzoin** is also a pathological product that is obtained by incising a deep-cut in the bark. It has been observed that after a span of about eight weeks, the exudating **balsamic resin** tends to become less sticky in nature and firm enough to collect. The entire exudate is usually collected in *two* stages, namely:

Stage 1: *First tapping*-yields *almond tears*, and

Stage 2: *Second tapping*-yields a more fluid material.


Characteristic Features

Sumatra Benzoin: It is pertinent to mention here that in pharmacy, only the **Sumatra Benzoin** is used. It occurs as blocks or irregular masses of tears having variable sizes usually imbedded either in an opaque or translucent matrix. It is rather brittle, and from within the tears are milky white in appearance. It generally becomes soft when warmed and gritty when chewed. The matrix is grayish brown to reddish in colour. Its taste is quite agreeable, balsamic and resembles to that of **storax**. It has a resinous and aromatic taste.

Siam Benzoin: The smaller tears of **Siam Benzoin** are darker in colour. It occurs largely in separate concavo-convex tears which are yellowish brown to rusty brown externally, whereas milky white internally. The tears are fairly brittle but normally become soft and plastic like on being chewed. It has a vanilla-like fragrance.

Chemical Constituents The chemical constituents of the *two* types of **Benzoin** are given below:

- (a) **Sumatra Benzoin:** It contains free balsamic acids, largely cinnamic acid (10%), benzoic acid (6%)-along with their corresponding ester derivatives. Besides, it also contains **teriterpene acids**, namely: **19-hydroxyoleanolic** and **6-hydroxyoleanolic acids, cinnamyl cinnamate, phenyl propyl cinnamate, phenylethylene** and lastly the traces of vanillin. It yields not less than 75% of alcohol-soluble extractives.
- (b) **Siam Benzoin:** It chiefly comprises of **coniferyl benzoate** (60-70%), benzoic acid (10%), **triterpene siaresinol** (6%) and traces of vanillin. It yields not less than 90% of alcohol-soluble extractives.



Coniferyl Benzoate

Chemical Tests

1. When 0.5 g of **Sumatra Benzoin** powder is warmed with 10 ml of KMnO_4 solution (5% w/v) in a test tube, a faint and distinct odour of benzaldehyde is developed. **Siam Benzoin** gives a negative test.
2. When 0.2 g of **Siam Benzoin** powder is digested with 5 ml of ether for 5 minutes and filtered; 1 ml of the filtrate is poured into a clean china-dish containing 2-3 drops of concentrated H_2SO_4 and mixed carefully, a deep purplish red colouration is developed instantly. **Sumatra Benzoin** gives a negative test.

Uses

1. Compound benzoin tincture is frequently employed as a topical protectant.
2. It is valuable as an expectorant when vapourized.
3. It finds its usage as a cosmetic lotion usually prepared from a simple tincture.
4. **Siam Benzoin** has been proved to be a better preservative for lard than the **Sumatra Benzoin**.

FURTHER READING REFERENCES

1. Agurell, S., Dewey, W.L., Willette, R.E., eds.: **The Cannabinoids: Chemical, Pharmacologic and Therapeutic Aspects**, Academic Press Inc., Orlando, Florida, (1984).
2. Atal, C.K. and Kapoor, B.M., eds., '**Cultivation and Utilization of Medicinal Plants**', Regional Research Laboratory, Jammu-Tawi, India, (1982).
3. Brown, R.G., and Brown, M.L., '**Woody Plants of Maryland**', Port City Press, Baltimore, (1972).
4. Council of Scientific and Industrial Research, **The Wealth of India**, 11 vols., New Delhi, (1948-1976).

5. Cutler, S.J., and Cutler, H.G. eds, **'Biologically Active Natural Products: Pharmaceuticals'**, CRC Press, London, (1999).
6. Duke, J.A., **'Handbook of Legumes of World Economic Importance'**, Plenum Press, New York, (1981).
7. Duke, J.A., **'Handbook of Medicinal Herbs'**, CRC Press, New York, (2001).
8. Dewick, P.M., **'Medicinal Natural Products: A Biosynthetic Approach'**, John Wiley & Sons, Ltd., 2nd, edn., New York, (2001).
9. Earle, F.R. and Jones, Q., *Analyses of Seed Samples from 113 Plant Families*, *Econ. Bot.* **16** (4), (1962).
10. Erichsen-Brown, C., **'Use of Plants for the Past 500 Years'**, Breezy Creeks Press, Aurora. Can., (1979).
11. Guenther, E., **'The Essential Oils'**, Vol. 1-6, Van Nostrand, New York, (1948-1952).
12. Hagen-Smit, A.J., **'Progress in the Chemistry of Natural Products'**, Vol. 1-12, Springer, Vienna, (1955).
13. Harborne, J.B., Tomas-Barberan, F.A., eds.: **'Ecological Chemistry and Biochemistry Plant Terpenoids'**, Oxford, Clarendon Press, London, (1991).
14. Herbert RB: *The biosynthesis of plant alkaloids and nitrogenous microbial metabolites*, *Nat. Prod. Rep.* **18**, 50-65, 2001; **Earlier Reviews:** 1999, **16**, 199-208; 1997, **14**, 359-372.
15. Hibino S., and Choshi T., *Simple indole alkaloids and those with a nonrearranged monoterpenoid unit*. *Nat Prod Rep.* **18**, 66-87, **Earlier Review:** Lounasmaa M and Tolvanen A, **17**, 175-191, 2000.
16. Irvine, F.R., **'Woody Plants of Ghana'**, Oxford University Press, London, (1961).
17. Keys, J.D., **'Chinese Herbs: Their Botany, Chemistry and Pharmacodynamics'**, Chas. E. Tuttle, Tokyo, (1976).
18. Kirtikar, K.R., Basu, B.D., and I.C.S. **'Indian Medicinal Plants'**, Vol. 1-4, 2nd. edn. reprint, Jayyed Press, New Delhi, (1975).
19. Kutchan T.M., **Molecular Genetics of Plant Alkaloid Biosynthesis**, In: **The Alkaloids, Chemistry and Pharmacology** (ed. Cordell GA). Vol. 50, Academic, San Diego, pp 257-316, 1998.
20. Misra N., Luthra R., Singh K.L., and Kumar S., **Recent Advances in Biosynthesis of Alkaloids**, In: **Comprehensive Natural Products Chemistry**, Vol. 4, Elsevier, Amsterdam, pp 25-59, 1999.
21. Purseglove, J.W., Brown, E.G., Green, C.L., and Robbins, S.R.J., **'Spices'**, Vol. 1-2, Longman, London, (1981).
22. Ramstad, E., **'Modern Pharmacognosy'**, McGraw Hill, London, (1956).
23. Robinson, R., **'The Structural Relation of Natural Products'**, Oxford-Clarendon Press, London, (1955).
24. Stöckigt J. and Ruppert M., **Striclosidine—the biosynthetic key to monoterpenoid indole alkaloids**, In: **Comprehensive Natural Products Chemistry**, Vol. 4, Elsevier, Amsterdam, pp 109-138, 1999.
25. Teranishi, R., Buttery, R.G., Sugisawa, H., Eds., **'Bioactive Volatile Compounds from Plants'**, Washington, DC, American Chemical Society, (1993).
26. Tyler, V.E., **'The Honest Herbal—A Sensible Guide to the Use of Herbs and Related Remedies'**, George F. Stickely, Philadelphia, (1982).

6

Phenylpropanoids

- Introduction
- Classification
- Biosynthesis of Phenylpropanoids
- Further Reading References

6.1 INTRODUCTION

Phenylpropanoids represent a large conglomerate of naturally occurring phenolic compounds essentially derived from the aromatic amino acids **phenylalanine** and **tyrosine** or in certain specific instances the intermediates obtained from **Shikimic Acid Biosynthetic Pathway**. In other words, these compounds comprise of a phenylring to which is attached a 3C-side chain; and may also contain one or more C₆—C₃ residues.

Interestingly, the unique combination of the phenyl-propane side chain (*i.e.*, 3C-atom) evidently present in '**phenylpropanoids**' are absolutely devoid of nitrogen atom, which is observed to be in contradiction to such other vital class of natural products, namely: **alkaloids**, **cyanogenic glycosides**, and **glucosinolates**. Obviously, the **phenylpropanoids** are distinctly phenolic in character by virtue of the presence of one or several hydroxyl groups attached to the aromatic ring (C₆ H₆), they are more often known among the phytochemists as '**plant phenolics**'.

6.2 CLASSIFICATION

The **phenylpropanoids** may be classified on the basis of their basic chemical moieties as enumerated below:

- (i) Hydroxycinnamic Acids
- (ii) Phenylpropenes
- (iii) Coumarins
- (iv) Abridged phenylpropanoids
- (v) Biphenylpropenoid derivatives
- (vi) High molecular weight phenylpropanoids

The above different categories of compounds belonging to the **phenyl propanoids** shall be discussed separately with the help of certain important examples of natural products in a systematic manner in the sections that follow:


6.2.1 Hydroxycinnamic Acids

The typical examples of **hydroxycinnamic acids** are, namely: *p*-coumaric acid, caffeic acid, ferulic acid, and sinapic acid, which shall be enumerated in the sections that follow:

6.2.1.1 Para-Coumaric Acid

Synonym *p*-Hydroxycinnamic acid.

Chemical Structure



para-Coumaric Acid


Biological Sources It is present in a variety of medicinal plant, namely: *Aloe barbadensis* Mill. (*Liliaceae*)-**Barbados Aloe**, Mediterranean Aloe, Curacao Aloe; *Euphorbia lathyris* L. (*Euphorbiaceae*)-**Mole plant**, Petroleum plant **Caper spurge**; *Hedra helix* L. (*Araliaceae*)-Ivy; *Hura crepitans* L. (*Euphorbiaceae*)-**Sandhox Tree**; *Malus sylvestris* Mill. (*Rosaceae*)-**Apple**; *Melilotus officinalis* Lam. (*Fabaceae*)-**Yellow Sweetlover**; *Trifolium pratense* L. (*Fabaceae*)-**Red Clover, Pavine Clover, Cowgrass**.

Characteristic Features It occurs as needles having mp 210-213°C. It may be crystallized in its anhydrous form from concentrated hot aqueous solution, but as the monohydrate from dilute aqueous solution on gradual chilling. Its uv_{max} (in 95% ethanol) are 223 and 286 nm (ϵ 14,450, 19,000). It is practically insoluble in ligroin and benzene, slightly soluble in cold water, and freely soluble in ethanol, ether and hot water.

6.2.1.2 Caffeic Acid

Synonym 3, 4-Dihydroxycinnamic acid.

Chemical Structure




Caffeic Acid

Biological Source It occurs widely in more than twenty different species of plants as detailed below: *Aconitum napellus* L. (*Ranunculaceae*)-**Aconite, Monkshood, Blue Rocket**; *Arctium lappa* L. (*Asteraceae*)-**Edible Burdock, Great Burdock, Lappa**; *Arnica montana* L. (*Asteraceae*)-**Mountain Tobacco, Leopard's-bane**; *Cinnamomum camphora* (L.) J.S. Presl. (*Lauraceae*)-**Camphor, Hon-Sho**; *Citrullus coloeynthis* (L.) Sehrad. (*Cucurbitaceae*)-**Colocynth, Bitter Apple, Wild Gourd**; *Clematis vitalba* L. (*Ranunculaceae*)-**Traveler's Joy**; *Coniun maculatum* L. (*Apiaceae*)-Hemlock; *Convalaria majalis* L. (*Liliaceae*)-**Lily of the Valley**; *Crataegus oxycantha* L. (*Rosaceae*)-Howthorn; *Digitalis purpurea* L. (*Serophulariaceae*)-**Common Foxglove, Digitalis**; *Equisetum hyemale* L.

(*Equisetaceae*)-Shavegrass, **Great Scouring Rush**; *Euphorbia pulcherrima* Wild ex Klotsch (*Euphorbiaceae*)-**Poinsettia**; *Euphrasia officinalis* L. (*Scrophulariaceae*)-**Eyebright**; *Gaultheria procumbens* L. (*Ericaceae*)-**Wintergreen, Teaberry, Boxberry**; *Leonurus cardiaca* L. (*Lamiaceae*)-Motherwort, *Santolina charnaecyparissus* L. (*Asteraceae*)-**Lavender-Cotton**, *Seopolia carniolica* Jacq. (*Solanaceae*)-**Seopolia**; *Solanum tuberosum* L. (*Solanaceae*)-**Potato**; *Solidago virgaurea* L. (*Asteraceae*)-**European Goldenrod, Woundwort**; *Stachys officinalis* (L.) Trevisan (*Lamiaceae*)-Betony; *Trifolium pratense* L. (*Fabaceae*)-**Red Clover, Pavine Clover, Cowgrass**; *Valeriana officinalis* L. (*Valerianaceae*)-**Valerian**; and *Viscum album* L. (*Loranthaceae*)-**European Mistletoe**.

Preparation It occurs in plants only in conjugated forms e.g., **chlorogenic acid**. It has also been isolated from **green coffee**,* and from **roasted coffee****

It can also be prepared by the hydrolysis of **chlorogenic acid** in an acidic medium as shown below:***



Characteristic Features Caffeic acid has yellow crystals obtained from concentrated aqueous solutions and the corresponding monohydrate from dilute solutions. It gets softened at 194°C and decomposes at 223-225°C. It is sparingly soluble in cold water, but freely soluble in cold ethanol and hot water.

Chemical Tests

1. It changes colour from yellow to orange in an alkaline medium.
2. It readily forms the methyl ester ($C_{10}H_{10}O_4$) which are obtained as colourless crystals from water (mp 152-153°C).

6.2.1.3 Ferulic Acid

Synonyms Caffeic acid 3-methyl ether; 4-Hydroxy-3-methoxycinnamic acid.

Biological Sources Ferulic acid is widely distributed in small amounts in a variety of plants, namely: seeds of *Citrullus colocynthis* (L.) Schrad. (*Cucurbitaceae*)-**Colocynth, Bilter Apple, Wild Gourd**; flowers of *Convallaria majalis* L. (*Liliaceae*)-**Lily-of-the-Valley**; leaves of *Digitalis purpurea* L. (*Scrophulariaceae*)-**Common Foxglove, Digitalis**; young shoots of *Equisetum hyemale* L. (*Equisetaceae*)-**Shavegrass, Great Scouring Rush**; leaves of *Euphorbia lathyris* L. (*Euphorbiaceae*)-

* Wolfrom *et al.* *J. Agr. Food Chem.*, **8**, 58 (1960).

** Krasemann, *Arch. Pharm.*, **293**, 721 (1960).


*** Fiedler, *Arzneimittel-Forsch.*, **4**, 41 (1954); Whiting, Carr, *Nature* **180**, 1479 (1957), Guren, *Chemical Abstracts*, **61**, 9965h (1964).

Mole Plant, Petroleum Plant, Caper Spurge; dried herb of *Euphrasia officinalis* L. (*Scrophulariaceae*)-**Eyebright;** gum-resin of *Ferula assafoetida* L. (*Apiaceae*)-**Asafoetida;** volatile oil of *Gaultheria procumbens* L. (*Ericaceae*)-**Wintergreen, Teaberry, Boxberry;** twigs of *Hedera helix* L. (*Araliaceae*)-**Ivy;** leaves of *Hura crepitans* L. (*Euphorbiaceae*)-**Sandbox Tree;** leaves of *Plantago major* L. (*Plantaginaceae*)-**Plantain;** volatile oil of *Rheum officinale* Baill. (*Polygonaceae*)-**Chinese Rhubarb, Canton Rhubarb, Shensi Rhubarb;** shrubs of *Serenoa repens* (Bartel.) Small (*Arecaceae*)-**Saw Palmetto.**

Preparation It has been isolated from *Ferula foetida* Reg. (*Umbelliferae*)* and from *Pirus laricio* Poir. (*Abietineae***).

It may also be prepared by the interaction of vanillin, malonic acid and piperidine in pyridine for three weeks and then precipitating **ferulic acid** with dilute HCl.

Chemical Structure



trans-Ferulic Acid

Characteristic Features

cis-form : Yellow oil; uv_{\max} (in ethanol): 316 nm.

trans-form : Orthorhombic needles obtained from water; mp 174°C uv_{\max} (in ethanol): 236, 322 nm. It is soluble in hot water, ethanol and ethyl acetate; moderately soluble in ether; and sparingly soluble in benzene and petroleum ether.

Identification Test It forms the corresponding sodium salt by treatment with NaOH solution whereby the solubility gets enhanced appreciably.

Uses It is used as a preservative of food products.

6.2.1.4 Sinapic Acid

Biological Source It is obtained from the leaves and twigs of *Viscum album* L. (*Loranthaceae*)-**European Mistletoe.**

Preparation It may be prepared by the hydrolysis of **sinapic acid choline ester** obtained from the **black mustard seeds** of *Brassica nigra* Koch (*Cruciferae*) either in acidic medium or by enzymatic hydrolysis as given below:

* H. Hlasiwetz, L. Barth, *Ann.*, **138**, 61 (1966)

** M. Bamberger, *Monatsh.*, **12**, 441 (1891).

Characteristic Features **Cinnamaldehyde** is a yellowish oily liquid having a strong odour of cinnamon. Its physical parameters are: d_{25}^{25} 1.048-1.052; bp_{100} 177.7°C, bp_{200} 199.3°C and bp_{760} 246°C, n_D^{20} 1.618-1.623. It dissolves in about 700 parts of water and in about 7 volumes of 60% ethanol. It is, however, miscible with ethanol, ether, chloroform and oils.

Chemical Test On addition of a drop of $FeCl_3$ (1% w/v) solution to a few drops of **cinnamaldehyde** a distinct brown colour is produced.

Uses

1. It is used extensively in the perfume industry.
2. It is employed for flavouring foods and beverages.

Interestingly, it has been observed that the pairs of the allyl ($CH_2=CH-CH_2-$) and propenyl ($CH_3CH=CH-$) isomers, such as: *eugenol* and *isoeugenol* invariably occur together in the same medicinal plant as stated below:

- (i) *Cananga odorata* (Lam.) Hook. f. & Thoms. (*Annonaceae*)-**Cananaga, Ylang-Ylang**; and
- (ii) *Myristica fragrans* Houtt. (*Myristicaceae*)-**Mael, Nutmeg**

Note **Isomerization of the allyl to the propenyl form may also be accomplished in the laboratory, but only under very drastic and specific experimental parameters i.e., in the presence of strong alkali. However, such isomerization rarely takes place under normal conditions of isolation from natural products, such as: solvent extraction with ether etc.**

6.2.3 Coumarins

Coumarin and its derivatives, such as: **hydroxy-coumarins** and **furanocoumarins** are present in a plethora of medicinal plants. However, the most common and the most widespread plant coumarin is the parent compound *i.e.*, coumarin itself, which is reported to occur in more than twenty-seven plant families *viz.*, *Caprifoliaceae*, *Leguminosae*, *Oleaceae*, *Rubiaceae*, *Solanaceae*, *Umbelliferae*-to name a few such families.


In a broader sense, the **coumarins** may be classified into *three* major categories, namely:

- (a) Coumarins,
- (b) Hydroxycoumarins, and
- (c) Furanocoumarins.

All these *three* classes of drugs shall be described with the help of some important examples from each class individually as below:

6.2.3.1 Coumarins

The chemistry of **coumarin** may be understood more vividly with the help of geometrical isomers of ***o*-hydroxycinnamic acids**, one of which instantly yields the **lactone coumarin** (or **benzopyran**), whereas the other fails to do so. Therefore, the former is the *cis*-isomer called **coumarinic acid**, and the latter the *trans*-isomer known as the **coumaric acid** as given below:



Coumarin

Synonyms *cis-o*-Coumarinic acid lactone; Coumarin; Coumarinic anhydride; Tonka bean comphor.

Biological Sources Coumarin is present in a large number of medicinal herbs, such as:


Acacia farnesiana (L.) Willd (*Fabaceae*)-**Cassie, Huisache**; *Apium graveolens* L. (*Apiceae*)-**Celery**; *Artemisia dracunculus* L. (*Asteraceae*)-**Tarragon**; *Chamaemelum nobile* (L.) All. (*Asteraceae*)-**Roman Camomile, English Camomile, Camomile**; *Cinnamomum verum* J.S. Presler (*Lauraceae*)-**Ceylon Cinnamon**; *Dipteryx odorata* (Aubl.) Willd (*Fabaceae*)-**Tonka Bean, Tonga, Cumaru**; *Hyoseyamus niger* L. (*Solanaceae*)-**Henbane, Henblain, Jusquaime**; *Myroxylon balsamum* var. *Pereirae* (Royle) Harms. (*Fabaceae*)-**Balsam of Peru**; *Peumus boldus* Molina (*Monimiaceae*)-**Boldo**; *Pimpinella anisum* L. (*Apiaceae*)-**Anise**; and *Trilisa odoratissima* (J.F. Gemel.) Cass (*Asteraceae*)-**Deertongue, Deer's Tongue**.

Characteristic Features Coumarin crystals have an orthorhombic and rectangular plates. They have a pleasant, fragrant odour resembling to that of the vanilla beans and a burning taste. The physical characteristics are, namely: mp 68-70°C and bp 297-299°C. Its solubility in water is very poor, viz., 1g dissolves in m 400 ml of cold and 50 ml of boiling water. However, it is freely soluble in ethanol, chloroform, ether, oils and also in alkaline solutions of NaOH or KOH.

Uses It is used extensively as a flavouring agent in pharmaceutical formulations.

6.2.3.2 Hydroxycoumarins

Hydroxycoumarins are invariably found in a large number of plant families. However, the relatively more common ones are based upon the following substances, such as: *umbelliferone* (**7-hydroxy coumarin**), *aesculetin* (**6, 7-dihydroxy-coumarin**) and *scopoletin* (**6-methoxy-7-hydroxy coumarin**) as given below.



Umbelliferone : R = H;
Aesculetin : R = OH;
Scopoletin : R = OCH₃;

Interestingly, some rarer **hydroxycoumarins** are, namely, **dephentin** (7, 8-dihydroxy coumarin) and **fraxetin** (6-methoxy, 7-8-dihydroxy coumarin) are both obtained from plant sources.


A few typical examples of **hydroxycoumarins** shall be described in the sections that follow, e.g., **Umbelliferone, Esculetin, and Scopoletin**.

6.2.3.2.1 Umbelliferone

Synonyms Hydrangin; Skimmetin.

Biological Sources **Umbelliferone** is present in a variety of medicinal plants, for instance: *Apium graveolens* L. (*Apiaceae*) **Celery**; *Artemisia abrotanum* L. (*Asteraceae*)-**Southernwood, Old Man**; *Daphne mezereum* L. (*Thymelaeaceae*)-**Mezereon**; *Dipteryx odorata* (Aubl.) Willd. (*Fabaceae*)-**Tonka Bean, Tonga, Cumaru**; *Ferula sumbul* Hook. (*Apiaceae*)-**Sumbul, Mask Root**; *Hydrangea paniculata* Seib. (*Saxifragaceae*)-**Peegee**; *Lavandule angustifolia* Mill. (*Lamiaceae*)-**Lavender**; *Matricaria chamomilla* L. (*Asteraceae*)-**Hungarian Camomile, German Camomile, Manzanilla**; and *Pimpinella anisum* L. (*Apiaceae*)-**Anise**.

Preparation **Asafoetida** contains resin (40-65%) which consists of chiefly a resin-alcohol **asaresinotannol** both in the free or combined form with ferulic acid, and of course, free **umbelliferone** is totally *absent* in the drug. Thus, umbelliferone is prepared by treating **ferulic acid** with HCl which gets converted to **umbellic acid** and the latter loses a molecule of water to give rise to **umbelliferone** as given below:



Umbelliferone may also be obtained by distillation of resin from **Umbelliferae**.*

Characteristic Features It is obtained as needles from water. It develops the characteristic odour of coumarin on heating. Its mp is 225-228°C. It usually sublimes. Its solubility in water is very poor *i.e.*, it dissolves 1 g in nearly 100 ml of boiling water. It is freely soluble in ethanol, chloroform, acetic acid and dilute alkaline solution. It is sparingly soluble in ether and the solutions exhibit a distinct blue fluorescence.

Identification Test When 0.5 g of **umbelliferone** is triturated with pure sand (SiO_2) and 5 ml of HCl, added 5 ml of water, filtered and to the filtrate added an equal volume of ammonia solution, it gives a beautiful blue fluorescence.

Uses

1. It is an important ingredient in most sunscreen lotions and creams.
2. It is most importantly used as an intracellular and pH sensitive fluorescent indicator and blood-brain-barrier (BBB) probe.

* Z wenger, *Ann.*, **115**, 1, 15 (1860).

6.2.3.2.2 Esculetin


Synonyms Aesculetin; Chicorigenin; 6, 7-Dihydroxy-coumarin.

Biological Source It is the *aglucon* of **esculin** and **cichorlin**: Esculin is derived from two different plant sources, namely: (a) the barks of *Crataegus oxyacantha* L. (*Rosaceae*)-**Hawthorn**; and (b) the flowers of *Centarea cyanus* Linn., (*Compositae*). It is a glucoside which upon hydrolysis gives the aglucon **esculetin**.


Esculetin is also obtained from **cichorlin**, which is a glucoside and found to be isomeric with esculin. **Cichorlin** is present in the flowers of the **chicory plant** (*Chichorium intybus* L., family: *Compositae*).

Preparation It is obtained by the hydrolysis of the following *two* glucosides, namely:

(a) **From Esculin:**



(b) **From Cichorlin:**



Characteristic Features It is obtained as prisms from glacial acetic acid and as leaflets by vacuum sublimation. Its mp is 268-270°C. It is soluble in dilute alkalis (2M solution) with the emission of blue fluorescence. It is almost insoluble in ether and in boiling water, but moderately soluble in hot ethanol and in glacial acetic acid.

Uses It is mostly in filters for absorption of uv-light


6.2.3.2.3 Scopoletin

Synonyms Chrysotropic acid; Gelseminic acid; 6-Methoxyumbelliferone; β -Methylesculetin; 7-Hydroxy 6-methoxycoumarin.

Biological Sources It is the *aglucone* of **scopolin**. **Scopoletin** occurs in the roots of *Arnica montana* L. (*Asteraceae*)-**Mountain Tobacco, Leopard's-bane**; leaves of *Artemisia abrotanum* L. (*Asteraceae*)-**Southernwood, Old Man**; roots and leaves of *Atropa belladonna* L. (*Solanaceae*)-**Belladonna, Deadly Nightshade**; barks of *Brunfelsia uniflora* (Phol.) D. Don. (*Solanaceae*)-**Manaca, Manacan**; fruits of *Capsicum annum* L. (*Solanaceae*)-**Chili, Sweet Peppers, Paprika**;

oil of the plant *Chamaemelum nobile* (L.) All. (*Asteraceae*)-**Roman Comomile; English Camomile, Comomile**; and roots of *Withania somniferum* (L.) Dunal (*Solanaceae*)-**Ashwagandha**.

Preparation It is obtained by the hydrolysis of the glucoside scopolin *i.e.*, 7-(β -D glucopyranosyloxy)-6-methoxy-2H-1-benzopyran-2-one as follows:



Characteristic Features **Scopoletin** occurs as prisms or needles from either acetic acid or chloroform. It melts at 204°C and has a uv_{\max} : 230, 254, 260, 298, 346 nm (log ϵ 4.11, 3.68, 3.63, 3.68, 4.07). It is slightly soluble in water or cold ethanol and quite soluble in hot ethanol and hot glacial acetic acid. It is moderately soluble in chloroform, but practically insoluble in the non-polar solvent benzene.

Identification Tests

1. Dissolve 0.1 g in ethanol and warm it in an electric water-bath to affect dissolution. The resulting solution gives a blue fluorescence.
2. A solution of 0.1 g in 3 ml of hot ethanol reduces the Fehling's solution thereby leaving behind a brick-red precipitate of cupric oxide (CuO).


It is pertinent to mention here that there exists some rarer species of hydroxycoumarins, such as: **daphentin** and **fraxetin**, which shall now be discussed in the sections that follows:

6.2.3.2.4 Daphentin

Synonyms 7, 8-Dihydroxycoumarin;

Biological Sources It is the aglucon of **daphnin**. It is obtained from the seeds and fruits of *Daphne mezereum* L. (*Thymelaeaceae*)-**Mezereon**; and the seeds of *Euphorbia lathyris* L. (*Euphorbiaceae*)-**Mole Plant, Petroleum Plant, Capers Spurge**.

Preparation **Daphentin** is prepared conveniently from its glucoside known as **daphnin** *i.e.*, 7, 8-dihydroxycoumarin 7- β -D-glucoside by treating the latter in *three* different ways, namely: (i) By boiling with dilute mineral acids; (ii) By enzymatic hydrolysis; and (iii) By sublimation as given below:



Characteristic Features The crystals obtained from dilute ethanol has a mp 256°C (decomposes). It undergoes sublimation on heating. It is soluble in boiling water, hot dilute alcohol and hot glacial acetic acid. It is found to be sparingly soluble in ether, CS₂, chloroform, and benzene.

Identification Tests


1. An aqueous solution of daphentin gives a green colouration with FeCl₃ solution, which turns red on the addition of sodium carbonate.
2. An alkaline solution of daphentin in alkali carbonate or alkali gives a yellow colour.

6.2.3.2.5 Fraxetin

Synonyms 7, 8-Dihydroxy-6-methoxycoumarin.

Biological Source It is the aglucon of **fraxin**. **Fraxin** is present in the seeds of *Acsculus hippocastanum* L. (*Hippocastanaceae*)-**Horse Chestnut**.

Preparation Fraxetin is obtained by heating **fraxin** with dilute sulphuric acid to affect the hydrolysis of glucoside and get the desired aglucon residue as shown here under:



Characteristic Features **Fraxetin** is obtained as plates from ethanol having mp 228°C. It has been observed that it turns first yellow at 150°C and subsequently brown at mp. It is soluble in 10 L of cold water, but in 300 ml of boiling water. It is somewhat more soluble in alcohol and practically insoluble in ether.

Identification Tests It forms the corresponding dimethyl ether termed as 6,7,8-trimethoxycoumarin (C₁₂H₁₂O₅) which has a mp 104°C and bp_{0.2} 90-100°C.


6.2.3.3 Furanocoumarins

Furanocoumarins, represent a class of relatively more complex coumarins that occur in various natural plant products. A few important members of this particular class are, namely: **Psoralen**; **Methoxsalen**; **Bergapten**; and **Imperatorin**, which shall be discussed below in an elaborated manner.

6.2.3.3.1 Psoralen

Synonyms Ficusin; 6-Hydroxy-5-Benzofuranacrylic acid δ-lactone; Furo (3, 2-δ)-coumarin.

Biological Source **Psoralen** belongs to one of a group of furanocoumarins occurring naturally in more than two dozen different plant sources, for instance: *Rutaceae* (e.g., **Bergamot**, **Limes**, **Cloves**); *Umbelliferae* (**Celery**; **Parsnips**); *Leguminosae* (e.g., **Psoralen coryfolia**); and *Moraceae* (e.g., **Figs**). It is also found in the **Rue Oil** obtained from *Ruta graveolens* L. (*Rutaceae*)-known as **Rue**, **Garden Rue** or **German Rue**. It is obtained from the leaves of *Ficus carica* Linn. (*Moraceae*)-**Figs**, **Anjir**.

Chemical Structure

Psoralen

Characteristic Features **Psoralen** crystals from ether have two sets of melting points *e.g.*, 163-164°C and 169-179°C (Spath). It is very soluble in chloroform, less soluble in alcohol, sparingly soluble in ether and practically insoluble in petroleum ether (60-80°C).

Identification Tests

1. Dissolve 1 mg of **psoralen** in 5 ml of ethanol and add to it 15 ml of a mixture made up of 3 parts of propylene glycol, 5 parts of acetic acid and 43 parts of water. The resulting mixture on being exposed to the uv-light in a uv-chamber, gives a distinct *blue-fluorescence*.
2. When 1 mg is dissolved in 2 ml ethanol, mixed with two drops of NaOH solution (0.1 M) and the resulting solution is subjected to uv-light, it emits a yellow fluorescence.


Uses

1. It is used in the treatment of leucoderma patches.
2. Psoralens have also exhibited photosensitizing and phototoxic effects in animals and human beings and, hence have been employed extensively in *photochemotherapy* for the treatment and management of **vitiligo***, **psoriasis**** and **mycosis fungoides*****.

6.2.3.3.2 Methoxsalen

Synonyms Xanthotoxin; Meloxine; Ammoidin; Meladinine; 8-Methoxypsoralen; 8-MOP; 8-MP; Oxsoralen.

Biological Source **Methoxsalen** is a naturally occurring analogue of **psoralen**, found in various species of *Rutaceae*, *Leguminosae*, and *Umbelliferae*. It is obtained from the fruits of *Fragara xanthoxyloides* and the fruits of *Ammi majus* belonging to the natural order *Umbelliferae*. It is also found in the herb *Ruta graveolens* (*Rutaceae*).

Chemical Structure

Methoxsalen

9-Methoxy-7H-furo[3,2-g][1]benzopyran-7-one; (C₁₂ H₈ O₄).

* T.F. Anderson, J.J. Voorhees, *Ann. Rev. Pharmacol. Toxicol.*, **20**, 235 (1980); *Vitiligo*: An acquired cutaneous disorder characterised by white patches, surrounded by areas of normal pigmentation.

** A. Kornhauser *et al.*, *Science*, **217**, 733 (1982); *Psoriasis*: A common chronic disease of the skin consisting of erythematous papules that coalesce to form plaques with distinct borders.

*** B.J. Parsons, *Photochem. Photobiol.*, **32**, 813-821 (1980); *Mycosis Fungoides*: A non-Hodgkin's form of cutaneous T-cell lymphoma of unknown etiology caused by a fungus.

Isolation The various steps involved are as under:

1. The *A. majus* fruits are dried, powdered, sieved and extracted with petroleum ether to complete exhaustion.
2. The petroleum ether extract is filtered and concentrated to obtain a dark green semi-crystalline solid mass (crude methoxalen) crystallizes out.

Note: The petroleum ether layer is carefully decanted off while hot and reserved separately for the isolation of imperatorin.

3. The residual dark-green solid mass is dissolved in minimum quantity of ethanol and boiled over an electric water-bath for 45-60 minutes. The contents are filtered immediately and the filtrate is concentrated under vacuo. It is cooled in a refrigerator overnight when pale-green crystals separate out. The crystals of **methoxsalen** thus obtained are purified first by washing with boiling water and finally recrystallizing from ethyl acetate.

Characteristic Features

1. It is obtained in two forms: *first*—as silky needles either from hot water or benzene + petroleum ether; *secondly*—as long rhombic prisms from ethanol + ether, having mp 148°C.
2. It is odourless *but* has a distinct bitter taste followed by tingling sensation.
3. It has uv_{max} : 219, 249, 300 nm ($\log \epsilon$ 4.32, 4.35, 4.06).
4. It has a pH 5.5.
5. *Solubility Profile*: It is practically insoluble in cold water; sparingly soluble in boiling water, liquid petroleum, ether; soluble in boiling ethanol, acetone, acetic acid, vegetable fixed oils, propylene glycol, benzene; freely soluble in chloroform; and soluble in aqueous alkalies with ring cleavage, but is reconstituted upon neutralization.

Identification Tests These are as follows:

1. A few crystals of **methoxsalen** on being triturated with little sulphuric acid in a porcelain dish produces an orange-yellow colour that gets changed to light green finally.
2. **Wagner's Reagent Test: Xanthotoxin** gives an instant precipitate with **Wagner's Reagent** ($I_2 + KI$).
3. **HNO₃ Test**: It gives a distinct yellow colouration with dilute HNO₃, which on rendering to alkaline with KOH or NaOH, changes to crimson colour.

Uses

1. It is used extensively in the treatment of leukoderma.
2. It is employed as a pigmentation agent.
3. It is also used in the treatment of psoriasis and mycosis fungoides.

6.2.3.3.3 Bergapten (e)


Synonyms Bergapten; Heraclin; Majudin; Psoraderm; 5-Methoxypsoralen; 5-MOP.

Biological Source **Bergapten** is the naturally occurring analogue of psoralen and an isomer of **methoxsalen**, mostly found in a wide variety of plants, such as: roots and fruits of *Angelica archangelica* L. (*Apiaceae*)-**Angelica, Garden Angelica, European Angelica**; seeds of *Apium*

graveolens L. (*Apiaceae*)-Celery; leaves, stems and fruits of *Petroselinum crispum* (Mill.) Nym. (*Apiaceae*)-Parsley; **Rue Oil** of *Ruta graveolens* L. (*Rutaceae*)-**Rue, Garden Rue, German Rue.**

Preparation **Bergapten** was first isolated from the oil of bergamot from *Citrus bergamia* Risso., belonging to the natural order *Aurantiodia**. It was also isolated from *Fagara xanthoxyloides* Lam., belonging to family *Rutaceae***.

Chemical Structure



Bergapten

Characteristic Features The crystals obtained from ethanol are needle-shaped having mp 188°C. It sublimes on heating. It is practically insoluble in boiling water, slightly soluble in glacial acetic acid, chloroform, benzene, and warm phenol. It is soluble in absolute ethanol (1 part in 60).

Identification Test It gives a distinct yellow-gold colouration when its solution is treated with a few drops of concentrate H_2SO_4 .

Uses


1. It is used in photochemotherapy of psoriasis.
2. It has been used to promote tanning in *suntan preparations e.g.*, creams and lotions.

6.2.3.3.4 Imperatorin

Synonyms Ammidin; Pentosalen; Marmelosin;

Biological Sources It is obtained from the roots and fruits of *Angelica archangelica* L. (*Apiaceae*) (**Angelica, Garden Angelica, European Angelica**); from the roots of *Imperatoria ostruthium* L. (*Umbelliferae*); from fruit of *Pastinaca sativa* L. (*Umbelliferae*); and also in the fruits of *Ammimajus* (*Umbelliferae*). However, the seed oil of *A. archangelica* is said to contain upto 0.5% **imperatorin**.

Chemical Structure



Imperatorin

9-[(3-Methyl-2-butenyl)oxy]-7H-furo [3, 2-g] [1] benzopyran-7-one; ($C_{16}H_{14}O_4$).

* Pomeranz, *Monatsh*, **12**, 379 (1891), **14**, 28 (1893).

** H. Thoms., E. Baeteke, *Ber.*, **44**, 3326 (1911).

Isolation The various steps involved are as follows:

1. The petroleum ether mother liquor left after the separation of **methoxsalen (Xanthotoxin)**, is concentrated under vacuo and allowed to cool in a refrigerator overnight when the crude imperatorin separates out.
2. The crude product is collected, dissolved in ether, filtered and concentrated under reduced pressure. It is kept in a refrigerator, and the crystals separating out are purified subsequently by recrystallization from ethanol.

Characteristic Features These are as given below:

1. It is obtained in *two* forms: *First*—as prisms from ether, and *secondly*—as long fine needles from hot water, having mp 102°C.
2. It has uv_{max} : 302, 265, 250 nm (log ϵ 3.95, 4.00, 4.24).
3. *Solubility Profile*: It is practically insoluble in cold water; very sparingly soluble in boiling water; freely soluble in chloroform; and soluble in benzene, ethanol, ether, petroleum, ether alkali hydroxides.

Identification Tests These are as stated below:

1. **Sulphuric Acid Test**: Imperatorin gives an intense deep orange colouration with a few drops of sulphuric acid which ultimately changes to brown colour.
2. **Marqui's Reagent**: It gives an orange colouration with Marqui's Reagent that rapidly changes to brown.
3. **Tollen's Reagent (Ammoniacal AgNO₃)**: It reduces Tollen's Reagent to produce a silver mirror.
4. **Fehlings Test**: It reduces Fehling's solution to give a brick-red precipitate of cupric oxide.
5. **Nitric Acid Test**: It gives a distinct yellow colour on boiling with dilute HNO₃, and this colour changes to purple on being treated with strong alkali *e.g.* NaOH or KOH.

6.2.4 Abridged Phenylpropanoids

Abridged phenylpropanoids are invariably acids and phenols, and quite rarely alcohol and aldehydes, which are attributed due to the β -oxidation of the C₃-side chain of: (a) **para-Coumaroyl CoA**, and (b) **para-Cinnamoyl CoA** followed by oxidative decarboxylation.

The various **abridged phenylpropanoids** present in a large number of **medicinal herbs** are usually classified into *three* major heads, namely:

- (a) With no side-chain,
- (b) With side-chain having one C-atom, and
- (c) With side-chain having two C-atoms.

All these *three* classes of compounds occurring in natural plants shall be discussed separately with the help of certain appropriate examples as stated below:


6.2.4.1 With No side-Chain

The most glaring example of an **abridged phenylpropenoid** that has no side-chain is **catechol** which shall be treated more explicitly as follows:

Catechol

Synonyms Pyrocatechol; Pyrocatechin; 1, 2-Dihydroxybenzene; 1, 2-Benzenediol.

Biological Sources It occurs naturally in various plant species, such as: whole plant of *Anandenathera peregrina* L. Speg. (*Mimosaceae*)-**Niopo, Cohoba, Yope, Yupa**; cortex of *Melia azedaraeh* L. (*Meliaceae*)-Chinaberry; and plant of *Rumex crispus* L. (*Polygonaceae*)-**Yellow Dock**.


Chemical Structure

Catechol


Preparation Being phenolic in character the aqueous extract may be treated with dilute alkalis carefully, and the resulting sodium salts are neutralized to yield the desired **catechol** from the natural plant sources.

It may also be obtained by several other methods as stated below:


- (a) **Decarboxylation of Protocatechuic Acid:** Protocatechuic acid is found in minute quantities in wheat grains, in wheat seedlings and in many other plants.



- (b) **From Salicylaldehyde:** Catechol is also obtained by the interaction of salicylaldehyde with hydrogen peroxide as follows:



- (c) **From Guaiacol:** It may also be prepared by treating guaiacol with hydrobromic acid as given below:



Characteristic Features **Catechol** is obtained as the monoclinic tablets or prisms from toluene. It usually undergoes discolouration on exposure to air or light. Its physical characteristics are: mp

105°C, d 1.344, bp₇₆₀ 245.5°C, bp₁₀₀ 176°C, bp₄₀ 150.6°C. It is steam volatile and sublimes on heating. Its dissociation constant K at 18° = 3.3×10^{-10} . It is soluble in 2.3 parts of water, ethanol, benzene, chloroform and ether; and very soluble in aqueous alkali solutions and pyridine. It has been observed that its aqueous solution soon turns brown.

Identification Test Dissolve 0.2 g of **catechol** in water and add to it a few drops of FeCl₃ (0.1% w/v) aqueous solution. The appearance of a green colour confirms the presence of catechol.

Uses

1. It is used as an antiseptic agent.
2. It finds its application in photography.
3. It is also employed for dyeing fur.

6.2.4.2 With Side-Chain Having One Carbon Atom

The **abridged phenylpropenoids** having a side chain with one carbon atom represent an important group of naturally occurring plant products, such as: **Benzoic acid, Gallic acid, Methyl salicylate, Salicin** and **Vanillin**.

These compounds shall be discussed here under in a concise descriptive manner.

6.2.4.2.1 Benzoic Acid

Synonyms Dracylic acid: Phenylformic acid, Benzene carboxylic acid.


Biological Source It mostly occurs in nature in free and combined forms. **Gum benzoin** may contain as much as 20% of **benzoic acid**, whereas most berries contain appreciable amounts *i.e.*, upto 0.05%.

Benzoic acid is found in a large number of medicinal herbs, namely: plant of *Aeacia farnesiana* (L.) Willd. (*Fabaceae*)-**Cassie, Huisache**; oil of *Cananga odorata* (Lam.) Hook. f. & Thoms. (*Annonaceae*)-**Cananga, Ylang-Ylang**; latex of *Daemonorops draco* bl. (*Arecaceae*)-**Dragon's Blood**; tubers of *Gloriosa superba* L. (*Liliaceae*)-**Glory Lilly**; plant of *Illicium verum* Hook. f. (*Magnoliaceae*)-**Star-Anise, Chinese Anise**; volatile of *Narcissus tazetta* L. (*Amaryllidaceae*)-**Daffodil, Chinese Sacred Lilly, Polyanthus Narcissus**; roots of *Paeonia officinalis* L. (*Ranunculaceae*)-**Peony**; Plant of *Piper methysticum* Forst. (*Piperaceae*)-**Kava-Kava**; leaves of *Plantago major* L. (*Plantaginaceae*)-**Plantain**; gum of *Styrax benzoin* Dryander (*Styracaceae*)-**Benzoin, Sumatra Benzoin, Styrax**; and pods of *Vanilla planifolia* Andr. (*Orchidaceae*)-**Vanilla**.


Preparation The alcoholic extract of the plant is concentrated cooled and treated with dilute mineral acid. The solid residue thus obtained is further recrystallized from hot alcohol.

It is also obtained synthetically in several ways as stated below:

(a) **Oxidation of Toluene:** Toluene when oxidized by air, it yields **benzoic acid**:



(b) **Decarboxylation of Phthalic Anhydride:** The decarboxylation of phthalic anhydride gives rise to **benzoic acid**:



Characteristic Features **Benzoic acid** has been obtained as monoclinic plates or tablets or leaflets. Its density ranges between 1.266-1.321. It has mp 122.4°C; and it sublimes at nearly 100°C. Its bp₇₆₀ 249.2°C, bp₁₀₀ 186.2°C, bp₄₀ 162.6°C. It is found to be steam-volatile. It has a flash point ranging between 121-131°C and dissociation constant K at 25°C = 6.40×10^5 . The pH of its saturated solution is 2.8. Its solubility in water at 25°C is 3.4 g/L and at 95°C is 68 g/L. Its solubility in other organic solvents are: cold ethanol 1 g/2.3 ml; boiling ethanol 1 g/1.5 ml; chloroform 1 g/4.5 ml; ether 1 g/3 ml; acetone 1 g/3 ml; carbon tetrachloride 1 g/30 ml; benzene 1 g/10 ml. carbon disulphide 1 g/30 ml. It is also soluble in fixed oils and volatile oils. It is slightly soluble in petroleum ether. The solubility of **benzoic acid** is enhanced by the presence of alkaline substances e.g., trisodium phosphate (Na₃PO₄) and borax.

Identification Test The corresponding calcium benzoate trihydrate salt gives an orthorhombic crystal or powder having a density of 1.44. It is highly soluble in boiling water but sparingly soluble in cold water *i.e.*, 1 g/25 ml.

Uses

1. It has been used in conjunction with salicylic acid in creams and ointments as an effective topical antifungal agent.
2. It is used extensively for the preservation of foods, fats, fruit juices, alkaloidal solutions.
3. It is employed as a *mordant* in calico printing.
4. It is also used for curing tobacco.

6.2.4.2.2 Gallic Acid

Synonym 3, 4, 5-Trihydroxybenzoic Acid.


Biological Sources **Gallic acid** is present in a very large cross-section of medicinal plants. A few such species are as follows, namely: seeds of *Abrus precatorius* L. (*Fabaceae*)-**Jequerity**; berries of *Aretostaphylos uva-ursi* (L.)-Spreng. (*Ericaceae*)-**Bearberry**; seeds of *Cimicifuga racemosa* (L.) Nutt. (*Ranunculaceae*)-**Black Cohosh, Black Snakeroot**; fruits of *Coriaria thymifolia* Humb. & Bonpl. (*Coriariaceae*)-**Shanshi**; resinoid substance (cypripedin) obtained from the rhizome of *Cypripedium* sp. (*Orchidaceae*) **Yellow Lady-slipper**; green branches of *Ephedra gerardiana* Wall. ex Staph (*Ephadraceae*)-**Pakistani Ephedra**; plant of *Eupatorium pertolatum* L. (*Asteraceae*)-**Boneset, Ague Weed**; roots of *Geranium maculatum* L. (*Geraniaceae*)-**Cranebill**; plant of *Juniperus sabina* L. (*Cupressaceae*)-**Sabine, Savin**; leaves of *Lawsonia inermis* L. (*Lythraceae*)-**Henna, Egyptian Privet, Mignonette**; root bark of *Quassia amara* L. (*Simaroubaceae*)-**Surinam Quassia, Bitterwood**; leaves

of *Tanacetum vulgare* L. (*Asteraceae*)-**Tansy**; and plant of *Tussilago farfara* L. (*Asteraceae*)-**Coltsfoot, Coughwort, Horse-Hoof**.

Preparation The two important methods of preparation of **gallic acid** from natural sources are given below:

- (a) **From Tannings of Nutgalls:** It is obtained either by alkaline or acid hydrolysis of the tannins from Nutgalls.
- (b) **From Spent Broths of *Penicillium glaucum* or *Aspergillus niger*:** It may also be prepared by carrying out the enzymatic hydrolysis from the spent broths of *P. glaucum* and *A. niger* which contains the enzyme *tannase*.

Chemical Structure




Gallic Acid

Characteristic Features **Gallic acid** is obtained as needles either from methanol or chloroform. It sublimes at 210°C that yields a fairly stable form which melts at 258-265°C (decomposed) and also an unstable form having mp 225-230°C. Its solubility in water in 1 g/87 ml, boiling water 1 g/3 ml, ethane 1 g/6 ml, ether 1 g/100 ml, glycerol 1 g/10 ml, and acetone 1 g/5 ml. It is found to be practically insoluble in benzene, chloroform and petroleum ether.

Identification Tests

1. **Gallic acid** is first converted to pyrogallol by means of the decarboxylation of the latter, which gives a distinct blue colour with FeCl_3 solution (0.1% w/v).



2. It forms its corresponding methyl ester with methanol which gives a sharp mp 202°C.

Uses It was used formerly as an astringent and styptic.

6.2.4.2.3 Methyl Salicylate It has been discussed in details under section 5.2.6.5.8.2 (c) in Chapter-5 on 'Terpenoids'.

6.2.4.2.4 Salicin


Synonyms Salicoside; Salicyl alcohol glucoside; Saligenin- β -D-glucopyranoside; 2-(Hydroxymethyl) phenyl- β -D-glucopyranoside.

Biological Sources It is obtained from the volatile oil of *Filipendula ulmaria* (L.) Maxim. (*Rosaceae*)-**Meadosweet, Queen-of-the Meadow**. It is also found in leaves and female flowers of the willow (*Salix*).

Preparation **Salicin** is prepared by several methods, such as:

- (a) **Bark of Poplar (Populus):** It is usually prepared by making the hot water extracts obtained either from the ground barks of poplar or willow.
- (b) **Root Bark of Viburnum prunifolium L. (Caprifoliaceae):** It may also be isolated from the root barks of *V. prunifolium* by means of the hot water extracts.*

Chemical Structure



Salicin

Characteristic Features **Salicin** has orthorhombic crystals from water with mp 199-202°C. Its physical parameters are: $[\alpha]_D^{25} - 62^\circ\text{C}$ to $- 67^\circ\text{C}$ (c = 3) and $[\alpha]_D^{20} - 45.6^\circ\text{C}$ (c = 0.6 in absolute ethanol). It is soluble in water at ambient temperature 1 g/23 ml, in boiling water 1 g/3 ml, in cold alcohol 1 g/90 ml and in alcohol at 60°C (1 g/30 ml). It is freely soluble in alkaline solutions, pyridine, and glacial acetic acid. It is found to be practically insoluble in chloroform and ether. The aqueous solutions are neutral to litmus and possesses a distinct bitter taste.

Uses

1. It is widely used as an analgesic
2. It is employed as a standard substrate in evaluating enzyme preparations containing β-glucosidase.


6.2.4.2.5 Vanillin

Synonyms Vanillic aldehyde; 3-Methoxy-4-hydroxybenzaldehyde; 4-Hydroxy-3-methoxy benzaldehyde.

Biological Sources **Vanillin** is found in a plethora of medicinal herbs, such as: fruits of *Ananas comosus* (L.) Merr. (*Bromeliaceae*)-**Pineapple**; volatile oil of *Croton eleutheria* Sw. (*Euphorbiaceae*)-**Cascarilla**; oleo-gum-resin of *Ferula asafoetida* L. (*Apiaceae*)-**Asafbetida**; flowerbuds of *Filipendula ulmaria* (L.) Maxim. (*Rosaceae*)-**Meadow-sweet, Queen of the Meadow**; leaves of *Ilex paragua-riensis* St. Hil. (*Aquifoliaceae*)-**Yerba Mate, Paraguay tea, South American Holly**; seeds of *Myroxylon balsamum* var. *Pereirae* (Royle) Harms. (*Fabiaceae*)-**Balsam of Peru**; essential oil of *Serenoa repens* (Bartel.) Small (*Arecaceae*)-**Saw Palmetto**; leaves of *Tilia europaea* L. (*Tiliaceae*)-**Linden Tree** (America), **Lime Tree** (Europe), and beans of *Vanilla planifolia* Andr. (*Orchidaceae*)-**Vanilla**.

Preparation **Vanillin** is prepared by the hydrolysis of the aldehyde glycoside *vanilloside* obtained from the unripe vanilla fruit to give rise to the desired aglycone residue (vanillin) as given below:

* Evans et al., *J. Am. Pharm. Assoc.*, **34**, 207 (1945).



It may also be obtained synthetically from eugenol or guaiacol; and also from the waste (lignin) of the wood-pulp industry.

Characteristic Features **Vanillin** has either a white or very slightly yellow needle-like appearance. It possesses a pleasant aromatic vanilla odour and taste. It undergoes gradual oxidation on exposure to humid and moist air. It gets affected by uv-light. Its physical characteristics are: mp 80-81°C; d 1.056 and bp₇₆₀ 285°C. Its solubility in water at ambient temperature is very low (1 g/100 ml), in hot water at 80°C (1 g/16 ml), in glycerol (1 g/20 ml); but freely soluble in ethanol, chloroform, ether, carbon disulfide, glacial acetic acid, pyridine and also soluble in oils and aqueous solutions of alkali hydroxides (NaOH, KOH). The aqueous solution of **vanillin** is acidic to litmus. It must be stored in air-tight and light-resistant containers.

Chemical Test **Vanillin** reduces the Tollen's Reagent (*i.e.*, ammoniacal silver nitrate solution) to give rise to the silver-mirror on warming in a water-bath thereby showing the presence of aldehyde moiety present in it.

Uses

1. It is used extensively as a pharmaceutical aid for flavouring pharmaceutical formulations *e.g.*, cough mixture, syrups and elixirs.
2. It is also used as a flavouring agent in beverages, malted-milk-foods, confectionery and in perfumery.
3. It is also employed in manufacture of 'liqueurs'.
4. It has more or less replaced *Vanilla pod* and *tincture vanilla* by virtue of the fact that 1 part of vanillin equals 400 parts of the former and 2.5-3 parts of vanillin equals 500 parts of the latter.
5. It is also used as a reagent in *Analytical Chemistry*.

6.2.4.3 With Side-Chain Having Two Carbon Atoms

The most glaring example of an **abridged-phenylpropanoid** is that of phenyl ethanol which shall be discussed here under:


6.2.4.3.1 Phenyl Ethanol

Synonyms 2-Phenylethanol; β-Phenylethyl alcohol; Benzyl carbinol; β-Hydroxyethylbenzenc; Benzeneethanol.

Biological Sources Phenyl ethanol is present in variety of essential oils in medicinal plants, namely: *Tillia europaea* L. (*Tiliaceae*)—**Linden Tree** (America), **Lime Tree** (Europe) and other volatile oils

of *viz.*, rose, carnation, hyacinth, aleppo pine, orange blossom, geranium Bourbon, champaea and neroli.

Preparation It is obtained by the fractional distillation of volatile oils stated above and collecting the fractions at 219-221°C. It may also be prepared by the reduction of ethylphenyl acetate in the presence pure sodium metal and absolute alcohol in a perfectly dry reaction flask as shown below:



Characteristic Features Phenyl alcohol is a colourless liquid having floral odour resembling to that of rose. Its physical characteristics are, namely: mp -27°C ; d_{25}^{25} 1.017-1.019; bp_{14} 104°C and n_D^{20} 1.530-1.533. Its solubility in water is very low *i.e.*, 2 ml gets dissolved in 100 ml water after thorough shaking; 1 ml is rapidly soluble in 1 ml of 50% ethanol; and completely miscible with ether and ethanol.

Uses

1. It is used in flavouring foods and beverages.
2. It is extensively employed in perfumery especially for making *rose perfumes*.
3. It is used as a *pharmaceutical aid* to combat microbial infections.

6.2.5 Biphenylpropanoid Derivatives

In this particular class of compounds, the side chains from two **phenylpropanoids** interact with each other to yield **biphenylpropanoid derivatives** that are commonly termed as *Lignans* or *Neolignans*.

(i) Lignans: **Lignans**, the plant products with *low molecular weight* that are accomplished by the oxidative coupling of *para*-hydroxyphenylpropene units wherein the two units may be linked by an oxygen bridge. Furthermore, the monomeric precursor units are, namely: cinnamic acid, cinnamyl alcohol, propenylbenzene and allylbenzene. However, the terminology **Lignan** or more precisely **Haworth Lignan** is generally applied to such compounds that are derived by coupling acid and/or alcohol exclusively; whereas, the compounds which are derived by coupling propenyl and/or allyl derivatives are known as **Neolignans**.*

Biological Source **Lignans** occur widely and have been obtained from roots, heart wood, foliage, fruit and resinous exudates of plants. They represent the dimer stage intermediate between the monomeric propylphenol units and lignin. However, the naturally occurring trimers and tetramers have not so far been reported. Nevertheless, the occurrence of **lignans** both in man and animal

* Gottlieb, O.R., *Fortschr. Chem. Org. Naturst*, **35**, 1-72 (1978)

species have been reported.* The α -lignan has been found in the roots and rhizomes of *Podophyllum hexandrum* Royle. (*Berberidaceae*).

Preparation Generally, the **lignans** are formed by the reduction of ferulic acid to coniferyl alcohol as its first and foremost step; and subsequently *via* the oxidative dimerization of the coniferyl alcohol units and the establishment of linkage through the β -carbon atom of the C_3 side-chain.


Characteristic Features **Lignans** are typically found as single enantiomeric forms *i.e.*, either as *d*- or *l*-isomers. However, these also occur as their racemic products *i.e.*, *dl*-forms. It has been observed that the **lignans** vary to a large extent with regard to their respective oxidation levels, degree of substitution, and above all the structural complexity.

Examples: **Podophyllum** and its chemical constituent *podophyllotoxin*. This has been discussed at length under Section 5.2.7 related to 'resin' in Chapter 5 on '**Terpenoids**'. The two important examples of **lignan** are that of **etoposide** and **teriposide** which shall be discussed in details below:

A. Etoposide

Synonyms Lastet, Vepesid, VP-16-213, NSC-141540, EPEG, 4'-Demethylepipodophyllotoxin 9-[4, 6-O-ethylidene- β -D-glucopyranoside.

Chemical Structure



Etoposide

Characteristic Features **Etoposide** is a semisynthetic **podophyllotoxin** structural analogue used as an '*antineoplastic agent*'. It essentially *differs* structurally from **podophyllotoxin** in the following manners, namely:

- (a) It has an ethylidene glucoside moiety attached at the C—1 position.
- (b) It has a epimeric configuration at the C—4 position of ring C, and
- (c) It possesses a hydroxyl function at the C—4' position rather than a methoxy moiety.

However, the hydroxyl (-OH) moiety at C-4' exerts *two* important properties to *Etoposide*, namely:

* Stitch S.R. *et al.*, *Nature*, **287**, 238 (1980); Setchel K.D.R., *ibid.*, 740

- (i) It is associated with etoposide's ability to induce *single-stranded DNA breaks*, and
- (ii) The ethylidene glucoside function is associated with etoposide's inability to *inhibit microtubule assembly*, an important specific property that may decrease the inherent toxic effects associated with **podophyllotoxin**.

Characteristic Features **Etoposide** is obtained as crystals from methanol having mp 236-251°C. Its physical parameters are $[\alpha]_D^{20} - 110.5^\circ\text{C}$ ($c = 0.6$ in chloroform), uv_{\max} (in absolute methanol) is 283 nm (ϵ 4245) and pKa 9.8.


Uses

1. It is employed in combination with other chemotherapeutic agents for refractory testicular tumours.
2. It is also used as a first line treatment in small cell lung cancer.
3. It has also been used extensively in the treatment of acute nonlymphocytic leukemias, non-Hodgkin's lymphomas, Hodgkin's disease, Kaposi's sarcoma, and neuroblastoma.

B. Teniposide

Synonyms Vumon, ETP, VM-26, Vehem-Sandoz, NSC-122819, 4'-Demethy-lepipodophyllotoxin- β -D-thenylidene glucoside.

Chemical Structure



Teniposide

Characteristic Features The characteristics features of the semi-synthetic derivative of podophyllotoxin are as follows the crystals obtained from absolute ethanol has mp 242-246°C. $[\alpha]_D^{20} - 107^\circ$ (in 9 : 1 chloroform/methanol), pKa 10.13 and uv_{\max} (in methanol): 283 nm ($E_{1\text{cm}}^{1\%}$ 64.1).

It differs from **etoposide** in the following respects:

- (i) It has an additional thenylidene ring on the glucopyranoside ring.
- (ii) Its pKa value is higher than that of **etoposide**.

Uses It is used as component of multiple-drug antineoplastic regimens for induction therapy in childhood acute lymphoblastic leukemia that is refractory to induction with other therapy.

(ii) Flavonoids: Interestingly, the *second* important class of the **biphenylpropanoid derivatives** is known as the **Flavonoids**. In general, **flavonoids** are amongst the most abundantly distributed natural

product compounds from the medicinal herbs having an enormous range of more than 2000 different compounds reported to be present either in the *free state* or as the *glycosides*. However, the ‘**flavonoid glycosides**’ have been described explicitly under Section 4.2.4 of Chapter 4 on ‘**Glycosides**’.

Mixed Biogenesis in Flavonoids **Flavonoids**, the aromatic compounds occurring in plants are usually biosynthesized by *three* different routes namely: (a) **acetate-malonate pathway**; (b) **acetate-mevalonate pathway**; and (c) **shikimic acid pathway**. Flavonoids have a mixed biogenesis, as evidenced by the fact that they are obtained from products of two or more of the main pathway.

The **flavonoid** and **isoflavonoid** ring structures are of mixed biosynthetic origin as depicted below in Fig. 6.1.




Fig. 6.1 Derivation of Flavonoid and Isoflavonoid Ring Systems.

From Figure 6.1, it may be observed that the ring A has been derived from *three* acetate units joined head-to-tail, whereas the ring B and the three carbon atoms of the pyran ring (*i.e.*, the central ring) are derived from cinnamic acid. It may, however, be observed that the acetate units are first and foremost get converted to CoA, wherein both the acetate-malonate and the **shikimic acid pathways** contribute to **flavonoid biosynthesis** exclusively.

Chalcones may be regarded as the precursors of all other classes of **flavonoids**. In fact, they have been isolated from a large number of plants, particularly members of the *Acanthaceae*, *Compositae*, *Gesneriaceae*, *Liliaceae*, *Oxalidaceae*, and *Scrophulariaceae*, wherein their presence can be apparently observed by their bright-yellow colouration to flower pigmentation. Fig. 6.2 represents the biogenetic relationship of the flavonoids.




Fig. 6.2 Biogenetic Relationship of Flavonoids.

* Hansel et al. *Deut Apotheken Ztg.*, **108**, 198 (1968).

Silybin is the active chemical constituent belonging to the **flavonoids**, which shall be discussed in details here under:


Silybin

Synonyms Silymarin, Apihepar, Laragon, Pluropon, Silarine, Silepan, Silirex, Silliver, Silmar.

Biological Source **Silybin** is obtained from the seeds of milk thistle, *Silybum marianum* (L.) Gaertn. (*Carduus marianus* L.) belonging to the natural order *Asteraceae*.

Chemical Constituents The seeds of milk thistle is chiefly comprised of *three* isomers, namely: **silidianin**, **silicristin** and the major component **silybin** (formerly known as **silymarin**). It has been recently characterized as a new class of substances termed as the **flavonolignans**. It has been more or less established beyond any reasonable doubt that **silybin** is produced in the plant by means of a radical **coupling of a flavonoid and coniferyl alcohol**.*

Chemical Structure



Silybin

Isolation A crude mixture of antihepatotoxic principle was first isolated from the plant (milk thistle) and designated as **silybin**.

Characteristic Features The anhydrous **silybin** has mp 158°C and it decomposes at 180°C. Its physical characteristics are as follows: $[\alpha]_D^{20} + 11^\circ$ (c = 0.25 in acetone + alcohol). Its uv_{max} (in methanol): 288 nm (log ϵ 4.33). It is soluble in acetone, ethyl acetate, methanol, ethanol and found to be sparingly soluble in chloroform. It is practically insoluble in water. It also occurs as the monohydrate crystals from a mixture of acetone and petroleum ether having mp 167°C (decomposes at 180°C); and from a mixture of methanol and water having mp 180°C.

Uses

1. **Silybin** is most importantly and widely employed as a therapeutic agent for protecting liver cells *in situ* or cell not yet irreversibly damaged by acting directly on the cell membranes (*i.e.* the targetted site) so as to prevent the entry of toxic substances.
2. It also augments and stimulates the 'protein synthesis' *i.e.*, anabolism of protein, thereby accelerating the process of regeneration and the production of *hepatocytes*.
3. It has also been experimentally proven that **silybin** binds specifically to a *regulative subunit of the DNA-dependent RNA polymerase I* at a particular site by mimicking a natural steroidal effector and thereby causing an activation of this enzyme. Consequently, the synthetic rate of ribosomal RNAs is increased considerably, thus leading to an enhanced formation of intact ribosomes that ultimately gives rise to an increased protein synthesis.

* Wagner *et al. Arzneimittel-Forsch* **18**, 688 (1968); and **24**, 466 (1974).

4. **Silybin** may also be employed as a supportive treatment for the management and cure of chronic inflammatory liver conditions and cirrhosis.*

General Biosynthesis of Flavonoids

The general biosynthesis of **flavonoids** essentially comprises of the interaction amongst the central intermediate *para*-coumaroyl CoA and three malonyl CoA units to elongate the side chain of the initial and original **phenylpropanoid** unit. It has been observed that the closure of the ring A yields the chalcone structure, while the follow up reaction ultimately closes the ring B as shown below in Fig. 6.3.




Fig. 6.3 General Flavonoid Biosynthetic Pathway
(Adapted from Robbers, J.E., *et al. Pharmacognosy and Pharmacobiotechnology*, Williams and Wilkins, London, 1996.)

6.2.6 High Molecular Weight Phenylpropanoids

Nevertheless, **phenylpropanoids**, play a vital role as building units in the formation of high-molecular weight polymers in plants. In general, these polymers are broadly classified into *two* major heads, namely:

- (i) Lignins, and
- (ii) Tannins

These two different categories of a **high-molecular weight phenylpropanoids** shall be discussed in details as under.

* It is marketed in the form of capsules which contains a concentrated extract equivalent to 140 mg of **silybin**.

6.2.6.1 Lignins

Lignins are the most abundant natural aromatic organic polymers found in virtually all vascular plants. It has been observed that the lignins together with cellulose, *q.v.*, and hemicellulose, *q.v.*, are the major cell wall components of the fibers of all *wood* and *grass* species in the plant kingdom. In fact, the **lignins** are sequestered in the secondary layer of the cell wall in close association with the cellulose matrix wherein the *phenolic hydroxy moieties* present in the lignins may be either hydrogen bonded or covalently attached to hemicellulose. **Lignin** is considered to be a sole factor in the contribution towards the strengthening of the cell-wall, and consequently for its synthesis. It is further regarded as a decisive factor in the adaptation of plants to a terrestrial habit in the process of evolution. It could only be possible by virtue of the fact that the lignified cell walls help to build the rigid and strong stems of the woody plants and the tress in general.

Composition **Lignin** is usually composed of coniferyl, *p*-coumaryl and sinapyl alcohols in varying proportion in a variety of plant species.

Uses

1. It is used as a source of vanillin, syringic aldehyde and dimethyl sulphoxide.
2. It is also employed as an extender for phenolic plastics.
3. It is used to strengthen rubber for shoe-soles.
4. It is used as an oil mud additive.
5. It also finds it application as a stablizer for asphalt emulsions.
6. It is employed to precipitate proteins.

6.2.6.2 Tannins

Synonyms Tannic Acid; Gallotannin; Gallotannic acid; *Acidum tannicum*.

Biological Sources **Tannic acid** usually occurs in the bark and fruit of a large number of plants, such as: roots of *Cimicifuga racemosa* (L.) Nutt. (*Ranunculaceae*)-**Black Cohosh, Black Snakeroot**; dried beans of *Coffea arabica* L. (*Rubiaceae*)-**Arabica Coffee, Arabian Coffee, Abyssinian Coffee**; barks of *Cornus florida* L. (*Cornaceae*)-**Dogwood, American Boxwood**; fresh forage (fodder) of *Equisetum arvense* L. (*Equisetaceae*)-**Field Horsetail**; leaves of *Eupatorium perfoliatum* L. (*Asteraceae*)-**Boneset, Ague Wood**; seeds of *Frangula alnus* Mill. (*Rhamnaceae*)-**Buckthorn**; roots of *Glycyrrhiza glabra* L. (*Fabaceae*)-**Common Licorice, Licorice Root, Spanish Licorice Root**; roots of *Paeonia officinals* L. (*Ranunculaceae*)-Peony; leaves of *Pilacarpus spp.* (*Rutaceae*)-**Jaborandi**; weeds of *Polygonum aviculare* L. (*Polygonaceae*)-**Prostrate Knotweed**; juice of the plant of *Rhamnus purshianus* DC. (*Rhamnaceae*)-**Cascara Sagrada, Cascara buckthorn**; flowers of *Tussilago farfara* L. (*Asteraceae*) **Coltsfoot, Coughwort, Horse-Hoof**; and plants of *Verbena officinalis* L. (*Verbenaceae*)-**Vervain, Verbena**.


Preparation **Tannic acid** is produced from Turkish or Chinese Nutgall usually formed by an aphid, *Schlechtendalia chinensis* found on the trees of *Rhus chinensis* belonging to family *Anacardiaceae*.

It may also be obtained by extraction from specially fermented oak **galls*** that are normally

* **Galls:** These are formed by virtue of the deposition of eggs by *gall-wasp viz., Adleria gallaetinctorial*.

grown on the young and tender twigs of *Quercus infectoria* (Oak Tree), belonging to the natural order *Fagaceae*.

Chemical Structure




Corilagin
(A Tannic Acid)

Characteristics Features **Tannic acid** is a yellowish-white to light brown, amorphous bulky powder or flakes, or spongy masses. It has a faint characteristic odour with a distinct astringent taste. It has a tendency to darken gradually on being exposed to air and light. It decomposes at 210-215°C mostly into pyrogallol and CO₂. It is highly soluble in water (1 g in 0.35 ml of water), 1 g in 1 ml of warm glycerol, and very soluble in acetone and ethanol. It is practically insoluble in chloroform, benzene, ether, petroleum ether, carbon disulphide and carbon tetrachloride.

Identification Tests

1. **Tannic acid**, when heated to 210-215°C, gets decomposed to yield **pyrogallol** and CO₂. The evolution of CO₂ may be confirmed when it turns freshly prepared lime-water milky.



2. It instantly gives rise to insoluble precipitates with albumin, starch, gelatin and a host of alkaloidal and metallic salts.
3. It readily forms a bluish-black colour or precipitate with ferric salts *e.g.*, FeCl₃.

Storage **Tannic acid** must be kept in well closed and protected from light containers.

Uses

1. Tannic acid with ferric salts are invariably used in the manufacture of inks.
2. It is used for tanning *i.e.*, making leather from hides of cow, goat, sheep and buffalo skin.
3. It is employed as a *pharmaceutical aid* due to its astringent and antiseptic actions.
4. It is used as a mordant in dyeing.
5. It is employed for sizing paper and silk.

6. It is also used for printing fabrics.
7. It finds its application to make imitation horn and tortoise shell when mixed with gelatin and albumin.
8. It is invariably employed to clarify beer and wine.
9. It is also used in 'photography'.
10. It is employed as a coagulant in the manufacture of rubber.
11. It is used in the large scale production of **gallic acid** and **pyrogallol**.
12. It is employed as a reagent in '**analytical chemistry**'.

6.3 BIOSYNTHESIS OF PHENYLPROPANOIDS

It has been observed that there are **two major precursors** for the **biosynthesis of phenylpropanoids**, namely: (a) **cinnamic acid**, and (b) **p-hydroxy-cinnamic acid (or p-coumaric acid)**. However, in plants these two chemical compounds are exclusively produced from the two aromatic amino acids phenylalanine and tyrosine, respectively, which are subsequently synthesised *via* the **Shikimic Acid Pathway** as depicted in Fig. 6.4.

Salient Features The salient features of the **biosynthesis of phenylpropanoids** *via* the **shikimic acid pathway** are enumerated below:

1. The biosynthetic pathway has been described explicitly in microorganisms by employing **auxotrophic mutants** of *Escherichia coli* and *Enterobacter aerogenes* which essentially require the aromatic amino acids for their normal growth.
2. Two *glucose metabolites*, namely: *erythrose 4-phosphate* and *phosphoenolpyruvate*, are found to react to give rise to a phosphorylated 7-carbon keto sugar, called 3-deoxy D-arabinoheptulosonic acid 7 phosphate (DAHP).
3. DAHP gets cyclized to 3-dehydroquinic acid, which is subsequently converted to **shikimic acid**.
4. The resulting **shikimic acid**, *via* a series of phosphorylated intermediates, gives rise to *chorismic acid*, which represents a vital branch-point-intermediate.
5. One of the branches leads to the formation of anthranilic acid and finally to tryptophan (an aromatic amino acid).
6. The second branch leads to the production of **prephenic acid**, which represents the last non-aromatic compound in the sequence.
7. **Prephenic acid** may be aromatized in *two* different manners, namely:
 - (a) First proceeds by dehydration and simultaneous decarboxylation to produce **phenylpyruvic acid**, which is the direct precursor of phenylalanine.
 - (b) Second takes place by dehydrogenation and decarboxylation to produce **p-hydroxy-phenylpyruvic acid**, which is the precursor of tyrosine.
8. **Cinnamic acid**, the phenylpropanoid precursor, is produced by the direct enzymatic deamination of phenylalanine.




Fig. 6.4 Biosynthesis of Phenylpropanoids via the Shikimic Acid Pathway
 (Adapted from Rebbers, J.E. *et al.* Pharmacognosy and Pharmacobiotechnology, Williams & Wilkins, London, 1996.)

9. ***para*-Coumaric acid** may be obtained in two different ways, namely:
- (a) By hydroxylation of cinnamic acid at the *para*-position, and
 - (b) By an analogous manner from tyrosine.

FURTHER READING REFERENCES

1. Canel C. *et al.*: **Molecules of Interest: Podophyllotoxin**, *Phytochemistry*, **54**: 115-120, 2000.
2. Cheeke, P.R., (ed.): **Toxicants of Plant Origin**, Vol. IV, Boca Raton, Florida, CRC Press Inc., 1989.
3. Davin L.B. and Lewis N.G.: **Dirigent Proteins and Dirigent Sites Explain the Mystery of Specificity of Radical Precursor Coupling in Lignan and Lignin Biosynthesis**, *Plant Physiol*, **123**: 453-461, 2000.
4. Forkmann G. and Heller W.: **Biosynthesis of Flavonoids**, *Comprehensive Natural Product Chemistry*, Vol. 1, Elsevier, Amsterdam, pp. 713-748, 1999.
5. Hahlbrock, K., School, D.: **Physiology and Molecular Biology of Phenylpropanoid Metabolism**, *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, **40**: 347, 1989.
6. Haslam, E.: **Plant Polyphenols: Vegetable Tannins Revisited**. Cambridge, Great Britain, Cambridge University Press, 1989.
7. Harborne, J.B., Mobry, T.J. (eds.): **The Flavonoids: Advances in Research Since 1980**, Chapman and Hill, Ltd., London, 1988.
8. Harborne J.B. and Williams C.A.: **Advances in Flavonoid Research Since 1992**, *Phytochemistry*, **55**: 481-504, 2000.
9. Hemingway, R.W., Lake, P.E., (eds.): **Plant Polyphenols: Synthesis Properties, Significance**, Plenum Press, New York, 1992.
10. Lewis, N.G., Davin, L.B., **Evolution of Lignan and Neolignan Biochemical Pathways: In: Isopentenoids and other Natural Products: Evolution and Function**, Nes, W.D., (ed.) American Chemical Society, Washington, D.C., 1994.
11. Lewis N.G. and Davis L.B.: **Lignans: Biosynthesis and Function**, *Comprehensive Natural Products Chemistry*, Vol. 1, Elsevier, Amsterdam, pp. 639-712, 1999.
12. Matern U. *et al.*: **Biosynthesis of Coumarins**, *Comprehensive Natural Products Chemistry*. Vol. 1, Elsevier, Amsterdam, pp. 623-637, 1999.
13. Murray, R.D.H., Mendez, J., Brown, S.A, **The Natural Coumarins**. John Wiley & Sons Ltd., Chichester, England, 1982.
14. Stafford, H.A., Ibrahim, R.K. (eds.): **Recent Advances in Phytochemistry**, Vol. 26, **Phenolic Metabolism in Plants**, Plenum Press, New York, 1992.

7

Alkaloids

- Introduction
- Classification of Alkaloids
- Alkaloids in Tissue Cultures
- Alkaloids in Chemosystematics
- Further Reading References

7.1 INTRODUCTION

The term **alkaloids** (or alkali-like) was first and foremost proposed by the pharmacist, W. Meissner, in 1819, for the basic nitrogen-containing compounds of plant origin.

Ladenburg defined **alkaloids**,—‘**as naturally occurring plant compounds having a basic character and containing at least one nitrogen in a heterocyclic ring.**’

With the advent of recent advanced knowledge in the chemistry of various alkaloids *two* more inevitable characteristic features were logically and justifiably added to the definition of **alkaloids**, namely:

- (a) Complex molecular structure, and
- (b) Significant pharmacological activity.

Furthermore, it was broadly observed that the basic properties of the **alkaloids** is solely by virtue of the presence of N-atom embedded into the *five*-or *six*- membered ring.

Therefore, the **alkaloids** are now generally defined as,—‘**physiologically active basic compounds of plant origin, in which at least one nitrogen atom forms part of a cyclic system.**’

Even this definition has a few anomalies as stated below, namely:

- (i) **Cholines and Betaines:** These two substances have the N-atom in the side chain and not in the aromatic ring as shown below:




Choline



Betaine

The **cholines** and **betaines** are regarded as simple alkylamines and not classified as **alkaloids**. They are designated by some school of thoughts as ‘**biological-amines**’ or ‘**proto-alkaloids**’.

- (ii) **Ephedrine:** It has the N-atom only in the side chain and not embedded in the aromatic ring as given below:




Ephedrine

- (iii) **Piperidine:** It is obtained from *Piper nigrum* (**Black Pepper**) and does not possess any pharmacological activity, but has a N-atom in a heterocyclic ring as given below:




Piperidine

- (iv) **Colchicine:** It is found to be neither basic nor it contains the N-atom in a heterocyclic ring, whereas it is considered as an alkaloid due to the fact it possesses distinct pharmacological activity as shown below:



Colchicine

- (v) **Thiamine (Vitamin B₁):** It conforms to the definition of **alkaloids** but is not regarded as an 'alkaloid' because of its almost universal distribution in living matter.



Thiamine Monochloride

Interestingly, **alkaloids** represent one of the most important group of chemical constituents occurring in the entire plant kingdom which exert extremely potent and vital physiological and pharmacological activities in the human beings. Therefore, it will be worthwhile to study the **alkaloids** with regard to the following aspects, namely:

- (i) Nomenclature
- (ii) Occurrence and distribution in different organs of the plant
- (iii) Site of formation of alkaloids in plants
- (iv) Functions of alkaloids in plants
- (v) Isomerism
- (vi) General characteristic features of alkaloids:
 - (a) Physical characteristics
 - (b) Chemical characteristics
- (vii) General methods of extraction and isolation of alkaloids.

These various aspects of **alkaloids** shall now be discussed adequately in a sequential manner so as to have a better in-depth of knowledge.

7.1.1 Nomenclature

The major characteristic of the nomenclature of **alkaloids** is the lack of any agreed systematic prevailing system. Therefore, by a general agreement, the chemical rules designate that the names of all alkaloids must end with the *suffix (-ine)*. The latin names end with (*-ina*). Thus, the names of the **alkaloids** are usually obtained in a number of ways, namely:

- (a) From the *generic name* of the plant producing them:
Examples: **Atropine** from *Atropa belladonna* Linn., (*Solanaceae*); and **Hydrastine** from *Hydrastis canadensis* L. (*Ranunculaceae*).
- (b) From the specific name of the plant yielding them:
Examples: **Belladonine** from *Atropa belladonna* L. (*Solanaceae*); and **Cocaine** from *Erythroxylum coca* Lam. (*Erythroxylaceae*).
- (c) From the common name of the drug producing them:
Example: **Ergotamine** from *Claviceps purpurea* (Er.) Tul. (*Hypocreales*) commonly known as **ergot**.
- (d) From their specific physiological activity:
Examples: **Emetine** from *Hedera helix* L. (*Araliaceae*) called **Ivy**; **Narcotine** from *Papaver somniferum* L. (*Papaveraceae*) known as **Opium Poppy**; and **Morphine** from *P. somniferum* L.
- (e) From the name of the discoverer:
Example: **Pelletierine** from the barks of *Punica granatum* Linn., (*Punicaceae*).
- (f) From their physical property:
Example: **Hygrine** from the roots of *Withania somniferum* (L.) Dunal (*Solanaceae*) called **Ashwagandha** (Hygro = moist).

7.1.2 Occurrence and Distribution in Different Organ's of Plant

McKee* (1962) reported that about 1000 **alkaloids**, which are known, belong to almost 100 families, 500 genera and spread over 1200 species. However, it has been observed beyond reasonable doubt

* McKee, H.S., *Nitrogen Metabolism in Plants*, (1962).

that the **alkaloids** are not evenly distributed amongst the plant kingdom. They have been found to be absent in *Algae* and in the lower groups of plants with the exception of one or two families of the fungi. The glaring examples of fungal alkaloids include those of ergot alkaloids.

The seeds of **papaya**, *Carica papaya* L. (*Caricaceae*), yield 660 to 760 mg BITC which is a bactericidal aglycone of **glucotropaeolin benzyl isothiocyanate**.

However, in the plant kingdom, the alkaloids generally, seem to get confined to a certain families and genera with regard to their distribution. For instance, amongst the angiosperms the families which have been recognized as outstanding for alkaloidal-yielding plants are, namely: *Apocynaceae*, *Berberidaceae*, *Papaveraceae*, *Ranunculaceae*, *Rubiaceae* and *Solanaceae*.

Monocotyledons, generally do not produce alkaloids, but investigation have revealed that *two* of the most promising families *viz.*, *Amaryllidaceae* and *Liliaceae* do contain alkaloid-containing plants.

Dicotyledons, mostly contain the **alkaloids**. It has been observed that neither *Labiatae* nor *Rosaceae* contain any alkaloid. Furthermore, almost 15% of all vascular plants contain alkaloids.





Alkaloids may occur in various parts of the plant. It may, however, be pointed out that in a particular species, normally only one or two specific organs and not all organs, essentially afford the function of alkaloidal formation. For instance, the **alkaloids** of the **tobacco plant**, *Nicotiana tabacum* Linn., (*Solanaceae*), are produced in the root and are subsequently translocated to the leaves where they usually accumulate, whereas the seeds are completely devoid of alkaloid. In another glaring example the **opium poppy**, *Papaver somniferum*, the alkaloids solely occur in the fresh latex of the fruit, while the seeds of poppy are virtually devoid of alkaloids. Likewise, the **colchicum*** corm. *Colchicum autumnale* Linn. (*Liliaceae*), the alkaloids are found both in the seed and in the corm. Interestingly, the bark of cinchona tree, *Cinchona officinalis* Linn., (*Rubiaceae*) contain the alkaloids (*viz.*, **quinine**) exclusively.

In some instances, there are noticeable fluctuations in the alkaloidal content in various organs of the plant during the different stages of its growth, during different seasons, and lastly between day and night. In certain perennials, the localization and accumulation of the alkaloids in one or two particular organs, appears to be more marked and pronounced with the advancement in the age of the plant.

In a broader sense, the particular **alkaloids** of complex structures are normally confined to specific plant families, such as: **hyoscyamine** in *Solanaceae*; and **colchicine** in *Liliaceae*. More importantly, a specific family may also contain several structurally non-related alkaloids *i.e.*, the basic-structure of alkaloids are altogether different, as exemplified under.


Salts of Alkaloids It has been found that a plethora of **alkaloids** occurring in various plant species are in the form of salts of organic acids, such as: acetic acid, malic acid, oxalic acid, succinic acid, tartaric acid, tannic acid or some other specific plant acids. In certain instances, the **alkaloids** are found to be in combination with **special plant acids**, for examples:

* **Colchicum**: It is derived from *Colchiss*, a port on the Black Sea where the plant was first found to be growing.


Family	Common Name	Botanical Name	Chemical Structure
Solanaceae	Nicotine	<i>Nicotiana tabacum</i> , or <i>N.rustica</i>	
	Hyoscyamine	<i>Hyoscyamus niger</i> L.; <i>Atropa belladonna</i> L.; <i>Datura Stramonium</i> L.,	
	Solanine	<i>Solanum tuberosum</i> L., (potato); <i>S. nigrum</i> L., <i>Lycopersicon esculentum</i> Mill. (tomato)	
	Capsaicin	<i>Capsicum annum</i> Linn, var (Chillies)	

Aconitine


associated with

**Morphine**


associated with

**Quinine**

associated with



Chelidonium associated with




Chelidonic acid

Rarely, alkaloidal salts with inorganic acids may also be present in plant products, such as: **morphine sulphate** in **opium poppy**.

Gluco-Alkaloids A few typical **alkaloids** are found in glycosidal combination with sugar moieties there by yielding the **gluco-alkaloids**.

Example: Solanidine (aglycone) obtained from the hydrolysis of the toxic glycoside **solanine** found in sprouts of potato tubers as given below:



7.1.3 Site of Formation of Alkaloids in Plants

The naturally occurring **alkaloids** that are found to be present in particular organs or parts of a specific plant, it does not logically suggest that they are either synthesized or formed in those particular organs. It may be further expatiated by the typical example of the **alkaloids** found in several **Datura** species and **Nicotiana** species, already discussed earlier, are invariably formed in the roots, but are rapidly translocated to the leaves. This glaring fact has been legitimately and explicitly demonstrated by the help of various experimental techniques innovated by researchers, namely: grafting techniques, labelling with radio-isotopes. Consequently, the leaves of such medicinal herbs, where the alkaloids usually get accumulated, is the ideal and vital part (organ) to be used for the subsequent extraction and isolation of relatively appreciable quantities of the alkaloids.

7.1.4 Function of Alkaloids in Plants

A good number of logical explanations, theories and principles have been put forward with regard to the possible function of **alkaloids** in plants or the probable reasons why they are present in them. It would be worthwhile to have a closer look and perhaps a better insight about certain possibilities that have gained cognizance over the years are described below along with their functions, namely:

- (a) As strategically located *poisonous agents* in plants thereby protecting them either against herbivorous animals or insects,
- (b) As probable by-products of various detoxification reactions representing a metabolic locking-up of compounds, otherwise harmful or detrimental to the plant,
- (c) As pronounced regulatory growth factors, and
- (d) As reserve substances in plant capable of supplying nitrogen or other necessary elements to its economy.

7.1.5 Isomerism

Generally, **isomerization** is the process of involving the change of one structure into another having the same empirical formula but with different properties in one or more respects.

A plethora of **alkaloids** contain one or more asymmetric carbon atoms in the molecule, and hence exhibit **optical activity**. It has been observed that in the majority of instances only the (–)-isomer (*i.e.*, the *levorotatory* component) has appreciable and distinct pharmacological activity than the corresponding (+)-isomer (*i.e.*, the *dextrorotatory* component) of the same alkaloidal species.

At this juncture, one needs to understand clearly the traditional designations *l*- and *d*- for the *levo*- and *dextro*- rotatory isomers respectively; and these are to be distinguished from the designations L- and D- which refer not to the *optical activity*, but to the *steric configuration* with regard to a conventionally accepted reference compound.


In fact, the optical activity is invariably associated with the **alkaloids** and their respective salts. However, the optical activity and the specific rotation usually varies with the solvent used, the temperature, the wave length of light and other minor factors.

There are quite a few typical and glaring examples that may serve to illustrate the considerable difference in the pharmacological activity observed amongst the different isomers of an **alkaloid**.


Examples

(a) Showing *d*- and *l*-isomers with distinct pharmacological activities, such as:

- (i) **Relative pressor activities* of D(–)-ephedrine and D(+)-ephedrine:** The relative pressor activities of **D(–)-ephedrine** is found to be 36 with regard to its **D(+)-ephedrine** isomer at 11 *i.e.*, the former is almost 3½ times more active than the latter as shown below:




D (–)-Ephedrine
(Relative pressure activity = 36)



D (+)-Ephedrine
(Relative pressure activity = 11)


* Increase of arterial blood-pressure.

- (ii) **Antimigraine activity of (-)-ergotamine and (+)-ergotamine:** It has been observed that the antimigraine activity of **(-)-ergotamine** possesses 3-4 times more activity than its corresponding (+)-ergotamine isomer:




(-)-Ergotamine

- (b) **Showing both (-)-and (+)-forms active pharmacologically:** In certain alkaloids, the (-) form as well as the (+) form are medicinally useful. **Examples:** The (-)-Quinine is primarily employed as a potent antimalarial agent; whereas the (+)-Quinine, also known as **quinidine**, is solely used in restoring cardiac arrhythmia to normal rhythm, as given below:



- (c) **Exception:** The (+) (+)-*d*-tubocurarine of *d*-tubocurarine is the only isomer that exhibits muscle relaxant properties, as shown below:

(+)-*d*-Tubocurarine Chloride

7.1.6 General Characteristics of Alkaloids

The general characteristics of **alkaloids** may be grouped together in *two* categories, namely:

- (a) Physical characteristics, and
- (b) Chemical characteristics.

These *two* categories shall now be discussed individually in the sections that follows:

7.1.6.1 Physical Characteristics

First and foremost, let us consider the solubility of alkaloids both in water and organic solvents along with some typical examples. In fact, a comprehensive knowledge of the solubility of complete range of alkaloids and their corresponding salts is of utmost pharmaceutical importance because of their extremely specific and potent pharmacological actions.

It is pertinent to mention here that in general the solubilities of different alkaloids and their respective salts usually exhibit considerable variation, which may be attributed from their extremely complex and varied chemical structures. However, it has been observed that the free alkaloid bases as such are invariably found to be fairly soluble in organic solvents, such as: ether, chloroform, relatively non-polar solvents (hexane, benzene, petroleum ether), immiscible solvent, lower alcohols (methanol, ethanol); but they are either practically insoluble or very sparingly soluble in water.

Interestingly, the alkaloidal salts are almost freely soluble in water, relatively less soluble in alcohol and mostly either insoluble or sparingly soluble in organic solvents:

Examples **Atropine sulphate** and **morphine hydrochloride** are much more soluble in water than their corresponding bases *i.e.*, **atropine** and **morphine**.

However, there are a few exceptions to the above stated generalizations, namely:

- (i) Certain **alkaloid** bases are water soluble, but these may be solely regarded as exceptions rather than any specific rule, such as: **ephedrine**, **colchicine**, **pilocarpine**; the **quaternary alkaloid-base** like **berberine** and **tubocurarine**; **caffeine-base** readily extracted from tea with water.
- (ii) **Narceine** and **pilocarpine** are insoluble in organic solvents, whereas morphine is sparingly soluble in organic solvents *viz.*, solubility in either 1:5000.
- (iii) Certain alkaloidal salts, for instance: **lobeline hydrochloride** and **aprotropine hydrochloride** are found to be soluble in organic solvent like chloroform.
- (iv) Some alkaloidal salts are sparingly soluble in water whereas others are extremely water-soluble, such as: **Quinine sulphate**-soluble in 1:1000 parts of water, **Quinine hydrochloride**-soluble in 1:1 part of water.

The physical characteristics of some potent **alkaloids**, such as: mp, optical rotation and solubility are enlisted below so as to have a glimpse of the distinct variation in the observed parameters:

S.No.	Alkaloid	mp (°C)	Optical rotation	Solubility
1.	Ajmaline	150-160	$[\alpha]_D^{20} + 144^\circ$	Chloroform, ether, ethanol, methanol
2.	Atropine	144-116	–	Benzene, chloroform, ether
3.	Berberine	145	–	Water
4.	Colchicine	142-150	$[\alpha]_D^{17} - 429^\circ$	Water, chloroform, ethanol
5.	Ephedrine	79	–	Water, ethanol, ether, chloroform, oils
6.	Hyoscyamine	108.5	$[\alpha]_D^{20} - 21.0^\circ$	Ethanol, dilute acids
7.	Morphine	197	$[\alpha]_D^{25} - 132^\circ$	Sparingly soluble in ethanol, chloroform, amyl alcohol,
8.	Physostigmine	105-106	$[\alpha]_D^{25} - 120^\circ$	Benzene, chloroform, oils
9.	Quinine	177	$[\alpha]_D^{17} - 117^\circ$	Chloroform, ether
10.	Reserpine	264-265 (dec.)	$[\alpha]_D^{23} - 118^\circ$	Chloroform, ethyl acetate, benzene.
11.	Strychnine	275-285	$[\alpha]_D^{18} - 104.3^\circ$	Chloroform, methanol, benzene
12.	Taxol	213-216 (dec.)	$[\alpha]_D^{20} - 49^\circ$	Methanol
13.	Vinblastine	211-216	$[\alpha]_D^{20} + 42^\circ$	Chloroform, ethanol
14.	Yohimbine	234	$[\alpha]_D^{20} + 108^\circ$	Chloroform, ethanol, benzene

7.1.6.2 Chemical Characteristics

The general chemical characteristics of the **alkaloids** are so broadly spread out, therefore, they shall be treated individually under the following heads, namely.

[A] N-in the Molecule Besides, the other normal elements *e.g.*, carbon, hydrogen, oxygen, the alkaloids must essentially contain at least one N-atom. The number of N-atoms vary from the bear minimum one in a molecule *e.g.*, **cocaine**, to even five in a molecule *e.g.*, **ergotamine**. It has been observed that these N-atoms are normally present as a part of the **heterocyclic ring** in the alkaloid molecule *e.g.*, **quinine, reserpine, strychnine, vinblastine** and **yohimbine**; whereas there are certain alkaloids that contain the N-atom in the aliphatic side chain *e.g.*, **ephedrine, mescaline**.

Invariably, the alkaloids contain the N-atom in the **tertiary-amine form** (R_3N) *e.g.*, **morphine, reserpine**; lesser in the **secondary-amine form** (R_2NH) *e.g.*, **ephedrine**; and very rarely in the **primary-amine form** (RNH_2) *e.g.*, **nor-pseudo-ephedrine**. Furthermore, whenever N-atom occurs either in the *tertiary-* or *secondary-*form, it essentially constitutes as an integral part of the ring-system, precisely the heterocyclic ring system.

Noticeably, the tertiary N-atoms wherein only two of the bonds are involved in a ring, the methyl moiety is usually found as the third component, for instance: N-methyl group in **morphine, cocaine, colchicine, dextro methorphan, codeine, physostigmine, vinblastine, vindesine** etc.

Hence, methyl moiety seems to be the only **alkyl group** that has been found to be substituted on the **N-atom**.


However, in some very specific cases, the N-atom occurs in the **quaternary ammonium form** ($R_4N^+ \cdot X^-$) *e.g.*, **tubocurarine chloride** [see section 7.1.5 (c)]. Nevertheless, the quaternary ammonium compounds are logically and technically not regarded as alkaloids by virtue of the following *two* particular reasons, namely:

- (i) N-atom does not possess a H-atom, and
- (ii) Chemical properties are quite different.


As a matter of convenience, they are legitimately grouped along with the *alkaloids*.

[B] O-in the Molecule In addition to the common elements C, H and N, a variety of **alkaloids** normally contains O-atom. Invariably, these specific alkaloids are found in the solid state, with a few exceptions where the oxygenated alkaloids usually occur as non-volatile liquids, such as: **pilocarpine**.


[C] Basicity (Alkalinity) In general, the **alkaloids** are basic (alkaline) in reaction, by virtue of the presence of N-atom present in the molecule. Hence, these are prone to the formation of their respective salts with various acids.



Pilocarpine
(Liquid, mp34°C)



Morphine
(Solid, mp197°C)



Strychnine
(Solid, mp 268–270°)

Degree of Basicity: The degree of basicity of the **alkaloids** mostly depends upon the prevailing influence caused due to the electrostatic status of the N-atom present in the alkaloid molecule, for instance, the number of N-atom present in the alkaloid, whether the N-atom is located in the ring or in the side-chain, the presence of alkyl group (*e.g.*, methyl) to the N-atom etc.

Another vital factor, which establishes the degree of basicity of an alkaloid, is the presence of *pri*-, *sec*-, *tert*-, or *quaternary* N-atom or atoms in it. In fact, such apparent differences in the degree of basicity arising from the various structural features, are eventually reflected by the different dissociation constant values (*i.e.*, pKa values) with regard to various alkaloids as stated below:


S.No.	Alkaloid	pKa Values
1	Berberine	2.47 ($K = 3.35 \times 10^{-3}$)
2	Colchicine	12.35 (at 20°C)
3	Emetine	$pK_1 = 5.77$; $pK_2 = 6.64$
4	Morphine	9.85
5	Papaverine	8.07 (at 25°C)
6	Physostigmine	$pK_{a1} = 6.12$; $pK_{a2} = 12.24$;
7	Quinine	$pK_1 = 5.07$ (at 18°C); $pK_2 = 9.7$;
8	Reserpine	6.6
9	Vinblastine sulphate	$pK_{a1} = 5.4$; $pK_{a2} = 7.4$;
10	Vincristine	5.0 ; 7.4 (in 33% DMF)

Salient Features

1. The weaker bases, *i.e.*, **alkaloids** having low pKa values, shall require a more acidic medium to form their respective salts with the corresponding acid.
2. The strongly basic **alkaloids** *i.e.*, those possessing high pKa values, shall require comparatively low acidic medium to form their respective salts with the acid.


Note: In a medium at a weakly acidic pH certain strongly basic alkaloids would be easily converted to their respective salt by interaction with the corresponding acid, whereas the alkaloids which are relatively weaker bases having lower pKa values shall still remain in their free-base form. Such a critical situation is skillfully exploited for the separation of a specific alkaloid or a group of alkaloids having closely identical pKa values, from other alkaloids that essentially possess either very low or very high pKa values.

3. The **alkaloids** are usually neutralized with acids to form salts that may be converted to the corresponding **free-base** by the cautious addition of selective weak bases, such as, ammonia, calcium hydroxide or sodium carbonate. The usage of either NaOH or KOH solutions must be avoided so as to prevent the decomposition or destruction of highly sensitive alkaloids.
4. **Amphoteric alkaloids:** There are some **alkaloids** which are amphoteric in nature *i.e.*, they are neither acidic nor basic in character; this is due to the presence of phenolic (–OH) moiety in **Morphine**, or the presence of carboxylic (–COOH) function in **Narceine**, as shown below:




Narceine

5. **Unstable alkaloidal salts:** There exists some specific **alkaloids** that inherently possess *weak-basic properties* and their salts are not so stable, for instance: **piperine**, **papaverine**, **narceine**, **narcotine**, and **caffeine**.




Piperine

Papaverine




Narcotine




Caffeine

6. **Neutral or slightly acidic alkaloids:** There are a few typical naturally occurring **alkaloids** that almost behave as either neutral or slightly acidic character, namely: ricinine and theophylline, as depicted below:



Ricinine



Theophylline

[D] Precipitation by Specific Reagents A good number of **alkaloids** obtained from various plant sources invariably give a distinct precipitate with certain specific reagents to an extent as small as *one microgram*. Based on these observations, these **alkaloid-precipitating reagents** are sometimes employed for either detecting the presence or absence of alkaloids in:

- Crude extracts or plant materials, and
- For ascertaining whether a specific extraction procedure has exhausted completely the alkaloidal contents or not.

However, a negative test *i.e.*, the absence of precipitation, may infer that the alkaloids are absent. It is pertinent to mention here that a positive test may not always indicate the presence of alkaloids, but may also be due to the presence of other plant constituents, such as: purines, proteins, betaines and ammonium salts etc. Therefore, it is always desired to rule out the possibility of a *false-test* by alkalifying the acidic solution with dilute ammonium hydroxide and subsequently extracting the liberated alkaloid with chloroform. The residue thus obtained, after the removal of the solvent (chloroform), is tested with the **alkaloid-precipitating reagents**. Now, if the test is positive, the presence of an alkaloid is almost confirmed.

Microcrystalline precipitates of alkaloids: **Alkaloids**, alike other amines, usually form *double-salts* with salts of heavy metals, such as, gold (Au), mercury (Hg) and platinum (Pt). The resulting double salts are found to be possessing characteristic microcrystalline structures. It has been observed

that under controlled and specific experimental parameters *viz.*, profile of mixing and gradual evaporation, a drop of an alkaloidal solution reacting with a drop of an appropriate alkaloidal-precipitating reagent, such as: chloroplatinic acid (H_2PtCl_6) or chlorauric acid ($\text{HAu} \cdot \text{Cl}_4$), on a microscopic-glass slide, gives rise to microcrystalline products having specific and characteristic shapes and structures solely based upon the manner of aggregation.* It may, however, be exploited skillfully as a convenient means of rapid-microscopical identification of an **alkaloid**.

The various reagents that are invariably used either for the testing of **alkaloids** by precipitation or by the formation of microcrystalline complexes (salts) are as stated below along with their individual compositions, namely:

(i) **Mayer's Reagent (Potassium-Mercuric Iodide Test Solution):**

Mercuric chloride	=	1.36 g
Potassium Iodide	=	3.00 g
Distilled water to make	=	100.00 ml

(ii) **Wagner's Reagent (Potassium Triiodide):**

Iodine	=	1.3 g
Potassium	=	2.0 g
Distilled water to make	=	100.00 ml

(iii) **Kraut's Reagent (Modified Dragendorff's Reagent or Potassium Bismuth Iodide):**

Bismuth Nitrate	=	8.0 g
Nitric Acid	=	20.0 ml
Potassium Iodide	=	27.2 g
Distilled water to make	=	100.00 ml

(iv) **Marme's Reagent (Potassium-Cadmium Iodide Reagent):**

Cadmium Iodide	=	10.0 g
Potassium Iodide	=	20.0 g
Distilled water to make	=	100.00 ml

(v) **Scheibler's Reagent (Phosphotungstic Acid Reagent):**

Sodium Tungstate	=	20.0 g
Disodium Phosphate	=	70.0 g
Distilled water to make	=	100.00 ml

Note: Acidify with nitric acid to litmus paper.

(vi) **Hager's Reagent:**

A saturated solution of *Picric Acid*.

(vii) **Sonnenschein's Reagent (Phosphomolybdic Acid):**

A 1% (w/v) solution of phosphomolybdic acid in ethanol.

(viii) **Bertrand's Reagent (Silicotungstic Acid):**

A 1% (w/v) solution of silicotungstic acid in distilled water.

(ix) **Reineckate salt solution:**

Ammonium Reineckate	=	1.0 g
$\text{NH}_4 [\text{Cr} (\text{NH}_3)_2 (\text{SCN})_4]$		

* A whole combination of several components.

Hydroxylamine HCl = 0.3 g
Ethanol = 100.00 ml

Note: Filter and store in a refrigerator.

[E] Colour Reactions with Specific Reagents Broadly speaking the colour reactions of the **alkaloids** are rather unspecific; however, they are certainly very sensitive so much so that even alkaloids present in microgram quantities invariably afford immediate and instant response. The ultimate development of a characteristic colour reaction is solely dependent upon either the dehydration or the oxidation of the alkaloid. Generally, a large number of these reagents essentially consist of concentrated sulphuric acid along with certain specific added compounds, such as, sulphomolybdic acid, formaldehyde, sulphovanadic acid, potassium arsenate, hydrogen peroxide, and selenious acid.

A number of such **specific reagents** shall be described in the section that follows:

(a) **Froehd's reagent:** Dissolve 5 mg of molybdic acid or sodium molybdate in 5 ml of pure concentrated H_2SO_4 .

Note: The reagent should be freshly prepared before use.

(b) **Erdmann's reagent:** A mixture of 10 drops of concentrated HNO_3 , and 100 ml of water are added to 20 ml of pure concentrated H_2SO_4 .

(c) **Marqui's reagent:** A mixture of 2-3 drops of formaldehyde solution (40%) with 3 ml of concentrated H_2SO_4 .

(d) **Mandalin's reagent:** Dissolve 1 g of finely powdered ammonium vanadate in 200 g of pure conc. H_2SO_4 .

(e) **Mecke's Reagent:** Dissolve 1 g of selenious acid in 200 g of pure concentrated H_2SO_4 .

(f) **Modified Dragendorff's reagent:** Dissolve 1.6 g of bismuth subnitrate in 60 ml of 20% glacial acetic acid, add to it 5 ml of 40% aqueous solution of KI, 5ml of glacial acetic acid and make up the volume to 100 ml of water.

(g) **Rosenthaler's reagent:** Dissolve 1 g of potassium arsenates in 100 g of pure concentrated H_2SO_4 .

(h) **Schaer's reagent:** Mix carefully 1 volume of pure 30% H_2O_2 with 10 volumes of concentrated H_2SO_4 .

Note: The reagent is always prepared afresh, before use.

Interestingly, there are some instances where in the intensity of the colour so produced is in *linear proportion* under standardized experimental parameters. Therefore, such specific colour reactions may be used exclusively for the quantitative determination of certain groups of **alkaloids**, such as:

- (i) **For Ergot Alkaloids:** The blue colour produced by the ergot alkaloids with the **Van Urk Reagent** (or **Ehrlich Reagent**) *i.e.*, *para*-dimethylaminobenzaldehyde in 65% H_2SO_4 , is employed for the quantitative estimation of **ergot alkaloids**.
- (ii) **For Belladonna Alkaloids:** The violet colour caused by the belladonna alkaloids with fuming HNO_3 and alcoholic KOH solution is employed for their assay.

[F] Stability of Alkaloids Alkaloids, in general, are not very stable. They normally undergo degradation or decomposition on being exposed to air, light, moisture and heat, besides chemical reagents. A few typical examples of alkaloids *vis-a-vis* their stability are stated below, namely:

- (i) Ergotamine gets destroyed by prolonged treatment with alkali, whereas strychnine can stand such vigorous action.
- (ii) An aqueous solution of alkaloids undergo rapid decomposition or degradation as compared to their solid forms.
- (iii) Storage of **alkaloids** in pure form or their dry extracts is usually done in a vacuum desiccator over a dehydrating agent *e.g.*, phosphorous pentoxide (P_2O_5) or calcium chloride ($CaCl_2$) anhydrous for their better stability.
- (iv) During the course of extraction of **alkaloids** followed by isolation, the solvent is preferably removed effectively by distillation under vacuum* (or reduced atmospheric pressure) or by subjecting it to evaporation in a Rotary Thin-Film Evaporator under vacuum so that the desired product is not exposed to excessive heat, thus avoiding decomposition.
- (v) **Alkaloids**, are stored in amber-coloured glass bottles preferably in a vacuum desiccator.

[G] Acid salts of Alkaloids A plethora of **alkaloids** are strongly alkaline in nature and most of them form well-defined salts. However, in certain instances the basicity of an **alkaloid** is quite weak and feeble, and hence the formation of the corresponding salts with either acetic or other weak acids is practically insignificant and rare. The salts formed with stronger acids *e.g.*, HCl, H_2SO_4 etc., get decomposed in the presence of water to liberate the free base and the acid. It has been observed that only a few of the alkaloids form carbonates, and consequently either the alkali carbonates or the alkali hydrogen carbonates are invariably used to liberate them from the aqueous solutions of their corresponding salts.

Alkaloids, in general, containing either one or more than one N-atom usually behave as **monoacidic bases**; and, therefore, form only one series of salts with acids as designated by 'BA' (where: B = base; and A = acid). It is pertinent to mention here that **quinine** in particular and the **cinchona alkaloids** in general are an exception to the earlier concept and found to behave as *diacidic bases*. Besides, a number of alkaloids to behave as monoacidic bases, even though they contain two N-atoms in their molecule. It is worthwhile to mention here that the basicities of the alkaloids is of utmost importance with regard to their quantitative volumetric estimation.

In common practice the salts of alkaloids are prepared by using cold and dilute solutions of the mineral acid specifically, *e.g.*, **morphine hydrochloride**, **atropine sulphate**, **quinine sulphate**, **ephedrine hydrochloride** etc. It may be pointed out that the use of concentrated mineral acids, or heating an alkaloid even with a dilute acid under pressure may ultimately lead to profound changes in them. Noticeably, the concentrated mineral acids invariably give rise to characteristic colour changes, that are usually used as a means of identification and characterization of the alkaloids. In addition to the complete decomposition of **alkaloids** by strong acids to result the various colour changes, the chemical changes caused by the mineral acids on them may be categorized into *three* different types, namely:

* Under vacuum (or reduced atmospheric pressure) the boiling point of solvent is lowered significantly *e.g.*, alcohol bp 78.5°C boils in vacuum at 40°C.

(a) **Dehydration:** Dehydration of **alkaloids** give rise to either *anhydro-* or *apo-* alkaloids, such as:

Apomorphine obtained from **Morphine**


Apoatropine obtained from **Atropine**

(b) **Demethoxylation:** The removal or elimination of the methoxyl groups from the alkaloids by treatment with either concentrated HCl or HI to produce methyl chloride (CH₃Cl) or methyl iodide (CH₃I) while giving rise to the corresponding *hydroxy base*. The methoxyl group (s) are present in a variety of alkaloids, for instance: **codeine, quinine, narcotine** and **papaverine**.

Example:



(c) **Hydrolysis:** A good number of **naturally occurring alkaloids** are obtained as *esters*. They easily undergo hydrolysis on being heated with either alkalis or mineral acids thereby resulting into the formation of the corresponding acids along with respective alcohols or phenols of the alkaloids. A few typical examples are as give below:



[H] Action of Alkalies The action of alkalies *e.g.*, NaOH and KOH on the **alkaloids** are found to be varying in nature as enumerated below:

- (a) Dilute alkaline solutions of KOH or NaOH normally decompose most alkaloidal salts and finally liberate the free alkaloids.
- (b) Certain alkaloids containing phenolic hydroxyl groups *e.g.*, **morphine**, on being treated with alkaline solutions yield, their corresponding soluble sodium or potassium salts.
- (c) The ester alkaloids usually undergo hydrolysis on being treated with dilute alkalies, such as: atropine, cocaine.
- (d) **Racemic Isomeride:** The action of alkali hydroxides on **hyoscyamine** in alcohol gives rise to the racemic isomeride atropine.
- (e) Fusion of alkaloids with dry KOH or NaOH by the application of heat ultimately leads to drastic decomposition of the former thereby yielding ultimately the simple heterocyclic bases, for instance: pyridine, quinoline, pyrrolidine etc.
- (f) Simple fusion of **alkaloids** with alkali hydroxides may give rise to distinct and visible colour changes.

[I] Pharmacological Activity The **alkaloids** exhibit a wide-spectrum and complete diversity of complex structures which ultimately is responsible for their extra ordinary broad-range of pharmacological activities covering both the cardio-vascular and central nervous system. It has been observed beyond any reasonable doubt that most alkaloids usually exert certain specific and definite pharmacological action. Moreover, a small quantity of an alkaloid (0.1–1.0 mg) may bring about a marked and pronounced pharmacological action on various organs and tissues both of animal and human origin. However, the potency of an individual alkaloid varies from one another widely and profusely.

A few typical pharmacological actions of some **alkaloids** are stated below showing their broad-spectrum of activities, namely:

S.No.	Alkaloid	Pharmacological Action
1	Morphine	Narcotic analgesic
2	Codeine	Expectorant, analgesic
3	Brucine	CNS-Stimulants
4	Strychnine	CNS-Stimulants
5	Ergotamine	Uterine muscle contractions
6	Atropine	Mydriatics
7	Homotropine	Mydriatics
8	Pilocarpine	Myotics
9	Physostigmine	Myotics
10	Ephedrine	Hypertensive
11	Reserpine	Hypotensive
12	Quinine	Antimalarial
13	Caffeine	CNS-stimulant
14	Tubocurarine	Neuromuscular blocking action
15	Emetine	Antiprotozoal action
16	Hyoscyamine	Relief of spasms of urinary tract
17	Cocaine	CNS-stimulant
18	Colchicine	Anti-gout
19	Lobeline	Treatment of asthma
20	Arecoline	Parasympathomimetic action
21	Protoberatrine A	For management of hypertension in pregnancy
22	Conessine	Antiprotozoal and antiamebic
23	Vasicine	Expectorant and bronchodilator
24	Vinblastine	Antineoplastic
25	Vincristine	Antineoplastic
26	Piperine	Carminative, stomachic
27	Heroin	Narcotic analgesic
28	Hyoscyne	Motion sickness (sedation)
29	Theophylline	Smooth muscle relaxant
30	Aconitine	Treatment of neuralgia, sciatica, rheumatism and inflammation.

7.1.7 General Methods of Extraction and Isolation of Alkaloids

The general methods of extraction and isolation of the alkaloids from the plant sources one has to take into consideration the following steps in a sequential manner, namely:

- (i) Separation of the alkaloid(s) from the main bulk of the non-alkaloidal substances,
- (ii) Most of the alkaloid-containing plants, several alkaloids having closely related chemical structures are normally present, such as: the cinchona alkaloids consist of more than twenty-five alkaloids. There is hardly any known plant source that contains only one alkaloid exclusively,
- (iii) Separation of each individual alkaloid from the mixture of alkaloids obtained from a particular plant source (*e.g.*, **cinchona bark**) using latest separation techniques, for instance, preparative **high-performances liquid chromatography, (HPLC)** column chromatography, by the help of chromatotron, and **high-performance thin-layer chromatography (HPTLC)**.

Nevertheless, the general methods of isolation of alkaloids largely depend upon several vital factors, for instance: (a) the alkaline nature of most alkaloids, (b) the ability and ease of formation of alkaloidal salts with acids, and (c) the relative solubilities of the resulting alkaloidal salts either in polar organic solvents *e.g.*, ethanol, chloroform, isopropanol etc., or in aqueous medium.

The general methods of **extraction of alkaloids** from the plant sources solely depend upon the purpose and scale of the operation (*e.g.*, pilot scale or commercial scale). It is also based on the quantum and bulk of the raw material to be employed in the operation. Of course, for research purposes column chromatography using ion-exchange resins have been used successfully and effectively to strip the plant materials of their alkaloidal contents. However, in the commercial scale large volumes of aqueous extracts of plant materials are normally pumped through huge metallic columns packed with cationic resins, which in turn pick up all basic components (cations). Subsequently, the alkaloids (*i.e.*, the basic components are conveniently washed off by flushing the column with a moderately strong acid. The column having the cationic resins can be reused once again for the next drug substances.

By the advent of the latest separation techniques and the copious volume of informations accumulated through the intensive and extensive research carried out with regard to the conventional processes essentially associated with the separation as well as isolation of the hundreds of **alkaloids** from the natural plant sources, the following *five* steps are most important and vital, namely:

- (i) Sample preparation
- (ii) Liberation of free alkaloidal base
- (iii) Extraction of alkaloidal base with organic solvent
- (iv) Purification of crude alkaloidal extract
- (v) Fractionation of crude alkaloids

All these *five* steps shall be discussed individually as under:

7.1.7.1 Sample Preparation

The first and foremost step is the sample preparation. The plant material is reduced to a moderately coarse powder by appropriate means using grinders and sieves, to facilitate maximum effective contact of the solvent with the ruptured alkaloid bearing tissues and cells. In the case of plant substances that are rich in oils and fats, such as: seeds, kernels, these non-alkaloidal chemical components need to be eliminated completely by extraction with a suitable non-polar solvent like n-hexane, light petroleum ether, in a soxhlet apparatus, which would not extract the alkaloids in question. However, it is always advisable to shake the light-petroleum ether or n-hexane fraction with a dilute mineral acid and subsequently test the acidic solution for the presence of alkaloids.

7.1.7.2 Liberation of Free Alkaloidal Base

It has been observed that the **alkaloids** invariably occur in the plant sources as the salt of acids, such as: oxalates, tannates etc. Therefore, when the plant substance is exposed to an alkaline medium, the alkaloidal salts are readily converted to the corresponding alkaloid bases.

Choice of Alkali Indeed, the choice of a suitable *mineral base* (alkali) for the ease of liberation of the alkaloid from the salts is not only very vital but also equally significant and largely depend on the following factors, namely:

(a) **Natural state of the alkaloids:** It has been observed that the salt of a *strongly basic alkaloid* with a mineral acid usually tends to undergo cleavage under the influences of a stronger base. Likewise, the corresponding salt of a *weakly basic alkaloid* and a relatively weak organic acid shall require a rather weaker base for its cleavage.

(b) **Chemical characteristics of the alkaloidal base:** The usage of strong alkali *e.g.*, NaOH or KOH should be avoided as far as possible by virtue of the fact that certain alkaloids undergo hydrolysis on prolonged contact with a strong base.

Example

- (i) **Hydrolysis of ester-alkaloids, *e.g.*, cocaine, hyoscyamine;**
- (ii) **Phenolic alkaloids *e.g.*, cephaeline, morphine.** These **alkaloids** normally get solubilized while in contact with a strong alkali and, therefore milder alkaline reagents *e.g.*, dilute ammonia solution are necessary for their liberation.

(c) **Presence of fatty substances:** The usage of strong alkali is strictly prohibited in the case of fat containing plant materials because of the formation of saponified products causing troublesome emulsions. In such cases, it is always preferred to defat the plant substance before proceeding for the liberation of free alkaloids.

Ammonium Hydroxide Solution Dilute aqueous ammonium hydroxide solution is one of the choicest alkali most frequently used for the liberation of alkaloids from the plant sources. It enjoys a two-fold advantage. First, being its adequate alkalinity to liberate most of the common alkaloids, and second by, its volatile nature so that it may be removed by evaporation of the solvent. As it has a tendency to be extracted by solvent ether from the aqueous solution, therefore, it is almost necessary to get rid of it by evaporation and subsequent washing repeatedly. In normal practice, usually even the last traces of ammonia are removed when the combined ethereal extract is reduced to half of its original volume under vacuum.

NaOH or KOH Solution The **alkaloids** that occur naturally as their tannate salts specially require either NaOH or KOH solution for their subsequent liberation. In certain typical instance even the use of KOH or NaOH fails to cleave the tannate salts because of their intimately strong bondage with the alkaloid and extremely insoluble nature.

Example

- (i) **Cinchona Bark:** It has got to be treated first by heating with dilute HCl so as to decompose the salts and liberate the alkaloids in the form of water soluble hydrochlorides, and
- (ii) **Pomegranate Bark:** It does not have the tannin so tenaciously bound to the alkaloids as in the case of cinchona bark. Hence, NaOH solution is strong enough to cause an effective split of the alkaloidal salts. It also acts to control the solubility of the water-soluble pomegranate alkaloids by preventing their dissociation.

7.1.7.3 Extraction of Alkaloidal Base

The extraction of alkaloidal base may be accomplished by *three* different types of solvents that are discussed below, namely:

[A] Extraction with Water-Miscible Solvents A plethora of **alkaloids** and their respective salts are soluble in alcohols, such as: methanol, ethanol, isopropanol; therefore, these very solvents may also be employed for the extraction of the plant substances. The usual pretreatment of the crude drug with alkali may be avoided completely, because alcohol appears to affect dissolution of not only the *alkaloidal salts* but also the *free bases* found in the plant substances. It is, however, believed that alcohol predominantly exerts a *hydrolyzing effect* upon the alkaloidal tannates and other salts. In

actual practice, neither pretreatment of the crude drug with an alkali nor acidification of the alcohol with a small amount of a mineral acid or an organic acid is required.

Note

1. **The penetration and hence the subsequent extraction of the crude drug is almost complete with the help of four successive extractions with an alcohol. Further, the loss of solvent is comparatively less than the chlorinated solvents e.g., chloroform.**
2. **The extraction of total alkaloids with alcohol is highly recommended because of its maximum efficiency and economical viability.**

[B] Extraction with Water-Immiscible Solvents In reality, the most widely used water-immiscible solvents for the extraction of alkaloids are: chloroform, diethyl ether (solvent ether) and isopropyl ether. However, a few other specific organic solvents, namely: ethylene chloride, carbon tetrachloride and **benzene*** may be employed with an evident advantage for certain specific alkaloids. Interestingly, *chloroform* is regarded as the choicest water-immiscible solvent for a broad-spectrum of alkaloids present in the plant kingdom and extracts them with varying degrees of ease.

Note: Chloroform is not suitable for the extraction of quaternary alkaloids e.g., tubocurarine.

[C] Extraction with Water The crude drug is subjected to extraction with water previously acidified with dilute solution of HCl, H₂SO₄ or CH₃COOH, which is subsequently rendered alkaline, preferably with dilute NH₄OH solution and finally extracted with a water-immiscible solvent as stated in [B] above.

Undoubtedly, water being an excellent and absolutely inexpensive polar solvent for the extraction of alkaloids, but it offers an enormous volume of disadvantages because it carries along with it a large number of other plant components, for instance: sugar, pigments (e.g., **chlorophylls**), starches, tannins, proteins etc., which ultimately puts across a colossal waste of time, energy and chemicals. Hence, its usage has been resulting to a bear minimum level.

In general, the alkaloids may be extracted by any of the following *three* well-defined and widely accepted processes, namely:

- (a) Soxhlet Extraction Process
- (b) Stas-Otto Process, and
- (c) Kippenberger's Process.

All these three processes shall now be discussed briefly in the sections that follows:

(a) Soxhlet Extraction Process: The soxhlet assembly is a continuous extractor which is generally suitable for the extraction of alkaloids from powdered plant materials with the help of organic solvents. In this instance, the powdered drug is usually moistened with dilute ammonia solution and then packed loosely in the thimble of the Soxhlet apparatus; and the organic solvent affords a deep penetration of the moist drug thereby allowing the greatest possible extraction of the alkaloids from the exposed surfaces of the cells and tissues of the crude drug. Once, the extraction is ascertained to have completed, the solvent is filtered and evaporated in a **Rotary Thin-Film Evaporator** and the residue is treated further for the isolation of individual alkaloids.

* **Benzene:** It is a carcinogenic chemical and hence its use may be avoided or done in a highly efficient fume cupboard.

(b) Stas-Otto Process: The **Stas-Otto process** essentially consists of treating the powdered and sieved drug substance with 90–95% (v/v) ethanol, previously acidified with tartaric acid. The proportion of crude drug to solvent should be maintained as 1 Kg to 1 L. The alcohol is distilled off under vacuum and the resulting aqueous residue is treated with petroleum-ether (60-80°C) to remove the fatty components completely. If any alkaloid is removed by the petroleum ether, it must be recovered by treating it with dilute mineral acid. Thus, the resulting aqueous extract is mixed with the main bulk of aqueous extract. The combined aqueous extract is filtered and evaporated to dryness preferably in a **Rotary Thin-Film Evaporator** under vacuum. The residue is extracted with absolute ethanol thereby dissolving the total alkaloids.

(c) Kippenberger's Process: In Kippenberger's process the powdered and sieved plant substance is first and foremost digested with solution of tannin (100 g) in glycerol (500 g) at a constant temperature of 40°C for a duration of 48 hours. The resulting mixture is further heated to 50°C so as to help in the complete coagulation of proteinous substances, cooled to ambient temperature and finally filtered. The resulting filtrate is thoroughly shaken with petroleum ether to get rid of faulty materials (oils, fats and waxes), and the last traces of petroleum ether is removed from the extract by heating either on a water-bath (electric) or exposure to Infra-Red Lamp. The fat-free crude plant extract is subsequently acidified and shaken with chloroform, successively to remove the bulk of the alkaloids, namely, atropine, codeine, colchicine, narcotine, nicotine, papaverine, spartenine and thebaine.

The resulting residual extract may still contain narceine, curarine and morphine. However, narceine and morphine may be isolated by passing freshly generated CO₂ directly into extract so as to convert the alkali hydroxide into their corresponding carbonate, which is then ultimately subjected to solvent extraction using a mixture of alcohol and chloroform. Finally, the third alkaloid, curarine, may be extracted by agitation with a mixture of equal volumes of ether and chloroform.

However, a combination of **Kippenberger's process** and **Stas-Otto process** by its application to the final alcoholic extract obtained by the latter process is found to give better separation of alkaloids.

7.1.7.4 Purification of Alkaloidal Extract

The main bulk of the **crude alkaloidal extract** is invariably subjected to further purification by means of either anyone or combination of the following methods:

(a) Extraction with Acid Solution The extraction of the **alkaloid** from the bulk of the crude alkaloid solution in immiscible organic solvent is invariably carried out by shaking with an acid solution. In usual practice, the use of HCl is restricted when chloroform remains as the solvent because of the fact that quite a few alkaloidal hydrochlorides are distinctly soluble in the latter. However, dilute H₂SO₄ is always preferred over HCl for general use in the extraction of alkaloids. Subsequently, the acid solution is rendered alkaline with dilute NH₄OH solution to liberate the alkaloids which is then extracted with an organic solvent. The solvent is removed under reduced pressure and the traces of moisture is removed with anhydrous sodium sulphate.

Note: The following two precautions may be observed, namely


- (i) To avoid the formation of stubborn and troublesome emulsions a solution of gum-tragacanth is often added to the aqueous-phase. In case, it still persists the two phases may be got separated by centrifugation, and**

(ii) To discard the presence of foreign interfering extractive components present in plant substances, such as: pigments, resins, waxes, oils and fats, the use of a 2.5-5% (w/v) solution of lead acetate is made to the alkaloidal extract which precipitates them effectively. The excess of lead present in the filtrate is removed by either passing H₂S gas through the Kipp's Apparatus or by adding sodium phosphate.


(b) **Precipitation of Alkaloid with Precipitating Reagent** The usual precipitation of the alkaloid as a complex compound is accomplished by the addition of a suitable precipitating reagent. The resulting alkaloidal complex is further purified by filtration, recrystallization and ultimately decomposed to obtain the desired free alkaloid(s).

Example

- (i) **Tannic-acid Complex:** It is normally decomposed by treatment with freshly prepared Pb(OH)₂ or Pb(CO₃)₂.
- (ii) **Precipitates obtained with HgCl₂, AuCl₃, PtCl₄, Mayer's Reagent:** These precipitates are decomposed by passing a stream of H₂S gas through its suspension.
- (iii) **Precipitates with Double Salts:** The double salt obtained with Dragendorff's Reagent is quickly boiled with 5% (w/v) BaCO₃ solution.
- (iv) **Precipitates with Nitrogenous Acids:** The precipitates obtained with nitrogenous acids like picric acid and picrolonic acid are normally decomposed by treatment with either NH₄OH or NaOH.



Picric acid



Picrolonic acid

- (v) **Reineckate Complex:** The complex obtained from alkaloid with *Reinecke Salt*, NH₄ [Cr (NH₃)₂ (SCN)₄], is normally decomposed by treating its solution in a mixture of acetone and water (1:1) with a silver sulphate solution.

It is pertinent to mention here that the **free liberated alkaloid** from the complexes stated above, (i) through (v), may be further extracted for its final recovery with an appropriate organic solvent, such as: chloroform.

(c) The purification of **alkaloids** may also be accomplished by the formation of its crystallised alkaloidal salt by the addition of an appropriate mineral or organic acid, such as: hydrochloric, hydrobromic, perchloric, sulphuric, oxalic and tartaric acids.

(d) Various known **separation techniques**, namely: **partition**, **ion-exchange** and **column chromatography** are invariably used for the purification of a host of alkaloids.

Besides, various physical parameters like: specific rotation, melting point, solubility are frequently used as a definite criteria of ascertaining the purity of alkaloids.

7.1.7.5 Fractionation of Crude Alkaloids

It has been observed largely that most of the alkaloid-bearing plant materials usually contain a mixture of closely-related alkaloids. Therefore, it has become almost necessary to carry out an effective fractionation of crude alkaloids from the extract or solution of total crude alkaloids.

However, the traditional and orthodox methods of separation are not only difficult but also tedious and cumbersome. The commonly employed techniques of separation that were found to be reliable and dependable may be short-listed as follows:

- (i) Fractional crystallization,
- (ii) Fractional distillation, and
- (iii) Derivatization with low solubility products.

The latest methods employed for the separation of **alkaloids** are the preparative **high performance liquid chromatography (HPLC)**, **high performance thin-layer chromatography (HPTLC)**, chromatotron, counter-current distribution and other chromatographic techniques including column chromatography, ion-exchange chromatography.

Following are some of the typical situations whereby the mixture of **alkaloids** may be separated effectively, such as:

- (a) A larger section of the **alkaloids** are easily soluble in chloroform and relatively less soluble in other organic solvents. In general, the order of solubility is as stated below chloroform > acetone > ethanol > methanol > ethyl acetate > ether > n-hexane. Keeping in view the above solubility profile of alkaloids in organic solvents, if one of the alkaloids is much less soluble in ethanol than chloroform, the fractional crystallization of this alkaloid is possible. In this particular instance the chloroform-fraction is concentrated to an appropriate level, and hot ethanol added in small proportions at intervals. Thus, upon cooling the alkaloid, which is less soluble in ethanol, separates out conveniently.
- (b) In case, the fractional crystallization of the mixture of closely related alkaloids become tedious and ineffective, one may try to form their respective salts,* and then carry out the separation indicated above.
- (c) The various acids, namely: HCl, HBr, HI, HClO₄, HNO₃, C₂H₂O₄, and C₆H₃N₃O₇, may either be employed in aqueous or methanolic solution. Thus, from the resulting methanolic solution, the salts of the respective alkaloids may be precipitated by the addition of ether. The precipitated crude alkaloidal salts may be further recrystallized from hot acetone containing a small proportion of methanol.
- (d) In certain other specific instances, the salts of the respective oxalates, picrates and perchlorates may be precipitated from their solutions in acetone, by the addition of ethyl acetate.

7.2 CLASSIFICATION OF ALKALOIDS

The **alkaloids**, as an important and enormously large conglomerate of naturally occurring nitrogen-containing plant substances having very specific as well as most diversified pharmacological properties may be classified in a number of modes and means.

* **Salts of Alkaloids:** that are used frequently are hydrochloride, hydrobromide, hydroiodide perchlorate, nitrate, oxalate and picrate.

Hegnauer* (1963) conveniently classified alkaloids into *six* important groups, corresponding to the six amino-acids legitimately considered as the starting points for their biosynthesis, such as: anthranilic acid, histidine, lysine, ornithine phenylalanine and tryptophan. Price* (1963) further took a leading clue from the earlier observation and considered in details the alkaloids present in one of the families, (*Rutaceae*) and logically placed them in the following *nine* chemical-structural categories, namely: **acridines, amides, amines, benzyloquinolines, canthinones, imidazoles, indolquinazolines, furoquinolines, and quinazolines.**

Another school of thought classifies **alkaloids** in the following *four* heads, namely:

(a) Biosynthetic Classification In this particular instance the significance solely lies to the precursor from which the alkaloids in question are produced in the plant biosynthetically. Therefore, it is quite convenient and also logical to group together all alkaloids having been derived from the same precursor but possessing different taxonomic distribution and pharmacological activities.

Examples

- (i) Indole alkaloids derived from *tryptophan*.
- (ii) Piperidine alkaloids derived from *lysine*.
- (iii) Pyrrolidine alkaloids derived from *ornithine*.
- (iv) Phenylethylamine alkaloids derived from *tyrosine*.
- (v) Imidazole alkaloids derived from *histidine*.

(b) Chemical Classification It is probably the most widely accepted and common mode of classification of **alkaloids** for which the main criterion is the presence of the basic heterocyclic nucleus (*i.e.*, the **chemical entity**).

Examples

- (i) Pyrrolidine alkaloids *e.g.*, **Hygrine**;
- (ii) Piperidine alkaloids *e.g.*, **Lobeline**;
- (iii) Pyrrolizidine alkaloids *e.g.*, **Senecionine**;
- (iv) Tropane alkaloids *e.g.*, **Atropine**;
- (v) Quinoline alkaloids *e.g.*, **Quinine**;
- (vi) Isoquinoline alkaloids *e.g.*, **Morphine**;
- (vii) Aporphine alkaloids *e.g.*, **Boldine**;
- (viii) Indole alkaloids *e.g.*, **Ergometrine**;
- (ix) Imidazole alkaloids *e.g.*, **Pilocarpine**;
- (x) Diazocin alkaloids *e.g.*, **Lupanine**;
- (xi) Purine alkaloids *e.g.*, **Caffeine**;
- (xii) Steroidal alkaloids *e.g.*, **Solanidine**;
- (xiii) Amino alkaloids *e.g.*, **Ephedrine**;
- (xiv) Diterpene alkaloids *e.g.*, **Aconitine**.

(c) Pharmacological Classification Interestingly, the **alkaloids** exhibit a broad range of very specific pharmacological characteristics. Perhaps this might also be used as a strong basis for the general classification of the wide-spectrum of alkaloids derived from the plant kingdom, such as:

* Swain, T. (ed), 'Chemical Plant Taxonomy', Academic Press, London, (1963).

analgesics, cardio-vascular drugs, CNS-stimulants and depressants, dilation of pupil of eye, mydriatics, anticholinergics, sympathomimetics, antimalarials, purgatives, and the like. However, such a classification is not quite common and broadly known.

Examples

- (i) **Morphine** as Narcotic analgesic;
- (ii) **Quinine** as Antimalarial;
- (iii) **Strychnine** as Reflex excitability;
- (iv) **Lobeline** as Respiratory stimulant;
- (v) **Boldine** as Cholaretics and laxatives;
- (vi) **Aconitine** as Neuralgia;
- (vii) **Pilocarpine** as Antiglaucoma agent and miotic;
- (viii) **Ergonovine** as Oxytocic;
- (ix) **Ephedrine** as Bronchodilator;
- (x) **Narceine** as Analgesic (narcotic) and antitussive.

(d) Taxonomic Classification This particular classification essentially deals with the ‘**Taxon**’ *i.e.*, the **taxonomic category**. The most common *taxa* are the genus, subgenus, species, subspecies, and variety. Therefore, the taxonomic classification encompasses the plethora of alkaloids exclusively based on their respective distribution in a variety of **Plant Families**, sometimes also referred to as the ‘**Natural order**’. A few typical examples of plant families and the various species associated with them are stated below, namely:


- (i) **Cannabinaceous Alkaloids:** *e.g.*, *Cannabis sativa* Linn., (**Hemp, Marijuana**).
- (ii) **Rubiaceous Alkaloids:** *e.g.*, *Cinchona Sp.* (**Quinine**); *Mitragyna speciosa* Korth (**Katum, Kratum, Kutum**); *Pausinystalia johimbe* (K. Schum) (**Yohimbe**).
- (iii) **Solanaceous Alkaloids:** *e.g.*, *Atropa belladonna* L., (**Deadly Nightshade, Belladonna**); *Brunfelsia uniflorus* (Pohl) D. Don (**Manaca, Manacan**); *Capsicum annum* L., (**Sweet Peppers, Paprika**); *Datura candida* (Pers.) Saff. (**Borrachero, Floripondio**); *Duboisia myoporoides* R. Br. (**Corkwood Tree, Pituri**); *Hyoscyamus niger* L. (**Henbane, Henblain, Jusquaim**); *Mandragora officinarum* L. (**Mandrake, Loveapple**); *Nicotiana glauca* R. Grah. (**Tree Tobacco**); *Seopolia carniolica* Jacq. (**Scopolia**); *Solanum dulcamara* L., (**Bittersweet, Bitter Nightshade, Felonwood**); *Withania somniferum* (L.) Dunal (**Ashwagandha**), etc.

Invariably, they are grouped together according to the name of the *genus* wherein they belong to, such as: **coca, cinchona, ephedra**.




Some ‘**phytochemists**’ have even gone a step further and classified the **alkaloids** based on their **chemotaxonomic classification**.

In the recent past, the **alkaloids** have been divided into *two* major categories based on the analogy that one containing a *non-heterocyclic nucleus*, while the other having the *heterocyclic nucleus*. These *two* classes of alkaloids shall be discussed briefly as under.

(a) Non-heterocyclic Alkaloids A few typical **alkaloids** having non-heterocyclic nucleus are enumerated below:





S.No.	Basic Ring Structure	Alkaloid	Botanical Origin	Family
1	Phenylethylamine 	Ephedrine Hordenine Capsaicin Mescaline Narceine	<i>Ephedra vulgaris</i> <i>Hordeum vulgare</i> <i>Capsicum annuum</i> <i>Laphophora williamsii</i> <i>Papaver somniferum</i>	<i>Gnetaceae</i> <i>Graminae</i> <i>Solanaceae</i> <i>Cactaceae</i> <i>Papaveraceae</i>

(b) **Heterocyclic Alkaloids** A large number of specific **alkaloids** possessing heterocyclic nucleus are stated below:

S.No.	Basic Ring Structure	Alkaloid	Botanical Origin	Family
1	Pyrrolidine 	Hygrine Stachydrine	<i>Erythroxylon coca</i> <i>Stachys tubrifera</i>	<i>Erythroxylaceae</i> <i>Labiatae</i>
2	Pyridine 	Arecoline Ricinine Trigenelline	<i>Areca catchu</i> <i>Ricinus communis</i> <i>Trigonella foenumgraecum</i>	<i>Palmaceae</i> <i>Euphorbiaceae</i> <i>Leguminosae</i>
3	Piperidine 	Conine Lobeline Pelletierine	<i>Conium maculatum</i> <i>Lobelia inflata</i> <i>Punica granatum</i>	<i>Umbelliferae</i> <i>Lobeliaceae</i> <i>Punicaceae</i>
4	Tropane [Piperidine-Pyrrolidine (N-Methyl)] 	Atropine Cocaine Hyoscyamine	<i>Atropa belladone</i> <i>Datura stramonium</i> <i>Erythroxylon coca</i> <i>Atropa belladona</i>	<i>Solanaceae</i> <i>Solanaceae</i> <i>Erythroxylaceae</i> <i>Solanaceae</i>
5	Quinoline 	Quinine, Quinidine Cuspareine	<i>Cinchona officinalis</i> <i>Cusparia trifoliata</i>	<i>Rubiaceae</i> <i>Rutaceae</i>
6	Isoquinoline 	Papaverine Berberine Emetine Corydaline Tubocurarine	<i>Papaver somniferum</i> <i>Hydrastis canadensis</i> <i>Uragoga ipecacuanha</i> <i>Corydalis aurea</i> <i>Corydalis solida</i> <i>Chondodendron tomentosum</i>	<i>Papaveraceae</i> <i>Berberidaceae</i> <i>Rubiaceae</i> <i>Fumariaceae</i> <i>Fumariaceae</i> <i>Menispermaceae</i>


(Contd.)

(Contd.)

7	<p>Aporphine Isoquinoline Phenanthrene</p> 	Boldine	<i>Peumus boldus</i>	<i>Monimiaceae</i>
8	<p>Norlupinane</p> 	Sparteine	<i>Lupinus luteus</i> , <i>Lupinus niger</i> , <i>Cytisus scoparius</i> , <i>Anagyris foetida</i>	<i>Leguminosae</i>
		Lupinine	<i>Lupinus luteus</i> <i>Anabasis aphylla</i>	<i>Leguminosae</i> <i>Chenopodiaceae</i>
9	<p>Indole (Benzopyrrole)</p> 	Ergotamine, Ergometrine Physostigmine	<i>Claviceps purpurea</i> <i>Physostigma</i> <i>Venenosum</i>	<i>Hypocreales</i> <i>Leguminosae</i>
		Reserpine	<i>Rauwolfia</i> <i>serpentina</i>	<i>Apocynaceae</i>
		Yohimbine	<i>Coryanthe</i> <i>johimbe</i> <i>Rauwolfia</i> <i>serpentina</i>	<i>Rubiaceae</i> <i>Apocynaceae</i>
		Vinblastine (Vincalukoblastine)	<i>Vince rosea</i>	<i>Apocynaceae</i>
		Strychnine	<i>Strychnos</i> <i>nux-vomica</i>	<i>Loganiaceae</i>
10	<p>Imidazole (Glyoxaline)</p> 	Pilocarpine	<i>Pilocarpus</i> <i>jaborandi</i>	<i>Rutaceae</i>
11	<p>Purine (Pyrimidine-Imidazole)</p> 	Caffeine	<i>Thea sinensis</i> <i>Camellia sinensis</i> <i>Coffea arabica</i> <i>Theobroma cacao</i>	<i>Ternstroemiaceae</i> <i>Rubiaceae</i> <i>Sterculiaceae</i>

(Contd.)

(Contd.)

12	Tropolone 	Colchicine	<i>Colchicum autumnale</i>	<i>Liliaceae</i>
13	Steroid 	Connesine Funtumine Solanidine Veratramine	<i>Holarrhena anti-dysenterica</i> <i>Funtumia latifolia</i> <i>Solanum spp.</i> <i>Veratrum grandiflorum,</i> <i>Veratrum viride</i>	<i>Apocynaceae</i> <i>Apocynaceae</i> <i>Solanaceae</i> <i>Liliaceae</i>
14	Terpenoid (Diterpene)	Aconine Aconitine (Glycoside) Atisine Lycoetionine	<i>Aconitum napellus</i> -do- <i>Aconitum heterophyllum,</i> <i>Aconitum anthora,</i> <i>Aconitum lycoctonum,</i>	<i>Ranunculaceae</i> -do- <i>Ranunculaceae</i> <i>Ranunculaceae</i>
15	Pyrrolizidine 	Senneccionine Senneciphylline	<i>Senecio vulgaris</i> <i>Senecio platyphillus</i>	<i>Compositae</i> <i>Compositae</i>

It is, however, pertinent to mention at this juncture that the enormous volume of authentic information accumulated so far with regard to the isolation of **alkaloids** from a variety of plant species and their subsequent characterization by the help of latest analytical techniques they may be classified as follows:

A. Alkaloids derived from Amination Reactions

- (i) Acetate-derived Alkaloids
- (ii) Phenylalanine-derived Alkaloids
- (iii) Terpenoid Alkaloids
- (iv) Steroidal Alkaloids

B. Alkaloids derived from Anthranilic Acid

- (i) Quinazoline Alkaloids
- (ii) Quinoline Alkaloids
- (iii) Acridine Alkaloids

C. Alkaloids derived from Histidine

Imidazole Alkaloids

D. Alkaloids derived from Lysine

- (i) Piperidine Alkaloids

- (ii) Quinolizidine Alkaloids
- (iii) Indolizidine Alkaloids
- E. **Alkaloids derived from Nicotinic Acid**
 - Pyridine Alkaloids
- F. **Alkaloids derived from Ornithine**
 - (i) Pyrrolidine Alkaloids
 - (ii) Tropane Alkaloids
 - (iii) Pyrrolizidine Alkaloids
- G. **Alkaloids derived from Tyrosine**
 - (i) Phenylethylamine Alkaloids
 - (ii) Simple Tetrahydro iso-quinoline Alkaloids
 - (iii) Modified Benzyl Tetrahydro iso-quinoline Alkaloids
- H. **Alkaloids derived from Tryptophan**
 - (i) Simple Indole Alkaloids
 - (ii) Simple β -Carboline Alkaloids
 - (iii) Terpenoid Indole Alkaloids
 - (iv) Quinoline Alkaloids
 - (v) Pyrroloindole Alkaloids
 - (vi) Ergot Alkaloids
- I. **Purine Alkaloids**

These broad and elaborated classification of the alkaloids shall now be treated individually at length in the sections that follows:

7.2.1 Alkaloids Derived from Amination Reactions

It has been duly established that the larger section of **alkaloids** are virtually derived from amino acid precursors by the help of certain specific processes that essentially introduce into the final structure not only a N-atom but also an amino acid carbon skeleton or a major part of it. However, a good number of **alkaloids** do not essentially conform with this analogy. They are usually synthesized primarily from non-amino acid precursors having the N-atom inserted into the structure at a comparatively latter stage. Interestingly, such structures are predominantly based on both steroidal and terpenoid skeletons. Besides, a few comparatively simpler **alkaloids** also appear to be derived exclusively with the help of similar late amination processes. An extensive and intensive studies on certain alkaloids it has been observed that the N-atom is specifically donated from an **amino acid source** through a *transamination reaction* using an appropriate ketone or aldehyde.

7.2.1.1 Acetate-Derived Alkaloids

Socrates was made to drink the decoction of the *Hemlock* plant and died soonafter. Thus, the poison present in it is really too dangerous for herbal administration by the uninitiated. The Hemlock plant is comprised of several potent alkaloids, such as: **coniine**, **γ -coniceine**, **conhydrine**, **N-methyl conine** and **pseudoconhydrine**. These **alkaloids** shall now be discussed as under:


A. Coniine

Synonyms Cicutine, Conicine.

Biological Sources It is obtained from the unripe, fully grown dried fruits of *Conium maculatum* L. (*Umbelliferae*).

It also occurs in the plant *Aethusa cynapium* L. (*Apiaceae*) (Fool's Parsley); *Cicuta maculata* L. (*Apiaceae*) (Water Hemlock).

Chemical Structure



Coniine

(S)-2-Propylpiperidine. It occurs naturally as the (S)-(+)- isomer.

Isolation Coniine may be isolated by adopting the various following steps, namely:

- (i) The powdered unripe, fully grown dried fruits of hemlock are mixed with a dilute solution of KOH and then subjected to steam distillation. The distillate is collected and neutralized carefully with dilute HCl and evaporated to dryness preferably under vacuum.
- (ii) The residue obtained as stated in (i) above is extracted with alcohol, filtered and the alcohol evaporated to dryness under vacuum. The alcohol helps in extracting the alkaloidal salts that are dissolved in water; it is then rendered alkaline either with diluted KOH solution or with dilute NH_4OH and finally extracted with ether successively.
- (iii) The ether from the combined ethereal layer is evaporated completely, when an oily liquid consisting of the free bases remains in the residue.
- (iv) Finally, the residue is subjected to fractional distillation in a current of H_2 -gas when the alkaloids could be broadly separated and a mixture containing *coniine* and γ -*coniceine* shall pass over as the first fraction at 171-172°C. These two alkaloids are consequently made to their corresponding hydrochloride salts, evaporated to dryness and extracted with acetone. Thus, coniine hydrochloride would be separated as an insoluble product, while the γ -*coniceine* may be recovered by evaporating acetone under vacuum.

Note: Coniine enjoys the unique distinction of being the First Alkaloid produced synthetically.

Characteristic Features

- (i) It is a colourless alkaline liquid.
- (ii) It darkens and polymerizes on being exposed to air and light.
- (iii) It has a mousy odour.
- (iv) Its physical parameters are as follows: mp $\sim -2^\circ\text{C}$; bp 166-166.5°C; bp₂₀ 65-66°C.
 d_4^{20} 0.844 – 0.848; n_D^{23} 1.4505; $[\alpha]_D^{25} + 8.4^\circ\text{C}$ (c = 4.0 in CHCl_3); $[\alpha]_D^{23} + 14.6^\circ$ (heat)
 $\text{pK}_a = 3.1$.
- (v) It is steam volatile.
- (vi) **Solubility:** 1 ml dissolves in 90 ml of water, less soluble in hot water. The base dissolves in about 25% water at room temperature. It is found to be soluble in alcohol, ether, acetone, benzene, amyl alcohol, and slightly soluble in chloroform.

Identification Tests


- (i) It readily forms the corresponding hydrobromide ($C_8H_{17}N.HBr$), obtained as prisms, mp $211^\circ C$, 1 g dissolves in 2 ml water, 3 ml alcohol, and soluble freely in ether and chloroform.
- (ii) Its hydrochloride ($C_8H_{17}N.HCl$) forms rhomboids, mp $221^\circ C$, freely soluble in water, alcohol and chloroform.
- (iii) It gives a red colouration with sodium nitroprusside slowly, which on addition of acetaldehyde changes to violet or blue.

Caution It exhibits potential symptoms of over exposure as: weakness, drowsiness, parasthesias, ataxia, nausea, excessive salivation, and bradycardia followed by tachycardia.*

Uses Externally, the coniine salts are used as ointments and infrequently employed for their local analgesic action in the symptomatic relief of pruritis, hemorrhoids and fissures.

B. γ -Coniceine

Biological Source It is obtained from the seeds of *Conium maculatum* L. (*Umbelliferae*).

Chemical Structure γ -Coniceine

2, 3, 4, 5-Tetrahydro-6-propylpyridine.

Characteristic Features

- (i) It is a colourless liquid alkaloid.
- (ii) It possesses a distinct mousy odour.
- (iii) It is steam volatile.
- (iv) Its physical parameters are: bp $171^\circ C$; d_4^{15} 0.8753; n_D^{16} 1.4661.
- (v) It is slightly soluble in water, but freely soluble in ethanol, chloroform and ether.


Identification Test

- (i) γ -Coniceine when subjected to reduction, it gives rise to a racemic mixture of *dl*-coniine.
- (ii) It forms γ -coniceine hydrochloride ($C_8H_{15}N.HCl$) which gives hygroscopic crystals from ether mp $143^\circ C$.

C. Conhydrine

Biological Source It is obtained from the seeds of *Conium maculatum* L. (*Umbelliferae*).

* Gosselin *et al.* Eds. **Clinical Toxicology of Commercial Products**, Williams and Wilkins, Baltimore, 5th ed., Sec-II., pp. 249-250 (1984).

Chemical Structure

Conhydrine


[R-(R*, S*)- α -Ethyl-2-piperidine methanol.

Characteristic Features

- (i) The crystals obtained from ether has mp 121°C, bp 226°C and $[\alpha]_D + 10^\circ\text{C}$.
- (ii) It is slightly soluble in water, but easily soluble in ethanol, ether and chloroform.

D. N-Methylconiine

Biological Source It is same as for (C) above.

Chemical Structure

N-Methylconiine

1-Methyl-2-propylpiperidine.


Isolation The *d*-form is stated to occur in Hemlock in small quantities, while the *l*-form may be isolated from residues left in the preparation of **coniine** by crystallization of the hydrobromides.

Characteristic Features The physical characteristic features of *dl*, *d*- and *l*-forms are given below:

Form	Nature	bp (°C)	d^{24}	$[\alpha]_D^{24}$	n_D^{13}	Solubility
<i>dl</i> -	–	bp 10.5 56.6	–	–	–	–
<i>d</i> -	Oily liquid	173-174	0.8318	+ 81°	1.4538	Slightly in water, soluble in organic solvents
<i>l</i> -	–	–	0.8318	– 84°	1.4538	Slightly in water, soluble in organic solvents

E. Pseudoconhydrine

Biological Source Its biological source is same as for (A) through (D) above.

Chemical Structure


(3*S-trans*)-6-Propyl-3-piperidinol.

Characteristic Features

- (i) It gives hygroscopic needles from absolute ether.
- (ii) Its mp stands at 106°C, whereas its monohydrate, scales, gives mp 60°C from moist ether.
- (iii) Its physical parameters are: bp 236°C ; $[\alpha]_D^{20} + 11^\circ$ ($c = 10$ in alcohol); pK (18°C): 3.70
- (iv) It is soluble in water and

Identification Tests It readily forms the hydrochloride salt ($C_8H_{17}NO.HCl$) as the crystals from ethanol having mp 213°C.

Biosynthesis of γ -Coniceine and Coniine A fatty acid precursor **octanoic acid (capric acid)** is employed, which is subsequently transformed into the ketoaldehyde through successive oxidation and reduction steps. The resulting ketoaldehyde acts as a substrate for a transamination reaction, the amino moiety is derived from L-alanine. The ultimate transformation lead to the formation of imine giving the heterocyclic ring present in **γ -coniceine**, and then reduction the **coniine** as shown below:

**7.2.1.2 Phenylalanine-Derived Alkaloids**


It has been observed that the aromatic amino acid **L-tyrosine** is not only a common but also an extremely vital precursor of **alkaloids**; whereas, **L-phenylalanine** is found to be much less frequently employed, and normally it specifically contributes *carbon atoms only*, such as: C_6C_1 , C_6C_2 or C_6C_3 units, without making available a N-atom from its amino function *e.g.*, as in the biosynthesis of **colchicine** and **lobeline**.

The various typical examples of phenylalanine-derived alkaloids are: **ephedrine**, **norpseudoephedrine (cathine)** and **capsaicin**, which shall be described hereunder:

A. Ephedrine

Biological Source It occurs in the dried young stems of the Chinese wonder drug **Ma Huang**, *Emhedra vulgaris*, *Ephedra sinica* Stapf., *Ephedra equisetina* Bunge belonging to family *Ginetaceae*, and also in several other Ephedra species. This is also found in *Ephedra geradiana* Wall ex. Stapf. (*Ephedraceae*) (Pakistani Ephedra). There are *two* most important forage ephedras in the United States, namely: *E. nevadensis* and *E. viridis*. The former are is *E. nevadensis* S. Wats (*Ephedraceae*) and known as **Mormon Tea** and **Nevada Jointfir**.

Chemical Structure



Ephedrine

α -[1-(Methylamino)ethyl] benzene methanol ($C_{10}H_{15}HO$).


Isolation Ephedrine usually exists singly in *Ephedra sinica* (1-3%) and *E. equisetina* (2%). However, it occurs in association with \pm -**Ephedrine** (*i.e.*, **pseudoephedrine**) in *E. vulgaris*.

However, the ephedrine and **pseudoephedrine** may be extracted conveniently from the dried young stems of the plant material by adopting the 'general procedures for alkaloid extraction' (section 1.7.3), by the help of successive benzene and dilute HCl extractions.

Preparation Ephedrine may be prepared by *two* methods, namely:


- (i) Fermentation method, and
- (ii) Synthetic method.

(a) **Fermentation Method:** It can be prepared commercially by fermenting a mixture of molasses* and benzaldehyde. The reaction product *i.e.*, methyl benzyl alcohol ketone *i.e.*, $C_6H_5-CH(OH)COCH_3$, a keto-alcohol is subsequently mixed with a solution of methyl amine and freshly prepared H_2 -gas is made to pass through it. Thus, we have:




* **Molasses:** A thick brown viscous liquid obtained as a by product of 'Sugar Industry' containing 8-10% cane sugar.


(b) **Synthetic Method:** Manske *et al.** (1929) synthesized (\pm)-**Ephedrine** by the catalytic reduction of 1-phenylpropane-1, 2-dione (or benzoylacetyl) in the presence of methylamine in methanol solution as given below:




Stereochemistry Since the **ephedrine** molecule contains *two* dissimilar chiral centres, four optically active isomers (or two pairs of enantiomers) are possible theoretically. Freudenberg (1932) put forward the following configurations of **ephedrine** and ψ -**ephedrine** (mp 118°C, $[\alpha]_D \pm 51.2^\circ$) are as follows:



Foder *et al.* (1949, 1950) confirmed that the ephedrine has the *erythro*-configuration, and ψ -ephedrine the *threo*-configuration as stated below:



The carbobenzoxy derivative of **nor- ψ -ephedrine** undergoes intramolecular rearrangement to the *O*-derivative in an acidic medium. In case, **nor- ψ -ephedrine** possesses the *threo*-configuration, then this ultimately gives rise to the favourable *trans*-orientation of the phenyl and methyl groups in the cyclic intermediate *i.e.*, the **steric repulsions** are at a bear minimum level. Likewise, the **nor-ephedrine** shall, therefore, exhibit essentially the *crythro*-configuration; and it was further revealed that its corresponding N-carbobenzoxy derivative does not undergo any molecular rearrangement whatsoever in an acidic environment to produce the *O*-derivative. Therefore, one may infer that the steric repulsions that would take place between the phenyl and methyl groups in



* Manske and Holmes (eds). **The Alkaloids**, Academic Press. N. York. Vol. 1 (1950)

the cyclic intermediate is evidently too high to allow its subsequent formation. Thus, it is absolutely possible, on this basis, to differentiate and distinguish between the stereoisomers of **ephedrine** and **Ψ-ephedrine**.

Characteristic Features The characteristic features of various forms of **ephedrine** and its salts are as stated under:

S. No.	Form	Nature	Name	mp (°C)	bp (°C)	[μ]D ₂₅	Solubility
1.	<i>dl</i> -	Crystals	Racephedrine, Racemic ephedrine	79	–	–	Soluble in water, ethanol, ether, chloroform, oils.
2.	<i>dl</i> -HCl	-do-	Ephedrin, Racephedrine HCl	187–188	–	–	1 g dissolves in 4 ml water, in 40 ml of 95% alcohol at 20°C. Insoluble in ether
3.	<i>dl</i> - Sulphate	-do-	Racephedrine Sulphate	247	–	–	Soluble in water and ethanol
4.	<i>l</i> -HCl	Crystals, waxy solid, granules	L-Erythro- 2-(methylamino) -1-phenylpropan- 1-ol.	34	255	–	1 g dissolves in 20 ml H ₂ O, 0.2 ml ethanol, soluble in chloroform, ether, oils
5.	<i>l</i> -HCl	Ortho- rhombic needles	Ephedral, Sanedrine	216–220	–	–30 to –35.5° (c = 5)	1 g dissolves in 3 ml water, 14 ml ethanol; In soluble in ether and chloroform
6.	<i>l</i> -Sulphate	-do-	–	245 (dec.)	–	–29.5 to –32.0° (c = 5)	1 g dissolves in 1.2 ml water, 95 ml ethanol; Freely soluble in hot alcohol.
7.	<i>d</i> -Pseudo- ephedrine	Rhombic tablets	<i>d</i> -Ψ- ephedrine, <i>d</i> -iso- ephedrine	119	–	+51° (c = 0.6 in alc.)	Sparingly soluble in water (differs from <i>l</i> -ephedrine). Freely soluble in alc. or ether.
8.	<i>d</i> -Pseudo- ephedrine hydrochloride	Needles	Galseud, Novafed, Rhinalair, Otrinol, Sinufed, Sudafed, Symptom-2,	181 to 182	–	+ 62° (c = 0.8)	Soluble in water, alcohol and chloroform

Special Features


- (i) **Ephedrine** does not yield a precipitate with Mayer's Reagent except in concentrated solution.
- (ii) Ephedrine in chloroform solution after long standing or on evaporation usually forms ephedrine hydrochloride and phosgene.
- (iii) Both **ephedrine** and **pseudoephedrine** are fairly stable to heat and when heated at 100°C for several hours does not undergo any decomposition.
- (iv) **Ephedrine** hydrochloride on being heated with 25% HCl gets partially converted to **pseudoephedrine**; and this conversion is reversible and soon attains on equilibrium.

Identification Tests

- (i) **Colour Test:** Dissolve 0.1 g ephedrine in 1 ml water with the addition of a few drops of dilute HCl. Add to it two drops of CuSO₄ solution (5% w/v) followed by a few-drops of NaOH (1N) solution when a reddish colour is obtained. Add to it 2-3 ml of ether and shake vigorously, the ethereal layer becomes purple and the aqueous layer turns blue.

- (ii) **Formation of Ephedrine Hydrochloride:** Dissolve 0.2-0.3g of ephedrine in 35 ml of chloroform in a stoppered test tube and shake vigorously. Allow it to stand for 12 hours and evaporate the chloroform, when crystals of ephedrine HCl are obtained, and
- (iii) **Formation of Benzaldehyde Odour:** Take 0.05 g of ephedrine in a small porcelain dish and triturate it with a few crystals of pure potassium ferricyanide, $[K_3Fe(CN)_6]$, add a few drops of water and heat on a water-bath, it gives rise to a distinct odour of *benzaldehyde*.

Biosynthesis of Ephedrine Alkaloids Interestingly, phenylalanine and ephedrine not only have the same carbon and nitrogen atoms but also have the same arrangement of C and N-atoms *i.e.*, the skeleton of atoms. Noticeably, L-phenylalanine is a precursor, possessing only seven carbons, a C_6C_1 fragment, gets actually incorporated. It has been observed that phenylalanine undergoes metabolism, probably *via* cinnamic acid to benzoic acid; and this perhaps in the form of its coenzyme-A ester, which is acylated with pyruvic acid and undergoes decarboxylation during the addition as shown below.



Biosynthesis of Ephedrine Alkaloids

A thiamine PP-mediated mechanism is put forward for the formation of the diketone, and a transamination reaction shall give rise to **cathinone**. Further reduction of the carbonyl moiety from either face yields the diastereomeric **norephedrine** or **norpseudoephedrine (Cathine)**. Ultimately, N-methylation would give rise to **ephedrine** or **pseudoephedrine**.

Uses


1. The *l*-**ephedrine** is extensively used as a bronchodilator.
2. The *d*-**psuedoephedrine** is employed widely as a decongestant.

B. Norpseudoephedrine

Synonyms Cathine; Katine; Nor- ψ -ephedrine.

Biological Sources It occurs naturally as the *D-threo*-form in the leaves of the **khat plant**, *Catha edulis* Forsk. (*Celastraceae*), which is widely found as an evergreen shrub native to Southern Arabia and Ethiopia. It is also found in relatively smaller amounts in the South American tree *Maytenus krukovii* A.C. Smith (*Celastraceae*); and in the mother liquors obtained from **Ma Huang** after the recovery of **ephedrine**.

Chemical Structure



Norpseudoephedrine

(R^* , R^*)- α -(1-Aminoethyl)-benzenemethanol.

Isolation It is isolated from the plant material as described under (A) in this section.

Characteristic Features The various physical parameters of different forms of **norpseudoephedrine** are summarized below:

S.No.	Form	Name	Nature	mp ($^{\circ}\text{C}$)	$[\alpha]_D^{20}$	Solubility
1.	D-	Norpseudo-ephedrine	Plates from methanol	77.5–78	$[\alpha]_{546}^{20}$ + 37.9 $^{\circ}$ (c = 3 in methanol)	Soluble in alcohol, chloroform, ether, and dilute acids
2.	Hydrochloride	Amorphan, Adiposetten, Exponcit N, Fasupond, Fugoa, Minusin	Prisms	180–181	+43.2 (H ₂ O)	Soluble in water
3.	Sulphate	–	Hexagonal plates.	298	$[\alpha]_{546}^{20}$ + 48.7 $^{\circ}$ (c = 1.4 in H ₂ O)	Soluble in water
4.	d/-Hydrochloride	–	Crystals	169–171	–	Soluble in water

Uses


1. It is widely employed as an anorexic.
2. It is also used in the optical resolution of externally compensated acids.

C. Capsaicin

Synonyms Axsain; Mioton; Zostrix.

Biological Source It is the pungent principle obtained in the fruit of various species of **Capsicum**, viz., *Capsicum annum* L. (*Solanaceae*) (Chilli, **Sweet Peppers**, **Paprika**).

Chemical Structure



Capsaicin

(E)-N-[4-Hydroxy-3-methoxyphenyl]-methyl]-8-methyl-6-nonenamide. It is phenolic in nature.

Isolation The **capsicum fruits** are crushed and extracted with either hot acetone or ethanol by using the method of percolation. The solvent *i.e.*, hot acetone or ethanol is evaporated under vacuum. The residue is extracted once again with successive quantities of warm acetone or ethanol until and unless the marc is completely free from any pungent principles. It contains approximately not less than 8% of **capsaicin**.

Characteristic Features

1. **Capsaicin** gives a distinct burning taste even when diluted to the extent of one part in one million parts of water. However, its pungency is destroyed by oxidation.
2. It is obtained as monoclinic, rectangular plates, scales from petroleum ether, having mp 65°C.
3. It has bp_{0.01} 210-220°C (air-bath temperature).
4. It has uv maximum: 227, 281 nm (ϵ 7000, 2500).
5. It is freely soluble in ether, benzene, chloroform; slightly soluble in CS₂; and practically in soluble in water.


Identification Tests

1. An alcoholic solution of **capsaicin** gives rise to a distinct bluish green colour upon adding a few drops of FeCl₃ solution (0.5% w/v).
2. When capsaicin is dissolved in a few drops of concentrated H₂SO₄ and a few crystals of sucrose is added, it yields a violet colour after a few hours.

Uses

1. It is used as a topical analgesic.
2. It is often employed as a tool in neurobiological research.
3. It is used in creams to counter neuralgia caused by herpes infections and in other pain-relieving formulations.

Biosynthesis of Capsaicin The aromatic fragment of the **capsaicin** molecule is derived solely from phenylalanine through chemical entities, viz., **ferulic acid** and **vanillin**. The later compound, an aldehyde, is actually the substrate for transamination to yield **vanillylamine**. However, the acid part of the resulting amide structure is of polypeptide origin having essentially a branched-chain fatty acyl-CoA which is produced by chain extension of isobutyryl-CoA. The aforesaid source of reactions are as given under:



Biosynthesis of Capsaicin

7.2.1.3 Terpenoid Alkaloids

A plethora of **alkaloids** solely based on mono-, sesqui-, di-, and tri-terpenoid skeletons have been isolated and characterized. However, logistic and scientific information (s) with regard to their actual formation in nature is more or less sparse. It has been observed that the monoterpene alkaloids are derived from the structurally related **iridoid materials**, wherein the O-atom in the heterocyclic ring is replaced by a N-containing ring as depicted below.



Terpenoid Alkaloids


A few typical examples of the **terpenoid alkaloids** are, namely: **aconine** and **actinidine**, which shall be discussed in the sections that follow:

A. Aconine

Biological Source **Aconine** is the hydrolyzed product of aconitine which is obtained from the dried roots of *Aconitum napellus* Linn. (*Ranunculaceae*) and other aconites. *A. napellus* is also known as **aconite**, **blue rocket** and **monkshood**. Usually it contains upto 0.6% of the total alkaloids of aconite, of which approximately one third is the alkaloid **aconitine**.

Chemical Structure (1 α , 3 α , 6 α , 14 α , 15 α , 16 α)-20-Ethyl-1, 6-16-trimethoxy-4-(methoxymethyl) aconitane-3, 8, 13, 14, 15-pentol.

Isolation The alkaloid **aconitine** is subjected to hydrolysis which yields benzoyl aconine and acetic acid. The resulting benzoyl aconine is further hydrolyzed to yield aconine and benzoic acid. **Aconine** being very soluble in water may be separated easily from the less water-soluble by product, *i.e.*, benzoic acid.



Characteristic Features

- (i) It is an amorphous powder with a bitter taste.
- (ii) It has mp 132°C, $[\alpha]_D + 23^\circ$ and pK_a 9.52.
- (iii) It is extremely soluble in water, alcohol; moderately soluble in chloroform and slightly soluble in benzene. It is practically insoluble in ether and petroleum ether.

Identification Tests It forms *two* distinct derivatives as given below:

- (a) **Aconine Hydrochloride Dihydrate ($C_{25}H_{42}ClNO_9 \cdot 2H_2O$):** It is obtained as crystals having mp 175-176°C and $[\alpha]_D - 8^\circ$.
- (b) **Aconine Hydrobromide Sesquihydrate ($C_{25}H_{42}BrNO_9 \cdot 1\frac{1}{2} H_2O$):** It is obtained as crystals from water with mp 225°C.


Uses

1. It is used in the treatment of neuralgia, sciatica, rheumatism and inflammation.
2. It is employed occasionally as analgesic and cardiac depressant.

B. Aconitine

Biological Source The botanical source is the same as described under (A) above.

Chemical Structure



(1 α , 3 α , 6 α , 14 α , 15 α , 16(β))-20-Ethyl-1, 6,-16-trimethoxy-4-(methoxymethyl) aconitane-3, 8, 13, 14, 15,-pento 8-acetate 14-benzoate (C₃₄H₄₇NO₁₁).

Characteristic Features

1. It occurs as hexagonal plates having mp 204°C.
2. Its pK_a value stands at 5.88.
3. Its specific rotation $[\alpha]_D + 17.3^\circ$ (Chloroform).
4. It is slightly soluble in petroleum ether; but 1 g dissolves in 2 ml chloroform, 7 ml benzene, 28 ml absolute ethanol, 50 ml ether, 3300 ml water.

Identification Tests Aconitine forms specific salts with HBr and HNO₃ having the following physical parameters.


(a) **Aconitine Hydrobromide Hemipentahydrate (C₃₄H₄₇O₁₁.HBr.2½ H₂O):** The hexagonal tablets mp 200-207°C and the dried substance mp 115-120°C. Its crystals obtained from ethanol and ether with ½ H₂O has mp 206-207°C. Its specific rotation $[\alpha]_D - 30.9^\circ$.

(b) **Aconitine Nitrate (C₃₄H₄₇NO₁₁.HNO₃):** The crystals have mp about 200°C (decomposes), $[\alpha]_D^{20} - 35^\circ$ (c = 2 in H₂O).

Uses

1. It is exclusively used in producing heart arrhythmia in experimental animals.
2. It has also been used topically in neuralgia.

Biosynthesis of Aconitine-Type Alkaloids Aconite is particularly regarded as extremely toxic, due to the presence of **aconitine**, and closely related C₁₉ **nonditerpenoid alkaloids**. It has been observed that the species of *Delphinium* accumulate **diterpenoid alkaloids**, for instance: **atisine**, which proved to be much less toxic when compared to aconitine. A vivid close resemblance of their structural relationship to diterpenes, such as: **ent-kaurene**, of course, little experimental evidence is available.



Biosynthesis of Aconitine-Type Alkaloids

From the above course of reactions it appears quite feasible that:

- (a) A **pre-ent-kaurene** carbocation usually undergoes **Wagner-Meerwein Rearrangements**,
- (b) The **atisine-skeleton** is produced subsequently by incorporating an N-CH₂-CH₂-O fragment (e.g., from 2-aminoethanol) to form the resulting heterocyclic rings,
- (c) The **aconitine-skeleton** is perhaps formed from the **atisine-skeleton** by further modifications as stated above,
- (d) A rearrangement process converts two fused 6-membered rings into a (7 + 5)-membered bicyclic system, and
- (e) One carbon from the exocyclic double bond is eliminated.

7.2.1.4 Steroidal Alkaloids

In general, the **steroidal alkaloids** represent an important class of alkaloids that essentially afford a close structural relationship to sterols *i.e.*, they contain a perhydro-1, 2-cyclopentanophenanthrene nucleus. Interestingly, these group of alkaloids invariably occur in the plant kingdom as glycosidal combination with carbohydrate moieties.

The **steroidal alkaloids** may be broadly classified into *two* major groups, namely:

- (a) Solanum Alkaloids, and
- (b) Veratrum Alkaloids.

These two class of alkaloids shall now be discussed in an elaborated fashion hereunder:

A. Solanum Alkaloids

A good number of plants belonging to the natural order *Solanaceae* have been found to accumulate favourably several **steroidal alkaloids** based on a C₂₇ *cholestane skeleton*, such as: **solasodine**, **tomatidine**, **solanidine**. These alkaloids usually occur in a wide variety of the genus *Solanum*, for instance: *Solanum laciniatum*; *S. dulcamara* Linn.; *S. nigrum* Linn.; *S. torvum* Swartz.; *S. lycopersicum* Linn.; (*Lycopersicon exculentum* Mill); *S. tuberosum*; *S. aviculare* etc. The three above mentioned alkaloids normally occur naturally in the plant as their corresponding glycosides. However, the two species of *Solanum*, namely: *S. laciniatum* and *S. aviculare* are considered to be a rich source of alkaloids (*i.e.*, the aglycone moieties) that are employed exclusively as the starting materials for the synthesis of several hormones and adreno-cortical steroids.

The **solanum alkaloids**, stated above are essentially the nitrogen-analogues of steroidal saponins. Unlike, their *oxygen* counterparts, all these N-containing alkaloids exhibit the same stereochemistry at C-25 (methyl being equatorial always), but C-22 isomers do exist, such as: solasodine and tomatidine.

The above cited *three* members of the **solanum alkaloids** shall be discussed as under:

A.1 Solasodine


Synonyms Solanocarpidine; Solanidine-S; Purapuridine.

Biological Sources It is obtained from the fruits of *Capsicum annuum* L. (*Solanaceae*) (**Chili, Paprika, Sweet Peppers**); shoots and berries of *S. dulcamara* L. (*Solanaceae*) (**Bittersweet, Bitter**

* GGPP = Geranylgeranyl diphosphate

Nightshade, Felonwood); leaves of *S. nigrum* L. (*Solanaceae*) (**Wonderberry, Black Nightshade, Prairie Huckleberry**).

Chemical Structure





Solasodine

(3 β , 22 α , 25R)-Spirosol-5-en-3-ol; (C₂₇H₄₃NO₂).

Isolation It is obtained by the hydrolysis of **solasonine** which yields **solasodine**, L-rhamnose, D-galactose and D-glucose respectively. It is the dehydrated product.

Characteristic Features

(i) It is obtained as hexagonal plates from methanol or by sublimation under high vacuum.



- (ii) It has mp 200-202°C; $[\alpha]_D^{25} - 98^\circ \text{C}$ [c = 0.14 in methanol]; $[\alpha]_D - 113^\circ$ (CHCl₃); pK_b 6.30.
 (iii) It is freely soluble in benzene, pyridine, and chloroform; moderately soluble in ethanol, methanol, and acetone; slightly soluble in water and practically insoluble in ether.

Identification Tests (for Solanum Alkaloids)


1. Dissolve 5-10 mg of the alkaloid in a few drops of hot amyl alcohol or ethanol and allow it cool gradually. The appearance of jelly-like product gives the characteristic test of the solanum alkaloids.
2. When a few mg of the alkaloids is treated with antimony trichloride solution in dry chloroform, it gives rise to a distinct red colouration.
3. The **solanum alkaloids**, in general, produces an instant red-violet colour with formaldehyde (HCHO) and sulphuric acid (H₂SO₄). This particular test is so distinct and sensitive that it is used for the quantitative estimation of these alkaloids colorimetrically.

Uses It is invariably used as a starting material for steroidal drugs.

A.2 Tomatidine


Biological Source It is obtained from the roots of Rutgers tomato plant [*Lycopersicon esculentum* Mill., cultivar. "Rutgers"] (*Solanaceae*) (Tomato).

Chemical Structure (3β, 5α, 22β, 25 S)-Spirosolan-3-ol; (C₂₇H₄₅NO₂).

**Characteristic Features**

1. It is obtained as plates from ethyl acetate having mp 202-206°C.
2. Its specific rotation $[\alpha]_D^{25} + 8^\circ$ (chloroform).


Isolation It is obtained by the hydrolysis of **tomatine** to yield a molecule of **tomatidine** along with 2 moles of D-glucose, 1-mole of D-xylose and 1-mole of D-galactose as depicted below:



Identification Test Its hydrochloride derivative (C₂₇H₄₅NO₂·HCl) is obtained as crystals from absolute ethanol having mp 265-270°C and $[\alpha]_D^{25} - 5^\circ$ (methanol).

A.3 Solanidine


Synonym Solatubine.




Biosynthesis of Solasodine, Tomatidine and Solanidine

- (ii) Another substitution affords 26-amino-22-hydroxycholesterol to cyclize thereby forming a heterocyclic piperidine ring,
- (iii) After 16 β -hydroxylation, the secondary amine is oxidized to an imine, and the ultimate *spiro-system* may be envisaged by virtue of a nucleophilic addition of the 16 β -hydroxyl on to the imine, and
- (iv) This specific reaction, however, establishes the configurations, *viz*: 22R-as in the case of **Solasodine**, and 22S-as in the case of **Tomatidine**.

B. Veratrum Alkaloids The **Veratrum alkaloids** represent the most important and medicinally significant class of steroidal alkaloids. It is, however, pertinent to mention here that the *basic ring systems* present in the Veratrum alkaloids are not quite the same as seen in the usual steroidal nucleus, as present either in the cholesterol or in the aglycone residues of the cardiac glycosides (A). Interestingly, one may observe in the structures of Veratrum alkaloids that the ring 'C' is a five-membered ring while ring 'D' is a six-membered ring (B) which apparently is just the reverse of the pattern in the regular steroidal nucleus as depicted in next page.



(A)



(B)

Examples

- (a) Alkamine portion of the ester alkaloids of **Veratrum**, viz., **Protoverine**, **Veracevine**, **Germine**.
 (b) Alkamine aglycones of glycosidic veratrum alkaloids, viz., **Veratramine**.

In general, the majority of **Veratrum alkaloids** may be classified into *two* categories solely based on their characteristic structural features, namely:

- (i) Ceveratrum alkaloids, and
 (ii) Jeveratrum alkaloids

These *two* categories of **Veratrum alkaloids** shall now be discussed individually in the sections that follows:

B.1 Ceveratrum Alkaloids The important alkaloids belonging to this group of alkaloids are, namely: **Protoveratrines**; **Veratridine**, **Cevadine**, **Germine** etc., which shall be treated separately hereunder:

B.1.1 Protoveratrines


Biological Sources It is obtained from the rhizome of *Veratrum album* L. (*Liliaceae*) and *Veratrum viride* Ait. (*Liliaceae*) (**American Hellebore**).

However, the alkaloids present in the rhizomes of *V. viride* are placed in *three* groups, such as:

Group-‘A’: Alkamines (esters of the steroidal bases) with organic acids, including **germidine**, **germitrine**, most valued therapeutically; besides, **cevadine**, **neogermitrine**, **neoprotoveratine**, **protoveratrines** and **veratridine**,

Group-‘B’: (Glycosides of the alkamines), mainly **pseudojervine** and **veratrosine**, and

Group-‘C’: (Alkamines), **germine**, **jervine**, **rubijervine**, and **veratramine**.

Chemical Structure

Isolation **Protoveratrine A and B** are usually extracted together and referred to as ‘**protoveratrines**’. About 2 kg of dried rhizomes of *V. album* is powdered and then extracted with benzene and ammonia. The total alkaloids are purified by extraction with acetic acid, re-extracted

into benzene. The solvent is removed under vacuum, the residue is dissolved in ether from which the crystalline powder of the crude protoveratrine separates out. The crude product is recrystallized from alcohol-acetic acid and upon subsequent alkanization of the solution with dilute ammonia. By this method one may obtain 8-10 g of **protoveratrine** powder from 8 kg of *V. album* rhizomes.

Consequently, **protaverine A and B** may be separated by the help of counter current distribution of the “**protaverine**” between benzene and acetate buffer (pH 5.5) and ultimately subjected to column chromatography on acid aluminium oxide (Al₂O₃).

Characteristic Features The characteristic features of the **protoveratrine**s are as follows:

- (i) The sternutative crystals obtained from ethanol have a slightly bitter taste.
- (ii) It decomposes at 266-267°C.
- (iii) Its specific rotation $[\alpha]_D^{25} - 38.6^\circ$ (pyridine), and $[\alpha]_D^{25} - 8.5^\circ$ (C = 1.99 in chloroform).
- (iv) It is soluble in chloroform, dilute aqueous acidic solutions and slightly soluble in ether. It is practically insoluble in water and petroleum ether.

However, the characteristic features of **protoveratrine A and B** are as stated below:

S. No.	Characteristic Features	Protoveratrine-A (Protalpa)	Protoveratrine-B (Veratrine; Neoprotoveratrine)
1.	Nature	Crystals obtained from acetone	Crystals obtained from acetone
2.	Decomposition temperature/mp	267-269° (dec.)	268-270° (dec.)
3.	Specific rotation	$[\alpha]_D^{25} - 40.5^\circ$ (pyridine); $[\alpha]_D^{25} - 10.5^\circ$ (chloroform);	$[\alpha]_D^{25} - 37^\circ$ (dec.); $[\alpha]_D^{25} - 3.5^\circ$ (chloroform);
4.	Solubility	Soluble in chloroform, hot ethanol and pyridine	Soluble in hot ethanol, pyridine and chloroform
5.	Stability in alkaline medium	Decomposes	Decomposes

Uses

1. It is used as an antihypertensive agent which exerts its action through reflex inhibition of pressor receptors in the heart and carotid sinus.
2. It also possesses emetic action.
3. It is used in the treatment of toxemia of pregnancy.


B.1.2 Veratridine

Biological Sources It is obtained from the seeds of *Schoenocaulon officinale* (Schelecht. and Cham.) A. Gray and also from the rhizome of *Veratrum album* L. (*Liliaceae*).

Chemical Structure (3β, 4α, 16β)-4, 9-Epoxyceveane-3, 4, 12, 14, 16, 17, 20-heptol 3-(3, 4-dimethoxybenzoate) as given under.

Isolation **Veratridine** can be isolated as the commercial **veratrine** (mixture) *i.e.*, the mixture of alkaloids **cevadine**, **veratridine**, **cevadiline**, **sabadine** and **cevine** obtained from the seeds of *S. officinale* stated above, as its sparingly soluble nitrate derivative.*

* Blount, *J. Chem. Soc.*, 122, (1935); Vejdelek *et. al. Chem. Listy* **153**, 33, (1956); Coll. *Czech. Chem. Commun.*, **22**, 98 (1957).



Characteristic Features

1. It is yellowish-white amorphous powder.
2. It tenaciously retains water.
3. It has mp 180°C after drying at 130°C.
4. Its specific rotation is $[\alpha]_D^{20} + 8.0^\circ$ (ethanol) and $pK_a 9.54 \pm 0.02$.
5. It is insoluble in water but slightly soluble in ether.

Identification Tests

1. It readily forms its nitrate derivative which is an amorphous powder and sparingly soluble in water.
2. Its sulphate salt is formed as its needles which happens to be very hygroscopic.
3. It readily forms its perchlorate derivative as long needles from water having mp 259-260°C (after drying at 120°C *in Vacuo*).

B.1.3 Cevadine

Synonym Veratrine

Biological Source It is obtained from the seeds of *Schoenocaulon officinale* (Schlecht and Cham.) A. Gray (*Sabadilla officinarum* Brandt.) belonging to family *Liliaceae*.


Chemical Structure $[3\beta(Z), 4\alpha, 16\beta]$ -4, 9-Epoxycevan-3, 4, 12, 14, 16, 17, 20-heptol 3-(2-methyl-2-butenate) as stated below.

Characteristic Features

1. It gives rise to flat needles from ether which decomposes at 213-214.5°C.
2. It has specific rotation $[\alpha]_D^{20} + 12.8^\circ$ (C = 3.2 in ethanol).
3. **Solubility:** 1 g dissolves in 15 ml ether or ethanol and is very slightly soluble in water.

Identification Tests

1. It forms **aurichloride derivative** which are obtained as fine yellow needles from ethanol that gets decomposed at 190°C.




- It readily produces **mercurichloride derivative** ($C_{32}H_{49}NO_9 \cdot HCl \cdot HgCl_2$) as silvery scales which decomposes at $172^\circ C$.

Caution Cevadine is extremely irritating locally particularly to the mucous membranes. Caution must be used in handling.

B.1.4 Germine

Biological Source Germine (an **alkamine**) is present in a plethora of *polyester alkaloids* that occur in *Veratrum* and *Zygadenus* species, such as: *Veratrum viride* Ait. (*Liliaceae*).

Chemical Structure ($3\alpha, 4\alpha, 7\alpha, 15\alpha, 16\beta$)-4, 9-Epoxyceevane-3, 4, 7, 14, 15, 16, 20-heptol ($C_{27}H_{43}NO_8$).



Characteristic Features

- It is obtained as crystals from methanol mp $221.5-223^\circ C$.
- It has specific rotation $[\alpha]_D^{25} + 4.5^\circ$ (95% ethanol) and $[\alpha]_D^{16} + 23.1^\circ$ ($C = 1.13$ in 10% acetic acid).


3. *Solubility*: It is soluble in chloroform, methanol, ethanol, acetone and water; and slightly soluble in ether.

Identification Tests It forms *three* different types of ‘acetates’ having specific characteristic features as stated below:


1. **3-Acetate derivative of Germinine ($C_{29}H_{45}NO_9$)**: It forms needles from ether having mp 219-221°C and $[\alpha]_D^{23} + 10^\circ$ (C = 1.05 in pyridine).
2. **16-Acetate derivative of Germinine ($C_{29}H_{45}NO_9$)**: It forms crystals from chloroform having mp 225-227°C and $[\alpha]_D^{23} - 19^\circ$ (C = 0.93 in pyridine).
3. **3, 4, 7, 15, 16-Pentaacetate derivative of Germinine ($C_{37}H_{53}NO_{13}$)**: It yields prisms from acetone + petroleum ether which decomposes at 285-287°C and $[\alpha]_D^{23} - 65^\circ$ (C = 0.65 in pyridine).

B.2 Jeveratrum Alkaloids The **Jeveratrum group of alkaloids** is usually represented by the structure of **veratramine**, **jervine** and **pseudojervine** etc., which essentially have the following salient features showing the points of difference in comparison to the *Ceveratrum alkaloids*:

The *three* important members of this particular category of alkaloids shall be treated separately in sections that follows:



Veratramine
[Jeveratrum Alkaloid]




Cevadine
[Ceveratrum Alkaloid]

S.No.	Jeveratrum Alkaloids	S.No.	Ceveratrum Alkaloids
1	The ring system beyond C-17 <i>i.e.</i> , the last two of the 6-membered rings, are altogether different from those in the Ceveratrum group of alkaloids	1	The alkamine portion has several –OH groups, (normally 6 to 9 in number), which are linked to various acids <i>e.g.</i> , vanillic acid, veratric acid, tiglic acid (angelic acid) to form their corresponding ester alkaloids.
2	The absence of the oxygen-bridge between C-4 and C-9.	2	An oxygen-bridge exists between C-4 and C-9.
3	The absence of several –OH moieties at C-4, C-12, C-14, C-17, C-20.	3	It has a chain of six cyclic rings of which the last two have a common N-atom.
4	The presence of a double-bond between C-5 and C-6.	4	At C-3 the H is replaced by a C ₅ H ₇ O moiety.

B.2.1 Veratramine

Biological Sources It is obtained in the rhizomes of *Viratrum viride* Ait. (*Liliaceae*) (**American Hellebore**); and also from *Veratrum grandiflorum* (Maxim.) Loes. F. (*Liliaceae*).

Chemical Structure The chemical structure of **veratramine** has also been referred to as **azasteroid**, wherein the N-atom is present in one or more side chains.



(3 β , 23 β)-14, 15, 16, 17-Tetrahydro-veratraman-3, 23-diol (C₂₇H₃₉NO₂).

Characteristic Features

1. It is obtained as crystals having mp 206-207°C.
2. It is slightly soluble in water, but soluble in ethanol and methanol.


Identification Tests

1. It forms a complex with digitonin (1:1) that has uv_{max} : 268 nm and $[\alpha]_D^{25} - 71.8^\circ$ (C = 1.21; $[\alpha]_D^{25} - 70^\circ$ (C = 1.56 in methanol).
2. **Dihydroveratramine Derivative:** The crystals of dihydroveratramine derivative has mp 192.5-194°C ; $[\alpha]_D^{25} + 26^\circ$ (C = 1.26 in acetic acid).

B.2.2 Jervine

Biological Sources It is obtained in the rhizomes of *Veratrum grandiflorum* (Maxim.) Loes F. *Veratrum album* L., and *Veratrum viride* Sol. (*Liliaceae*).

Chemical Structure



(3 β , 23 β)-17, 23-Epoxy-3-hydroxyveratraman-11-one (C₂₇H₃₉NO₃).

Characteristic Features

1. The needles obtained from methanol and water has mp 243.5-244°C (Saito).
2. Its specific rotation $[\alpha]_D^{20} - 150^\circ$ (ethanol) (Saito); and $[\alpha]_D^{20} - 167.6^\circ$ (chloroform) (Poethke).
3. It has uv_{max} : 250, 360 nm (ϵ 1500, 60).


Identification Tests

1. **Diacetyljervine (C₃₁H₄₃NO₅):** The diacetyljervine has mp 173-175°C; $[\alpha]_D - 112^\circ$; uv_{max} (ethanol): 250, 360 nm (ϵ 16400, 80).
2. **Jervine Hydrochloride** has mp 300-302°C.

B.2.3 Pseudojervine

Biological Sources It is obtained from the rhizomes of *Veratrum viride* Ait (*Liliaceae*) (**American Hellebore**); *V. album* L. (*Liliaceae*); and *V. eschscholtzii* Gray. (*Liliaceae*).

Chemical Structure It is the glucoside of **jervine** as given below.



(3 β , 23 β)-17, 23-Epoxy-3-(β -D-glucopyranosyloxy) veratraman-11-one. (C₃₃H₄₉NO₈).

Characteristic Features

1. It is obtained as lustrous leaflets having mp 300-301°C (dec.).
2. Its specific rotation $[\alpha]_D^{25} - 133^\circ$ (C = 0.48 in 1.3 ethanol-chloroform).
3. **Solubility:** It is soluble in benzene, chloroform; slightly soluble in ethanol and almost insoluble in ether.

Note: It is, however, pertinent to observe here that the *Zygadenus* species and the *Schoenocaulon* species appear to have only the *Ceveratrum* alkaloids and practically no *Jeveratrum* alkaloids. Interestingly, the large number of *Veratrum* species seem to contain both these type of steroidal alkaloids.

7.2.2 Alkaloids Derived from Anthranilic Acid

Anthranilic acid is found to be a key intermediate in the biosynthesis of L-tryptophan. Therefore, it has been established that this biotransformation ultimately is solely responsible to the elaboration of the **indole alkaloids**. In the course of this conversion, the anthranilic acid residue is specifically decarboxylated, thus the C₆N skeleton is further utilized. In general, there are several such instances wherein the anthranilic acid itself serves as an *alkaloid precursor*, by employing various means and processes that essentially retain the full skeleton and further exploit the carboxyl function legitimately. Interestingly, in mammals, L-tryptophan gets degraded back to anthranilic acid. However, this particular route is of least importance in the plant kingdom.

The alkaloids derived from anthranilic acid may be classified into *three* major categories, namely:

- (i) Quinazoline alkaloids,
- (ii) Quinoline alkaloids, and
- (iii) Acridine alkaloids.

The aforesaid categories of alkaloids shall be discussed in an elaborated fashion hereunder individually.

7.2.2.1 Quinazoline Alkaloids


Vasicine is a **quinazoline alkaloid** which will be described below.

A. Vasicine

Synonym Peganine

Biological Sources It is obtained from the leaves of *Adhatoda vasica* (L.) Nees (*Acanthaceae*) (**Malabar Nut, Adotodai, Paveltia**); and the seeds of *Peganum harmala* L. (*Rutaceae*) (**Harmel, Syrian Rue, African Rue**).

Chemical Structure



Vasicine

1, 2, 3, 9-Tetrahydropyrrolo [2, 1-b] quinazoline-3-ol; (C₁₁H₁₂N₂O).

Isolation It is isolated from the leaves of *Adhatoda vasica** and also from the seeds of *Peganum harmala*** by adopting the standard methods of isolation described earlier in this chapter.

* Sen, Ghosh, *J. Indian Chem. Soc.*, **1**, 315 (1924);

** Späth, Nikawitz, *Ber.* **67**, 45, (1934);

Characteristic Features

- d,l*-Form:**
1. It is obtained as needles from ethanol having mp 210°C.
 2. It sublimes on being subjected to high vacuum.
 3. It is soluble in acetone, alcohol, chloroform; and slightly soluble in water, ether and benzene.
- l*-Form:**
1. It is obtained as needles from ethanol with mp 212°C.
 2. Its specific rotation $[\alpha]_D^{14} - 254^\circ$ (C = 2.4 in CHCl₃); $[\alpha]_D^{14} - 62^\circ$ (C = 2.4 in ethanol).

Note: In dilute HCl it is obtained as its dextrorotatory form.


Identification Tests

1. Hydrochloride dihydrate derivative is obtained as needles having mp 208°C (dry).
2. Hydroiodide dihydrate derivative is formed as needles with mp 195°C (dry).
3. Methiodide derivative is obtained as needles from methanol having mp 187°C.
4. Acetyl vasicine derivative (C₁₁H₁₁N₂O COCH₃) is formed as crystals having mp 123°C and bp_{0.01} 230-240°C.

Uses


1. It is mostly used as an expectorant and bronchodilator.
2. It also shows oxytocic properties very similar to those exhibited by **oxytocin** and **methyl ergometrine**.
3. **Vasicine** also shows abortifacient action which is due to the release of prostaglandins.

Biosynthesis of Vasicine Various studies in *Peganum harmala* have evidently revealed **vasicine** (**peganine**) to be derived from the anthranilic acid, while the remaining portion of the structure comprising of a pyrrolidine ring provided by ornithine. The probable mechanism of **vasicine** skeleton may be explained by virtue of the nucleophilic attack from the N-atom present in anthranilate upon the pyrrolidinium cation, ultimately followed by amide formation. However, interestingly this pathway is not being adopted in *Justicia adhatoda*. Thus, a comparatively less predictable sequence from N-acetylanthranilic acid and aspartic acid is observed as shown below:



B. Vasicinone

Biological Source The plant source remain the same as described under **vasicine**.

Chemical Structure

1, 2, 3, 9-Tetrahydropyrrolo [2, 1-b] quinazoline-6-one-3-ol ($C_{11}H_{10}N_2O_2$).




Uses It is used mainly as an expectorant which action is solely due to stimulation of the bronchial glands.


7.2.2.2 Quinoline Alkaloids

In general, the alkaloids containing essentially the '**quinoline**' nucleus include a series of alkaloids obtained exclusively from the **cinchona bark**, the major members of this particular group are, namely: **quinine**, **quinidine**, **cinchonine** and **cinchonidine**. Interestingly, more than twenty five alkaloids have been isolated and characterized either from the *Yellow Cinchona* i.e., *Cinchona calisaya* Wedd. and *Cinchona ledgeriana* Moens ex Trimen, or from the *Red Cinchona* i.e., *Cinchona succirubra* Pavon ex Klotzsch (Family: *Rubiaceae*). The aforesaid alkaloids are also found in their hybrids as well as in the *Cuprea Bark* obtained from *Remijia pedunculata* and *Remijia purdieana* belonging to the natural order *Rubiaceae*.


However, it has been revealed that an average commercial yield of the **cinchona alkaloids** in the dry bark materials from the said plant materials are as follows: **quinine** (5.7%); **quinidine** (0.1-0.3%); **cinchonine** and **cinchonidine** (0.2-0.4%). Nevertheless, the other closely related minor alkaloids are present in relatively smaller quantities.

Basic Structures of Cinchona Alkaloids The various **quinoline alkaloids**, which possess potent medicinal activities are, namely: **quinine**, **quinidine**, **cinchonine**, and **cinchonidine**. It is interesting to observe that these alkaloids not only have a closely related structure but also similar medicinal characteristics. These alkaloids possess the *basic skeleton* of 9'-rubanol that is derived from the parent compound known as ruban. Thus, ruban is obtained from the combination of *two* distinct heterocyclic nuclei, namely: (a) 4-methyl quinoline nucleus, and (b) quinuclidine nucleus. However, this particular nomenclature was suggested by Rabe so as to simplify the naming of such compounds and also to signify its origin from the natural order *Rubiaceae*.







Ruban
[8-9' Quinuclidyl Methyl
Quinoline nucleus]




9'-Rubanol




Quinine



Quinidine



Cinchonine




Cinchonidine

In this context, a few important drugs belonging to the **quinoline alkaloids** shall now be discussed in the sections that follows:

A. Quinine

Biological Sources It is obtained from the bark of *Cinchona calisaya* Wedd; *Cinchona ledgeriana* Moens ex Trimen; *Cinchona officinalis* Linn. f.; *Cinchona robusta* How; and *Cinchona succirubra* Pavon ex Klotzsch belonging to family *Rubiaceae*.

Chemical Structure



(8 α , 9R)-6'-Methoxycinchonan-9-ol, (C₂₀H₂₄N₂O₂).

Isolation The schematic method of isolation of the *Cinchona Alkaloids* in general and that of **quinine** in particular has been provided in the following *Flow-chart* in a sequential manner. Hence, this particular flow-chart also includes the method of isolation of other important members of this group *i.e.*, **quinidine**, **cinchonine** and **cinchonidine** as given under.


Notes

1. Bisulphates of cinchona as alkaloids [B.H₂SO₄] are readily soluble in water.
2. **Quinine sulphate** [Br.H₂SO₄] is sparingly soluble in water [1:720].
3. **Cinchonine** is practically insoluble in ether.
4. Tartrates of **Quinine** and **Cinchonidine** are insoluble, whereas the tartrates of **Cinchonine** and **Quinidine** are soluble in water.

Characteristic Features

1. The orthorhombic needles obtained from absolute ethanol are triboluminescent and having mp 177°C (with some decomposition).
2. It sublimes in high vacuum at 170-180°C.
3. Its specific rotation is $[\alpha]_D^{15} - 169^\circ$ (C = 2 in 97% ethanol); $[\alpha]_D^{17} - 117^\circ$ (C = 1.5 in chloroform); and $[\alpha]_D^{15} - 285^\circ$ (C = 0.4 M in 0.1 N H₂SO₄).
4. Its dissociation constant pK₁ (18°) is 5.07 and pK₂ 9.7.
5. **Neutral Salt of Quinine [(B)₂.H₂SO₄.8H₂O]**: It is formed by neutralization from boiling water, which is sparingly soluble in water (*viz.*, 1 in 720 at 25°C). The octahydrate neutral salts of quinine undergoes efflorescence on being exposed to air and gets converted to the corresponding dihydrate salt which is more stable.
6. **Acid Sulphate of Quinine [(B).H₂SO₄.7H₂O]**: The quinine bisulphate is soluble in water (1 in 8.5 at 25°C) and in ethanol (1 in 18). The aqueous solution is acidic to litmus.
7. **Tetrasulphate Salt of Quinine [(B)₂.2H₂SO₄.7H₂O]**: The tetrasulphate salt of quinine is very soluble in water.

Schematic Method of Isolation of Cinchona Alkaloids



Identification Tests

1. **Fluorescence Test:** Quinine gives a distinct and strong blue fluorescence when treated with an oxygenated acid, such as: acetic acid, sulphuric acid. This test is very marked and pronounced even to a few mg concentration of quinine.

Note: The hydrochloride and hydroiodide salts of quinine do not respond to this fluorescence test.

2. **Thalleioquin Test:** Add to 2-3 ml of a weakly acidic solution of a quinine salt a few drops of bromine-water followed by 0.5 ml of strong ammonia solution, a distinct and characteristic emerald green colour is produced. The coloured product is termed as **thalleioquin**, the chemical composition of which is yet to be established. This test is so sensitive that quinine may be detected to a concentration as low as 0.005%.

Notes: Quinidine and cupreine (a Remijia alkaloid) give also a positive response to this test; but cinchoninine and cinchonidine give a negative test.

3. **Erythroquinine Test (or Rosequin Test):** Add to a solution of quinine in dilute acetic acid 1-2 drops of bromine water, a drop of a solution of potassium ferrocyanide [$K_4(FeCN)_6$] (10% w/v), and to it add a drop of strong ammonia solution, the solution turns red instantly. In case, it is shaken immediately with 1 ml of chloroform, the red colour is taken up by the chloroform layer.

4. **Herpathite Test:** To a boiling mixture containing 0.25 g of quinine in 7.5 ml glacial acetic acid, 3 ml ethanol (90% v/v), 5 drops of conc. sulphuric acid and add to it 3.5 ml of 1% iodine solution in ethanol, the appearance of crystals of **iodosulphate of quinine** (*i.e.*, **sulphate of iodo-quinine**)-is known as **Herpathite** after the name of its discoverer. It has the chemical composition [$(B_4).3H_2SO_4.2HI.I_4.3H_2O$] which separates out as crystals (on cooling), having a metallic lustre that appears dark green in reflected light and olive green in transmitted light.

Uses

1. It is used as a flavour in carbonated beverages.
2. It is widely used as an antimalarial agent in tropical countries.
3. It is employed as a skeletal muscle relaxant.

Biosynthesis of Quinine The various steps whereby *Coryanthe-type* indole alkaloids are converted to quinoline derivatives have not yet been elucidated and hence established. Therefore, only a partial biosynthetic pathway may be written for **quinine** as given under.

B. Cinchonine

Biological Source **Cinchonine** is obtained from a variety of cinchona bark, especially in the bark of *Cinchona micrantha* R. and P. belonging to family *Rubiaceae*.

Chemical Structure Please see structure under Section 7.2.2.2, (9S-Cinchonan-9-ol) ($C_{19}H_{22}N_2O$).


Isolation It has already been described under **quinine** Section 'A' above.

Characteristic Features

1. Its prisms, needles are obtained from ether and ethanol having mp 265°C.
2. It begins to sublime at 220°C.
3. Its specific rotation is $[\alpha]_D + 229^\circ$ (in ethanol).
4. One gramme of it dissolves in 60 ml ethanol, 25 ml boiling ethanol, 110 ml chloroform and 500 ml ether. It is practically insoluble in water.

Identification Tests

1. **Cinchonine dihydrochloride ($C_{19}H_{22}N_2O.HCl$):** It is white or faintly yellow crystals or crystalline powder. It is freely soluble in water and ethanol.




Biosynthesis of Quinine

2. **Cinchonine hydrochloride dihydrate** ($C_{19}H_{22}N_2O \cdot HCl \cdot 2H_2O$): It is obtained as fine crystals having mp when anhydrous $215^\circ C$ with decomposition. One g dissolves in 20 ml of water, 3.5 ml of boiling water, 1.5 ml of ethanol, 20 ml of chloroform and slightly soluble in ether.
3. **Cinchonine sulphate dihydrate** [$(C_{19}H_{27}N_2O)_2 \cdot H_2SO_4 \cdot 2H_2O$]: It occurs as lustrous, very brittle crystals having mp $198^\circ C$ (when anhydrous). One g dissolves in 65 ml of water, 30 ml of hot water, 12.5 ml of ethanol, 7 ml of hot ethanol, 47 ml of chloroform and slightly soluble in ether.

7.2.2.3 Acridine Alkaloids

The origin of the **acridine-ring-system** is by virtue of an extension of the process that essentially involves the combination of anthranilic acid and acetate/malonate as shown in the following sequence of reactions; whereas, a rather more direct route to the above leads to the **quinoline-ring-system** discussed in Section 7.2.2.2 earlier.




There are a few typical examples of the **acridine alkaloids**, such as: **Rutacridone**, **Acronycine** and **Melicopicine**.

A. Rutacridone

Biological Source The fresh and dried leaves of *Ruta graveolens* L. (*Rutaceae*) (**Rue, Garden Rue, German Rue**).


Chemical Structure




Uses

1. In Chinese medicine rue is considered as an emmenagogue, hemostat, intestinal antispasmodic, sedative, uterine stimulant, vermifuge, rheumatism, cold and fever.
2. In Poland, it is used as an aphrodisiac and choleric.
3. The herb is used medicinally as a bitters, an aromatic stimulant, ecbolic and in suppression of the menses.

The chemical structures of **acronycine** and **melicopicine** are given below:




Acronycine



Melicopicine

Biosynthesis of Rutacridone, Acronycine and Melicopicine The anthraniloyl-CoA is observed to act as a *starter-unit* for the extension of chain *via* one molecule of **malonyl-CoA**, and formation of amide ultimately generates the heterocyclic system, that would adopt finally the more stable 4-hydroxy-2-quinolone form as shown in the following sequence of reactions. Interestingly, the position C-3 is highly nucleophilic; and, therefore, is susceptible to alkylation, especially *via* dimethylallyl diphosphate in the instance of all the three alkaloids, namely: **rutacridone**, **acronycine**, and **melicopicine**. This seems to allow the formation of additional six-membered oxygen containing heterocyclic ring system (**acronycine**); and five-membered oxygen containing heterocyclic ring system (**rutacridone**).



7.2.3 Alkaloids Derived from Histidine

The amino acid *L-histidine*, containing the heterocyclic imidazole ring, is considered to be the right precursor of alkaloids that essentially comprise of this ring-system.

A good number of *Pilocarpus species*, belonging to family *Rutaceae*, found to contain plethora of alkaloids with an *imidazole ring*, namely: **pilocarpine**, **isopilocarpine**, and **pilosene**. It has been observed that the alkaloids in these species invariably reside in the leaves. **Pilocarpine** constitutes 0.5-1.0% of the dried leaf material. **Isopilocarpine** appears to vary significantly within a range from 5 to 7.5% of the total alkaloids. Further, the alkaloids are located mostly in the upper epidermal leaves of the cells of the leaves, and also in the cells of the mesophyll bordering upon the lower epidermis.

The *three* major alkaloids derived from histidine shall be described in the sections that follows.


7.2.3.1 Pilocarpine

Synonyms Ocusert Pilo.

Biological Source **Pilocarpine** is obtained from the leaves of closely related plants of the genus *Pilocarpus*, belonging to the natural order *Rutaceae*. However, the genus comprised of a variety of species commonly known by various names, such as: *Pilocarpus jaborandi* (*Pernambuco Jaborandi*), (*Pilocarpus pennatifolius* (**Paraguay Jaborandi**); *Pilocarpus microphyllus* (**Maranhã Jaborandi**); *Pilocarpus selloanus* (**Rio Jaborandi**); *Pilocarpus trachylophus* (**Ceara Jaborandi**); *Pilocarpus spicatus* (**Aracati Jaborandi**); *Pilocarpus heterophyllus* (**Barqui Simento Jaborandi**); and *Pilocarpus racemosus* (**Guadeloupe**).


It is worthwhile to mention here that *P. microphyllus* is the major commercial source of this drug.

Chemical Structure



(3*S*-*cis*)-3-Ethyldihydro-4-[1-methyl-1*H*-imidazol-5-yl]methyl]-2(3*H*)-furanone (C₁₁H₁₆N₂O₂).

Pilocarpine is a monoacidic tertiary base comprising of a lactone ring and an imidazole nucleus. It is the lactone of **pilocarpic acid**, an acid with a **glyoxaline** nucleus, as given below:



Isolation The finely powdered leaves of **Jaborandi** is first extracted with ethanol (95% v/v) containing 1% HCl. The ethanol is distilled off under vacuo and the residue is taken up with a little

water and neutralized carefully by the addition of dilute ammonia. The resins separating out are filtered off and the filtrate is concentrated to a small volume. The resulting concentrated filtrate is alkali-fied with an excess of ammonia and the liberated alkaloids are shaken out with at least three successive portions of chloroform. The chloroform is removed from the combined extract under vacuo.


The residue is dissolved in a minimum volume of distilled water and neutralized with dilute HNO_3 (6N). The mixture of nitrates of **pilocarpine** and **isopilocarpine** crystallizes out upon cooling; which may be further separated by fractional crystallization from ethanol.

Characteristic Features

1. It is found as oil or crystals having mp 34°C .
2. It boils at bp₅ 260°C with partial conversion to its isomer isopilocarpine.
3. Its specific rotation is $[\alpha]_D^{18} + 106^\circ$ ($C = 2$) and dissociation constant pK_1 (20°C) 7.15; and pK_2 (20°C) 12.57.
4. It is soluble in water, alcohol, chloroform, sparingly soluble in ether and benzene; and practically insoluble in petroleum ether.
5. It exhibits an absorption maximum at 263 nm.
6. It behaves as a monoacidic base.
7. It usually gives distinct precipitates with a number of reagents, such as: **Wagner's Reagent**, **Mayer's Reagent**, **Hager's Reagent**, **silicotungstic acid**, **phosphomolybdic acid**, gold and platinum halides.


Note: Some of these precipitates do help in the identification of pilocarpine.

8. **Cessation of Lactone-Ring:** The lactone ring is opened-up (undergoes cessation) by treatment with strong alkalies like NaOH, KOH, which ultimately form salts with the formation of **pilocarpic acid** as given below:




Note: The cessation of the lactone-ring absolutely destroys the physiological activity of pilocarpine; and the lactone-ring is not affected by either ammonia or alkali carbonates.


9. **KMnO₄-Oxidation:** KMnO_4 oxidation destroys the imidazole ring in pilocarpine and yields ammonia, methyl amine, **pilopic acid**, **homopilopic acids** plus other products.



Isomerism **Pilocarpine** and **isopilocarpine** are stereoisomers, that essentially exhibit the stereochemical difference in the **lactone moiety** of the molecule as shown below:



Pilocarpine



Isopilocarpine

However, the above observation is based on the experimental evidence, which specifically depicts that the isomerism of the above two alkaloids still persists, even when the **imidazole moiety** undergoes destruction under mild experimental conditions.

Identification Tests

1. **Helch's Violet-Colour Test:** Pilocarpine readily forms a violet coloured compound when a solution of either the base or its salt is first treated with hydrogen peroxide (H_2O_2) and then with potassium dichromate ($\text{K}_2\text{Cr}_2\text{O}_7$) in the presence of few drops of dilute sulphuric acid (Helch, 1902).

Note:

- (i) **The violet-coloured compound (*i.e.*, pilocarpine perchromate) is soluble in chloroform and benzene. It was further characterized as pilocarpine perchromate by Biedebach (1933).**
 - (ii) **Shupe successfully employed the Helch's reaction to determine pilocarpine quantitatively by the colourimetric assay.**
2. **Ekkert's Colour Tests:** Add to 1 ml of 1% (w/v) solution of pilocarpine hydrochloride ($\text{C}_{11}\text{H}_{16}\text{N}_2\text{O}_2 \cdot \text{HCl}$) 1 ml of sodium nitroprusside solution (2% w/v) and 1 ml of NaOH solution (1N). Allow the reaction mixture to stand for 6-8 minutes and then acidify with dilute HCl when a wine or red colour develops.

(Note: Isopilocarpine hydrochloride also gives a similar colour test.)

Further, when a few drops of 0.1 N sodium thiosulphate solution are added to the wine or red colour solution, it changes to distinct green colouration.

Note: Elvidge (1947) put forward a method for the assay of the total alkaloids of *Pilocarpus* leaves based on the Ekkert's colour test.


Uses

1. Pilocarpine possesses miotic and diaphoretic actions.
2. Pilocarpine nitrate is used extensively as an ophthalmic drug having cholinergic action.
3. It is also employed to reduce the intra-ocular pressure in glaucoma patients.

7.2.3.2 Isopilocarpine

Synonym β -Pilocarpine.

Biological Source It is same as stated under Section 7.2.3.1 on **pilocarpine**.

Chemical Structure

Isolation It has been discussed under Section 7.2.3.1 above

Characteristic Features

1. It is a hygroscopic oily liquid or prisms.
2. It has the following physical parameters: bp₁₀ 261°C; $[\alpha]_D^{18} + 50^\circ$ (C = 2); pK₁ (18°C) 7.17.
3. It is miscible with water and alcohol; very soluble in chloroform; less soluble in ether and benzene; and almost insoluble in petroleum-ether.

Identification Tests Its derivatives have the specific physical parameters, namely:

1. **Isopilocarpine hydrochloride hemihydrate (C₁₁H₁₆N₂O₂ · HCl · ½H₂O):** It is obtained as scales from ethanol having mp 127°C; when anhydrous mp 161°C; $[\alpha]_D^{18} + 39^\circ$ (C = 5). It is soluble 0.27 part water and 2.1 parts ethanol.
2. **Isopilocarpine nitrate (C₁₁H₁₆N₂O · HNO₃):** It occurs as prisms from water, scales from ethanol, having mp 159°C; $[\alpha]_D^{18} + 39^\circ$ (C = 2). It is soluble in 8.4 parts of water and in 350 parts of absolute ethanol.


Uses

1. It is used as an antiglaucoma agent
2. It is also employed as miotic.

7.2.3.3 Isopilosine

Synonyms Carpiline; Carpidine; Pilosine (this compound was originally called pilosine *i.e.*, the *cis*-isomer of isopilosine).

Biological Source It is obtained from the dried leaflets of *Pilocarpus microphyllus* (Rutaceae), which has the total alkaloidal content (0.5-1.%) that consists principally **pilocarpine** along with small portion of **isopilosine**, **pilosine** and related structures.

Chemical Structure


[3S-[3 α (S*), 4 β]]-Dihydro-3-(hydroxyphenylmethyl)-4-[(1-methyl-1H-imidazol-5-yl) methyl]-2(3H)-furanone. (C₁₆H₁₈N₂O₃).

Isolation It is isolated from the leaves of *P. microphyllus* Stapf. (*Rutaceae*) by adopting standard procedures.*

Characteristic Features


1. It is obtained as needles from ethanol mp 182-182.5°C.
2. Its specific rotation $[\alpha]_D^{20} + 83.9^\circ$ (ethanol); uv_{max} (ethanol) : 210 nm ($\log \epsilon$ 4.10).

Biosynthesis of Imidazole Alkaloids *L-Histidine*, an amino acid contains an imidazole ring; and is, therefore, the most probable precursor of alkaloids containing this ring system. The **imidazole alkaloids** usually found in Jaborandi leaves (*P. microphyllus* and *P. jaborandi*; *Rutaceae*) are most likely derived from histidine; however, sufficient experimental data are lacking. Interestingly, the additional carbon atoms may originate from acetate or supposedly from threonine in the case of **pilocarpine**, whereas **pilosine** incorporates a phenylpropane C₆H₃ unit as shown below.



7.2.4 Alkaloids Derived from Lysine

The amino acid *L-lysine* happens to be the homologue of *L-ornithine*, and it also caters as an alkaloid precursor, employing pathways that are analogous to those known for ornithine. The 'additional methylene moiety' present in lysine affords the formation of six-membered piperidine ring systems, very similar to ornithine that provided five-membered ring systems, as shown below:



The various alkaloids that are derived from lysine are invariably grouped together under the following categories, namely:

- (a) Piperidine alkaloids
- (b) Quinolizidine alkaloids, and
- (c) Indolizidine alkaloids

The above cited categories of alkaloids shall now be discussed separately here under.

* F.L. Pyman *J. Chem. Soc.* **101**, 2260 (1912).

7.2.4.1 Piperidine Alkaloids


The various important alkaloids that essentially have the *piperidine nucleus* are, namely: **Coniine**, **Lobeline**, **Lobelanine** and **Piperine**, which shall be discussed individually in the sections that follows:

A. Coniine

Synonyms Cicutine; Conicine.

Biological Source It is the toxic principle of poison *Hemlock*, *Conium maculatum* L. (*Umbelliferae*). It is found in the seeds of *Cicuta maculata* L. (*Apiaceae*) (Water Hemlock).

Chemical Structure



Coniine

(S)-2-Propylpiperidine (C₈H₁₇N)

It occurs naturally as the (S)–(+) isomer.

Isolation It is isolated by standard procedures described earlier from the pitcher plant *Sarracenia flava**

The various steps involved in the isolation of **coniine** are as follows:

1. The powdered Hemlock fruits are mixed with KOH solution and then subjected to steam distillation. The distillate thus obtained is neutralized and evaporated to dryness.
2. The resulting residue is extracted with ethanol successively, filtered and the solvent evaporated under vacuo. The ethanol extracts the alkaloidal salts that are dissolved in water, which is subsequently rendered alkaline with KOH solution and finally extracted with ether at least 3-4 times.
3. Ether is evaporated under vacuo, when an oily liquid comprising of the free bases remains as the residue.
4. The fractional distillation of the oily liquid *under vacuo* or in a current of hydrogen gas, separates them into coniine and γ -coniceine at approximately 171-172°C. These two alkaloids are converted to their corresponding hydrochlorides, evaporated to dryness and extracted with acetone. Thus, coniine hydrochloride will be obtained as the insoluble substance, while the coniceine hydrochloride shall remain in acetone and recovered separately.

Note: Coniine enjoys the distinction of being the first ever alkaloid prepared synthetically.

Characteristic Features

1. It is a colourless alkaline liquid, which darkens and polymerizes on exposure to light and air.
2. It has a typical mousy odour.
3. It has mp $\sim -2^\circ\text{C}$, and bp 166-166.5°C; bp₂₀ 65-66°C.
4. It is a steam-volatile substance.

* N.V. Mody *et al. Experientia*, **32**, 829 (1976)

5. Its physical parameters are: d_4^{20} 0.8440-0.848; n_D^{23} 1.4505; $[\alpha]_D^{25} + 8.4^\circ$ (C = 4.0 in chloroform); $[\alpha]_D^{23} + 14.6^\circ$ (neat) and pK_a 3.1.
6. *Solubility*: 1 ml dissolves in 90 ml of water, and less soluble in hot water. The base dissolves about 25% of water of ambient temperature. It is freely soluble in ethanol, ether, benzene, acetone and amyl alcohol; and slightly soluble in chloroform.

Identification Tests

1. **Coniine Hydrobromide Derivative (C₈H₁₇N · HBr)**: Its prisms have mp 21°C ; 1 g dissolves in 2 ml water, 3 ml ethanol, and freely soluble in ether and chloroform.
2. **Coniine Hydrochloride Derivative (C₈H₁₇N · HCl)**: It occurs as rhomboids having mp 221°C, freely soluble in water, chloroform and ethanol.

(*R*)-(-) *Form*: It is a liquid, bp₇₅₆ 165°C; $[\alpha]_D^{25} - 8.1^\circ$ (C = 4.0 in chloroform); $[\alpha]_D^{23} - 14.2^\circ$ (neat).

(±)-*Form*: It has bp 200-210°C.

Uses


1. It has been used in convulsive and spasmodic diseases, such as: asthma, chorea, epilepsy, pertussis and tetanus.
2. **Coniine** has also been recommended in carditis, delirium, glandular swellings, jaundis, mania, nervous diseases, neuralgia, rheumatism, spasms and ulcers.

B. Lobeline

Synonyms α-Lobeline; Inflatine;

Biological Sources It is obtained from the herb and seeds of *Lobelia inflata* L., (*Lobeliaceae*) (**Indian Tobacco, Asthma Weed**); leaves of *Lobelia tupa* L. (*Campanulaceae*) (**Tupa, Devil's Tobacco**).

Chemical Structure



Lobeline

[2R-[2α, 6α (S*)]]-2-[6-(2-Hydroxy-2-phenylethyl)-1-methyl-2-piperidinyl]-1-phenylethanone (C₂₂H₂₇NO₂).

Isolation The various steps are as follows:

1. The powdered **lobelia herb** is moistened with water, acidified slightly with acetic acid and left as such for 3-4 hours. The resulting mass is then pressed and the process of moistening and pressing is repeated subsequently.

2. The acidic solutions thus collected are mixed and rendered alkaline with sodium bicarbonate carefully. The alkaline solution is extracted with ether successively. For purification, the ethereal extract is shaken with water, acidified with dilute sulphuric acid. The acidified liquid is again rendered alkaline with sodium bicarbonate solution and shaken with ether.
3. The combined ethereal extract is evaporated and the yellow oily liquid, comprising of the total alkaloids, is dissolved in water, acidified with HCl, filtered and then shaken with chloroform successively. Thus, the chloroform will exclusively extract the lobeline hydrochloride, while leaving the salts of the other alkaloids in the aqueous layer. The chloroform is then evaporated under vacuo to obtain the brownish oily residue.
4. The above residue is then taken up with double its volume of hot water at 80°C. The aqueous solution is kept in a vacuum desiccator over concentrated H₂SO₄ for several hours when lobeline hydrochloride separates out as crystals.
5. To recover the **lobeline** base, the resulting HCl salt is dissolved in warm water, rendered alkaline with dilute NaOH carefully and extracted with ether several times. The ethereal extract is evaporated and the residue is recrystallized from ethanol or benzene.

Characteristic Features

1. Lobeline is obtained as needles from ethanol, ether and benzene having mp 130-131°C, and specific rotation $[\alpha]_D^{15} - 43^\circ$ (ethanol).
2. It is freely soluble in chloroform, ether, benzene and hot ethanol; and very slightly soluble in water and petroleum ether.

Identification Tests


1. **Colour Test: Lobeline** on the addition of a few drops of concentrated sulphuric acid followed by a drop of formalin solution gives rise to a distinct red colouration.
2. **Froehd's Test:** It produces an instant rose red colouration with **Froehd's Reagent** that ultimately changes to blue.
3. **Erdmann's Reagent:** It develops a faint green colour which intensifies on slight warming.
4. **Lobeline Hydrochloride (C₂₂H₂₇NO₂·HCl) (Lobron, Zoolobelin):** It is obtained as rosettes of slender needles from ethanol with mp 178-180°C; $[\alpha]_D^{20} - 43^\circ$ (C = 2); and uv_{max} (methanol) 245, 280 nm (log ϵ 4.08, 3.05). Its solubility profile is as follows: 1 g dissolves in 40 ml of water, 12 ml of ethanol, very soluble in chloroform and very slightly soluble in ether. A 1% (w/v) solution in water has a pH of 4.0-6.0.
5. **Lobeline Sulphate [(C₂₂H₂₇NO₂)₂·H₂SO₄] (Lobeton, Unilobin, Bantron, Toban, Lobidan):** Its crystals obtained from ethanol exhibits specific rotation $[\alpha]_D^{20} - 25^\circ$ (C = 2). It is soluble in 30 parts of water and slightly in ethanol.

Uses

1. It is widely used as a respiratory stimulant.
2. Its effects resemble those of **nicotine** and hence used in lozenges or chewing tablets, containing 0.5-1.5 mg of **Lobeline Sulphate**, to help in breaking the tobacco habit, otherwise known as 'smoking deterrants'.

C. Lobelanine

Biological Source After **lobeline**, **lobelanine** is obtained as the most abundant alkaloid of *Lobelia inflata* L. (*Lobeliaceae*). (**Indian Tobacco, Asthma Weed**).

Chemical Structure

Lobelanine

cis-2, 2'-(1-Methyl-2, 6-piperidine-diyl) bis [1-phenylethanone], (C₂₂H₂₅NO₂).

Isolation The aqueous layer obtained in step (3), as stated under isolation of lobeline, is subjected to column chromatography and the **lobelanine** is collected as one of the major fractions.

Characteristic Features


1. It is obtained as rosettes of needles from ether or petroleum ether having mp 99°C.
2. It is freely soluble in acetone, benzene, ethanol, chloroform; and slightly soluble in water and ether.

Identification Tests

1. **Lobelanine Hydrochloride (C₂₂H₂₅NO₂.HCl)**: The crystals obtained from dilute ethanol decomposes at 188°C; it is soluble in chloroform; and slightly soluble in absolute ethanol and cold water.
2. **Lobelanine Hydrobromide (C₂₂H₂₅NO₂.HBr)**: The crystals do not give a sharp mp, but gets decomposed at 188°C.
3. **Lobelanine Nitrate (C₂₂H₂₅NO₂.HNO₃)**: The crystals obtained from dilute ethanol has mp 153-154°C.
4. It differs from **Lobeline** in lacking OH moiety; and therefore, does not react with nitrous acid nor with benzyl chloride.
5. It being a diketonic compound-forms a dioxime.

D. Lobelanidine

Biological Source It is same as that of lobelanine.

Chemical Structure

Lobelanidine

[2α(R*), 6α(S*)-Methyl-α, α'-diphenyl-2, 6-piperidine-diethanol (C₂₂H₂₉NO₂).

Isolation It is obtained as one of the fractions obtained from the column chromatography of the aqueous extract from step (3) under isolation of **lobeline**.

Characteristic Features

1. It is obtained as scales from ethanol with mp 150°C.
2. It distils unchanged in vacuo.
3. It is freely soluble in benzene, chloroform, acetone; slightly soluble in ether, petroleum ether; and almost insoluble in water.


Identification Tests

1. **Lobelanidine Hydrochloride ($C_{22}H_{29}NO_2 \cdot HCl$):** It is obtained as needles from ethanol having mp 135-138°C.
2. **Lobelanidine Hydrobromide ($C_{22}H_{29}NO_2 \cdot HBr$):** Its crystals have a mp 189°C.


Synthesis from Lobeline, Lobelanine and Lobelanidine First of all, **lobelanine** may be synthesized by the interaction of one molecule of glutaric dialdehyde, two moles of benzoyl acetic acid, and one mole of methylamine hydrochloride; allowing the reaction mixture to stand for 40 hours at 35°C and at pH 4.5. Thus, the resulting product **lobelanine** gives rise to:

- (a) **Lobeline:** When subjected to *partial reduction*, and
- (b) **Lobelanidine:** On being subjected to *complete reduction*.

All these reactions are summarized as given below.




Biosynthesis of Lobeline and Lobelanine The two above stated alkaloids, namely: **lobeline** and **lobelanine**, commonly found in the antiasthmatic medicinal plant *Lobelia inflata*, found to comprise of the piperidine rings with alternative C_6C_2 side-chains derived from phenylalanine *via* cinnamic acid. In fact, these alkaloids are formed as shown below wherein benzoylacetyl-CoA, an emerging intermediate in the β -oxidation of cinnamic acid helps to cater for the nucleophile engaged in the **Mannich reaction**. Thus, oxidation of the piperidine ring brings forth a new iminium species that can react further with a second mole of **benzoylacetyl-CoA**, again *via* **Mannich reaction**. Both **lobeline** and **lobelanine** are the resulting products obtained from further N-methylation and/or carbonyl reduction reactions.



E. Piperine

Biological Source It is obtained from the dried unripe fruit of *Piper nigrum* L. (Black Pepper), *Piper longum* L., *Piper retrofractum* Vahl. (*Piper officinarum* C.D.C.), and *Piper clusii* C.D.C.; and also in the root bark of *Piper geniculatum*. Sw. belonging to family *Piperaceae*.

Chemical Structure



Piperine

(E, E)-1-[5-(1, 3-Benzodioxol-5-yl)-1-oxo-2, 4-pentadienyl] piperidine ($C_{17}H_{19}NO_3$).

Isolation The dried unripe fruits are extracted with ethanol in a Soxhlet apparatus till extraction is complete. The solvent is evaporated under vacuo in a Rotary Thin Film Evaporator. The residue of the alcoholic extract is digested with dilute alkali to affect saponification, when piperine remains unaffected. The residue, thus obtained is decanted and washed with distilled water several times.

The resulting product is dissolved in hot ethanol and on cooling the crystalline **piperine** separates out.

Characteristic Features

1. Piperine is obtained as monoclinic prisms from alcohol having mp 130°C.
2. It is tasteless at first, but has a burning aftertaste.
3. Its dissociation constant pK (18°C) is 12.22.
4. **Solubility Profile:** 1 g of **piperine** dissolves in 15 ml ethanol, 1.7 ml chloroform, 36 ml ether; freely soluble in acetic acid and benzene; and almost insoluble in water (40 mg/L at 18°C), and petroleum ether.


Identification Tests

1. **Wagner's Reagent Test:** The addition of **Wagner's reagent** to an alcoholic solution of **piperine** gives rise to bluish needle like crystals having mp 145°C.
2. **Platinum Chloride Test [H₂PtCl₆]:** **Piperine** on treatment with platinum chloride solution (0.5% w/v) produces an instant orange red colouration, which upon standing gives needles of **piperine-H₂PtCl₆**.
3. **Piperine** reacts with a few drops of concentrated sulphuric acid yields a distinct red colouration.

Uses

1. It is used as an insecticide.
2. It is also employed extensively as condiment in food preparations.
3. It is used to give a 'pungent' taste to brandy.




Biosynthesis of Piperine In the **biosynthesis of piperine**, the piperidine ring forms part of a tertiary amide moiety which is incorporated *via* piperidine itself *i.e.*, the reduction product of Δ^1 -piperideine as shown under. Interestingly, the piperic acid residue, in obtained from a cinnamoyl-CoA precursor. The extension of chain is caused by virtue of acetate/malonate and ultimately combines as its CoA-ester with the previously obtained piperidine nucleus.



Biosynthesis of Piperine

7.2.4.2 Quinolizidine Alkaloids

The **quinolizidine alkaloids** comprise of **lupinine**, **lupanine** and **sparteine** which are responsible for the toxic properties are characterized by a quinolizidine skeleton. The bi-heterocyclic nucleus is closely related to the ornithine-derived pyrrolizidine system, but is believed to be formed from two molecules of lysine.




The aforesaid *three* alkaloids shall now be discussed individually in the pages that follows:

A. Lupinine

Synonyms *l*-Lupinine; (-)-Lupinine.

Biological Source The naturally occurring *l*-form is obtained from the seeds and herb of *Lupinus luteus* L. and other *Lupinus* species belonging to the natural order *Leguminosae*; and also found in *Anabasis aphylla* L. (*Chenopodiaceae*).

Chemical Structure



[1*R*-*trans*]-Octahydro-2*H*-quinolizidine-1-methanol (C₁₀H₁₉NO).

Isolation The isolation of **lupinine** from the seeds and herb of *Lupinus luteus* may be affected by the method evolved by Couch* (1934).

Characteristic Features

1. **Lupinine** is obtained as stout orthorhombic prisms from acetone having mp 68.5-69.2°C.
2. Its physical parameters are: bp₄ 160-164°C; bp₇₅₅ 269-270°C; [α]_D²⁶ - 25.9° (C = 3 in water); [α]_D²⁸ - 21° (C = 9.5 in ethanol);
3. It is soluble in water, ethanol, ether and chloroform.
4. It is a strong base.

Identification Tests

1. ***l*-Form Lupinine Hydrochloride Derivative (C₁₀H₂₀ClNO):** Its orthorhombic prisms have mp 208-213°C, and [α]_D-14°.


* Couch, J.F., *J. Am. Chem. Soc.*, **56**, 2434 (1934).

2. ***dl*-Form Lupinine:** The crystals obtained from acetone have mp 58.5-59.5°C.

B. Lupanine

Biological Source It is obtained from the herb of *Genista tinctoria* L. (*Fabaceae*) (**Dyer's Broom**).

Chemical Structure



Lupanine

(7 α , 7 α , 14 α , 14 β)-Dodecahydro-7, 14-methano-2H, 11H-dipyridiol [1, 2-a: 1', 2'-e] diazocin-11-one; (C₁₅H₂₄N₂O).

Isolation The racemic and optical isomers of **lupanine** have been duly isolated from various species of *Lupinus* (*Fabaceae/Leguminosae*) as stated below:

(\pm)-**Lupanine**—from white lupins;

***d*-Lupanine**—from blue lupins;

***l*-Lupanine**—from the natural racemic form;

Characteristic Features The physical parameters of the above *three* forms of **lupanine** are given below:

***dl*-Lupanine:** It is obtained as orthorhombic prisms obtained from acetone having mp 98-99°C; bp_{1.0} 185-195°C; It is soluble in ethanol, ether, chloroform and water; and insoluble in petroleum ether.

***d*-Lupanine (Synonym: 2-Oxosparteine):** It is obtained as syrup crystallizing difficultly in hygroscopic needles having mp 40-44°C; bp₃ 190-193°C; n_D²⁴ 1.5444; [α]_D²⁵ +84° (C = 4.8 in ethanol). It is found to be freely soluble in water, ethanol, ether and chloroform.

***l*-Lupanine (Synonym: Hydrorhombinine):** It is a viscous liquid having bp_{1.0} 186-188°C; [α]_D -61° in acetone.


Identification Test **Lupanine** forms the corresponding lupanine hydrochloride dihydrate (C₁₅H₂₄N₂O.HCl.2H₂O) which is obtained as rhombic crystals from water having mp 127°C (dry).

C. Sparteine

Synonyms *l*-Sparteine; Lupinidine.

Biological Sources It is obtained from yellow and black **lupin beans** *Lupinus luteus* L., and *Lupinus niger* Hort.; and also found in *Cytisus scoparius* (L.) Link. (*Fabaceae*) (**Scotch Broom**); *Anagyris foetida* L., belonging to natural order *Leguminosae*. Besides, it is also obtained from the roots of *Aconitum napellus* L. (*Ranunculaceae*) (**Aconite, Monkshood, Blue Rocket**); from the herbs of *Chelidonium majus* L. (*Papaveraceae*) (Celandine, Great Celandine, Nipplewort); from leaves of *Peumus boldus* Molina (*Monimiaceae*) (Boldo).

Chemical Structure



Sparteine

[7S-(7 α , 7 α , 14 α , 14 α β)]-Dodecahydro-7, 14-methano-2H, 6H-dipyrido [1, 2-a: 1', 2'-e] [1, 5] diazocine, (C₁₅H₂₆N₂).

Isolation It is isolated from yellow and black lupin beans by the method put forward by Clemo* (1949).

Characteristic Features

1. It is a viscous oily liquid having bp₈ 173°C.
2. It is volatile with steam.
3. Its physical parameters are: $[\alpha]_D^{21} - 16.4^\circ$ (C = 10 in absolute ethanol); n_D^{20} 1.5312; d_4^{20} 1.020; pK_1 at 20°C : 2.24; pK_2 :9.46; pH of 0.01 molar solution is 11.6.
4. **Solubility profile:** It is freely soluble in ethanol, ether and chloroform; and 1 g dissolves in 325 ml of water.

Identification Test


Sparteine Sulphate Pentahydrate (C₁₅H₂₆N₂.H₂SO₄.5H₂O): (Synonyms: Depasan; Tocosamine) It is obtained as columnar crystals which loses water of crystallization at 100°C turning brown and ultimately gets decomposed at 136°C. The pH of a 0.05 molar solution is 3.3. It is practically insoluble in ether and chloroform, and 1 g dissolves in 1.1 ml of water, 3 ml of ethanol.

Uses

1. It is used mostly as an oxytocic.
2. It is employed as a cardiac depressant, cathartic, diuretic and for stimulating uterine contractions.
3. Sparteine is used occasionally as a quinidine substitute in stubborn cases of atrial fibrillation.

Biosynthesis of Lupinine, Lupanine and Sparteine Experimental evidence reveals lysine to be incorporated into **lupinine** *via* **cadaverine**; however, the intermediate related to **homospermidine** is excluded. It has been observed that Δ^1 -piperideine happens to be an important intermediate after **cadaverine**. Thus, the proposed pathway given below suggests coupling of two such molecules. In fact, the two tautomers of Δ^1 -piperideine, as N-analogues of corresponding carbonyl compounds, are in a position to couple by an aldol-type mechanism. In reality, this coupling takes place in solution at physiological pHs, although the *stereospecific coupling* as shown in the proposed pathway shall evidently require the participation of suitable enzymes. After coupling, the imine system gets hydrolyzed, the resulting primary amine function undergoes oxidation, and ultimately the formation of the **quinolizidine nucleus** is accomplished by Schiff base formation. Thus, **lupinine** is then synthesized by two further reductive steps. Hence, the pathway to **sparteine** and **lupanine** eventually requires participation of another molecule of **cadaverine** or Δ^1 -piperideine.


* Clemo *et al. J. Chem. Soc.* 6.63, (1949)



Biosynthesis of Lupinine, Lupanine and Sparteine

7.2.4.3 Indolizidine Alkaloids

The **indolizidine alkaloids** are usually characterized by the presence of a 5-membered and a 6-membered cyclic ring with a N-atom fused in them as shown below:




The two typical examples of **indolizidine alkaloids** are, namely: **castanospermine** and **swainsonine**, which shall be discussed hereunder:

A. Castanospermine

Biological Source It is obtained from the seeds of the **Australian leguminous tree** *Castanospermum australe* A. Cunn. (*Leguminosae*) (**Moreton Bay Chestnut**).

Chemical Structure



[1S-(1 α , 6 β , 7 α , 8 β , 8a β)]-Octahydro-1, 6, 7, 8-indolizinetetrol; (C₈H₁₅NO₄). It is a polyhydroxy alkaloid.

Isolation The isolation of the naturally occurring (+)-form of **castanospermine** from the seeds of the Australian leguminous tree has been duly accomplished.*

Characteristic Features

1. The crystals obtained from aqueous ethanol have mp 212-215°C (decomposed).
2. Its specific optical rotation is $[\alpha]_D^{25} + 79.7^\circ$ (C = 0.93 in water); and dissociation constant pK 6.09.


Uses Its has demonstrated activity against the AIDS virus HIV, by virtue of their ability to inhibit ghyosidase enzymes involved in glycoprotein biosynthesis. However, the glycoprotein coating seems to be vital for the proliferation of the AIDS virus.

B. Swainsonine

Biological Source It is obtained from the plant *Swainsona canescens* (*Leguminosae/Fabaceae*).

* Hohenschutz *et al. Phytochemistry*, **20**, 811 (1981).

Chemical Structure



[1S-(1 α , 6 β , 7 α , 8 β , 8 $\alpha\beta$)-Octahydro-1, 2, 8-indolizinetriol; (C₈H₁₅NO₃).

Biosynthesis of Castanospermine and Swainsonine These two alkaloids are regarded as a hybrid between the **pyrrolizidine** and **quinolizidine** alkaloids. It is, however, pertinent to mention here that these two alkaloids though are derived from lysine, yet their origin entirely deviates from the usual and common lysine-derived moieties in that **L-pipecolic acid** is found to be an intermediate in the pathway. In fact, there are *two* established routes known to the formation of pipecolic acid in nature, as shown below; wherein the point of difference solely based on whether the N-atom is taken-up either from the α - or the ϵ -amino portion of lysine.

In short, for the **indolizidine-alkaloid biosynthesis** the following salient features may be observed, namely:

- Pipecolic acid is produced *via* the aldehyde and Schiff base by retaining the N-atom from the α -amino function,
- Indolizidine nucleus is formed subsequently by incorporating a C₂-acetate unit through simple reactions,
- The resulting compound leads to the formation of *castanospermine* through a sequential hydroxylation reactions,
- Also a branch point compound results into the formation of **swainsonine** that essentially possess the opposite configuration at the ring fusion.


7.2.5 Alkaloids Derived from Nicotinic Acid

The alkaloids derived from the **nicotinic acid** are commonly known as the '**Pyridine Alkaloids**'.

In general, the alkaloids found in tobacco (*Nicotiana tabacum*, *Solanaceae*) include a variety of alkaloids, such as: **nicotine**, **anabasine**, and **niacin (Vitamin B₃, nicotinic acid)**. Interestingly, the '*pyridine unit*' has its origins in **vitamin B₃ (nicotinic acid)**; whereas, a combination of a pyridine ring with a pyrrolidine ring gives rise to **nicotine**, or a combination of a pyridine ring with a piperidine unit forms **anabasine**.

7.2.5.1 Pyridine Alkaloids

The *three* above mentioned **pyridine alkaloids**, *viz.*, **nicotine**, **anabasine** and **niacin**, shall now be discussed individually in the sections that follows:




Biosynthesis of Castanospermine and Swainsonine

A. Nicotine

Synonyms Nicolan; Nicabate; Nicoderm; Nicotell TTS; Nicopatch; Nicotinell; Habitrol; Tabazur.

Biological Sources It is obtained from the dried leaves of *Nicotiana tabacum* Linn., (*Solanaceae*) (**Virginia Tobacco; Tobacco**); sprouts of *Asclepias syriaca* L. (*Asclepiadaceae*) (**Common Milkweed**); dried leaves and roots of *Datura metel* L. (*Solanaceae*) (**Unmatal, Metel, Hindu Datura**); leaves of *Duboisia myoporoides* R. Br. (*Solanaceae*) (**Corkwood Tree, Pituri**); fresh forage of *Equisetum arvense* L. (*Equisetaceae*) (Field Horsetail); herbs of *Equisetum hyemale* L. (*Equisetaceae*) (**Shavegrass, Great Scouring Rush**); leaves of *Erythroxylum coca* Lam., (*Erythroxylaceae*) (**Coca**); fruits and leaves of *Nicotiana glauca* R. Grah. (*Solanaceae*) (**Tree Tobacco**); and leaves of *Nicotiana rustica* Linn. (*Solanaceae*)—present upto 2-8%.

Chemical Structure



(S)-3-(1-Methyl-2-pyrrolidixyl) pyridine; (C₁₀H₁₄N₂).

Preparation Commercial **nicotine** is entirely a byproduct of the tobacco industry; and the extraction from *N. tabacum* has been described in literature.*

Characteristic Features

1. It is a colourless to pale yellow oily liquid, very hygroscopic in nature, and turns brown on exposure to air and light.
2. It has an inherent acrid burning taste.
3. It develops the odour of pyridine.
4. It has a bp₇₄₅ 247°C with partial decomposition; and bp₁₇ 123-125°C.
5. It is a steam-volatile product.
6. Its physical parameters are; n_D²⁰ 1.5282 ; d₄²⁰ 1.0097 ; [α]_D²⁰ -169° ; pK₁ (15°) 6.16 and pK₂ 10.96; and pH of 0.05 M solution 10.2.
7. It readily forms salts with almost any acid; and double salts with many metals and acids.
8. **Solubility:** It is miscible with water below 60 °C; and on mixing nicotine with water the volume contracts. However, it is found to be very soluble in chloroform, ethanol, ether, petroleum ether and kerosene oils.

Identification Tests

1. **Nicotine Hydrochloride (C₁₀H₁₄N₂.HCl):** It is obtained as deliquescent crystals having specific rotation [α]_D²⁰ +104° (p = 10).

* Gattermann, Wieland, 'Laboratory Methods of Organic Chemistry' New York, 24th edn., (1937).

- Nicotine Dihydrochloride ($C_{10}H_{14}N_2 \cdot 2HCl$):** The deliquescent crystals are extremely soluble in water and ethanol.
- Nicotine Sulphate [$(C_{10}H_{14}N_2)_2 \cdot H_2SO_4$] (Synonym: Nicotine neutral sulphate):** It is obtained as hexagonal tablets having optical rotation $[\alpha]_D^{20} + 88^\circ$ ($p = 70$); and is soluble in water and ethanol.
- Nicotine Bitartrate ($C_{10}H_{14}N_2 \cdot 2C_4H_6O_6$) (Synonym: Nicotine Tartrate):** It is obtained as the dihydrate, crystals having mp $90^\circ C$; $[\alpha]_D^{20} + 26^\circ$ ($C = 10$); and is found to be very soluble in ethanol and water.
- Nicotine Zinc Chloride Double Salt Monohydrate ($C_{10}H_{16}Cl_4N_2Zn \cdot H_2O$):** It is very soluble in water; and sparingly soluble in ether and ethanol.
- Nicotine Salicylate ($C_{17}H_{20}N_2O_3$): (Synonym: Eudermol):** It is obtained as hexagonal plates having mp $118^\circ C$; $[\alpha]_D^{20} + 13^\circ$ ($C = 9$); and is found to be freely soluble in ethanol and water.

Uses


- It is used extensively as an insecticide and fumigant.
- It finds its application as a '**contact poison**' in the form of soap *i.e.*, as its oleate, laurate and naphthenate salts.
- It is also used as a '**stomach poison**' in combination with bentonite.
- One of its recent applications nicotine is employed as chewable tablets of lozenges for the treatment of smoking withdrawal syndrome.
- It possesses a unique action on the autonomic ganglia which it first stimulates and subsequently depresses ultimately leading to *paralysis*.

B. Anabasine

Synonym Neonicotine;

Biological Sources It is obtained from the leaves of *Duboisia myoporoides* R. Br. (*Solanaceae*) (**Corkwood tree; Pituri**); fruits and leaves of *Nicotiana glauca* R. Grah. (*Solanaceae*) (**Tree Tobacco**—claimed to be the richest source of **anabasine** (1.2%); leaves of *Nicotiana tabacum* L. (*Solanaceae*) (**Tobacco, Tabac, Virginia Tobacco**); and also the leaves of *Anabasis aphylla* L. (*Chenopodiaceae*).

Chemical Structure 3-(2-Piperidixyl) pyridine; ($C_{10}H_{14}N_2$).



Isolation Anabasine is extracted on a large scale in Russia; and the industrial extraction processes have been reported by Sadykov and Timbekov* (1956).

Characteristic Features

1. It is a liquid freezing at 9°C; and boiling at 270-272°C; bp₁₄ 145-147°C; bp₂ 105°C.
2. Its physical parameters are: d₄²⁰ 1.0455; n_D²⁰ 1.5430; and [α]_D²⁰ - 83.1°.
3. It is soluble in most organic solvents and water.

Identification Tests

1. Being a secondary amine it can form a nitroso derivative.
2. **Anabasine Hydrochloride:** Its specific optical rotation is [α]_D + 16.5° (C = 10 in water).

Uses It is invariably employed as an effective insecticide.

C. Niacin

Synonyms Nicotinic Acid; Pellagra Preventive Factor (or P.P. Factor); Vitamin B₃; Akotin; Daskil; Nicacid; Niacor; Nicangin; Nicobid; Nicolar; Niconacid; Nico-Span; Wampocap. The term 'niacin' has also been applied to nicotinamide.


Biological Sources It is widely distributed in nature; and appreciable quantities are found in fish, yeast, liver, and cereal grains.

Chemical Structure



3-Pyridinecarboxylic acid; (C₆H₅NO₂).

Preparation It may be prepared by the **oxidation of nicotine**** as given below.



Characteristic Features

1. It is obtained as needles from ethanol and water having mp 236.6°C.


* Sadykov and Timbekov, *J. Appl. Chem.*, USSR, **29**, 148 (1956).

** MeElvain, S.M., *Org. Synth. Coll. Vol I*, 385 (1941).

2. It sublimes without any decomposition.
3. It is a nonhygroscopic substance and fairly stable in air.
4. It shows uv_{\max} : 263 nm; and pH 2-7 of a saturated solution.
5. *Solubility*: 1 g dissolves in 60 ml water; freely soluble in boiling water and ethanol; soluble in propylene glycol; and insoluble in ether.

Identification Test

1. **Niacin Sodium Salt Sesquihydrate ($C_6H_4NNaO_2 \cdot 1\frac{1}{2}H_2O$) (Synonym: Direktan)**: It is obtained as either white crystals or crystalline powder, which is stable in air. Its solubility profile are as follows: 1 g dissolves in ~ 1.4 ml of water, in 60 ml ethanol, in 10 ml glycerol; and insoluble in ether. The pH of aqueous solution is ~ 7.
2. **N-Oxide Derivative (Oxiniacetic Acid)**: It is obtained as needles mp 254-255°C (dec.) and uv_{\max} (0.1 N. H_2SO_4):220, 260 nm (ϵ 22400, 10200).




Uses


1. It is used as antihyperlipoproteinemic agent.
2. It is a vital vitamin (enzyme cofactor).

Biosynthesis of Nicotine, Anabasine and Niacin Interestingly, plants such as *Nicotiana* make use of an altogether different pathway employing glyceraldehyde 3-phosphate and L-aspartic acid precursors as given under. Thus, the dibasic acid **quinonilic acid** features in the aforesaid pathway which upon decarboxylation gives rise to **nicotinic acid**.

It is pertinent to mention here that the formation of **nicotine** caused by a pyrrolidine ring derived from ornithine, quite possibly as the N-methyl- Δ^1 -pyrrolinium cation gets hooked on to the pyridine ring present in nicotinic acid thereby displacing the carboxyl function during the course of reactions as depicted in (B). Further, a **dihydronicotinic acid** intermediate is most likely to be engaged permitting decarboxylation to the enamine 1, 2-dihydropyridine. It, therefore, allows an aldol-type interaction with the N-methylpyrrolinium cation, and ultimately undergoes dehydrogenation of the dihydropyridine ring reversed to a pyridine ring yields *nicotine*. In this fashion, **nornicotine** is derived by the oxidative demethylation of nicotine.


Finally, **anabasine** is generated from nicotinic acid and lysine *via* the Δ^1 -piperidinium cation in an effectively analogous sequence as shown in (C) below.





7.2.6 Alkaloids Derived from Ornithine

A non-protein amino acid, *L-ornithine*, usually constitutes an integral part of the 'urea-cycle' in animals, wherein it is eventually produced from *L-arginine* in a reaction sequence catalyzed by the enzyme *arginase* as given below:



Evidently, *L-ornithine* possesses δ - and α -amino moieties, and the N-atom from the former moiety which is eventually incorporated into the alkaloid structures along with the C-chain, except for the carboxyl function. Thus, the *L-ornithine* exclusively provides a C₄N building block to the alkaloid structure; not only as a **pyrrolidine ring system**, but also as a part of the **tropane alkaloids**. Nevertheless, the reactions of ornithine are fairly comparable to those of lysine, which in turn provides a C₅N unit bearing its ϵ -amino moiety.

The various **alkaloids** derived from ornithine may be categorized into *three* heads, namely:

- (i) Pyrrolidine Alkaloids,
- (ii) Tropane Alkaloids, and
- (iii) Pyrrolizidine Alkaloids.

The above categories of **alkaloids** shall be discussed separately hereunder.


7.2.6.1 Pyrrolidine Alkaloids

The *three* glaring examples of **pyrrolidine alkaloids** are, namely: **hygrine**, **cuscohygrine** and **stachydrine**, which would be discussed below:

A. Hygrine

Biological Sources It occurs in the leaves of *Erythroxylon coca* Lam., (*Erythroxylaceae*) (Coca); and the roots of *Withania somniferum* (L.) Dunal. (*Solanaceae*) (**Ashwagandha**).

Chemical Structure



Hygrine

(R)-1-(1-Methyl-2-pyrrolidinyl)-2-propanone; (C₈H₁₅NO).

Characteristic Features

1. It is a liquid having bp₁₁ 76.5°C; bp₁₄ 81°C; n_D²⁰ 1.4555.
2. It is soluble in dilute mineral acids, chloroform and ethanol; and slightly soluble in water.


Identification Test It forms oxime readily (C₈H₁₆N₂O) which is obtained as crystals from ether having mp 123-124°C.

Uses The drug is broadly used as a sedative, hypnotic laxative and diuretic.

B. Cuscohygrine

Synonyms Cuskhygrine; Bellaradine.

Biological Sources It is obtained from the roots of *Atropa belladonna* L. (*Solanaceae*) (**Belladonna**, **Deadly Nightshade**); roots of *Datura innoxia* Mill. (*Solanaceae*) (**Thorn Apple**) upto 5-30%; seeds of *Datura metal* L. (*Solanaceae*) (**Unmatal**, **Metel**, **Hindu Datura**); leaves of *Hyocyamus niger* L. (*Solanaceae*) (**Henbane**, **Henblain**, **Jusquaim**); herb of *Mandragora officinarum* L. (*Solanaceae*) (**Mandrake**, **Loveapple**); rhizome of *Scopolia carniolica* Jacq. (*Solanaceae*) (**Scopolia**); and the roots of *Withania somniferum* (L.) Dunal (*Solanaceae*) (**Ashwagandha**).

Chemical Structure

Cuscohygrine

1, 3-Bis (1-methyl-2-pyrrolidinyl)-2-propanone; ($C_{13}H_{24}N_2O$).

Isolation It is isolated from the naturally occurring plant sources by standard method.*

Characteristic Features

1. It is a oily liquid having bp_{23} 169-170°C; bp_{14} 152°C; bp_2 118-125°C; d_4^{20} 0.9733; n_D^{20} 1.4832.
2. It is found to be miscible with water; and freely soluble in ethanol, ether, and benzene.


Identification Tests

1. Cuscohygrine Hemiheptahydrate: Its needles have mp 40°C.
2. Cuscohygrine Hydrobromide ($C_{13}H_{24}N_2O \cdot 2HBr$): It forms prisms from ethanol having mp 234°C.

C. Stachydrine

Synonyms Methyl hygrate betaine; Hygric acid methylbetaine.

Biological Sources It is obtained from the forage of *Achillea millefolium* L. (*Asteraceae*) (**Yarrow**); flowers of *Chrysanthemum cinerarifolium* (**Trevir.**) Vis. (*Asteraceae*) (**Pyrethrum, Dalmatian Insect Flower**); branches of *Lagochilus inebrians* Bunge (*Lamiaceae*) (**Intoxicating Mint**); dry plant of, *Leonurus cardiaca* (L.) (*Lamiaceae*) (**Motherwort**); the 'betaine fraction' of alfalfa *Medicago sativa* L. (*Fabiaceae*) (**Alfalfa**) (0.785%); and herbage of *Stachys officinalis* (L.) Trevisan (*Lamiaceae*) (**Betony**).

Chemical Structure

Stachydrine

(S)-2-Carboxy-1, 1-dimethylpyrrolidinium inner salt; ($C_7H_{13}NO_2$).

Isolation It has been isolated by reported method by Schulze** and Jahns.***

* Liebermann, *Ber.*, **22**, 679 (1898)

** Schulze, *Ber.* **26**, 939 (1893);

*** Jahns, *Ber.* **29**, 2065 (1896);

Characteristic Features


1. It is obtained as monohydrate deliquescent crystals having mp 235°C (anhydrous).
2. It is sweetish in taste.
3. It is soluble in water, dilute mineral acids and ethanol;
4. It isomerizes at the mp to methyl hydrate.

Identification Tests

1. **Stachydrine Hydrochloride ($C_7H_{17}NO_2 \cdot HCl$):** Its large prisms are obtained from absolute ethanol which gets decomposed at 235°C. It is very soluble in water and soluble in 13 parts of ethanol.
2. **Stachydrine Acid Oxalate ($C_7H_{13}NO_2 \cdot C_2H_2O_4$):** Its needles have mp 106°C. It is practically insoluble in absolute ethanol.
3. **Stachydrine Aurichloride ($C_7H_{13}NO_2 \cdot HAuCl_4$):** Its yellow needles have mp 225°C (rapid heating). It is quite soluble in hot water, but practically insoluble in cold water.
4. **Stachydrine Platinichloride Tetrahydrate ($(C_7H_{17}NO_2)_2 \cdot H_2PtCl_6 \cdot 4H_2O$):** It is obtained as orange crystals decomposing at 210-220°C (rapid heating). It is found to be very soluble in dilute ethanol and water. It may also be obtained with two moles of water of crystallization.

7.2.6.2 Tropane Alkaloids

Tropane is a bicyclic compound obtained by the condensation of one mole each of **pyrrolidine** and **piperidine** as shown below.



Tropane is regarded as the principle base of a plethora of alkaloids obtained from various members of the natural order, viz., *Solanaceae*, *Erythroxylaceae*, *Convolvulaceae*, and *Dioscoreaceae*. It essentially consists of a 7-carbon bicyclic ring with a N-atom strategically bridged between C-1 and C-5 and providing a C_7N unit. It is, however, pertinent to mention here that the tropane base contains two chiral centres (*i.e.*, asymmetric C-atoms), namely: C-1 and C-5, but surprisingly it does not exhibit any optical activity (an exception) by virtue of the fact that *intramolecular compensation* prevails. It happens to be a *meso*-compound.


A few important members belonging to the **tropane alkaloids** are, namely: **atropine**, **cocaine**, **cinnamoyl cocaine**, **ecgonine** and **hyoscyamine**. These alkaloids shall now be treated individually in the sections that follows:

A. Atropine

Synonyms Tropine tropate; *dl*-Hyoscyamine; *dl*-Tropyl Tropate; Tropic acid ester with Tropine.

Biological Sources It is obtained from the roots and leaves of *Atropa belladonna* Linn. (*Solanaceae*) (**Belladonna**); and the seeds and leaves of *Datura stramonium* Linn. (*Syn.*: *Datura tatula* Linn.) (*Solanaceae*) (**Jimson Weed, Thorn Apple, Stramonium**), besides other species of *Solanaceae*, such as: *D. metel* Linn.; *D. innoxia* Mill., *D. alba* Nees.; and *D. fastuosa* Linn.

Chemical Structure



1 α H, 5 α H-Tropan-3 α -ol (\pm)-tropate (ester); (C₁₇H₂₃NO₃).

Characteristic Features

1. **Atropine** is obtained as long orthorhombic prisms from acetone having mp 114-116°C.
2. It usually sublimes in high vacuum at 93-110°C.
3. It has a dissociation constant pK 4.35; and the pH of a 0.0015 molar solution is 10.0.
4. **Solubility:** 1 g dissolves in 455 ml water; 90 ml water at 80°C; 2 ml ethanol; 1.2 ml ethanol at 60°C; 27 ml glycerol; 25 ml ether, 1 ml chloroform; and in benzene.

Identification Tests It forms various types of salts, namely:

1. **Atropine Hydrochloride (C₁₇H₂₃NO₃.CH₃NO₃):** The granular crystals have mp 165°C. It is soluble in water and ethanol. The pH of 0.05 molar solution is 5.8.
2. **Atropine Methyl Bromide (C₁₇H₂₃NO₃.CH₃Br) (Tropin):** Its crystals have mp 222-223°C. It is soluble in 1 part of water, slightly soluble in ethanol, and practically insoluble in ether and chloroform.
3. **Atropine Methylnitrate (C₁₇H₂₃NO₃.CH₃NO₃) (Methylatropine nitrate, Eumydrin, Metropine, Harvatrate, Metanite, Ekomin):** Its crystals have mp 163°C. It is found to be freely soluble in water or ethanol; and very slightly soluble in chloroform and ether.
4. **Atropine Sulphate Monohydrate [(C₁₇H₂₃NO₃)₂.H₂SO₄.H₂O] (Atropisol):** It is obtained as either crystals or powder with mp 190-194°C. It is *inactive* optically. It has a very bitter taste. It shows pH ~ 5.4. Its bitterness is threshold 1:10,000. It is found to be incompatible with a host of substances, such as, tannin, alkalies, salts of gold and mercury, borax, bromides, iodides, benzoates and vegetable decoctions or infusions.

Its solubility profile is: 1 g dissolves in 0.4 ml water; in 5 ml cold and 2.5 ml boiling ethanol; in 2.5 ml glycerol; 420 ml chloroform and 3000 ml ether.

Uses


1. It is used in preanaesthetic medication.
2. It is employed as an anticholinergic agent.
3. It is also used as a mydriatic.
4. It is employed as an antidote in opium and chloral hydrate poisoning.
5. It is frequently employed to minimize spasm in cases of intestinal gripping caused due to strong purgatives.
6. It also find its applications to reduce such secretions as: saliva, sweat, and gastric juice.

B. Cocaine

Synonyms 2β-Carbomethoxy-3β-benzoyloxytropane; *l*-Cocaine; β-Cocaine; Benzoylmethylecgonine; Ecgonine methyl ester benzoate.

Biological Sources It is obtained from the leaves of *Erythroxylon coca* Lam. and other species of *Erythroxylon*, (*Erythroxylaceae*); and leaves of *Erythroxylon truxillense* Rusby (*Erythroxylaceae*).

Chemical Structure



[1R-(exo, exo)]-3-(Benzoyloxy)-8-methyl-8-azabicyclo[3, 2, 1] octane-2-carboxylic acid methyl ester; (C₁₇H₂₁NO₄).

Isolation **Cocaine** is extracted from the plant by digestion either with sodium carbonate solution or with lime water and by subsequent solvent extraction using petroleum ether (bp 160-180°C; or 200-220°C). The combined petroleum ether extract is shaken up with dilute HCl. The solution of hydrochloride thus obtained is concentrated carefully in a thin-film evaporator. In case, the leaves are rich in cocaine content, as in the Peruvian coca leaves, a major portion of cocaine gets separated as crystals.

Characteristic Features

1. **Cocaine** is obtained as the monoclinic tablets from ethanol having mp 98°C.
2. It usually becomes volatile above 90°C; however, the resulting sublimate is *not* crystalline in nature.

- Its physical parameters are as follows; bp_{0.1}, 187-188°C; $[\alpha]_D^{18} - 35^\circ$ (50% ethanol); $[\alpha]_D^{20} - 16^\circ$ (C = 4 in chloroform); pKa (15°C) 8.61 and pKb (15°C) 5.59.
- Solubility Profile:** 1 g of cocaine dissolves in 600 ml of water; 270 ml of water at 80°C; 6.5 ml of ethanol; 0.7 ml of chloroform; 3.5 ml of ether; 12 ml of turpentine; 12 ml of pure olive oil; and 30-50 ml of liquid petrolatum. It is also soluble in acetone, carbon disulphide and ethyl acetate.

Identification Tests

- Cocaine Permanganate:** The addition of a drop of saturated solution of KMnO₄ to a solution of cocaine prepared in a saturated solution of alum gives rise to a violet crystalline precipitate due to the formation of cocaine permanganate. It clearly shows characteristic violet aggregates of plates when examined under the microscope.
- Cocaine Hydrochloride (C₁₇H₂₁NO₄·HCl) (Cocaine Muriate):** It is obtained as granules, crystals, or powder. It has a slightly bitter taste and usually numbs lips and tongue. Its physical characteristics are: mp ~ 195°C; $[\alpha]_D - 72^\circ$ (C = 2 in aqueous solution); 1 g dissolves in 0.4 ml of water; 3.2 ml cold and 2 ml hot alcohol; 12.5 ml chloroform. It is also soluble in glycerol and acetone; and insoluble in ether or oils.
- Cocaine Nitrate Dihydrate (C₁₇H₂₂N₂O₇·2H₂O):** Its crystals have mp 58-63°C. It is freely soluble in water or ethanol; and slightly soluble in ether.
- Cocaine Sulphate (C₁₇H₂₁NO₄·H₂SO₄):** It is obtained as white, crystalline or granular powder, which is soluble in ethanol and water.

Uses


- It is used as a local anaesthetic as it causes numbness.
- Its main action is a CNS-stimulant and, therefore, categorized as 'narcotic drugs'. It is a highly habit-forming drug.

C. Cinnamoyl Cocaine

Synonyms Ecgonine Methyl Ester; Cinnamoylcocaine; Cinnamoyl-methylecgonine; Ecgonine Cinnamate Methyl Ester.

Biological Source It is obtained from the leaves of *Erythroxylon coca* Lann. (*Erythroxylaceae*), particularly from the Javanese leaves.

Chemical Structure



Cinnamoylcocaine [(E)-Form]

[1R-(exo, exo)]-8-Methyl-3-[(1-oxo-3-phenyl-2-propenyl)oxy]-8-azabicyclol [3, 2, 1] octane-2-carboxylic acid methyl ester; (C₁₉H₂₃NO₄).

Isolation Instead of the Peruvian leaves the Java leaves of *E. coca* are treated in the same manner and fashion as described under cocaine earlier (section 'B'). It has been observed that the mixed hydrochlorides mostly comprise of cinnamoyl cocaine which gets separated as fine needles.

Characteristic Features

1. It is obtained as fine needles having mp 121°C.
2. Its specific optical rotation is $[\alpha]_D - 4.7^\circ$ (chloroform).
3. It is freely soluble in ether, ethanol and chloroform; and almost insoluble in water.


Identification Tests

1. It reduces an acidic solution of KMnO₄ in cold *i.e.*, at ambient temperature, which helps to detect the presence of this alkaloid in an admixture with cocaine.
2. It undergoes hydrolysis when warmed with HCl to yield ***l*-ecgonine**, cinnamic acid and methanol.

D. Ecgonine

Biological Source It is also obtained from the leaves of *Erythroxylum coca* Lam. (*Erythroxylaceae*) (**Coca**) as its *l*-form.

Chemical Structure



Ecgonine

[1R-(exo, exo)]-3-Hydroxy-8-methyl-8-azabicyclol [3, 2, 1] octane-2-carboxylic acid; (C₉H₁₅NO₃).

It is the principal part of the **cocaine** molecule.

Isolation Ecgonine may be obtained by the hydrolysis of **cocaine** as given below:



Characteristic Features

1. The *l*-form **ecgonine monohydrate** is obtained as triboluminescent, monoclinic prisms from ethanol having mp 198°C (anhydrous substance gets decomposed at 205°C).
2. Its specific optical rotation $[\alpha]_D^{15} - 45^\circ$ (C = 5); dissociation constants are: pKa 11.11, and pKb 11.22.
3. **Solubility Profile:** 1 g dissolves in 5 ml water, 67 ml ethanol, 20 ml ethanol, 75 ml ethyl acetate; sparingly soluble in ether, acetone, benzene, chloroform and petroleum ether.

Identification Tests


1. **Ecgonine Hydrochloride (C₉H₁₅NO₃.HCl):** It is obtained as the triclinic plates obtained from water having mp 246°C; $[\alpha]_D^{15} - 59^\circ$ (C = 10); soluble in water and slightly in ethanol.
2. ***dl*-Ecgonine Trihydrate:** It is obtained as plates from 90% ethanol having mp 93-118°C (anhydrous substance gets decomposed at 212°C).

Uses It is mostly used as a topical anaesthetic.

E. Hyoscyamine

Synonyms *l*-Tropine Tropate; Daturine; Duboisine; *l*-Hyoscyamine; Cystospaz; Levsin; *l*-Tropic acid ester with Tropine; 3 α -Tropanyl S-($-$)-Tropate.

Biological Sources It is obtained from the roots and leaves of *Atropa bella-dona* L. (*Solanaceae*) (0.21%) (**Thorn Apple**); fruits, roots and leaves of *Datura metel* L. (*Solanaceae*) (**Unmatal, Metel, Hindu Datura**); leaves and seeds of *Datura stramonium* L. (*Solanaceae*) (**Jimson Weed, Thorn Apple, Stramonium**); root bark of *Duboisia myoporoides* R. Br. (*Solanaceae*) (**Pituri, Corkwood Tree**); young plants of *Hyoscyamus niger* L. (*Solanaceae*) (**Henbane, Henblain Jusquaim**); seeds of *Lactuca virosa* L. (*Asteraceae*) (**Bitter Lettuce, Wild Lettuce**); and the herb *Mandragora officinarum* L. (*Solanaceae*) (**Mandrake, Loveapple**).

Chemical Structure

Hyoscyamine

1 α H, 5 α H-Tropan-3 α -ol ($-$)-tropate (ester); (C₁₇H₂₃NO₃).

Isolation **Hyoscyamine** may be isolated from the Belladonna leaves by adopting the following steps sequentially:

1. The finely powdered and sieved **Belladonna leaves** is extracted with 95% (v/v) ethanol in a Soxhlet Apparatus till no more alkaloids come out from the marc. The ethanolic extract is concentrated to a syrupy residue under vacuo and subsequently treated with dilute HCl. The resinous matter is separated by filtration and the resulting solution is further purified by shaking out with petroleum ether (40-60°C) several times.
2. The purified acidic solution thus obtained is made alkaline with ammonia solution (dilute) carefully and extracted with chloroform successively. The combined chloroform layer is once

again shaken with dilute HCl, and the acidic solution made alkaline with dilute ammonia solution and extracted with chloroform successively.

- The combined chloroform layer is removed by distillation under reduced pressure. The crude alkaloids thus obtained is neutralized with oxalic acid. The oxalates of atropine and hyoscyamine may be separated by fractional crystallization from acetone and ether wherein the **hyoscyamine oxalate** being more soluble gets separated as the second crop.

Characteristic Features

- Hyoscyamine** is obtained as silky tetragonal needles from evaporating ethanol having mp 108.5°C.
- The physical parameters are: $[\alpha]_D^{20} - 21^\circ$ (ethanol); and dissociation constant K at 19° is 1.9×10^{-12} .
- Solubility Profile:** 1 g dissolves in 281 ml water (pH 9.5), 69 ml ether, 150 ml benzene, and 1 ml chloroform. It is freely soluble in dilute mineral acids and ethanol.

Identification Tests The various identification tests for **hyoscyamine** are, namely:

- Gerrard Reaction: Hyoscyamine** (and also **atropine**) responds to the **Gerrard Reaction** wherein about 5-10 mg of it reacts with mercuric chloride solution (2% w/v) in 50% ethanol to give rise to an instant red colouration without warming.
- Schaer's Reagent:** A few mg of **hyoscyamine** when made to react with a few drops of the **Schaer's Reagent** *i.e.*, 1 volume of 30% H₂O₂ mixed with 10 volumes of concentrated sulphuric acid, produces a distinct green colouration.
- Vitali-Morin Colour Reaction:** A few mg of **hyoscyamine** (and also atropine) is treated with about 0.2 ml of fuming HNO₃, evaporated to dryness on the water-bath. To the residue is then added 0.5 ml of a 3% (w/v) solution of KOH in methanol, it gives a bright purple colouration, that changes to red and finally fades to colourless.

Note: (a) The 3% solution of KOH must be freshly prepared.

(b) The reaction is very sensitive *i.e.*, upto 0.0001 mg of any of the alkaloids *viz.*, strychnine, apomorphine, veratrine, physostigmine etc. give a positive test.

- para-Dimethylaminobenzaldehyde Reagent:** [Prepared by dissolving 2 g of *p*-Dimethylaminobenzaldehyde in 6 g of H₂SO₄ to which 0.4 ml of water is added previously]. Add to 5-10 mg of **hyoscyamine** in an evaporating dish 2-3 drops of this reagent and heat on a boiling water-bath for several minutes. A distinct red colouration is produced that ultimately gets changed to permanent cherry red upon cooling.
- Hyoscyamine Hydrobromide (C₁₇H₂₃NO₃.HBr):** It is obtained as deliquescent crystals having mp 152°C; very soluble in water; 1 g dissolves in 3 ml ethanol; 1.2 ml chloroform and 2260 ml ether.
- Hyoscyamine Hydrochloride (C₁₇H₂₃NO₃.HCl):** The crystals have mp 149-151°C; and freely soluble in water and ethanol.
- Hyoscyamine Methyl Bromide (C₁₇H₂₃NO₃.CH₃Br) (N-Methylhyo-scyaminium bromide):** The crystals have mp 210-212°C; and freely soluble in water, dilute ethanol; and slightly soluble in absolute ethanol.

8. **Hyoscyamine Sulphate Dihydrate** [(C₁₇H₂₃NO₃)₂·H₂SO₄·2H₂O] (**Egacene, Peptard, Egazil Duretter**): It is obtained as needles from ethanol having mp 206°C (when dry); [α]_D¹⁵ – 29° (C = 2); pH 5.3 (1 in 100); 1 g dissolves in 0.5 ml water and about 5.0 ml ethanol; and very slightly soluble in ether and chloroform.

Uses

1. It is mostly employed as an anticholinergic drug.
2. It exerts relaxation of bronchial and intestinal smooth muscles (*i.e.*, antispasmodic action).
3. It also inhibits contraction of the iris muscle of the eye to produce mydriasis.
4. It decreases significantly decreases the sweat gland and salivary gland secretions.

Biosynthesis of Hygrine, Cuscohygrine, Cocaine, Cinnamoyl Ecgonine (Methylecgonine) and Hyoscyamine The **pyrrolidine ring system**, present in **hygrine** and **cuscohygrine**, is formed initially as a Δ^1 -pyrrolinium cation. The extra C-atoms required for hygrine formation are derived from acetate *via* acetyl-CoA; and the sequence appears to involve stepwise addition of *two* acetyl-CoA units as shown below:

These *two* steps may be explained as under:

- (a) The enolate anion from acetyl-CoA serves as nucleophile for the pyrrolinium ion in a Mannich-like reaction, that may give rise to products having either R or S stereochemistry.
- (b) An addition is caused by virtue of a Claisen condensation which essentially extends the side-chain, and the product is 2-substituted pyrrolidine, thereby retaining the thioester moiety of the second acetyl-CoA.


It has been observed that **Hygrine** and most of the naturally occurring **tropane alkaloids** is devoid of this specific C-atom, which is subsequently eliminated by suitable decarboxylation/hydrolysis reactions. Interestingly, the genesis of the bicyclic structure of the tropane skeleton existing in either *cocaine* or **hyoscyamine** is accomplished due to the repetition of the Mannich-like reaction stated above. These reactions are summarized in the description given under.

7.2.6.2 Pyrrolizidine Alkaloids


The **bicyclic pyrrolizidine nucleus** is formed by the utilization of two moles of ornithine and this pathway is accomplished *via* the intermediate **putrescine**. However, it has been observed that the plant sources usually synthesizing the pyrrolizidine alkaloids seem to be devoid of the *decarboxylase enzyme* that helps in the transformation of ornithine into putrescine; in fact, *ornithine* is really incorporated by way of arginine.

In nature, the **pyrrolizidine alkaloids** have a relatively broad stretch of distribution, but are specifically present in certain genera of the *Leguminosae/Fabaceae* (*e.g.*, **Crotalaria**); the *Compositae/Asteraceae* (*e.g.*, **Senecio**); and the *Boraginaceae* (*e.g.*, **Heliotropium, Symphytum, and Cynoglossum**). Broadly speaking the pyrrolizidine bases do not occur in their free form, but are mostly found as esters with rare mono- or di-basic acids, the **necic acids**.

The *two* important alkaloids of this category are, namely: **Retronecine** and **Senecionine**, which shall be discussed as under:



(continued)




Biosynthesis of Hygrine, Cuscohygrine, Cocaine, Carbamoyl Ecgonine (Methyl-ecgonine) and Hyoscyamine

A. Retronecine The most common base portion of the **pyrrolizidine alkaloids** is **retronecine**. The ‘Necine’ bases are *1-methylpyrrolizidines* of different stereochemical configurations and degree of hydroxylation which invariably occur as esters in alkaloids of *Senecio*, *Crotalaria* and a plethora of genera of the *Boraginaceae* as stated earlier.

Biological Source It is obtained from the herbs of *Heliotropium europaeum* L. (*Boraginaceae*) (**Heliotrope, Turnsole**).

Chemical Structure



(1*R-trans*)-2, 3, 5, 7, a-Tetrahydro-1-hydroxy-1 H-pyrrolizine-7-methanol; (C₈H₁₃NO₂).

Characteristic Features

1. It is obtained as crystals from acetone having mp 119-120°C.
2. It has the specific optical rotation $[\alpha]_D^{20} + 4.95^\circ$ (C = 0.58 in ethanol).

Identification Test It gives the racemic mixture *i.e.*, (\pm) form as crystals from acetone having mp 130-131°C.

Uses


1. The plant is used for cancer and is popularly known as “**Herbe Du Cancer**” in Europe.
2. It is also used for snakebite and scorpion stings.

B. Senecionine

Synonym Aureine;

Biological Source The hepatotoxic alkaloid is obtained from the whole plant of *Senecio vulgaris* L. (*Compositae*); weed of *Senecio aureus* L. (*Asteraceae*) (**Squaw Weed, Liferoot, Golden Groundsel**); and preblossoming plant of *Tussilago farfara* L. (*Asteraceae*) (**Coltsfoot, Coughwort, Horse-Hoof**).

Chemical Structure



12-Hydroxysenecionan-11, 16-dione; (C₁₈H₂₅NO₅): is described by Barger and Blackie (1936).*

Characteristic Features

1. It is obtained as plates having mp 236°C and a bitter taste.
2. Its specific optical rotation $[\alpha]_D^{25} - 55.1^\circ$ (C = 0.034 in chloroform).
3. It is practically insoluble in water; freely soluble in chloroform; and slightly soluble in ether and ethanol.

Uses

1. It is used as an excellent drug to control pulmonary hemorrhage.
2. It is also used to hasten labour and check the pains of parturition.

Biosynthesis of Retronecine and Senecionine It has been observed that the plants synthesizing the above mentioned **pyrrolizidine alkaloids** seem to be devoid of the decarboxylase enzyme transforming ornithine into putrescine; in fact, ornithine is actually incorporated by way of arginine. The various steps involved essentially in the **biosynthesis of retronecine** and **senecionine** are summarized as below:

1. Two moles of putrescine are condensed in an NAD⁺-dependent oxidative deamination reaction to yield the corresponding imine, which is subsequently transformed into **homospermidine** by the aid of NADH reduction.
2. The genesis of the creation of the pyrrolizidine skeleton is on account of the **homospermidine** molecule by a sequential series of interactions, such as: oxidative deamination, imine formation, intramolecular **Mannich reaction**, that specifically exploits the enolate anion produced from the aldehyde.
3. The '*pyrrolizidine skeleton*' thus provides a C₄N unit from ornithine, together with an additional four C-atoms from the same amino acid precursor.
4. The **senecionine** is a diester of **retronecine** with **senecic acid**.

These steps have been depicted as under.

7.2.7 Alkaloids Derived from Tyrosine

The pyridoxal phosphate (PLP)-dependent decarboxylation of *L-Tyrosine* yields the simple phenylethylamine *tyramine*, that subsequently undergoes di-N-methylation thereby producing **hordenine**. **Hordenine** is regarded as a germination inhibitory alkaloid obtained from barely *viz.*, *Hordeum vulgare* (*Graminae/Poaceae*).

There are a number of alkaloids derived from tyrosine which may be classified as stated below:

- (i) Phenylethylamine alkaloids,
- (ii) Simple Tetrahydro iso-quinoline alkaloids,
- (iii) Modified tetrahydro iso-quinoline alkaloids,
- (iv) Phenylethylisoquinoline alkaloids, and
- (v) Amaryllidaceae alkaloids.

* Barger, Blackie, *J. Chem. Soc.* 743 (1936)

Characteristic Features The characteristic features of some racemic forms, optical isomers and their respective salts are enumerated below:

1. ***dl*-Ephedrine (Synonyms: Racephedrine; Racemic Ephedrine):** The crystals have mp 79°C; and are soluble in oils, chloroform, ether, water, and ethanol.
2. ***dl*-Ephedrine Hydrochloride (Synonyms: Ephetonin; Racephedrine Hydrochloride) (C₁₀H₁₅NO.HCl):** The crystals have mp 187–188°C; and pH 6.0. Its solubility profile are: 1 g dissolves 4 ml water, 40 ml of 95% ethanol at 20°C; and practically insoluble in ether.
3. ***dl*-Ephedrine Sulphate (Synonym: Racephedrine Sulphate) (C₁₀H₁₅NO.H₂SO₄):** The crystals have mp 247°C, and are soluble in ethanol and water. Its solution has a pH of 6.0.
4. ***l*-Ephedrine [L-Erythro-2(methylamino)-1-phenylpropan-1-ol]:** It is obtained as waxy solid, crystals or granules, having a soapy feel and the substance gradually decomposes on exposure to light. It may contain water upto ½ mole (5.2%). However, the anhydrous product is hygroscopic in nature having mp 34°C. Interestingly, the absorption of water enhances mp to 40°C; and bp 255°C. The pH of aqueous solution (1 in 200) is 10.8. 1 g of it dissolves in 20 ml water, 0.2 ml ethanol; and freely soluble in ether, chloroform and oils.
5. ***l*-Ephedrine Hydrochloride (Synonyms: Ephedral; Senedrine):** It is obtained as orthorhombic needles having mp 216-220°C, which are affected by light. Its specific optical rotation $[\alpha]_D^{25} - 33$ to -35.5° (C = 5). The pH of aqueous solution (1 in 200) is 5.9. 1 g dissolves in 3 ml water, 14 ml ethanol; and is found to be practically insoluble in chloroform and ether.
6. ***l*-Ephedrine Sulphate:** Its orthorhombic needles have mp 245°C (decomposed) and are affected by light. Its specific optical rotation $[\alpha]_D^{25} - 29.5$ to -32.0° (C = 5). 1 g dissolves in 1.2 ml water and 95 ml ethanol; and freely soluble in hot alcohol. Its pH is about 6.

Identification Tests

1. Dissolve 0.01 g of **ephedrine** in 1 ml water by adding a few drops of dilute HCl. To this add two drops of CuSO₄ solution (5% w/v) followed by a few-drops of NaOH solution when a reddish colour is developed. Now, add 2-3 ml ether and shake the contents thoroughly; the ethereal layer turns purple while the lower aqueous layer becomes blue.
2. Dissolve 0.2 g of **ephedrine** in 30 ml of chloroform in a stoppered flask and shake the contents vigorously. Allow the mixture to stand for at least 12 hours at room temperature and then remove the chloroform over an electric water bath. The crystals of ephedrine hydrochloride separate out.
3. Triturate 0.05 g of **ephedrine** with a few crystals of [K₃Fe(CN)₆] *i.e.*, potassium ferricyanide, followed by a few drops of water and heat on a water bath slowly when a distinct odour of benzaldehyde (*i.e.*, similar to the odour of bitter almonds) is given out.

Uses


1. *l*-Ehedrine is used extensively as a bronchodilator.
2. It also exerts excitatory action on the CNS and produces noticeable effects on skeletal muscles.
3. It is also employed as nasal decongestant.

B. Hordenine

Synonyms Anhaline; Eremursine; Peyocactine.

Biological Sources It is obtained from the plant of *Lophophora williamsii* (Lamair) Coult. (*Cactaceae*) (**Peyote**) and *Selenicereus grandiflorus* Britt and Rose (*Coctaceae*) (**Night Blooming Cereus**).

Chemical Structure



4-[2-Dimethylamino) ethyl] phenol; (C₁₀H₁₅NO).

Isolation It is isolated from barley germs by the method suggested by Erspamer and Falconieri* (1952).

Characteristic Features

1. It is obtained as orthorhombic prisms from ethanol or benzene + ether; as needles from water having mp 117-118°C.
2. It sublimes at 140-150°C and has a bp₁₁ 173°C.
3. **Solubility Profile:** It is very soluble in chloroform, ethanol and ether; 7 g dissolves in 1 L of water; practically insoluble in petroleum ether; and sparingly soluble in benzene, xylene and toluene.

Identification Test **Hordenine** readily forms its hydrochloride salt which is obtained as needles from ethanol having mp 177°C; and it is very soluble in water.


Uses It exhibits digitalis-like activity.

C. Mescaline

Synonym Mezcaline

Biological Sources It is obtained from *Peyote* (**Mescal Buttons**) the flowering heads of *Lophophora williamsii* (Lemaire) Coult. (*Coctaceae*) and the cactus *Trichocereus pachanoi* Britton and Rose (*Cactaceae*) (**Achuma, San Pedro Aguacoli**).

Chemical Structure



3, 4, 5-Trimethoxybenzeneethanamine; (C₁₁H₁₇NO₃).

* Erspamer and Falconieri, *Naturwiss*, **39**, 431 (1952).

Isolation Mescaline has been successfully isolated from the plant source by Banholzer *et al.** (1952).

Characteristic Features

1. The crystals have mp 35-36°C and bp₁₂ 180°C.
2. It is moderately soluble in water; freely soluble in ethanol, chloroform and benzene; and practically insoluble in ether and petroleum ether.

Identification Tests It forms readily a variety of salts, such as:


1. **Mescaline Hydrochloride** ($C_{11}H_{17}NO_3 \cdot C_{11}H_{17}NO_3 \cdot HCl$): The needles have mp 181°C and freely soluble both in ethanol and water.
2. **Mescaline Sulphate Dihydrate** [$(C_{11}H_{17}NO_3)_2 \cdot H_2SO_4 \cdot 2H_2O$]: It is obtained as prisms having mp 183-186°C; soluble freely in methanol and hot water; and sparingly soluble in ethanol and cold water.
3. **Mescaline Acid Sulphate** ($C_{11}H_{17}NO_3 \cdot H_2SO_4$): The crystals have mp 158°C.
4. **N-Acetylmescaline**: It mostly occurs naturally, mp 94°C.
5. **N-Methylmescaline**: It occurs naturally, bp 130-140°C.
6. **N-Benzoylmescaline**: It is obtained as needles from aqueous ethanol having mp 121°C; and is found to be very soluble in ether and ethanol.

Note: This is a controlled substance (hallucinogen) listed in the US code of Federal Regulations [Title 21 Part 1308.11 (1995)].

D. Narceine

Biological Source It is obtained from the dried latex (**opium**) by incision from the unripe capsule of *Papaver somniferum* Linn., (*Papaveraceae*) to the extent of 0.1-0.5%.

Chemical Structure



Narceine

6-[[6-[2-(Dimethylamino) ethyl]-4-methoxy-1, 3-benzodioxol-5 yl] acetyl]-2, 3-dimethoxy benzoic acid; ($C_{23}H_{24}NO_8$).

Isolation The isolation of **narceine** from **morphine** mother liquors is tedious.** It may also be prepared from **narcotine** or **gnoscopine*****

* Banholzer *et al. Helv. Chim. Acta*, **35**, 1577 (1952).

** Merek, *Chem. Ztg.*, **13**, 525 (1889)

*** Roser, *Ann.* **247**, 167 (1888).

Characteristic Features

1. The anhydrous material is very hygroscopic in nature having mp 138°C; and its uv_{\max} (ethanol) is 270 nm ($\log \epsilon = 3.98$).
2. Usually the alkaloid is obtained as the trihydrate.
3. The clusters of silky and prismatic needles are obtained from water having mp 176°C.
4. Its dissociation constants are pK_b at 20° = 10.7; $K_b = 2 \times 10^{-11}$; $pK_a = 9.3$; $K_a = 5 \times 10^{-10}$.
5. The pH of its saturated solution is 5.8.
6. **Solubility Profile:** 1g dissolves in 770 ml water; 220 ml boiling water; moderately soluble in hot alcohol; almost insoluble in benzene, chloroform, ether, petroleum ether.
7. It forms salts with solutions of alkali hydroxide and also with dilute mineral acids.

Identification Test Ethylнарсеine Hydrochloride (C₂₅H₃₂ClNO₈) (Synonym: Narcyl): It is obtained as plates from water having mp 208-210°C. It is slightly soluble in cold water, insoluble in ether; and freely soluble in hot water, ethanol and chloroform.

Uses

1. Narcyl is used as a narcotic analgesic.
2. Narcyl is also employed as an antitussive agent.

Biosynthesis of Hordenine and Mescaline Decarboxylation of L-tyrosine *via* pyridoxal phosphate (PLP) yields the simple phenylethylamine derivative *tyramine*, which on di-N-methylation gives rise to **hordenine**. Besides, phenylethylamine derivatives commonly exhibit either 3, 4-di- or 3, 4, 5-trihydroxylation reactions, and are subsequently derived *via dopamine i.e.*, the decarboxylation product obtained from *L-DOPA* (L-dihydroxyphenylalaline). The two variants of **catecholamines**, namely: *first*, a mammalian neurotransmitter **noradrenaline (norepinephrine)**, and *secondly*, the most common '*fight or flight*' hormone released in animals from the adrenal gland due to fear psychosis or stress **adrenaline (epinephrine)**. Furthermore, these two compounds are formed due to β -hydroxylation and N-methylation of dopamine.

Lastly, aromatic hydroxylation and O-methylation convert dopamine into *mescaline*. All these reactions have been shown sequentially as given below.


7.2.7.2 Simple Tetrahydro Isoquinoline Alkaloids


The typical representatives of the simple **tetrahydroisoquinoline derivatives** are the closely-related alkaloids occurring along with mescaline are, namely: **anhalamine**, **anhalonine** and **anhalonidine**.

These *three* alkaloids shall be discussed in the pages that follow:

A. Anhalamine

Biological Sources It is obtained from the plant *Lophophora williamsii* (Lemaire) Coult. (*Cactaceae*) (*Peyote*), and *Anhalonium lewinii*. Henn. (*Cactaceae*).

Chemical structures



Biosynthesis of Hordenine and Mescaline

1, 2, 3, 4-Tetrahydro-6, 7-dimethoxy-8-isoquinolinol, ($C_{11}H_{15}NO_3$).

Characteristic Features

1. The crystals have mp 189-191°C.
2. Its uv_{max} (ethanol) is 274 nm ($\log \epsilon$ 2.90).
3. **Solubility Profile:** It is found to be almost insoluble in cold water, cold ethanol, ether and freely soluble in hot water, ethanol, acetone and dilute acids.

Identification Test Anhalamine Hydrochloride Dihydrate ($C_{11}H_{15}NO_3 \cdot HCl \cdot 2H_2O$): It is obtained as crystals from water having mp 258°C.


Uses It may play a minor role in causing hallucinations.

B. Anhalonine

Synonym Anhalanine

Biological Sources It is obtained from the **mescal buttons** [*Lophophora williamsii* (Lemaire) Coult. (*Anhalonium lewinii* Henn). *Cactaceae*]; and also in *Ariocarpus*, in *Gymnocalycium gibbosum*.

Chemical Structure



6, 7, 8, 9-Tetrahydro-4-methoxy-9-methyl-1, 3-dioxolo [4, 5-h] isoquinoline, ($C_{12}H_{15}NO_3$).

Characteristic Features

1. It is obtained as rhombic needles from petroleum ether having mp 86°C and $bp_{0.02}$ 140°C.
2. Its specific optical rotation $[\alpha]_D^{25} - 63.8^\circ$ (methanol); and $[\alpha]_D^{25} - 56.3^\circ$ (chloroform).
3. It is found to be freely soluble in ethanol, ether, chloroform, benzene and petroleum ether.

Identification Test


Anhalonine Hydrochloride ($C_{12}H_{15}NO_3 \cdot HCl$): It is obtained as orthorhombic prisms decomposing at 255°C. Its aqueous solution is almost neutral. It is found to be freely soluble in hot water.

Uses It may be employed as a mild hallucinating agent.

C. Anhalonidine

Biological Source It is invariably obtained from the **mescal buttons**, the buds of *Lophophora williamsii* (Lemaire) Coult. (*Anhalonium lewinii* Henn.) belonging to the natural order *Coctaceae*.

Chemical Structure



1, 2, 3, 4-Tetrahydro-6, 7-dimethoxy-1-methyl-8-isoquinolinol; (C₁₂H₁₇NO₃).

Characteristic Features


1. It is mostly obtained as small octahedral crystals from benzene having mp 160-161°C.
2. Its uv_{max} (ethanol) is 270 nm ($\log \epsilon$ 2.81).
3. Its aqueous solution acts as a strong base.
4. It is freely soluble in water, ethanol, chloroform and hot benzene; sparingly soluble in ether; and practically insoluble in petroleum ether.
5. It has been observed that the solutions of anhalonidine acquire a reddish colouration on standing.

Uses It may be used as a mild hallucinogen.

Biosynthesis of Anhalamine, Anhalonine and Anhalonidine Interestingly, the two additional C-atoms present in **anhalonidine** and **anhalonine** are provided by pyruvate; whereas, the C-atom for **anhalamine** is supplied by glyoxylate, as shown below. However, in each instance, a carboxyl group is lost from this aforesaid additional precursor. The *pyruvate i.e.*, the keto-acid eventually reacts with an appropriate phenylethylamine, in this particular instance the *dimethoxy-hydroxy derivative*, thereby yielding a **Schiff Base**. Further, a **Mannich-like mechanism** helps in the cyclization to produce the heterocyclic isoquinoline nucleus, whereby the mesomeric effect of an oxygen substituent caters for the nucleophilic site on the aromatic ring. Evidently, restoration of aromaticity *via* proton loss yields the tetrahydroquinoline nucleus, thus representing overall a biosynthetic equivalent of the **Pictet-Spengler Isoquinoline Synthesis**.* Subsequently, the carboxyl group is eliminated, not by means of a simple decarboxylation process, but *via* an unusual oxidative decarboxylation process that essentially involves the following steps, namely:

- (i) *First*, producing the intermediate *imine*,
- (ii) *Secondly*, subjecting to *reduction* yielding *anhalonidine*,
- (iii) *Thirdly*, subjecting to *methylation* giving rise to *anhalonine*,
- (iv) *Fourthly*, subjecting phenylethylamine precursor employing the *glyoxylic acid* instead of *pyruvic acid* generating *anhalamine*.

* **Pictet-Spengler Isoquinoline Synthesis:** Formation of tetrahydro-isoquinoline derivatives by condensation of β -arylethylamines with carbonyl compounds and cyclization of the Schiff bases formed:





It is, however, pertinent to mention here that the aforesaid mode of coupling is perhaps less frequently found than that involving carbon-carbon bonding between aromatic rings. The major **opium alkaloids** viz., **morphine**, **codeine** and **thebaine** are obtained through this mode of coupling. (R)-*Reticuline* has been established beyond any reasonable doubt as the precursor of the above **three morphinan alkaloids**. Interestingly, there exist an ample evidence to show that the later stages of the proposed biosynthetic pathway undergo *modifications* in certain strains of *opium poppy*. Thus, in such modified strains of opium poppy **thebaine** is being converted to **oripavine** and **morphinone**, whereby the phenolic O-methyl moiety is removed before that of the ether, i.e., the same steps are carried out but in an altogether different order.


The various alkaloids belonging to this category, namely: **morphine**, **codeine**, **thebaine**, **reticuline**, **oripavine** and **morphinone** shall be discussed separately in the following sections:

A. Morphine

Synonyms Morhium; Morphia; Dolcontin; Duromorph; Morphina; Nepenthe.

Biological Sources **Morphine** is obtained from a variety of medicinal plants, such as: *Argemone mexicana* L. (*Papaveraceae*) (**Prickly Poppy**); *Eschscholzia californica* Cham. (*Papaveraceae*) (**California Poppy**); *Papaver bracteatum* Lindl. (*Papaveraceae*) (**Great Scarlet Poppy**; Thebaine Poppy); *Papaver somniferum* L. (*Papaveraceae*) (**Opium Poppy**; and Poppysed Poppy Keshi).



Chemical Structure

(5 α , 6 α)-7, 8-Didehydro-4, 5-epoxy-17-methylmorphinan-3, 6-diol; (C₁₇H₁₉NO₃).

Isolation The latex obtained by incision on the unripe capsule of opium poppy is first collected in clean, plastic containers, and the process of incision is repeated at least four times on the same capsule after an interval of two days. Care must be taken to make the incisions on the superficial surface only so as to collect exclusively the external exudation of latex. Subsequently, the latex is dried carefully either by exposing to air on metallic shallow plates or by passing a stream of hot air. Thus the 'opium' or the dried latex is stored for the isolation of **morphine**. It is found to contain usually 9.5% **morphine** when calculated as anhydrous morphine.


The morphine may be isolated from '**Powdered Opium**' by adopting the following steps sequentially:

- Step-1:** The powdered opium is shaken with calcium chloride solution and filtered.
- Step-2:** The resulting filtrate is concentrated and to it is added 10% w/v sodium hydroxide solution carefully *i.e.*, to solubilize morphine, codeine and narceine. It is now filtered.
- Step-3:** The filtrate containing morphine, codeine and narceine is extracted with chloroform. The resulting mixture is separated.
- Step-4:** The lower chloroform layer contains codeine, whereas the upper aqueous layer comprises of **morphine** and **narceine**.
- Step-5:** The aqueous layer is first acidified and subsequently made alkaline with ammonia, whereby morphine gets precipitated and collected as a white solid residue (Yield = 9.5%).


Characteristic Features

1. **Morphine** is obtained as short, orthorhombic, columnar prisms from anisole that gets decomposed at 254°C. It also occurs in its metastable phase having mp 197°C. However, the high melting form sublimates at 190-200°C (0.2 mm pressure at 2 mm distance).
2. It has a bitter taste.
3. **Morphine** (free-base) unlike most other alkaloids in their free-base forms is found to be sparingly soluble in chloroform and nearly insoluble in ether or benzene.
4. **Morphine** gets dissolved in caustic alkalis by virtue of the fact that the OH moiety at C-3 is phenolic in nature and the other OH function at C-6 is a secondary alcoholic group.
5. **Morphine** is a monoacidic base and hence, forms salts that crystallizes rapidly. These are found to be neutral to litmus and methyl orange.
6. The average pH of a saturated solution of morphine salt is found to be 4.68.

7. **Morphine** is levorotatory.
8. **Morphine** upon oxidation converts the secondary alcoholic function to a ketonic group and thus **codeinone** is formed.



9. **Morphine** upon being heated in a sealed tube with concentrated HCl at 140°C eliminates a molecule of water to yield apomorphine as given below:



Identification Tests There are several colour tests that are employed for the identification of **morphine**, namely:

1. **Brouadrel-Boutmy Test:** Add to an aqueous solution of a salt of morphine (*e.g.*, morphine hydrochloride, morphine acetate trihydrate, morphine tartrate trihydrate) a few drops of a mixture of dilute solutions of potassium ferricyanide [K₃Fe(CN)₆] (0.1% w/v) and ferric chloride [FeCl₃] (0.1% w/v) when a deep blue colouration is produced.
Note: (1) Morphine causes reduction of K₃Fe(CN)₆ to K₄Fe(CN)₆ i.e., potassium ferrocyanide which gives a deep blue colouration with FeCl₃.
(2) Codeine fails to give this test; and hence it may be employed to differentiate between morphine and codeine.
2. **Schneider-Weppen Test:** **Morphine** when mixed with 6 to 8 portions of powdered cane sugar (C₆H₁₂O₆) intimately and treated with a few drops of concentrated H₂SO₄, the development of a purple-red colour takes place. However, the addition of bromine water before the addition of H₂SO₄ renders the test rather more sensitive.
3. **Iodic Acid Test:** **Morphine** (10 mg) when dissolved in dilute H₂SO₄ (2N) is treated with a solution of pure iodic acid or potassium iodate; and the resulting mixture is shaken with chloroform (5 ml), a distinct violet colouration is imparted to the chloroform layer.

Note: Morphine reduces iodic acid and potassium iodate.

4. **Sodium Nitrite Test:** To a solution of **morphine** in dilute HCl add a few drops of sodium nitrite solution (1% w/v). Allow the reaction mixture to stand for 5-8 minutes and then make it alkaline with dilute ammonia solution, the development of a red colour confirms the presence of morphine.

Note: (1) It is a non-specific test for morphine and is also given by other phenolic substances.

(2) It legitimately distinguishes morphine from codeine.

5. **Nitric Acid Test: Morphine** readily gives an orange-red colouration when a few mg of it is treated with a few drops of concentrated nitric acid.
- (a) The resulting orange-red colouration rapidly changes to yellow on heating.
- (b) The orange-red colouration gets easily disappeared on the addition of a few drops of stannous chloride solution (SnCl_2) (1% w/v).
6. **Ferric Chloride Test:** When a neutral solution of **morphine** is treated with a few drops of ferric-chloride solution (1% w/v), a greenish-blue colour is produced.

Derivatives of Morphine A number of derivatives of morphine are produced that essentially have distinct characteristic features as enumerated below:

1. **Morphine Monohydrate ($\text{C}_{17}\text{H}_{19}\text{NO}_3 \cdot \text{H}_2\text{O}$):**

- (i) It is obtained as orthorhombic, sphenoidal prisms, or needles from methanol that gets decomposed at 254-256°C with rapid heating.
- (ii) It darkens on exposure to light and also loses water of crystallization at 130°C.
- (iii) Its physical parameters are: d_D^{20} 1.32; $[\alpha]_D^{25} - 132^\circ$ (methanol); pK_b at 20°C = 6.13, pK_a 9.85; pH of a saturated solution 8.5; and uv_{max} in acid: 2.85 nm, in alkali: 298 nm.
- (iv) **Solubility Profile:** 1 g dissolves in about 5000 ml of water, 1100 of boiling water, 210 ml of ethanol, 98 ml of boiling ethanol, 1220 ml of chloroform, 6250 ml of ether, 114 ml of amyl alcohol, 10 ml of boiling methanol, 525 ml of ethyl acetate; freely soluble in solutions of fixed alkali and other alkaline earth hydroxides, in phenols, cresols; moderately soluble in mixtures of chloroform with alcohols; and slightly soluble in ammonia benzene.

2. **Morphine Acetate Trihydrate ($\text{C}_{19}\text{H}_{23}\text{NO}_5 \cdot 3\text{H}_2\text{O}$):**

- (i) It is a yellowish-white powder.
- (ii) It has a slight acetic odour.
- (iii) Its specific optical rotation $[\alpha]_D^{15} - 77^\circ$ (water).
- (iv) It dissolves 1 g in 2.25 ml of water, 2 ml of boiling water, 22 ml of ethanol, 2 ml of ethanol at 60°C, 4.5 ml of glycerol, 4.75 ml of chloroform; and practically insoluble in ether.

3. **Morphine Tartrate Trihydrate [$(\text{C}_{17}\text{H}_{19}\text{NO}_3)_2 \cdot \text{C}_4\text{H}_6\text{O}_6 \cdot 3\text{H}_2\text{O}$]:** It is obtained as a crystalline powder. It is soluble in 11 parts of water; slightly soluble in alcohol; and practically insoluble in ether, chloroform and carbon disulphide.

Uses

1. It is used as a potent narcotic analgesic.
2. It is usually given in severe pains and also in such instances where patient fails to show positive response to other analgesics.

3. It exerts a biphasic action on the CNS.
4. It is found to sedate the respiratory centre, emetic centre and the cough centre through its action in the medulla.
5. It stimulates the chemoreceptor-trigger-zone located in the medulla that ultimately causes nausea and vomiting; and this is perhaps regarded as a side-effect.
6. It also exerts sedative and hypnotic actions.


Note: Morphine and its salts are habit forming drugs. Hence, its use must be done under the strict observation of a physician.

B. Codeine

Synonyms Codicept; Morphine monomethyl ether; Morphine 3-methyl ether; Methyilmorphine.

Biological Sources It is obtained from the plant *Argemone mexicana* L. (*Papaveraceae*) (**Prickly Poppy**); *Eschscholzia californica* Cham. (*Papaveraceae*) (**California Poppy**); *Papaver bracteatum* Lindl. (*Papaveraceae*) (**Great scarlet poppy, Thebaine Poppy**); and *Papaver somniferum* L. (*Papaveraceae*) (**Opium Poppy, Poppyseed Poppy Keshi**).

Chemical Structure



(5 α , 6 α)-7, 8-Didehydro-4, 5-epoxy-3-methoxy-17-methyl-morphinan-6-ol; (C₁₈H₂₁NO₃).

Preparation It is invariably present in opium from 0.7 to 2.5% depending on the sources of plant substances. However, mostly it is prepared by carrying out the methylation of **morphine**.

Characteristic Features

1. It is obtained as monohydrate orthorhombic sphenoidal rods or tablets (octahedra) from water or dilute ethanol having mp 154-156°C (after drying at 80°C).
2. It is found to sublime (when anhydrous) at 140-145°C under 1.5 mm reduced pressure.
3. It is observed to melt to oily drops when heated in an amount of water is sufficient for complete solution, and subsequently crystallizes on cooling.
4. Its physical parameters are: d_4^{20} 1.32; $[\alpha]_D^{15}$ -136° (C = 2 in ethanol); $[\alpha]_D^{15}$ -112° (C = 2 in chloroform); pK (15°) 6.05; pH of a saturated solution 9.8.
5. **Solubility Profile:** 1 g dissolves in 120 ml water, 60 ml water at 80°C, 2 ml ethanol, 1.2 ml hot ethanol, 13 ml benzene, 18 ml ether, 0.5 ml chloroform; freely soluble in methanol, dilute acids and amyl alcohol; and almost insoluble in solutions of alkali hydroxides and in petroleum ether.

Identification Test It forms various types of salts, namely:

1. **Codeine Acetate (C₂₀H₂₅NO₅):** The dihydrate is obtained as crystals having an acetic acid odour. It is found to be soluble in water and ethanol. It loses acetic acid on keeping and subsequently turns into a product which is incompletely soluble in water.
2. **Codeine Hydrobromide (C₁₈H₂₁NO₃.HBr):** The dihydrate is obtained as crystals and the anhydrous product shows a mp 190-192°C; $[\alpha]_D^{22} - 96.6^\circ$; 1 g dissolves in 60 ml water, 110 ml ethanol; and pH about 5.
3. **Codeine Hydrochloride (C₁₈H₂₁NO₃.HCl):** Its dihydrate salt is obtained as small needles having mp ~ 280°C with some decomposition; $[\alpha]_D^{22} - 108^\circ$; 1 g dissolves in 20 ml of water, 1 ml boiling water, 180 ml ethanol; and pH about 5.
4. **Codeine Salicylate (C₂₅H₂₇NO₆):** It is obtained as white crystalline powder; slightly soluble in water; and freely soluble in ethanol or ether.
5. **Codeine Phosphate (C₁₈H₂₄NO₇.P) (Galcodine):** The hemihydrate salt (USP) is obtained as fine, white, needle-shaped crystals or crystalline powder. It is odorless and affected by light. The solution is acidic to litmus. It is freely soluble in water; very soluble in hot water; slightly soluble in ethanol; and more soluble in boiling ethanol.
6. **Codeine Sulphate (C₃₆H₄₄N₂O₁₀.S):** The trihydrate is obtained as crystals or crystalline powder; 1 g dissolves in 30 ml water; 6.5 ml water at 80°C; 1300 ml ethanol; insoluble in chloroform or ether; pH 5.0.
7. **Codeine Methyl Bromide (C₁₉H₂₄Br.NO₃) (Eucodin) :** Its crystals have mp ~ 260°C; soluble in 2-3 parts of water, in hot methanol; sparingly soluble in ethanol; and insoluble in chloroform and ether.

Uses


1. It is mostly used as a narcotic analgesic.
2. It is invariably employed as an antitussive.

C. Thebaine

Synonym Paramorphine;

Biological Sources It is obtained from the fresh capsule latex (0.125%), dried 0.25 to 0.26% of *Papaver bracteatum* Lindl. (*Papaveraceae*) (**Great Scarlet Poppy, Thebaine Poppy**); and the air-dried milky exudation obtained from excised unripe fruits of *Papaver somniferum* L. (*Papaveraceae*) (**Opium Poppy, Poppyseed Poppy Keshi**).

Chemical Structure



(5 α)-6, 7, 8, 14-Tetrahydro-4, 5-epoxy-3, 6-dimethoxy-17-methylmorphinan; (C₁₉H₂₁NO₃).

Isolation Thebaine may be isolated from opium by means of the following steps, namely:

Step-1: Opium (dried latex) is treated with calcium chloride solution and then extracted with warm water. Allow it to remain as such for 24 hours.

Step-2: Filter the resulting product and collect the residue and filtrate separately.

Residue— contains the salts of calcium as lactate, sulphate, resinate and meconate (To be discarded).

Filtrate— contains the hydrochloride of various alkaloids present in opium.

Step-3: Add dilute NaOH solution (2N) carefully to the resulting filtrate and allow it to stand for 4-6 hours. Filter the contents of the flask:

Filtrate— contains morphine, codeine and narceine

Residue— contains thebaine, papaverine and narcotine

Step-4: Dissolve the residue or precipitate in dilute ethanol (50% v/v), make slightly acidic with the addition of dilute glacial acetic acid and finally add to it approximately *three volumes* of boiling distilled water.

Step-5: Filter the above reaction product:

Filtrate— contains thebaine

Residue— contains papaverine and narcotine

Step-6: Concentrate the filtrate obtained in Step-5 under reduced pressure and add to it dilute NH₄OH solution to make it alkaline; and extract the liberated alkaloid thebaine successively with chloroform. Thebaine is obtained after evaporation of chloroform under vacuo.

Characteristic Features

1. It is obtained as orthorhombic, rectangular plates by sublimation at 170-180°C under atmospheric pressure and a 1 mm distance mp 193°C (rapid heating).
2. Its physical parameters are: $[\alpha]_D^{15} - 219^\circ$ (p = 2 in ethanol); $[\alpha]_D^{23}$ (p = 5 in chloroform); pK at 15°C = 6.05; and pH of a saturated solution is 7.6.
3. **Solubility Profile:** 1 g dissolves in 1460 ml water at 15°C, in about 15 ml hot ethanol, 13 ml chloroform, 200 ml ether, 25 ml benzene, 12 ml pyridine; and not very soluble in petroleum ether.

Identification Tests Thebaine forms a number of salt derivatives which have specific characteristic features, such as:

1. **Thebaine Salicylate (C₁₉H₂₁NO₃.C₇H₆O₃):** It is obtained as crystals which are soluble in 750 parts of water. Thus, thebaine may be separated from other major alkaloids of opium by forming its salicylate derivative which is sparingly soluble in water.
2. **Thebaine Hydrochloride Monohydrate (C₁₉H₂₁NO₃.HCl.H₂O):** It is obtained as orthorhombic prisms from alcohol having $[\alpha]_D^{23} - 164^\circ$ (p = 2). It is found to be soluble in about 12 parts of water and in ethanol. The pH of a 0.05 molar solution is 4.95.
3. **Thebaine Oxalate Hexahydrate (2 C₁₉H₂₁NO₃.C₂H₂O₄.6 H₂O):** It is obtained as prisms. It is soluble in 10 parts of water and also in ethanol; and is almost insoluble in ether.
4. **Thebaine Binopalate Monohydrate (C₁₉H₂₁NO₃.C₂H₂O₄.H₂O):** It is obtained as prisms and found to be soluble in 45 parts of water.

5. **Thebaine Bitartrate Monohydrate** ($C_{19}H_{21}NO_3 \cdot C_4H_6O_6 \cdot H_2O$): It is obtained as prisms, soluble in 130 parts of water, quite soluble in both hot water and hot ethanol.
6. It gives a red colour on the addition of a few drops of cold sulphuric acid which ultimately changes to orange yellow.


Uses It is an opiate analgesic.

D. Reticuline

Synonym Coclanoline.

Biological Sources It is obtained from the plant *Hydratis canadensis* L. (*Ranunculaceae*) (Goldenseal); the leaves of *Laurus nobilis* L. (*Lauraceae*) (**Bay, Grecian Laurel, Green Bay**); the air-dried milky exudation obtained from excised unripe fruits of *Papaver somiferum* L. (*Papaveraceae*) (**Opium Poppy, Poppyseed Poppy Keshi**); and the leaves of *Sassafras albidum* (Nutt.) Nees (*Lauraceae*) (**Sassafras**).

Chemical Structure



1, 2, 3, 4-Tetrahydro-1-[(3-hydroxy-4-methoxyphenyl) methyl]-6-methoxy-2-methyl-7-isoquinolinol;
($C_{19}H_{23}NO_4$).

Isolation Gopinath *et al.*,* has described the isolation of *d*-form of reticuline from *Anona reticulata* Linn., (*Annonaceae*).

Characteristic Features

1. The *dl*-form of **reticuline** is obtained as pink crystals having mp 146°C.
2. The uv_{max} : 284 nm ($\log \epsilon$ 3.85).
3. **Solubility Profile**: It is soluble in aqueous buffer of pH < 7.5 or > 11; and is practically insoluble in water at pH 8-10.

Identification Tests

(S)-Form Reticuline Perchlorate ($C_{19}H_{23}NO_4 \cdot HClO_4$): It is obtained as colourless prisms from ethanol having mp 203-204°C. Its specific optical rotation $[\alpha]_D^{18} + 88.3^\circ$ (C = 0.21 in ethanol).


E. Oripavine

Synonym O³-Demethylthebaine.

* Gopinath *et al.*, *Ber.* **92**, 776 (1959).

Biological Sources It is obtained from the plant *Papaver bracteatum* Lindl. (*Papaveraceae*) (**Great Scarlet Poppy, Thebaine Poppy**); and *Papaver orientale* Linn. (*Papaveraceae*).

Chemical Structure



(5 α)-6, 7, 8, 14-Tetrahydro-4, 5-epoxy-6-methoxy-17-methyl-morphinan-3-ol; (C₁₈H₁₉NO₃).


Isolation It has been isolated from plant source by Kiselev and Konovalova.*

Characteristic Features The crystals have mp 200-201°C; and $[\alpha]_D^{20} - 211.8^\circ$.

Identification Tests

1. **Oripavine Hydrochloride (C₁₈H₁₉NO₃.HCl):** It is obtained as crystals which decompose at 244-245°C.
2. **Oripavine Methiodide (C₁₈H₁₉NO₃.CH₃I):** The crystals decompose at 207-208°C.

F. Morphinone It has been observed that the later stages of the biosynthetic pathway starting from reticuline leading to **thebaine** and **morphine** are strategically modified in some strains of opium poppy. Therefore, in such strains, thebaine is converted by way of **oripavine** and **morphinone**. In this pathway the phenolic *O*-methyl function is removed before that of the enol ether, *i.e.*, accomplishing the same steps but in a different order. In other words, **morphinone** is obtained by the demethylation of **oripavine** as shown below:



Biosynthesis of Morphine, Codeine, Thebaine, Oripavine and Morphinone The various steps involved are as follows:

* Kiselev and Konovalova *J. Gen. Chem. USSR*, **18**, 142 (1948).

1. **(R)-Reticuline**, may be redrawn as shown in page 495 following pathway is found to be the substrate for one-electron oxidation *via* the phenol moiety present in each ring thereby yielding the diradical.
2. Subsequent coupling *ortho* to the phenol group in the tetrahydroisoquinoline nucleus, and *para* to the phenol in the benzyl substituent, gives rise to **salutaridine**—a *dienone* which is found as minor alkaloidal component in the **opium poppy** *Papaver somniferum*.
3. **Thebaine** is achieved *via* **salutaridinol** produced from salutaridine by means of the stereospecific reduction of the carbonyl group.
4. In **thebaine** the ring closure to form the ether linkage is caused due to the nucleophilic attack of the phenol moiety on the dienol system followed by a displacement of the hydroxyl group.
5. Future reactions essentially involve conversion of **thebaine** into *morphine* *via* *codeine* by virtue of a process that exclusively modifies the oxidation state of the diene ring, but apparently removes *two* *O*-methyl groups.
6. One is evidently present as an enol ether, removal of which yields *neopinone*, that subsequently gives rise to **codeinone** and then **sodeine** by the help of allylic isomerisation and reduction respectively.
7. In certain specific strains of **opium poppy**, **thebaine** is changed to **oripavine** and **morphinone** by virtue of the pathway that essentially removes the phenolic *O*-methyl function before that of the enol ether.

7.2.8 Alkaloids Derived from Tryptophan

L-Tryptophan is a neutral heterocyclic amino acid containing essentially an indole ring system. It has been observed that it serves as a precursor for a wide spectrum of indole alkaloids. However, there exists an ample concrete evidence that major rearrangement reaction may convert the predominant **indole-ring system** into a **quinoline-ring system** thereby enhancing further the overall ability of tryptophan to act broadly as an alkaloid precursor.


The various alkaloids derived from tryptophan are conveniently classified into the following categories, namely: (i) **Simple Indole Alkaloids**; (ii) **Simple β -Carboline Alkaloids**; (iii) **Terpenoid Indole Alkaloids**; (iv) **Quinoline Alkaloids**; (v) **Pyrroloindole Alkaloids**; (vi) **Ergot Alkaloids**.


These aforesaid categories of alkaloids shall be discussed separately with typical important examples followed by the possible biosynthetic pathways, wherever necessary.

7.2.8.1 Simple Indole Alkaloids

L-Tryptophan (*i.e.*, α -aminoindole-3-propanoic acid) on decarboxylation yields *tryptamine*. The *N*-methyl and *N,N*-dimethyl derivatives of the latter are broadly distributed in the plant kingdom as *serotonin*—a simple hydroxylated derivative. Sequential biotransformation *viz.*, decarboxylation, *N*-methylation and hydroxylation gives rise to the formation of **psilocin**; whereas, phosphorylation of the OH group in psilocin yields *psilocybin*.

The *three* alkaloids, namely: **serotonin**, **psilocin** and **psilocybin** shall be discussed in the sections that follow:






Biosynthesis of Morphine, Codeine, Thebaine, Oriparine and Morphinone

A. Serotonin

Synonyms 5-Hydroxytryptamine; 5-HT; Enteramine; Thrombocytin; Thrombotonin;

Biological Sources The root bark of *Gossypium hirsutum* L. (*Malvaceae*) (**American Unplanted Cotton**) contains serotonin.

Chemical Structure



Serotonin

3-(2-Aminoethyl)-1H-indol-5-ol; (C₁₀H₁₂N₂O).

Identification Tests

1. **Serotonin Hydrochloride (C₁₀H₁₂N₂O.HCl)** It is obtained as hygroscopic crystals, sensitive to light having mp 167-168°C. It is water soluble and the aqueous solutions are found to be stable at pH 2-6.4.
2. **Serotonin complex with Creatinine Sulphate Monohydrate (C₁₄H₂₁N₅O₆S.H₂O) (Antemovis)** It is obtained as plates which decomposes at 215°C. Its uv_{max} (water at pH 3.5): is 275 nm (ϵ 15,000). It has two dissociation constants $pK_1' = 4.9$ and $pK_2' = 9.8$. The pH of a 0.01 molar aqueous solution is 3.6. It is found to be soluble in glacial acetic acid; very sparingly soluble in methanol and ethanol (95%); and insoluble in absolute ethanol, acetone, pyridine, ethyl acetate, chloroform, benzene and ether.

Uses


1. It is a potent vasoconstrictor.
2. It is also a neurotransmitter in the CNS and is important in sleep-walking-cycles.

B. Psilocin

Synonyms Psilocyn.

Biological Sources It is obtained from the sacred mushroom of Mexico known as **Teonanacatl**. It is also found in the fruiting bodies of *Psilocybe maxicana* Heim, (*Agaricaceae*).

Chemical Structure



Psilocin

3-[2-(Dimethylamino) ethyl]-1H-indol-4-ol; (C₁₂H₁₆N₂O);

Isolation It has been successfully isolated in trace amounts from the fruiting bodies of *Psilocybe mexicana**.

Characteristic Features

1. It is obtained as plates from methanol having mp 173–176°C.
2. It is an amphoteric substance.
3. It is unstable in solution, more precisely in an alkaline solution.
4. It is very slightly soluble in water.
5. Its uv_{\max} : 222, 260, 267, 283, 293 nm (log ϵ 4.6, 3.7, 3.8, 3.7, 3.6).

Uses It is a hallucinogenic substance


Note It is a controlled substance listed in the U.S. Code of Federal Regulations, Title 21 Part 1308, 11 (1995).

C. Psilocybin

Synonym Indocybin;

Biological Sources These are same as mentioned in **psilocin 'B'** above.

Chemical Structure



3-[2-(Dimethylamino) ethyl]-1H-indol-4-ol dihydrogen phosphate ester; (C₁₂H₁₇N₂O₄P).

Isolation The method of isolation of **psilocybin** is the same as stated under **psilocin**.

Characteristic Features


1. **Psilocybin** is obtained as crystals from boiling water having mp 220-228°C; and from boiling methanol mp 185-195°C.
2. It has uv_{\max} (methanol): 220, 267, 290 nm (log ϵ 4.6, 3.8, 3.6).
3. The pH of a saturated solution in 50% aqueous ethanol is 5.2.
4. **Solubility Profile:** It is soluble in 20 parts of boiling water, 120-parts of boiling methanol; sparingly soluble in ethanol; and practically insoluble in chloroform, benzene.

Uses It is a hallucinogenic substance and exerts its action at a dose level of 6-20 mg.

Biosynthesis of Serotonin, Psilocin and Psilocybin The different steps involved in the biosynthesis of **serotonin**, **psilocin** and **psilocybin** may be summarized as stated below:


* Hofmann *et al.*, *Experientia*, **14**, 107 (1958); Heim *et al.*, *Helv. Chim. Acta*, **42**, 1557 (1959).

1. L-Tryptophan upon oxidation gives rise to the corresponding hydroxylated derivative known as 5-hydroxyl-L-tryptophan, which further undergoes decarboxylation to yield **serotonin** also termed as **5-hydroxytryptamine (or 5-HT)**.
2. L-Tryptophan undergoes decarboxylation to yield the tryptamine, which affords N-Methylation and N,N-dimethylation in the presence of S-adenosylmethionine (SAM). The resulting dimethyl derivative upon oxidation gives rise to the product **psilocin** another hydroxylated derivative.
3. Phosphorylation of the hydroxyl function in psilocin affords **psilocybin**.
4. Interestingly, both **psilocin** and **psilocybin** are solely responsible for attributing the hallucinogenic properties of the so-called '*magic mushrooms*', that include species of **Psilocybe**, **Panaeolus** and the like.



7.2.8.2 Simple β -Carboline Alkaloids

The alkaloids based on the **β -carboline ring system** obviously suggest the formation of a new six-membered heterocyclic ring employing the *ethylamine side-chain present in tryptamine* exactly in the same manner to the evolution of tetrahydroisoquinoline alkaloids (see Section 7.2.7.2). The exact mechanism whereby the above rearrangement is accomplished may be explained by virtue of the fact that C-2 of the indole nucleus is *nucleophilic* due to the adjacent nitrogen atom. Therefore, C-2 can conveniently participate in a **Mannich/Pictet-Spengler type reaction**, thereby enabling it to attack a **Schiff base** produced from tryptamine and either an aldehyde (or keto acid) as given below:



It has been observed that relatively *simpler structures* make use of *keto-acids*, such as: **harman**, **harmaline**, **harmine** and **elaegnine**. These alkaloids shall be treated individually in the sections that follow.


Interestingly, the comparatively complex carbolines, for instance: the **terpenoid indole alkaloids** e.g., **ajmaline** are usually generated by the help of a pathway that specifically utilize *an aldehyde*, such as: secologanin. This particular section shall be dealt with separately under Section 7.2.8.3.

A. Harman

Synonyms Aribine; Loturine; Passiflorin; 2-Methyl- β -carboline; 3-Methyl-4-carboline;

Biological Sources It is obtained from the bark fruit of *Passiflora incarnata* L. (*Passifloraceae*) (**May pop, Passion flower**); seed of *Peganum harmala* L. (*Rutaceae*) (**Harmel, Syrian Rue, African Rue**), bark of *Sickingia rubra* (Mart.) K. Schum. (**Arariba rubra Mart.**), (*Rubiaceae*); and bark of *Symplocos racemosa* Roxb. (*Symplocaceae*).

Chemical Structure



Harman

1-Methyl-9H-pyrido [3, 4, b] indole; (C₁₂ H₁₀ N₂).

Isolation Poindexter and Carpenter* isolated this alkaloid from the cigarette smoke.

Characteristic Features

1. It is obtained as orthorhombic crystals from heptane and cyclohexane having mp 237-238°C.
2. It has a bitter taste.
3. It exhibits distinct bright blue fluorescence in uv light.
4. Its pKa's are 7.37 and 144.6.
5. It has uv_{max} (methanol): 234, 287, 347 nm (log ϵ 4.57, 4.21, 3.66).
6. It is practically insoluble in water and freely soluble in dilute acids.

Identification Test **Harman Hydrochloride (C₁₂H₁₀N₂.HCl)** It is obtained as rosettes of needles from a mixture of ethanol + 20% HCl in water which sublimes at 120-130°C.

Uses It is a narcotic hallucinogen.


B. Harmaline

Synonyms Harmidine; Harmalol Methyl Ether; O-Methyl-harmalol; 3, 4-Dihydroharmine;

Biological Sources It is obtained from the seeds of *Peganum harmala* L. (*Zugophyllaceae*); and *Banisteria cappi* Spruce (*Malpighiaceae*). It is also obtained from the fruit of *Passiflora incarnata* L. (*Passifloraceae*) (**Passionflower, Maypop**).

* Poindexter and Carpenter, *Chem. & Ind.* (London), **1962**, 176.

Chemical Structure



Harmaline

4, 9-Dihydro-7-methoxy-1-methyl-3H-pyridol [3, 4,-b] indole; ($C_{13} H_{14} N_2 O$).

Characteristic Features

1. It is obtained as orthorhombic bipyramidal prisms, or tablets from methanol; and as rhombic octahedra from ethanol having the same mp 229-231°C.
2. Its solutions give a blue fluorescence.
3. Its dissociation constant pK_a 4.2.
4. It has uv_{max} (methanol): 218, 260, 376 nm ($\log \epsilon$ 4.27, 3.90 and 4.02)
5. It is found to be slightly soluble in water, ethanol, ether; and very soluble in dilute acids and hot ethanol.

Identification Tests Harmaline forms definite derivatives as shown below:

1. **Harmaline Hydrochloride Dihydrate ($C_{13}H_{14}N_2O.HCl.2H_2O$):** It is obtained as slender, yellow needles that are found to be moderately soluble in ethanol and water.
2. **N-Acetylharmaline:** It is obtained as needles having mp 204-205°C.

Uses


1. It is recognized as a narcotic hallucinogen.
2. It is used as a CNS-stimulant.

C. Harmine

Synonyms Telepathine; Leucoharmine; Yageine; Banisterine;

Biological Sources It is obtained from the seeds of *Peganum harmala* L. (*Zygophyllaceae*); *Banisteria caapi* Spruce. (*Malpighiaceae*); and *Banisteriopsis inebrians* Morton. (*Malpighiaceae*). It is also obtained from the fruit of *Passiflora incarnata* L. (*Passifloraceae*).

Chemical Structure



Harmine

7-Methoxy-1-methyl-9H-pyrido [3, 4-b] indole; ($C_{13} H_{12} N_2 O$).

* Rainhard *et al.* *Phytochemistry*, 7, 503, (1968).

Isolation Harmin may be isolated from the seeds of *Peganum harmala* L. (*Zygophyllaceae*) by the method suggested by Reinhard *et al.*

Characteristic Features

1. It is obtained as slender, orthorhombic prisms from methanol having mp 261°C (decomposition).
2. It sublimes and has pKa value of 7.70.
3. It has uv_{max} (methanol): 241, 301, 336 nm (log ϵ 4.61, 4.21, 3.69).
4. It is found to be slightly soluble in water, ethanol, ether and chloroform.

Identification Tests


Harmine Hydrochloride Dihydrate (C₁₃H₁₂N₂O.HCl.2H₂O) It is obtained as crystals having mp 262°C (decomposition), but when anhydrous mp 321°C (decomposition), but when anhydrous mp 321°C. The aqueous solution exhibits a distinct blue fluorescence. It is found to be soluble in 40 parts of water and freely soluble in hot water.

Uses It finds its usage as a CNS-stimulant and also as a narcotic hallucinogen.

D. Elaeagnine

Biological Source It is obtained from the bark of *Elaeagnus angustifolia* Linn., (*Synonyms: E. hortensis* Bieb.) (*Elaeagnaceae*).

Chemical Structure




Elaeagnine

Biosynthesis of Elaeagnine, Harman, Harmaline and Harmine The various steps involved in the biosynthesis of the above mentioned *four* alkaloids are briefly summarized as under:

1. Tryptamine and acetyl carboxylic acid (*i.e.*, keto acid) undergoes a **Mannich-like reaction** to yield a **β -carboline carboxylic acid**, which on oxidative decarboxylation gives rise to 1-methyl β -carboline.
2. The resulting product on subsequent reduction gives rise to the alkaloid **elaeagnine**.
3. The **1-methyl β -carboline** upon mild oxidation yields the alkaloids *harman* with the elimination of a mole of water from the 6-membered heterocyclic nucleus.
4. The **1-methyl β -carboline** upon hydroxylation followed by methylation produces **harmaline**.
5. **Harmaline** on further oxidation generates **harmine** by the loss of a mole of water from the 6-membered pyridine ring at C-3 and C-4 positions.

The these steps are sequentially arranged in the following course of reactions:



7.2.8.3 Terpenoid Indole Alkaloids

Terpenoid indole alkaloids is perhaps one of the major groups of alkaloids in the plant kingdom which comprise of more than 3000 recognized alkaloids till date. Interestingly, they are found to be confined to *eight* different natural orders (*i.e.*, families), of which the *Apocynaceae*, the *Loganiaceae*, and the *Rubiaceae* are predominantly the best known sources.

However, it is pertinent to mention here that practically in all the structure a *tryptamine residue* is strategically located in the molecule; while the remaining fragment is invariably recognized as a C_9 or C_{10} residue.


The wisdom, relentless efforts and meticulous in-depth studies carried out by numerous groups of researchers dealing with plant substances across the globe ultimately led to *three* main *structural variants* entirely based on their good judgement and understanding namely:

- (a) *Corynanthe* Type *e.g.*, **ajmalicine** and **akuammicine**,
- (b) *Aspidosperma* Type *e.g.*, **tabersonine**, and
- (c) *Iboga* Type *e.g.*, **catharanthine**.

It has since been established beyond any reasonable doubt that the C_9 or C_{10} component present in the aforesaid *three* types of structural variants *i.e.*, *Corynanthe*, *Aspidosperma* and *Iboga* groups was definitely of the *terpenoid origin*. Besides, it was also confirmed that the *secoroid secologanin* was duly proclaimed to be the *terpenoid derivative*, which perhaps must have initially combined with the tryptamine residue of the molecule. From these scientific and logical evidences one may safely infer that the *three* above mentioned groups of alkaloids might be not only related but also rationalized in terms of rearrangements taking place exclusively in the *terpenoid portion* of the various structural variants as shown in the pathway given below.

Salient Features The salient features of the above pathway are as follows:

1. **Secologanin** (a *secoroid* and a *terpenoid* derivative) is formed through **geraniol** via **loganin**, which essentially contains the 10C-framework a typical characteristic feature of the *Corynanthe* moiety.
2. The resulting *Corynanthe* C-skeleton undergoes subsequent rearrangements to give rise to *Aspidosperma* and *Iboga* groups.



Pathways for Coryanthe, Aspidosperma and Iboga Type Alkaloids

- This intra-molecular rearrangement may be represented by detachment of a 3C-unit, which is subsequently reunited to the remaining C₇ fragment in one of the two different manners as shown in the pathway.
- Interestingly, where C₉ terpenoid units are complied with, the alkaloids usually, seem to have lost a C-atom marked in the circle, which exactly corresponds to the carboxylate function of secologanin molecule. Therefore, its ultimate elimination by way of hydrolysis/decarboxylation is now understood without any reasonable doubt.
- Thus, the *Coryanthe* type of C-skeleton yields *ajmalicine* and *akuammicine*.
- The *Aspidosperma* type of C-skeleton yields **tabersonine** and **vindoline**.
- The *Iboga* type of C-skeleton gives rise to *catharanthine*.

A few typical examples of terpenoid indole alkaloids, namely: **Ajmalicine** (Raubasine); Akuammicine; **Vindoline**; and **Catharanthine** shall be discussed below:


A. Ajmalicine

Synonyms Raubasine; Circolene; Hydrosarpan; Lamuran; Isoarteril;

Biological Sources It is obtained from the plants of *catharanthus lanceus* Pichon (Boj.) (*Apocynaceae*) (Lanceleaf Periwinkle); *Catharanthus roseus* (L.) G. Don (*Apocynaceae*) (**Periwinkle**, **Madagascar** or **Cape Periwinkle**, **Old Maid**]; leaves of *Mitragyna speciosa* Korth. (*Rubiaceae*)

(**Katum, Kutum, Krantum**); *Rauwolfia serpentina* (L.) Benth. (*Apocynaceae*) (**Rauwolfia, Chandra, Sarpaganda**); and bark of *Corynanthe johimbe* K. Schum., (*Rubiaceae*).

Chemical Structure



Ajmalicine

(19 α)-16, 17-Didehydro-19-methyl-oxayohimban-16-carboxylic acid methyl ester; (C₂₁H₂₄N₂O₃).

Isolation **Ajmalicine** may be isolated either from the bark of *Corynanthe johimbe* by the method suggested by Heinemann*, or from the roots of *Rauwolfia serpentina* by the procedure adopted by Hofmann.**

Characteristic Features

1. It is obtained as prisms from methanol which decompose at 257°C.
2. It has specific optical rotation $[\alpha]_D^{20} - 60^\circ$ (C = 0.5 in chloroform); $[\alpha]_D^{20} - 45^\circ$ (C = 0.5 in pyridine); and $[\alpha]_D^{20} - 39^\circ$ (C = 0.25 in methanol).
3. It exhibits uv_{max} (methanol): 227, 292 nm (log ϵ 4.61, 3.79).

Identification Tests

1. **Ajmalicine Hydrochloride (C₂₁H₂₄N₂O₃.HCl)**: It is obtained as leaflets from ethanol having mp 290°C (decomposed); $[\alpha]_D^{20} - 17^\circ$ (C = 0.5 in methanol); and is sparingly soluble in water or dilute HCl.
2. **Ajmalicine Hydrobromide (C₂₁H₂₄N₂O₃.HBr)**: It is obtained as diamond-shaped plates from methanol having mp 295-296°C.

Uses


1. It is mostly used as antihypertensive and anti-ischemic agent (both cerebral and peripheral).
2. It has a broad application in the relief of obstruction of normal cerebral blood flow.

B. Akuammicine

Biological Source It is obtained from the plant substance of *Catharanthus roseus* (L.) G. Don (*Apocyanaceae*) (**Periwinkle, Madagascar or Cape Periwinkle, Old Maid**); and also from the seeds of *Picralima klaineana*, Pierre, belonging to the natural order (*Apocyanaceae*).

* Heinemann, H., *Ber.* **67**, 15 (1934).

** Hofmann, A., *Helv. Chim. Acta.* **37**, 849, (1954).

Chemical Structure

Akuammicine

2, 16, 19-20-Tetradehydrocuran-17-oic acid methyl ester; ($C_{20}H_{22}N_2O_2$).

Characteristic Features

1. It is obtained as plates from a mixture of ethanol and water having mp 182°C.
2. Its physical parameters are: $[\alpha]_D^{16} - 745^\circ$ ($C = 0.994$ in ethanol); pKa 7.45; and uv_{max} (ethanol): 227, 330 and 330 nm ($\log \epsilon$ 4.09, 4.07, 4.24).


Identification Tests It forms the following derivatives:

1. **Akuaminicine Hydrochloride Dihydrate ($C_{20}H_{22}N_2O_2 \cdot HCl \cdot 2H_2O$):** It is obtained as leaflets from ethanol or water having mp 171°C; and has $[\alpha]_D^{21} - 610^\circ$ ($C = 1.430$ in ethanol).
2. **Akuaminicine Perchlorate Monohydrate ($C_{20}H_{22}N_2O_2 \cdot HClO_4 \cdot H_2O$):** It is obtained as needles from a mixture of ethanol and water having mp 134-136°C.
3. **Akuammicine Hydroiodide Monohydrate ($C_{20}H_{22}N_2O_2 \cdot HI, H_2O$):** It is obtained as square plates from water having mp 128°C.
4. **Akuammicine Methiodide:** It is obtained as crystals from water with mp 252°C.
5. **Akuammicine Nitrate:** It is obtained as needles from hot water having mp 182.5°C.

Uses The drug exhibits a slight digitalis-like reaction; and is, therefore, believed to act as a heart poison.

C. Vindoline

Biological Sources It is obtained from the plant *Catharanthus roseus* (L.) G. Don (*Apocynaceae*) (**Periwinkle, Madagascar or Cape Periwinkle; Old Maid**). It is found to be the major alkaloid from the leaves of *Vinca rosea* Linn. (*Apocynaceae*).

Chemical Structure

Vindoline

(2 β ,3 β ,4 β ,5 α ,12 β ,19 α)-4-(Acetyloxy)-6, 7-didehydro-3-hydroxy-16-methoxy-1 methylaspido-permidine-3-carboxylic acid methyl ester; (C₂₅H₃₂N₂O₆).

Isolation It is isolated from the leaves of *Vinca rosea* by the method suggested by Gorman *et al.**

Characteristic Features

1. **Vinodoline** is obtained in two forms: *first*, as needles from a mixture of acetone and petroleum ether having mp 164-165°C; and *secondly*, as prisms having mp 174-175°C.
2. It has $[\alpha]_D^{20}$ - 18° (chloroform) and dissociation constant pKa 5.5 in 66% DMF.
3. It has uv_{max} (ethanol): 212, 250, 304 nm (log ϵ 4.49, 3.74, 3.57).


Identification Tests It gives specific derivatives as.

1. **Vindoline Hydrochloride (C₂₅H₃₂N₂O₆.HCl)**: It is obtained as crystals from acetone having mp 161-164°C.
2. **Demethoxy Vindoline (C₂₄H₃₀N₂O₅) (Vindorosine, Vindolidine)**: It is obtained as needles from benzene and petroleum ether having mp 167°C. It has $[\alpha]_D^{16}$ -31° (Chloroform); and uv_{max} (methanol): 250, 302 nm (log ϵ 3.98, 3.52).

D. Catharanthine

Biological Sources It is obtained in the plant of *Catharanthus lanceus* Pichon (Boj.) (*Apocynaceae*) (**Lanceleaf Periwinkle**); and *Catharanthus roseus* (L.) G. Don (*Apocynaceae*) (**Periwinkle, Madagascar or Cape Periwinkle, Old Maid**). It is also found in *Vinca rosea* Linn. (*Apocynaceae*).

Chemical Structure



Cathranthine

3, 4-Didehydroibogamine-18-carboxylic acid methyl ester; (C₂₁ H₂₄ N₂ O₂).

Isolation It may be isolated from *Vinca rosea* Linn by the method recommended by Gorman *et al.***

Characteristic Feature

1. Its crystals obtained from methanol has mp 126-128°C.
2. It has uv_{max} (ethanol): 226, 284, 292 nm (log ϵ 4.56, 3.92, 3.88).
3. It has specific optical rotation $[\alpha]_D^{27}$ + 29.8° (CHCl₃); and dissociation constant pKa' 6.8.

* Gorman *et al.* *J. Am. Pharm. Assoc.* **48**, 256, (1959).

** M. Gorman *et al.*, *J. Arn. Pharm. Assoc. Sci. Ed.* **48**, 256 (1959).

Uses

1. Its pharmacological action resembles to that of *R. serpentina*.
2. It also shows beneficial growth inhibition effects in certain human tumors.
3. It is used as a diuretic.


Biosynthesis of Ajmalicine, Vindoline and Catharanthine The various steps involved in the biosynthesis of **ajmalicine**, **vindoline** and **catharanthine** are summarized below:

1. Condensation of secologanin with tryptamine in a Mannich-type reaction gives rise to the **tetrahydro- β -carboline system** and generates *strictosidine*.
2. The structural variations involved in converting the *Coryanthe* type skeleton into the corresponding *Aspidosperma* and *Iboga* types are evidently quite complex and are given in the pathway as under.
3. **Preakuammicine** is obtained from **strictosidine** via the *enol*-form of **dehydrogeissoschizine**.
4. **Preakuammicine** undergoes intramolecular rearrangement to produce **stemmadenine**, which subsequently gives rise to a hypothetical intermediate.
5. The hypothetical intermediate may be redrawn which undergoes Diel's-Alder type reaction to produce **catharanthine**.
6. Dehydrogeissoschizine yields **ajmalicine**.
7. The hypothetical intermediate gives rise to **vindoline** via **tabersonine**.

It is pertinent to mention here that the sequence of alkaloid formation has been proved initially by noting carefully which alkaloids become labelled as a feeding experiment progresses, but more recently it has been confirmed by suitable enzymatic experimental studies.

It is important to mention here that there exists a plethora of structural variants of terpenoid indole alkaloids which may be exemplified with the help of the following specific examples of certain potent alkaloids, namely:

- (i) **Yohimbine**: It is a carboxylic variant related to ajmalicine and appears to arise from dehydrogeissoschizine by an elaborated mechanism.
- (ii) **Reserpine**: It is a trimethoxybenzoyl ester of yohimbine-like alkaloid. It has an additional-OCH₃ moiety at C.-11 of the indole nucleus.
- (iii) **Rescinnamine**: It is a trimethoxycinnamoyl ester of yohimbine-like alkaloid. It also contains an additional methoxyl substituent on the indole-system at C-11.
- (iv) **Vinblastine**: The nucleophilic vindoline, C-5 of the indole nucleus is being activated adequately by the OMe at C-6, besides the N-atom of the indole moiety. The resulting adduct is subsequently reduced in the dihydropyridinium ring by the NADH-dependent 1, 4-addition, giving the substrate for hydroxylation. Its ultimate reduction gives rise to vinblastine.
- (v) **Vincristine**: It is the oxidized product of vinblastine whereby the inherent N-formyl group on the indoline fragment is transformed.
- (vi) **Strychnine**: The loss of one C from a preakuammicine-like structure via hydrolysis/ decarboxylation followed by an addition of the additional two C-atoms by means of aldol-condensation with the formyl moiety, complexed as a hemiacetal in the well-known **Wieland-Gumlich aldehyde**. The ultimate formation of strychnine from its hemiacetal is by virtue of the formation of both *ether* and *amide* linkages.



Pathways for Ajmalicine, Preakuammicine, Catharanthine and Vindoline


The above mentioned *six* structural variants of the **terpenoid indole alkaloids** shall now be discussed individually in the sections that follow.

A. Yohimbine

Synonyms Quebrachine; Corynine; Aphrodine;

Biological Sources It is found in the root bark of *Alchornea floribunda* Muell. Arg. (*Euphorbiaceae*) (**Niando**); plant* of *Catharanthus lanceus* Pichon (Boj.) (*Apocynaceae*) (**Lanceleaf Periwinkle**); bark of *Pausinystalia johimbe* (K. Schum.) (*Rubiaceae*) (**Yohimbe**); root of *Rauvolfia serpentina* (L.) Benth. (*Apocynaceae*) (**Rauvolfia, Chandra, Sarpaganda**); and plant of *Rauvolfia tetraphylla* L. (*Apocynaceae*) (**Pinque-Pinque**).

Chemical Structure



Yohimbine

(16a, 17a)-17-Hydroxy-yohimban-16-carboxylic acid methyl ester; (C₂₁H₂₆N₂O₃).

Characteristic Features

1. It is obtained as orthorhombic needles from dilute alcohol having mp 234°C.
2. Its specific optical rotations are: $[\alpha]_D^{20} + 50.9^\circ$ to $+ 62.2^\circ$ (ethanol); $[\alpha]_D^{20} + 108^\circ$ (pyridine); and $[\alpha]_{546}^{20} + 129^\circ$ (C = 0.5 in pyridine).
3. It has uv_{max} (methanol): 226, 280, 291 nm (log ϵ 4.56, 3.88, 3.80).
4. It is freely soluble in ethanol, chloroform, hot benzene; moderately soluble in ether; and sparingly soluble in water.

Identification Tests

Yohimbine Hydrochloride (C₂₁H₂₆N₂O₃.HCl) (Aphrodyne, Yocon, Yohimex, Yohidrol): It is obtained as orthorhombic plates or prisms from ethanol which decompose at 302°C. Its specific optical rotation $[\alpha]_D^{22} + 105^\circ$ (water). It is found to be soluble in nearly 120 ml water, 400 ml ethanol, and the aqueous solution is almost neutral.

Uses

1. It is an adrenergic blocking agent, which has been used extensively in angina pectoris and arteriosclerosis.
2. It has been used successfully for the treatment of impotency in patients with vascular or diabetic problems.
3. It is invariably employed as a pharmacological probe for the study of α_2 -adrenoreceptor.


* Emboden reported that this plant contains upto 5% **yohimbin**.

B. Reserpine

Synonyms Crystoserpine; Eskaserp; Rau-sed; Reserpoid; Rivasin; Serfin; Sandril; Sedaraupin; Serpasil; Serpine; Serpasol; Serpiloid.

Biological Sources It is obtained from the plant *Catharanthus roseus* (L.) G. Don (*Apocynaceae*) (**Periwinkle, Madagascar or Cape Periwinkle, Old Maid**); root of *Rauwolfia serpentina* (L.) Benth (*Apocynaceae*) (**Rauwolfia, Chandra, Sarpaganda**); root of *Rauwolfia tetraphylla* L. (*Apocynaceae*) (**Pinque-Pinque**); and from the plant of *Vinca minor* L. (*Apocynaceae*) (**Periwinkle**).

Chemical Structure



Reserpine

(3 β , 16 β , 17 α , 18 β , 20 α)-11, 17-Dimethoxy-18-[(3, 4, 5-trimethoxy benzoyl)oxy] yohimban-16-carboxylic acid methyl ester; (C₃₃ H₄₀ N₂ O₉);

Isolation Reserpine may be isolated by adopting the following steps in a sequential manner:

1. The powdered and sieved roots are allowed to swell in a NaHCO₃ solution (10% w/v) for a period of 10-12 hours. The resulting solution is extracted with benzene, until the extracts give a weak positive reaction with HgI₂.
2. The combined benzene extracts are concentrated and ether is added to the benzene solution. The resulting mixture is extracted with dilute HCl. The combined acidic solution is washed with ether, filtered and extracted with chloroform in a successive manner.

Note: The chloroform will specifically extract the weakly basic alkaloids, such as: **Reserpine and Rescinnamine**.

3. The combined chloroformic extract is washed subsequently with 10% (w/v) sodium carbonate solution and followed by water so as to get rid of any *free acids* present. The resulting extract is finally evaporated to dryness under vacuo.
4. The residue is dissolved in anhydrous methanol and seeded with a pure crystal of **reserpine** and allowed to cool gradually when reserpine will crystallize out.
5. However, **rescinnamine, deserpidine** and other minor weakly basic alkaloids could be obtained from the mother liquor conveniently.
6. The mother liquor is evaporated to dryness, and the residue is dissolved in the minimum quantity of benzene and subjected to column chromatography over a column packed with acid-washed

alumina. The alkaloids are eluted in the different fractions by making use of benzene, chloroform, methanol (10%) in a sequential manner.

Characteristic Features

1. It is obtained as long prisms from dilute acetone which get decomposed at 264-265°C; (decomposes at 277-277.5°C in an evac-tube).
2. Its specific optical rotations are: $[\alpha]_D^{23} - 118^\circ$ (CHCl₃); $[\alpha]_D^{26} - 164^\circ$ (C = 0.96 in pyridine); $[\alpha]_D^{26} - 168^\circ$ (C = 0.624 in DMF).
3. It has uv_{max} (CHCl₃): 216, 267, 295 nm (61700, 17000, 10200).
4. **Reserpine** is weakly basic in nature, pKa 6.6.
5. It is found to be freely soluble in chloroform (~ 1g/6 ml), glacial acetic acid, methylene chloride; soluble in benzene, ethyl acetate; slightly soluble in acetone, methanol, ethanol (1g/1800 ml), ether, in aqueous solutions of citric and acetic acids; and very sparingly soluble in water.

Identification Tests

1. Most solutions of reserpine upon standing acquire a distant yellow colouration and a marked and pronounced fluorescence; especially after the addition of an acid or upon exposure to light.
2. **Reserpine Hydrochloride Hydrate (C₃₃H₄₀N₂O₉.HCl.H₂O)**: It is obtained as crystals which decompose at 224°C.

Uses


1. It is a hypotensive drug which exhibits strong hypotensive and sedative activity.
2. It is also employed to alleviate mild anxiety conditions *i.e.*, the drug shows a mild tranquillizing effect.

C. Rescinnamine

Synonyms Reserpinine; Anaprel; Apoterin S; Cartric; Cinnaloid; Moderil;

Biological Sources It is obtained from the roots of *Rauvolfia serpentina* (L.) Benth. (*Apocynaceae*) (**Rauvolfia, Chandra, Sarpaganda**).

Chemical Structure



Rescinnamine

3, 4 5-Trimethoxy-cinnamic acid ester of methyl reserpate; (C₃₅ H₄₂ N₂ O₉).

Isolation Rescinnamine may be isolated from step (5) onwards as described under **Morphine**.

Characteristic Features


1. It is obtained as fine needles from benzene having mp 238-239°C (under vacuum).
2. Its specific optical rotation is $[\alpha]_D^{24} - 97^\circ$ (C = 1 in chloroform).
3. It has uv_{max} (methanol): 228, 302 nm (log ϵ 4.79, 4.48).
4. **Solubility Profile:** It is moderately soluble in methanol, benzene, chloroform and other organic solvents; and practically insoluble in water.

Uses It is mostly used as an antihypertensive.

D. Vinblastine

Synonyms Vincalukoblastine; VLB; 29060-LE;

Biological Source It is obtained from *Vinca rosea* Lin.. (*Apocynaceae*).

Chemical Structure

Isolation It may be isolated from *Vinca rosea* Linn., either by the method suggested by Noble *et al.** or by Gorman *et al.***

Characteristic Features

1. It is obtained as solvated needles from methanol having mp 211-216°C.
2. Its specific optical rotation $[\alpha]_D^{26} + 42^\circ$ (chloroform).
3. It has uv_{max} (ethanol): 214, 259 nm (log ϵ 4.73, 4.21).
4. It is soluble in alcohols, chloroform, acetone, ethyl acetate and is practically insoluble in water and petroleum ether.

Identification Tests It forms derivatives as given below:

1. **Vinblastine Sulphate (C₄₆H₅₈N₄O₉·H₂SO₄) (Exal, Vebe, Velban):** It is obtained as crystals mp 284-285°C. Its physical parameters are: $[\alpha]_D^{26} - 28^\circ$ (C = 1.01 in methanol); pKa₁ 5.4; pKa₂ 7.4. It has uv_{max} (methanol): 212, 262, 284, 292 nm (log ϵ 4.75, 4.28, 4.22, 4.18). One part is

* Noble *et al.* *Ann. N.Y. Acad. Soc.* **76**, Art 3, 892-894 (1958)

** Gorman *et al.* *J. Am. Chem. Soc.*, **81**, 4745, 4754, (1959).

soluble in 10 parts of water, 50 parts of chloroform; very slightly soluble in ethanol; and practically insoluble in ether.

2. **Vinblastine Dihydrochloride Dihydrate** ($C_{46}H_{58}N_4O_9 \cdot 2HCl \cdot 2H_2O$): It is obtained as crystals that decompose at 244-246°C.

Uses

1. The alkaloid is used for the treatment of a wide variety of neoplasms.
2. It is also recommended for generated Hodgkin's disease, lymphocytic lymphoma, hystiocytic lymphoma, mycosis fungoides, advanced testicular carcinoma, Kaposi's sarcoma, and choriocarcinoma and lastly the breast cancer unresponsive to other therapies.
3. It is effective as a single entity, however, it is normally given along with other neoplastic agents in combination therapy for the increased therapeutic effect without any noticeable additive toxicity.
4. It arrests mitosis at the metaphase.
5. It is found to be effective in the acute leukemia of children.

E. Vincristine

Synonyms Leurocristine; VCR; LCR.

Biological Sources It is also obtained from *Vinca rosea* Lin., (*Catharanthus roseus* G. Don) belonging to the natural order *Apocynaceae*.

Chemical Structure Please see the chemical structure under **Vinblastine**. It may also be named as: 22-Oxovincaleukoblastine.

Isolation **Vincristine** may be isolated from **Vinca rosea** Linn., by the method suggested by Svoboda.*

Characteristic Features

1. It is obtained as blades from methanol having mp 218-220°C.
2. Its specific optical rotation $[\alpha]_D^{25} + 17^\circ$; $[\alpha]_D^{25} + 26.2^\circ$ (ethylene chloride); pKa: 5.0, 7.4 in 33% DMF.
3. It has uv_{max} (ethanol): 220, 255, 296 nm ($\log a_m$ 4.65, 4.21, 4.18).

Identification Tests

Vincristine Sulphate ($C_{46}H_{56}N_4O_{10} \cdot H_2SO_4$) (**Vincrex, Oncovin, Vincosid, Kyocristine**): Its crystals are obtained from ethanol and is found to be unstable.

Uses

1. **Vincristine sulphate** is recommended for the treatment of acute lymphocytic leukemia, and in combination therapy in Hodgkin's disease, lymphosarcoma, reticulum cell sarcoma, neuroblastoma, Wilm's tumour and rhabdomyosarcoma.

Note: Viucristine sulphate being highly unstable; therefore, its refregerated storage in sealed ampules is absolutely essential.


2. It is broadly used as an antineoplastic agent.

* Svobada, *Lyoydia*, 24, 173 (1961)

F. Strychnine

Biological Sources It is abundantly found in the seeds of *Strychnos Nux Vomica* L. (*Loganiaceae*) (**Nux Vomica, Strychnine**); beans of *Strychnos ignatti* Berg. (*Loganiaceae*); roots of *S. cinnamomifolia* Thw.; seeds, bark and wood of *S. colubrina* Linn.; and plant of *S. malaccensis* Benth. (**Syn:** *S. gautheriana* Pierre).

Chemical Structure



Strychnine

Strychnidine-10-one; (C₂₁ H₂₂ N₂ O₂)

Salient Features

1. **Strychnine** contains *two* N-atoms even then it happens to be a mono-acidic base.
2. **Strychnine** readily forms a variety of salts, such as: nitrate, N₆-oxide, phosphate and sulphate. Interestingly, the N-atom which is specifically involved in the salt formation is the one that is located *farthest* from the aromatic benzene ring.
3. The second N-atom is strategically positioned as an amide nitrogen; and, therefore, it does not exhibit any basic characteristics.

Isolation **Strychnine** may be isolated from the seeds of *S. nux vomica* by adopting the following steps sequentially:

1. The seeds of nux vomica are dried, ground and sieved which are mixed with an adequate quantum of pure slaked lime and made into a paste by adding a requisite amount of water. The wet mass thus obtained is dried at 100°C and extracted with hot chloroform in a continuous extractor till the extraction is completed.
2. The alkaloids are subsequently removed from the chloroform solution by shaking with successive portions of dilute sulphuric acid (2N). The combined acid extracts are filtered to get rid of any foreign particles or residue.
3. To the resulting acidic filtrate added an excess of ammonia to precipitate the alkaloids (strychnine + brucine).
4. The precipitate is extracted with ethanol (25% v/v) several times which exclusively solubilizes *brucine*, and ultimately leaves *strychnine* as an insoluble residue.
5. The residue containing **strychnine** is filtered off and is finally purified by repeated recrystallization from ethanol.

Characteristic Features

1. It is obtained as brilliant, colourless cubes from a mixture of chloroform and ether having mp 275-285°C, and d¹⁸ 1.359.

2. Its specific optical rotation $[\alpha]_D^{18}$ -104.3° (C = 0.254 in ethanol); $[\alpha]_D^{25}$ -13° (C = 0.4 in chloroform).
3. Its dissociation constant pKa (25°) 8.26.
4. It has λ_{max} (95% ethanol); 2550, 2800, 2900 Å ($E_{1\text{cm}}^{1\%}$ 377, 130, 101).
5. **Solubility Profile:** 1g dissolves in 182 ml ethanol, 6.5 ml chloroform, 150 ml benzene, 250 ml methanol, 83 ml pyridine; and very slightly soluble in water and ether.
6. A solution of strychnine containing 1 part in 700,000 parts of water gives a distinct bitter taste.

Identification Tests Strychnine may be identified either by specific *colour tests* or by specific *derivatives*:

(a) Colour Tests

1. **Sulphuric Acid-Dichromate Test: Strychnine** (5-10 mg) when dissolved in a few drops of concentrated sulphuric acid and stirred with a crystal of pure potassium dichromate [$\text{K}_2\text{Cr}_2\text{O}_7$] it gives an instant reddish-violet to purple colouration.
2. **Mandelin's Reagent Test: Strychnine** or its corresponding salt when treated with **Mandelin's Reagent*** it gives rise to a violet to blue colouration.
3. **Ammonium Vanadate (V) Test: Strychnine** or its salt when treated with a saturated solution of ammonium vanadate, it produces a violet to blue colouration.
4. **Nitric Acid Test: Strychnine** on being treated with a trace of HNO_3 (conc.) yields an instant yellow colouration.

Note: Strychnine derivatives will also give this test except strychnine nitrate.
Note: A similar test with Brucine gives an intense orange-red colouration. It may be used to differentiate between strychnine and brucine.

(b) Strychnine Derivatives: The various important **strychnine** derivatives are as given under:

1. **Strychnine Nitrate ($\text{C}_{21}\text{H}_{23}\text{N}_3\text{O}_5$):** It is obtained as colourless, odourless needles or white crystalline powder 1g dissolves in 42 ml water, 10 ml boiling water, 150 ml ethanol, 80 ml ethanol at 60°C, 105 ml chloroform, 50 ml glycerol; and insoluble in ether. It shows a pH ~ 5.7.
2. **Strychnine N⁶ Oxide ($\text{C}_{21}\text{H}_{22}\text{N}_2\text{O}_3$):** It is obtained as monoclinic prisms from water which decompose at 207°C. It has pK value 5.17. It is found to be freely soluble in ethanol, glacial acetic acid, chloroform; fairly soluble in water; sparingly soluble in benzene; and practically insoluble in ether and petroleum ether.
3. **Strychnine Phosphate ($\text{C}_{21}\text{H}_{25}\text{N}_2\text{O}_6\text{P}$):** It is usually obtained as its dihydrate salt ($\text{C}_{21}\text{H}_{25}\text{N}_2\text{O}_6\text{P} \cdot 2\text{H}_2\text{O}$) which is colourless or white crystals or white powder. 1g dissolves in slowly in ~ 30 ml water, more soluble in hot water, and slightly soluble in ethanol. The aqueous solution is acidic to litmus.
4. **Strychnine Sulphate ($\text{C}_{42}\text{H}_{46}\text{N}_4\text{O}_8\text{S}$):** It normally crystallizes as pentahydrate [$2\text{C}_{21}\text{H}_{22}\text{N}_2\text{O}_2 \cdot \text{H}_2\text{SO}_4 \cdot 5\text{H}_2\text{O}$]. It is colourless, odourless, very bitter crystals or white crystalline powder. It effloresces in dry air and loses all its water of crystallization at 100°C. It shows mp

* **Mandelin's Reagent** Dissolve 0.1g of ammonium vanadate in 10 ml of hot water and add to it 1-2 ml of conc. H_2SO_4 . Filter and preserve the solution.


when anhydrous $\sim 200^{\circ}\text{C}$ with decomposition. 1g dissolves in 35 ml water, 7 ml boiling water, 81 ml ethanol, 26 ml ethanol at 60°C , 220 ml chloroform, 6 ml glycerol, and insoluble in ether. A 1 : 100 solution shows pH 5.5.

5. **Strychnine Gluconate Pentahydrate ($\text{C}_{27}\text{H}_{34}\text{N}_2\text{O}_9 \cdot 5\text{H}_2\text{O}$):** Its crystals darken above 80°C . It is soluble in 2 parts water ~ 40 parts ethanol. The aqueous solution is found to be neutral.
6. **Strychnine Glycerophosphate Hexahydrate ($\text{C}_{45}\text{H}_{53}\text{N}_4\text{O}_{10}\text{P} \cdot 6\text{H}_2\text{O}$):** 1g dissolves in ~ 350 ml water, ~ 310 ml ethanol; slightly soluble in chloroform; and very slightly soluble in ether.
7. **Strychnine Hydrochloride Dihydrate ($\text{C}_{21}\text{H}_{23}\text{ClN}_2\text{O}_2 \cdot 2\text{H}_2\text{O}$):** It is obtained as trimetric prisms which are efflorescent in nature. 1g dissolves in ~ 40 ml water, ~ 80 ml ethanol, and insoluble in ether. The pH of a 0.01 M solution is 5.4.


Uses

1. **Strychnine** is extremely interesting pharmacologically and is regarded as a valuable tool in both physiologic and neuroanatomic research.
2. It is extremely toxic, and functioning as a central stimulant.
3. It causes excitation of all parts of the central nervous system and blocks inhibitory spinal impulses at the post synaptic level. This may lead to an exaggeration in reflexes ultimately leading to *tonic convulsions*.
4. The drug is rarely used in modern medical practice but is utilized as a *vermin killer i.e.*, animal or insect killer.
5. It is used chiefly in poison baits for rodents.

Biosynthesis of Yohimbine, Reserpine, Rescinnamine, Vinblastine, Vincristine and Strychnine Dehydrogessoschizine (*keto-form*) undergoes isomerization by means of the nucleophilic attack on to carbonyl through a conjugated system, which subsequently forms an onium ion that upon reduction produces **yohimbine** as shown below:



Reserpine and **deserpindine** are essentially the trimethoxybenzoyl esters of yohimbine-type alkaloids; whereas, **rescinnamine** is a trimethoxycinnamoyl ester. Interestingly, both **reserpine** and **rescinnamine** contain an additional methoxyl moiety present strategically on the indole ring system at C-11, which is accomplished by virtue of hydroxylation and methylation at a late stage along the pathway. A predominant and characteristic feature of these alkaloids is that they exhibit the *opposite stereochemistry* at C-3 to **yohimbine** and **strictosidine** as depicted below:




The biosynthetic pathway leading to **vinblastine** and **vincristine** is supposedly involve the following vital steps:

1. An oxidative reaction on **catharanthine**, catalysed by an enzyme peroxidase, thereby producing a peroxide that aptly loses the peroxide as a leaving group, ultimately breaking a carbon-carbon covalent bond as shown in the diagram given below.
2. The intermediate electrophilic ion is attacked on to the conjugated iminium system by the **vindoline**, whereby C-5 of the indole nucleus being appropriately activated by the $-OCH_3$ moiety located at C-6, and also by the N-atom present in the indole ring.
3. The resulting adduct is subsequently reduced in the dihydropyridinium ring by $NADH^*$ -dependent 1, 4-addition thereby giving rise to the substrate for hydroxylation.
4. Ultimately, reduction of the above resulting product generates **vinblastine**.
5. The oxidized product from vinblastine, with its N-formyl moiety rather than N-methyl on the vindoline fragment, may finally yield **vincristine**.


The biosynthetic pathway leading to **strychnine** essentially comprise of the following steps, namely:

1. **Preakuammicine** loses one C-atom *via* hydrolysis followed by decarboxylation.
2. Addition of the *two* extra C-atoms is accomplished by means of Aldol-condensation reaction with acetyl-CoA, whereby it yields the **Wieland-Gumlich aldehyde** as a complexed hemiacetal form.

* **NADH** = Nicotinamide adenine dinucleotide (reduced form).




3. The subsequent construction of ether and amide linkages gives rise to the formation of **strychnine** from the above hemiacetal as shown below.



7.2.8.4 Quinoline Alkaloids

A good number of very prominent and remarkable examples of the ‘**quinoline-alkaloids**’ derived from *tryptophan* are nothing but the modifications of the terpenoid indole alkaloids commonly found in the genus *Cinchona* belonging to the natural order *Rubiaceae*.


Interestingly, more than twenty alkaloids have been isolated and characterized from the bark of *Cinchona calisaya* and *Cinchona ledgeriana*, very commonly known across the globe as the **Yellow Cinchona**; besides the other equally well-known species *Cinchona succirubra*, popularly known in trade as the **Red Cinchona**. However, the four long prized and most popular **quinoline alkaloids** known for their antimalarial activities are namely: **quinine**, **cinchonine**, **quinidine**, and **cinchonidine**. These alkaloids shall now be described individually in the sections that follow. It is worthwhile to state here that these structures are not only unique but also remarkable wherein the indole nucleus is replaced by a quinoline system through an intramolecular rearrangement as given below:



A. Quinine

Biological Sources The *cinchona* species (*Rubiaceae*) specifically contains **quinine** in the bark upto 16% (mostly 6-10%) in a variety of its species, namely: *Cinchona calisaya* Wedd.; *C. ledgeriana* Moens ex Trimen; *C. officinalis* Linn. f.; *C. robusta* How.; and *C. succirubra* Pavon ex Klotzsch. The representative samples of dried **cinchona**, **cinchona bark** or **peruvian bark** is found to contain nearly 0.4 to 4% **quinine**.

Chemical Structure



Quinine

(8a, 9R)-6'-Methoxycinchonan-9 ol; (C₂₀H₂₄N₂O₂).

Isolation of Quinine, Cinchonine, Cinchonidine and Quinidine The isolation of all the *four* important quinoline alkaloid, such as: **quine, cinchonine; cinchonidine** and **quinidine** may be accomplished by adopting the following steps carefully and sequentially.

Step I: The cinchona bark is dried, powdered, sieved and treated with calcium oxide (slaked lime), NaOH solution (10% w/v) and water and kept as such for 6-8 hours.

Step II: The resulting mixture is treated with benzene in sufficient quantity and refluxed for 12-16 hours. The mixture is then filtered while it is hot.

Step III: The hot filtrate is extracted successively with 6N. sulphuric acid. The mixture of alkaloidal bisulphate is heated upto 90°C and maintained at this temperature upto 20-30 minutes.

Step IV: The resulting solution is cooled to room temperature and made alkaline by the addition of solid pure sodium carbonate till a pH 6.5 is attained.

Step V: The alkaloidal sulphate solution thus obtained is treated with sufficient quantity of activated charcoal powder (1g per 1L), boil, shake vigorously and filter.

Step VI: Cool the hot filtrate slowly in a refrigerator (2-10°C) overnight and again filter. Collect the residue and the filtrate separately.

Step VII: The residue (or precipitate) of quinine sulphate is boiled with water and made alkaline by adding cautiously solid sodium carbonate. The resulting precipitate is that of **quinine**.

Step VIII: The filtrate obtained from step-VI comprises of **cinchonine, cinchonidine** and **quinidine**; which is treated with NaOH solution (10% w/v) very carefully to render it *just alkaline*. It is successively extracted with adequate quantity of ether. The lower (aqueous layer) and the upper(ethereal layer) are collected separately.

Step IX: The aqueous layer contains **cinchonine**. It is evaporated to dryness in a Rotary Film Evaporator, extracted with absolute ethanol, decolourized with activated charcoal powder and allow it to crystallize slowly in a refrigerator (2-10°C) overnight. The crystals of **cinchonine** are obtained.

Step X: The ethereal layer obtained in step-VIII contains **quinidine** and **cinchonidine**. It is extracted with dilute HCl (2N) several times till a drop of the extract on evaporation does not give a positive test for alkaloids. Neutralize the combined acidic extract by adding solid sodium potassium tartrate *carefully*. Filter the resulting mixture and collect the *precipitate* and the *filtrate* separately.

Step XI: The precipitate of **cinchonidine tartrate** is treated with dilute HCl carefully. The resulting solution of alkaloid hydrochloride is made alkaline by the addition of dilute ammonium hydroxide when **cinchonidine** is obtained as a precipitate.

Step XII: The filtrate obtained from Step-X contains quinidine tartrate which is treated with solid potassium iodide powder carefully till the whole of quinidine gets precipitated as quinidine hydroiodide salt. It is filtered and the solid residue is finally treated with dilute NH_4OH to obtain the precipitate of **quinidine**.

Characteristic Features

1. It is obtained as triboluminescent, orthorhombic needles from absolute ethanol having mp 177° (with some decomposition).
2. It sublimes in high vacuum at $170\text{-}180^\circ\text{C}$.
3. Its specific optical rotations are: $[\alpha]_D^{15} - 169^\circ$ ($C = 2$ in 97% ethanol); $[\alpha]_D^{17} - 117^\circ$ ($C = 1.5$ in chloroform); $[\alpha]_D^{15} - 285^\circ$ ($C = 0.4$ M in 0.1 N H_2SO_4).
4. Its dissociation constants are: pK_1 (18°) 5.07; and pK_2 9.7.
5. The pH of its saturated solution is 8.8.
6. It gives a distinct and characteristic blue fluorescence which is especially strong in dilute sulphuric acid.
7. **Solubility Profile:** 1 g dissolves in 1900 ml water; 760 ml boiling water; 0.8 ml ethanol; 80 ml benzene; 18 ml benzene at 50° ; 1.2 ml chloroform; 250 ml by ether; 20 ml glycerol; 1900 ml of 10% ammonia water; and almost insoluble in petroleum ether.

Identification Tests Quinine may be identified either by a series of **Colour Tests** or by the formation of several known derivatives having characteristic features; and these shall be discussed separately as under:

(a) Colour Tests: These are, namely

1. **Oxygenated Acids:** Oxygenated acids, such as: sulphuric acid or acetic acid gives a strong blue fluorescence with quinine. This test is very sensitive even in extremely dilute solutions.

Note: Halogen quinine compounds and hydrochloride salts of quinine do not give fluorescence in solution.

2. **Herpathite Test:** To a boiling mixture of quinine (0.3g) in 7.5 ml glacial acetic acid, 3 ml ethanol (90% v/v) and 5 drops of concentrated H_2SO_4 , add 3.5 ml of I_2 solution (1% w/v) in ethanol, crystals of *iodosulphate of quinine* or **Herpathite*** separates out on cooling. The crystals thus obtained exhibit metallic lustre, appears dark in reflected light and alive-green in transmitted light.
3. **Thalleioquin Test:** When a few drops of bromine water are added to 2 or 3 ml of a weakly acidic solution of quinine salt, followed by the addition of 0.5-1.0 ml of strong ammonia solution, it produces a distinct characteristic emerald green colouration. It is an extremely sensitive colour test which may detect quinine even upto a strength as low as 0.005% (w/v). The end coloured product is known as **thalleioquin** for which the exact chemical composition is not yet known.

* **Herpathite** The iodo sulphate of quinine (or sulphate of iodo-quinine) is now known as **Herpathite** after the name of its discoverer [Formula: $\text{B}_4 \cdot 3\text{H}_2\text{SO}_4 \cdot 2\text{HI} \cdot \text{I}_4 + 3\text{H}_2\text{O}$]

Note: (a) This test is given by quinidine and also by other Remijia alkaloids e.g., cupreine.
(b) Both cinchonine and cinchonidine do not respond to the Thalleioquin Test.

- Erythroquinine Test (or Rosequin Test):** Dissolve a few mg of quinine in dilute acetic acid, add to it a few drops of bromine water (freshly prepared), followed by a drop of a 10% (w/v) solution of potassium ferrocyanide [$K_4 Fe(CN)_6$]. Now, the addition of a drop of concentrated NH_4OH solution gives rise to a red colouration instantly. If shaken quickly with 1-2 ml of chloroform, the red colouration is taken up by the lower chloroform-layer.

(b) **Derivatives/Salts of Quinine:** These are as follows:


- Quinine Trihydrate:** It is obtained as a microcrystalline powder having mp $57^\circ C$. It effloresces and loses one mol of water in air, two moles of water over H_2SO_4 , and becomes anhydrous at $125^\circ C$.
- Quinine Bisulphate Heptahydrate ($C_{20}H_{24}N_2O_2 \cdot H_2SO_4 \cdot 7H_2O$)** [*Synonyms:* **Quinbisan, Dentojel, Biquinate**): It is obtained as very bitter crystals or crystalline powder. It effloresces on exposure to air and darkens on exposure to light. 1 g dissolves in 9 ml water, 0.7 ml boiling water, 23 ml ethanol, 0.7 ml ethanol at $60^\circ C$, 625 ml chloroform, 2500 ml ether, 15 ml glycerol and having a pH 3.5.
- Quinine Dihydrochloride ($C_{20}H_{24}N_2O_2 \cdot 2HCl$)** (*Synonyms:* **Quinine dichloride; Acid quinine hydrochloride; Quinine bimuriate**): It is obtained as a powder or crystals having a very bitter taste. 1g dissolves in about 0.6 ml water, 12 ml ethanol; slightly soluble in chloroform; and very slightly soluble in ether. The aqueous solutions are found to be strongly acidic to litmus paper (pH about 2.6).
- Quinine Hydrochloride Dihydrate ($C_{20}H_{24}N_2O_2 \cdot HCl \cdot 2H_2O$)**: It is obtained as silky needles having a bitter taste. It effloresces on exposure to warm air. It does not lose all its water below $120^\circ C$. 1 g dissolves in 16 ml water, in 0.5 ml boiling water, 1.0 ml ethanol, 7.0 ml glycerol, 1 ml chloroform, and in 350 ml ether. A 1% (w/v) aqueous solution shows a pH 6.0-7.0.
- Quinine Sulphate Dihydrate [$(C_{20}H_{24}N_2O_2)_2 \cdot H_2SO_4 \cdot 2H_2O$]** (*Synonyms:* **Quinamm; Quinsan; Quine, Quinate**): It is obtained as dull needles or rods, making a light and readily compressible mass. It loses its water of crystallization at about $110^\circ C$. It becomes brownish on exposure to light. Optical rotation $[\alpha]_D^{15} - 220^\circ$ (5% solution in about 0.5 N . HCl). 1g dissolves in 810 ml water, 32 ml boiling water, 120 ml ethanol, 10 ml ethanol at $78^\circ C$; slightly soluble in ether and chloroform, but freely soluble in a mixture of 2 vols. chloroform and 1 vol. absolute ethanol. Its aqueous solutions are neutral to litmus. The pH of a saturated solution in 6.2.

Uses

- It is frequently employed as a flavour in carbonated beverages.
- It is used as an antimalarial agent.
- It is also employed as a skeletal muscle relaxant.
- It has been used to treat hemorrhoids and varicose veins.
- Quinine** is also used as a oxytotic agent.
- Quinine** is supposed to be prophylactic for flu.

Biosynthesis of Quinine A survey of literature reveals that the intrinsic details of the biosynthetic pathways are lacking; however, an assumed biogenetic process essentially involving the following steps:

1. **L-Tryptophan and secologanin yields strictosidine**, which upon hydrolysis and decarboxylation produces **coryantheal**.
2. **Coryantheal** undergoes intramolecular changes, *first*-by cleavage of C-N bond (*via* iminium), and *secondly*-by formation of an altogether new C-N bond (again *via* iminium). This gives rise to an intermediate.
3. The resulting intermediate undergoes further intramolecular changes to yield **cinchoninone** having a quinoline nucleus.
4. **Cinchoninone** in the presence of **NADPH*** reduces the carbonyl function and generates **quinine**:




* NADPH = Nicotinamide adenine dinucleotide phosphate (reduced form).

B. Cinchonine

Biological Sources It occurs in most varieties of cinchona bark as mentioned under **quinine** (section 'A'). Besides, **cinchonine** especially occurs in the bark of *Cinchona micrantha* R & P. belonging to the natural order *Rubiaceae*.

Chemical Structure



Cinchonine

(9S) - Cinchonan-9-ol; (C₁₉ H₂₂ N₂O)

Isolation The detailed method of isolation has been given under **quinine** (section 'A'). Besides, Rabe* has put forward another method of isolation of **cinchonine**.

Characteristic Features

1. **Cinchonine** is obtained as needles from ethanol or ether having mp 265°C.
2. It begins to sublime at 220°C.
3. Its specific optical rotation is $[\alpha]_D + 229^\circ$ (in ethanol).
4. **Solubility Profile:** 1g dissolves in 60 ml ethanol, 25 ml boiling ethanol, 110 ml chloroform, 500 ml ether; and practically insoluble in water.
5. It has two distinct dissociation constants: pK₁ 5.85 and pK₂ 9.92.

Identification Tests **Cinchonine** may be identified by forming its specific derivatives, namely:

1. **Cinchonine Hydrochloride Dihydrate (C₁₉H₂₂N₂O.HCl.2H₂O):** It is obtained as fine crystals. The mp of its anhydrous salt is 215 °C with decomposition. 1g dissolves in 20 ml water, 3.5 ml boiling water 1.5 ml alcohol, 20 ml chloroform; and slightly soluble in ether. The aqueous solution is almost neutral.
2. **Cinchonine Dihydrochloride (C₉H₂₂N₂O.2HCl):** It is usually obtained as white or faintly yellow crystals or crystalline powder. It is found to be freely soluble in water or ethanol.
3. **Cinchonine Sulphate Dihydrate [(C₁₉H₂₂N₂O)₂.H₂SO₄.2H₂O]:** It is commonly obtained as lustrous extremely bitter crystals. Its anhydrous salt has mp 198°C. 1g dissolves in 65 ml water, 30 ml hot water, 12.5 ml ethanol, 7 ml hot ethanol, 47 ml chloroform; and slightly soluble in ether. The aqueous solution is practically neutral.
4. **Epichonine [Synonyms (9R)-Cinchonan-9-ol]:** It has mp 83°C; and $[\alpha]_D^{22} + 120.3^\circ$ (C = 0.806 in ethanol).

Uses

1. It is used as an antimalarial agent.

* Rabe, *Ber.* 41, 63 (1908)


- It is employed as a tonic in waters, bitters and liqueurs.
- It is broadly used for febrifuge, schizonticide, stomachic, amebiasis, dysentery, flu, fever, and as a mild stimulant of gastric mucosa.

C. Quinidine

Synonyms Conquinine; Pitayine; β -Quinine;

Biological Source **Quinidine** is obtained from the various species of *Cinchona* as described under quinine (section 'A'). It is reported to be present in cinchona barks ranging between 0.25-3.0%.

Chemical Structure It is the dextrorotatory *stereoisomer* of **quinine**



Quinidine

(9S)-6'-Methoxycinchonan-9-ol; (C₂₀H₂₄N₂O₂).

Isolation **Quinidine** may be isolated from the cinchona bark by the method stated under **quinine** (section 'A').

Characteristic Features

- Quinidine** is obtained as triboluminescent crystals having mp 174-175°C after drying of the solvated crystals.
- Its specific optical rotations are: $[\alpha]_D^{15} + 230^\circ$ (C = 1.8 in chloroform); $[\alpha]_D^{17} + 258^\circ$ (ethanol); and $[\alpha]_D^{17} + 322^\circ$ (C = 1.6 in 2m HCl).
- It has two dissociation constants, namely: pK_1 (20°) 5.4; and pK_2 10.0.
- It gives a distinct and characteristic blue fluorescence in dilute sulphuric acid (2N).
- The uv absorption spectrum is identical with that of **quinine**.
- Solubility Profile:** 1 g gets dissolved in 2000 ml cold water, 800 ml boiling water, 36 ml ethanol, 56 ml ether, 1.6 ml chloroform; very soluble in methanol; and practically insoluble in petroleum ether.

Identification Tests The various derivatives of **quinidine** have specific characteristic features as enumerated below:

- Quinidine Sulphate Dihydrate [(C₂₀H₂₄N₂O₂)₂·H₂SO₄·2H₂O] (Synonyms **Quinidex; Quinicardine; Quinora; Extentabs; Cin-Quin**):**
- It is mostly obtained as white, very bitter, odorless, fine crystals which is frequently cohering in masses. It does not lose all of its water of crystallization below 120°C. It has been found to darken on exposure to light. It has $[\alpha]_D^{25} \sim + 212^\circ$ (in 95% ethanol); and $\sim + 260^\circ$ (in dilute HCl). The pH of a 1% (w/v) solution between

6.0-6.8. Its pKa values are : 4.2 and 8.8. 1 g dissolves in 90 ml water, 15 ml boiling water, 10 ml ethanol, 3 ml methanol, 12 ml chloroform; and insoluble in ether and benzene.

Note: Quinidine sulphate dihydrate is the salt of an alkaloid obtained either from various species of Cinchona and their hybrids, or from Cuprea bark, obtained from Remijia pedunculata and Remijia purdieana belonging to the natural order Rubiaceae.

2. **Quinidine Gluconate ($C_{26}H_{36}N_2O_9$) (Synonyms Quinaglute; Duraquin; Gluconic acid quinidine salt):** It is obtained as crystals having mp 175-176.5°C; and soluble in 9 parts of water and 60 parts of ethanol.
3. **Quinidine Polygalacturonate ($C_{20}H_{24}N_2O_2 \cdot C_6H_{10}O_7 \cdot H_2O$) [Synonyms Galactoquin; Cardioquin; Naticardina):** It is obtained as an amorphous powder mp 180°C (decomposes). The anhydrous substance is found to be insoluble in methanol, ethanol, chloroform, ether, acetone, dioxane; and soluble in 40% methanol or ethanol: 12%; in water at 25°C: ~ 2%.
4. **Quinidine Hemipentahydrate:** It is obtained as prisms from dilute ethanol, mp ~ 168°C, and loses 1/2 H₂O on exposure to air.
5. **Quinidine Hydrogen Sulphate Tetrahydrate ($C_{20}H_{24}N_2O_2 \cdot H_2SO_4 \cdot 4H_2O$) (Synonyms Kiditard; Kinichron; Kinidin Durules; Quiniduran; Chinidin - Duriles; Quinidine Bisulphate):** It is obtained as rods which is soluble in 8 parts of water and emitting a distinct blue fluorescence.
6. **Neutral Hydroiodide of Quinidine ($C_{20}H_{24}N_2O_2 \cdot HI$):** It is obtained as a crystalline powder when KI is added to a neutral aqueous solution of a quinidine salt. It is very sparingly soluble in water (1 part in 1250 parts at 15°C). It is found to be much less soluble than that of the other cinchona alkaloids.

Quinidine also gives a specific colour test as given below:

Ferrocyanide Test for Quinidine A small quantum (10-15 mg) of a **quinidine salt** is mixed thoroughly with 0.5-1.0 ml of freshly prepared bromine water in an evaporating dish. The contents are transferred carefully into a test tube with the help of 1 ml of distilled water. To this is added 1 ml of chloroform, contents shaken and then allowed to stay for a few minutes. A few drops of a 10% (w/v) solution of potassium ferrocyanide [$K_4 Fe (CN)_6$] and 3 ml of a 5N. NaOH solution are added with continuous shaking. The chloroform layer attains a red colour.

Note: Quinine or its salt under identical treatment gives a negative test, and hence it may be used to distinguish between quinidine and quinine.

Uses

1. It is used as an antiarrhythmic agent (Class 1A)*.
2. It finds its applications as an antimalarial drug.
3. It is most commonly employed to treat various cardiac arrhythmias, namely: atrial flutter, AV junctional and ventricular contractions, atrial and ventricular tachycardia, atrial fibrillation, and premature atrial condition.


* **Class 1A Antiarrhythmic Agent** When the antiarrhythmic mechanisms is accomplished through membrane stabilization.

D. Cinchonidine

Synonyms Cinchovatine; α -Quinidine;

Biological Sources It is obtained in most varieties of the cinchona bark as described under quinine (section 'A'). It is, however, observed to be present especially in the bark of *Cinchona pubescens* Vahl. (*C. succirubra* Pav.) and *Cinchona pitayensis* Wedd., (*Rubiaceae*).

Chemical Structure



Cinchonidine

(8a, 9R)-Cinchonan-9-ol; (C₁₉ H₂₂ N₂ O).

Isolation Cinchonidine can be conveniently isolated from the bark of various species of *cinchona* as described explicitly under **quinine** (section 'A'). However, it may also be isolated by the method suggested by Leers.*

Characteristic Features

1. It is obtained as orthorhombic prisms or plates from ethanol having mp 210°C.
2. It has specific optical rotation $[\alpha]_D^{20} - 109.2^\circ$ (in ethanol).
3. **Solubility Profile:** It is found to be freely soluble in chloroform and ethanol; moderately soluble in ether; and practically insoluble in water.
4. It has two dissociation constants: pK₁ 5.80 and pK₂ 10.03.

Identification Tests Cinchonidine may be identified by preparing its specific derivatives that possess characteristic features, such as:

1. **Cinchonidine Dihydrochloride (C₁₉H₂₂H₂O.2HCl):** It is obtained as white or slightly yellow crystals or powder. It is freely soluble in ethanol and water.
2. **Cinchonidine Hydrochloride Dihydrate (C₁₉H₂₂N₂O.HCl.2H₂O):** It is obtained as a crystalline powder. It loses all of its water of crystallization at 120°C. It has $[\alpha]_D^{20} - 117.5^\circ$ (in water). It is soluble in 25 parts of cold water, more soluble in boiling water; soluble in chloroform and ethanol; and slightly soluble in ether. The aqueous solution is almost neutral in nature.
3. **Cinchonidine Sulphate Trihydrate [(C₁₉H₂₂N₂O)₂.H₂SO₄.3H₂O]:** It is obtained as silky, acicular crystals which effloresce on being exposed to air and get darkened in light. The mp of anhydrous salt is nearly 240°C with decomposition. 1g dissolves in 70 ml water, 20 ml hot water, 90 ml ethanol, 40 ml hot ethanol, 620 ml chloroform; practically insoluble in ether. The aqueous solution is more or less neutral.

** Leers, *Ann.*, **82**, 147 (1952)

4. **Epincinchonidine [Synonyms: (8 α , 9S)-Cinchonan-9-ol]**: It has mp 104°C; and $[\alpha]_D^{20} + 63^\circ$ (C = 0.804 in ethanol).

Uses It is mostly used as an antimalarial agent.

Totaquine **Totaquine** is nothing but a mixture of the total alkaloids of the well-known cinchona bark. It is invariably exploited as a 'cheap substitute' for quinine in an unethical practice in trade. It is found to contain not less than 7% and not more than 12% of quinine units anhydrous form; and not more than 80% of the total anhydrous crystallizable **cinchona alkaloids**.

The following table summarizes the characteristic features and specific tests for the *four* major **cinchona alkaloids**, namely: **Quinine, Quinidine, Cinchonine** and **Cinchonidine**.


Differences Among Four Major Cinchona Alkaloids

S. No.	Characteristics	Quinine	Quinidine	Cinchonine	Cinchonidine
1	Chemical Formula	Methoxy Vinyl rubanol	Methoxy Vinyl rubanol	Vinyl rubanol	Vinyl rubanol
2	Rotation of Alcoholic solution	(-)	(+)	(+)	(-)
3	Fluorescence in oxygenated acids	Blue	Blue	-	-
4	Thallioquin Test	+ve	+ve	-ve	-ve
5	Erythroquinine Test	+ve	+ve	-ve	-ve
6	Herrpathite Test	+ve	-ve	+ve	-ve
7	Solubility in (ml):				
	Water	1900	2000	-	-
	Chloroform	1.2	1.6	110	++
	Ethanol	0.8	36.0	60	++
	Glycerol	20.0	-	-	-
	Ether	250	56.0	500	+

Biosynthesis of Cinchonine, Quinidine and Cinchonidine The various sequential steps involved in the biosynthesis of **Cinchonine, Quinidine** and **Cinchonidine** are stated as under:

1. **Strictosidine** is obtained by the interaction of L-tryptophan and secologanin as already shown in the **Biosynthesis of Quinine**.
2. Strictosidine undergoes a molecular rearrangement to form an aldehyde which upon hydrolysis and decarboxylation yields *corynantheal*.
3. **Corynantheal** generates **cinchoninone** by virtue of *two* transformations; *first*: an intermediate formed due to the cleavage of C-N bond (*via* iminium) then formation of a new C-N bond (again *via* iminium); and *secondly*: cleavage of the indole C-N bond. The resulting product loses a molecule of water to yield **cinchoninone**.
4. **Cinchoninone** undergoes epimerization at C-8 *via* enol to form the stereoisomer, which upon interaction with NADPH gives rise to *chnchonine* and *quindine* respectively.
5. **Cinchonone** with direct interaction with NADPH gives rise to **cinchonidine** and **quinine** respectively.


The outline of the biosynthesis elaborated above from (1) through (5) may be summarized as depicted below:



Biosynthesis of Cinchonine, Quinidine and Cinchonidine


7.2.8.5 Pyrroloindole Alkaloids

The *indole* nucleus has two C-atoms in the heterocyclic portion, *viz.*, C-2 and C-3. Interestingly, both C-2 and C-3 may be regarded as nucleophilic in character. However, it has been established beyond any reasonable doubt that the reactions essentially involving C-2 appear to be the most common in **alkaloid biosynthesis**.



Indole

It is, however, pertinent to mention here that the nucleophilic character of C-3 has been duly exploited thereby generating the almost rare **pyrroloindole nucleus** as given below:



Pyrroloindole


Physostigmine is a typical example of this specific category of alkaloid which shall now be discussed in details as under:

Physostigmine

Synonyms Eserine; Cogmine;

Biological Sources It is obtained from the seeds of *Physostigma venenosum* Balf. (*Fabaceae*) (**Calabar Bean, Ordeal Bean**) yielding not less than 0.15% of the total alkaloids of **physostigma**.

Chemical Structure



Physostigmine

(3a*S-cis*)-1, 2, 3, 3α, 8, 8α-Hexahydro-1, 3α, 8-trimethylpyrrolol [2, 3-*b*] indol-5-yl methylcarbamate (ester); (C₁₅ H₂₁ N₃ O₂).

Isolation **Physostigmine** may be isolated by adopting the following *two* steps, namely:

Step I: The seeds are dried, powdered, sieved and extracted by continuous percolation with hot ethanol (95%) and the solvent is subsequently removed by distillation under vacuo. Water is added to the residue and the floating fatty layer is separated. The lower aqueous layer is subjected to alkalization with sodium carbonate and the liberated alkaloid is then extracted with ether successively.

Step II: The combined ethereal extract is then concentrated to a small volume and washed with 5% (w/v) sulphuric acid repeatedly unless and until the washings give a positive acidic reaction to litmus paper. To this aqueous acidic solution (containing the alkaloids as sulphates) is added an excess of a saturated solution of *sodium salicylate* when the physostigmine salicylate separates out as a crystalline product. The **physostigmine** may be recovered from the resulting salt by treating it with sodium carbonate followed by an immediate extraction with ether successively. The ether is evaporated in a Rotary Thin-Film Evaporator and the desired physostigmine is collected as prisms or clusters.

* Schwyzer, *Die Fabrikation Pharmazeutischer and Chemisch-Technischer Produkte* (Berlin, 1931) p 338.

However, **physostigmine** may also be isolated by the methods described by Schwyzer* and Cheminitius.*

Characteristic Features

1. It is obtained as orthorhombic sphenoidal prisms or clusters of leaflets from ether or benzene having mp 105-106°C. It is also available as an unstable, low melting form mp 86-87°C.
2. Its specific optical rotations are: $[\alpha]_D^{17} - 76^\circ$ ($C = 1.3$ in chloroform); and $[\alpha]_D^{25} - 120^\circ$ (benzene).
3. It has two dissociation constants: $pK_{a1} 6.12$, and $pK_{a2} 12.24$.
4. **Solubility Profile:** It is slightly soluble in water; soluble in ethanol, benzene, chloroform and oils.

Identification Tests **Physostigmine** may be identified either by specific colour tests or by preparing their derivatives as stated below:

(a) **Colour Tests:** These are as follows:

1. **Physostigmine** or its salts, a few mg, when warmed with 1 ml of strong ammonia solution it gives rise to a yellowish-red colouration. On further evaporation to dryness on a steam-bath, a bluish residue (eserine blue) is obtained that is soluble in ethanol forming a blue solution.
2. Both solid and solutions of **physostigmine** eventually turn red on being exposed to heat, light and air; and also on contact with traces of metals. This colour change indicates hydrolysis to **eseroline** and oxidation to **rubreserine**.
3. **Physostigmine** gives an instant blue colouration when treated with potassium ferricyanide $[K_3Fe(CN)_6]$ and a few drops of $FeCl_3$ solution (1% w/v).
4. **Physostigmine** produces a deep-yellow colouration on being heated with 0.5-1 ml of KOH solution (1% w/v).

Note: (a) This is a very sensitive test and can detect it upto 10 mcg level.

(b) Under controlled experimental parameters the intensity of the yellow colour produced may be measured spectrophotometrically at 470 nm and can serve as an assay method.

5. When a small quantity of **physostigmine** is heated in a porcelain basin on a steam bath with a drop or two of fuming HNO_3 , a yellow solution is obtained. The resulting solution on evaporation to dryness forms a green residue due to the formation of **chloreserine**, which is readily soluble in ethanol to give a green solution.
6. **Physostigmine** when treated with a solution of phosphomolybdic acid and ammonium meta vanadate in H_2SO_4 it gives rise to an emerald green colour.

(b) **Derivatives:** Major derivatives of **physostigmine** are:

1. **Physostigmine Salicylate ($C_{22}H_{27}N_3O_5$) (Antilirium):** It is obtained as acicular crystals having mp 185-187°C. It has uv_{max} (methanol): 239, 252, 303 nm ($\log \epsilon$ 4.09, 4.04, 3.78). 1g dissolves in 75 ml water at 25°C. The pH 0.5% (w/v) aqueous solution is 5.8. It is soluble in 16 ml ethanol; 5 ml of boiling ethanol; 6 ml of chloroform; and 250 ml of ether.
2. **Physostigmine Sulphate $[(C_{15}H_{21}N_3O_2)_2 \cdot H_2SO_4]$:** It is mostly obtained as deliquescent scales having mp 140°C (after drying at 100°C). 1g dissolves in 0.4 ml ethanol, 4 ml water, 1200 ml

* Cheminitius, *J. Prabt. Chem.* **116**, 59 (1927).

ether. The pH of 0.05 M aqueous solution is 4.7. The solutions of the sulphate salt are more prone to change colour than those of the corresponding salt of the salicylate.

3. **Physostigmine Sulphite** $[(C_{15}H_{21}N_3O_2)_2 \cdot H_2SO_3]$: The white powder is found to be freely soluble in ethanol and water. The aqueous solution is observed to remain colourless for a long duration.


Uses

1. It possesses a cholinergic (anticholinesterase) and miotic activities.
2. It was used earlier to treat myasthenia gravis; but now it is more frequently used for the eye.
3. It is employed as an antidote for reversing CNS and cardiovascular (*viz.*, arrhythmia and tachycardia) effects of excessive dosages with tricyclic antidepressants.
4. It helps in the contraction of the ciliary muscle of eye, and a decrease in the intraocular pressure produced by an increased out-flow of the aqueous humor.
5. Physostigmine is employed frequently in ophthalmology to treat glaucoma.

Biosynthesis of Physostigmine The various steps involved in the biosynthesis of **physostigmine** are as follows.

1. Tryptamine undergoes C-methylation at C-3 of the indole nucleus due to its nucleophilic character.
2. Formation of the '**third pyrrole**' ring takes place by virtue of the nucleophilic attack of the primary amine function on to the iminium ion.
3. Further substitution on the phenyl ring leads to the formation of physostigmine.

The above *three* steps are summarized as given below:



Biosynthesis of Physostigmine

7.2.8.6 Ergot Alkaloids


Ergot is a fungal disease very commonly and widely observed on a good number of wild as well as cultivated grasses, and is produced by different species of **claviceps**. This particular disease is usually

characterized by the formation of hard and seedlike ‘**ergots**’ in place of the normal seeds. However, these specific structures are frequently termed as **sclerotia**, which represent the ‘**resting stage**’ of the fungus.

The generic name, ‘**claviceps**’, usually refers to the club-like nature of the **sclerotium***, whereas *purpurea* signifies its purple colour. As these *sclerotia* are elongated and somewhat pointed in shape and appearance, hence the common name of *spurred rye* has been assigned to the drug.

Medicinal **ergot** is the dried sclerotium of the fungus *Claviceps purpurea* (Fries) belonging to the natural order *Clavicipitaceae* developed in the ovary of rye, *Secale cereale* (*Germinae/Poaceae*). There are certain other species of *Claviceps* which have been found to produce ergots in the ovaries of other member of *Graminae* and *Cyperaceae*.

In fact, there exist *four* main categories of **ergot alkaloids** which may be distinguished, namely: (a) **clavine alkaloids**, (b) **lysergic acids**, (c) **lysergic acid amides**, and (d) **ergot peptide alkaloids**. There are, in fact, *ten* **ergot peptide alkaloids** which are: **ergotamine**, **ergosine**, **ergocristine**, **ergocryptine**, **ergocornine**, **ergotaminine**, **ergosinine**, **ergocristinine**, **ergocryptinine**, and **ergocorninine**; however, the *last five* alkaloids being isomers of the *first five*. The aforesaid alkaloids are beautifully typified by a structure comprising of *four* components, *viz.*, lysergic acid, dimethylpyruvic acid, proline and phenylalanine strategically joined together in amide linkages as depicted below:



Interestingly, the poisonous properties of ergots in grain, specifically rye, for animal as well as human consumption, purposefully and unknowingly, have long been recognized. The dreadful causative agents are collectively termed as the ‘**ergot alkaloids**’, containing essentially an **indole nucleus**. These, group of alkaloids are also referred to as ‘**ergolines**’.

* **Sclerotium** A hardened mass formed by the growth of certain fungi. The sclerotium formed by ergot on rye is of medical importance due to its toxicity.

The *three* important and typical members of the ergot alkaloids (ergolines), namely: **ergonovine**, **ergotamine** and **lysergamde (ergine)** shall be discussed individually in the sections that follows:

A. Ergonovine


Synonyms Ergometrine; Ergobasine; Ergotocine; Ergostetrine; Ergotrate; Ergokline; Syntometrine.

Biological Sources It is obtained from the seeds of *Ipomea violaceae* Linn. (*Ipomea tricolor* Cav.) belonging to family *Convolvulaceae* (Morning glory, Tliltliltzen, Ololiuqui); and also from the dried seeds of *Rivea corymbosa* Hall. F. (*Convolvulaceae*) (Snakeplant).

The percentage of **Ergometrine** and **Ergine** present in the *Rivea* and *Ipomea* species are as given below:

Alkaloids	Rivea (%)	Ipomea (%)
Ergometrine	–	0.005
Ergine	0.0069	0.035

Chemical Structure



Ergonovine

[8β (s)]-9, 10-Diadehydro-N-(2-hydroxy-1-methylethyl)-6-methylergoline-8-carboxamide; (C₁₉H₂₃N₃O₂).

Isolation The following steps may be followed stepwise:

1. The seeds are dried, powdered, sieved and finally defatted with n-hexane in a Soxhlet apparatus.
2. The defatted material is extracted with hot dilute sulphuric acid (6N) successively. The acid extract is then treated with an excess of barium sulphate, and the barium is removed with CO₂ and subsequent filtration.
3. The resulting filtrate is then concentrated by evaporation under reduced pressure.
4. The concentrated solution is taken up in ethanol, made alkaline with NH₄OH and subjected to extraction with chloroform successively.
5. The resulting chloroform extract is further extracted with dilute H₂SO₄ (6N). The acidic solution is made alkaline with ammonia and saturated with NaCl and then extracted with ether several times.
6. The solvent is removed from the ether extract under vacuo leaving the alkaloidal residue.
7. **Ergonovine** may be recrystallized from acetone.

It may also be prepared from D-lysergic acid and L (+)-2-amino-1-propanol by the method of Stoll and Hofmann.*

Characteristic Features

1. **Ergonovine** is obtained as tetrahedral crystals from ethyl acetate, and as fine needles from benzene. It tends to form solvated crystals having mp 162°C.
2. It has specific optical rotation $[\alpha]_D^{20} + 90^\circ$ (in water).
3. Its dissociation constant is pK_a 6.8.
4. It is found to be freely soluble in lower alcohols, acetone and ethyl acetate; more soluble in water than the other principal alkaloids of ergot; and slightly soluble in chloroform.

Identification Tests *As per se* the **ergot alkaloids** may be identified either by general precipitation and colour reactions or by preparing their derivatives as stated below:

(a) Precipitation Reactions

- (i) The **ergot alkaloids** are readily precipitated by the alkaloidal reagents. However, Mayers reagent is regarded to be the most sensitive test whereby on opalescence in dilutions of 1 ppm can be obtained.
- (ii) Iodine solution in KI also gives an instant precipitate with very dilute solutions of **ergot alkaloids**.

(b) **Colour Tests:** The most vital colour tests are given as under:

- (i) **Keller's Test:** To a solution of the alkaloid in glacial acetic acid add a few mg of solid $FeCl_3$ and then add 1-2 ml of concentrated sulphuric acid along the side of the tube. The appearance of an intense blue colouration is accomplished at the junction of the two layers.
- (ii) **Van Urk Test:** When a solution containing an **ergot alkaloid** is mixed with **Van Urk Reagent****, it gives rise to a characteristic deep blue colouration.

Note: (a) **Van Urk Reagent may also be used in spraying developed paper chromatograms of the ergot alkaloids, and for this purpose 10% (v/v) HCl is used instead of H_2SO_4 .**

(b) **The spectrophotometric assay for total ergot alkaloids is also based on the blue colour given with Van Urk Reagent.**

- (iii) **Glyoxylic Acid Reagent Test:** **Ergot alkaloids** gives a blue colouration with the addition of Glyoxylic acid reagent and a few drops of concentrated H_2SO_4 .
- (iv) **Fluorescence Test:** The aqueous solution of the salts of ergot alkaloids produce a distinct blue fluorescence.

(c) **Derivatives of Ergonovine:** The various derivatives of **ergonovine** are as follows:

- (i) **Ergonovine Maleate (Ergometrine Maleate) ($C_{19}H_{23}N_2O_2 \cdot C_4H_4O_4$) [Synonyms Cornocentin; Ermetrine; Ergotrate Maleate]:** It is obtained as crystals that decompose at 167°C. It has specific optical rotation $[\alpha]_D^{25} + 48^\circ$ to $+ 57^\circ$. 1g dissolves in 36 ml water and 120 ml ethanol. It is almost insoluble in chloroform and ether.

* Stoll, A., and Hofmann, A., *Helv. Chim. Acta*, **26**, 944 (1943).

** **Van Urk Reagent** Mix together 0.125g of *para*-dimethylamino. benzaldehyde; 0.1 ml of $FeCl_3$ soln. (5% w/v), and 15% (v/v) H_2SO_4 to make 100 ml.

- (ii) **Methylergonovine Maleate** ($C_{20}H_{25}N_2O_2 \cdot C_4H_4O_4$): It is a semisynthetic homologue of ergonovine and prepared from lysergic acid and 2-aminobutanol. It is obtained as a white to pinkish-tan microcrystalline powder.
- (iii) **Ergonovine Tartrate Hydrate (Ergometrine Tartrate Hydrate)** [$(C_{19}H_{23}N_3O_2)_2 \cdot C_4H_6O_6 \cdot H_2O$] (**Basergin, Neofermergen**): It is obtained as crystals that are slightly soluble in water.

Uses

1. **Ergonovine** is used as an oxytocic.
2. **Ergonovine maleate** also acts as an oxytocic and produces much faster stimulation of the uterine muscles as compared to other ergot alkaloids.
3. **Methylergonovine maleate** is observed to act as an oxytocic whose actions are slightly more active and longer acting than **ergonovine**.

B. Ergotamine

Biological Source It is obtained from the seeds of *Claviceps purpurea* (Fr.) Tul. (*Hypocreales*) (**Ergot**).

Chemical Structure The chemical structure of **ergotamine** has been given in Section 7.2.8.6.

Isolation The method of Stoll* may be adopted as stated below:

1. The powdered dried ergot is first defatted with n-hexane or petroleum ether (40-60°)
2. The marc consisting of the defatted powdered ergot is thoroughly mixed with aluminium sulphate and water so as to fix the alkaloids by converting them into the double salts.
3. The resulting alkaloidal double salts are subjected to continuous extraction with hot benzene that removes the alkaloid exclusively on one hand; and the unwanted substances *e.g.*, **ergot oil**, soluble acid, neutral substances like-phytosterol, colouring matter and organic acids on the other.
4. The benzene is removed under vacuo and the residue thus obtained is stirred for several hours with a large volume of benzene and subsequently made alkaline by passing NH_3 gas.
5. The resulting solution is filtered and the benzene extract is concentrated under vacuo to approximately 1/50th of the original volume, whereupon ergotamine crystallizes out.
6. An additional quantity of ergotamine may also be crystallized from the mother liquor by treatment with petroleum ether.
7. Ergotamine may be further purified by crystallization from aqueous acetone.

Characteristic Features

1. It is obtained as elongated prisms from benzene that get decomposed at 212-214°C.
2. It usually becomes totally solvent-free only after prolonged heating in a high vacuum.
3. It is found to be highly hygroscopic in nature; and darkens and decomposes on exposure to air, heat and light.
4. It has specific optical rotation $[\alpha]_D^{20} - 160^\circ$ (chloroform).
5. It is soluble in 70 parts methanol, 150 parts acetone, 300 parts ethanol; freely soluble in chloroform, pyridine, glacial acetic acid; moderately soluble in ethyl acetate; slightly soluble in benzene; and practically insoluble in petroleum ether and water.

* Stoll, *Helv. Chim. Acta* **28**, 1283, (1945)

Identification Tests The precipitation reactions and the colour tests are the same as described under **ergonovine**. However, the specific derivatives of **ergotamine** are as stated below:

1. **Ergotamine Tartrate** [(C₃₃H₃₅N₅O₅)₂.C₄H₆O₆] (**Ergomar; Ergate; Ergotartrat; Ergostat; Exmigra; Fermergin; Lingraine; Gynergen; Lingran**): It is normally obtained as solvated crystals *e.g.*, the *dimethanolate*; also occurs as heavy rhombic plates from methanol having mp 203°C (decomposes). It has specific optical rotation $[\alpha]_D^{25} - 125^\circ$ to -155° (C = 0.4 in chloroform). One gram dissolves in either 500 ml of ethanol or water.
2. **Ergotamine Hydrochloride** [C₃₃H₃₅N₅O₅.HCl]: It is obtained as rectangular plates from 90% (v/v) ethanol which get decomposed at 212°C. It is found to be soluble in water-ethanol mixtures; and sparingly in water or ethanol alone.
3. **Dihydroergotamine Mesylate** (C₃₃H₃₇N₅O₅.CH₃SO₃H) (**Agit; 1 Dihydro-ergotamine methane sulphonate; Angionorm; DET MS; Dergotamine; D.H.E. 45; Diergo; Dihyergot; Dirgotarl; Endophleban; Ergomimet; Ikaran; Migranal; Morena; Ergont; Ergotonin; Orstanorm; Tonopres; Verladyn; Seglor**): It is obtained as large prisms from 95% (v/v) ethanol having mp 230-235°C; and moderately soluble in water.

Note: (a) It is the salt of a semisynthetic alkaloid prepared from ergotamine by hydrogenation of the Δ^9 double bond in the lysergic acid nucleus.

(b) It is mostly used in the treatment of migraine because it is found to be better in efficacy and more tolerated than the parent alkaloid.

Uses


1. It is employed as a potent antimigraine drug.
2. **Ergotamine** possesses oxytocic properties, but it is not employed for that effect.
3. **Ergotamine tartrate** is used invariably to prevent or abort vascular headaches, including *migraine* and *cluster headaches*. The mechanism of action is perhaps due to direct vasoconstriction of the dilated carotid artery bed with concomitant lowering in the amplitude of pulsations.
4. **Ergotamine tartrate** is also an antagonist of the serotonin activity.
5. **Ergotamine tartrate** is frequently used along with caffeine for the management and control of migraine headache. Both serve as cerebral vasoconstrictors; while the latter is considered to increase the action of the former.
6. **Methylergonovine maleate** is an oxytocic reported to be longer acting and more active than **ergonovine**.

C. Ergine

Synonyms Lysergamide; Lysergic acid amide;

Biological Sources It is obtained from the immature seeds of *Argyreja nervosa* (Burm.) Bojer (*Convolvulaceae*) (**Wood Rose, Silver Morning Glory**); Beeds of *Ipomea Violaceae* L. (*Convolvulaceae*) (**Tlitliltzen, Ololiuqui**); seeds of *Rivea corymbosa* Hall. F. (*Convolvulaceae*) (**Snakeplant**); and also from the seeds of *Ipomea tricolor* Cav (*Convolvulaceae*).

Chemical Structure



Ergine

9, 10-Didehydro-6-methylergoline-8β-carboxamide; (C₁₆ H₁₇ N₃O).

Isolation It is isolated from the seeds of *Rivea corymbosa* (L.) and from *Ipomea tricolor* Cav. by the method of Hofmann and Tschertter.*

Characteristic Features

1. It is obtained as prisms from methanol which get decomposed at 242°C.
2. It has a specific optical rotation of $[\alpha]_{5461}^{20} + 15^\circ$ (C = 0.5 in pyridine).

Identification Tests The precipitation reactions and the colour tests are the same as described under **ergonovine** (Section A).

Ergine may also be identified by forming its derivative as stated below:

Ergine Methane Sulphonate (C₁₆H₁₇N₃O·CH₃SO₃H) It is obtained as prisms from a mixture of methanol and acetone that get decomposed at 232°C.

Uses It has a pronounced depressant action.

Note: It is a controlled substance listed in the U.S. Code of Federal Regulations, Title 21 Part 1308, 13 (1995).


Biosynthesis of Ergotamine The various steps involved in the biosynthesis of **ergotamine** are as enumerated below:

1. Three amino acids, viz., L-alanine, L-phenylalanine, and L-proline in the presence of ATP and enzyme SH; or D-(+)-lysergic acid in the presence of ATP and enzyme SH undergo two steps: *first*-activation via AMP esters, and *secondly*-attachment to the respective enzymes, thereby giving rise to an intermediate. It is worthwhile to observe that the enzyme is comprised of *two subunits* that essentially bind the substrates as indicated in the biosynthetic pathway given below.
2. The comparatively more complex structures comprising of the peptide fragments, such as: **ergotamine** are eventually formed by sequential addition of amino acid residues to the *thioester-bound lysergic acid*, yielding a linear lysergyl-tripeptide covalently attached to the enzyme complex.

* Hofmann and Tschertter, *Experientia*, **16**, 414 (1964).

- The resulting complex undergoes lactam formation followed by release from enzyme. In other words, the cyclized tripeptide residue is rationalized instantly by the formation of a lactam (amide) that releases ultimately the product from the enzyme.
- This resulting product first affords hydroxylation then followed by generation of a hemiketal-like linkage to give rise to the formation of *ergotamine*.


All these aforesaid steps (1) through (4) have been duly depicted in the following biosynthetic pathway.



Biosynthesis of Ergotamine
(Adapted from - 'Medicinal Natural Products' Dewick P.M.)


Peptide Alkaloids in Ergot Interestingly, it has been observed critically that three amino acids, namely: alanine, phenylalanine and proline, actually form the basis for the various structures which are encountered in the domain of the '**ergot alkaloids**'. Therefore, these known and established structures may be subdivided into *three* major groups which are: ergotamine group, ergoxine group, and ergotoline group.

The various alkaloids having the peptide linkages found in '**ergot**' are depicted as under.


**Ergotamine group**

R = CH ₂ Ph	ergotamine
R = CH ₂ CHMe ₂ S	ergosine
R = $\overset{S}{\text{C}}\text{H}(\text{Me})\text{Et}$	(β -ergosine)
R = CHMe ₂	ergovaline
R = Et	ergobine

() Not yet known in nature

**Ergoxine group**

ergostine
ergoptine
(β -ergoptine)
ergonine
ergobutine

**Ergotoxine group**

ergocristine
α -ergocryptine
β -ergocryptine
ergocornine
ergobutyryne

7.3 ALKALOIDS IN TISSUE CULTURES

The quantum growth and progress in the past three decades especially with regard to the legitimate utilization of plant **tissue cultures** in the exclusive bioproduction of naturally occurring chemical compounds under specific aseptic conditions through various well established means and methods almost identical to those employed to culture microorganisms has virtually opened up an altogether new and virgin horizon in the latest field of biotechnology. Therefore, the application of tissue culture techniques in the context of the biosynthesis of important secondary metabolites from plants *viz.*, **alkaloids**, not only holds a well-deserved promise for the rational controlled production of plant constituents but also supports the fact that higher plants do provide an important source of medicinally active chemical entities.


Although it has been established beyond any reasonable doubt that most of the work carried but on '**alkaloid biosynthesis**' has been more or less directly concerned with the intact plants or parts thereof, for instance: leaves, roots or shoots, there have been certain evidence and investigations using tissue cultures. This type of work is particularly beneficial in locating and establishing the site of alkaloid synthesis. It is, however, pertinent to mention here that the inferences drawn from various experimental findings that *tobacco stem callus tissue* will not synthesize alkaloids unless and until the root formation has started either spontaneously or by means of chemical stimulation.

Likewise, it has been observed interestingly that the latex isolated from the capsule of the **opium poppy** (*Papaver somniferum*) will synthesize morphine either from dopa or tyrosine, but the latex obtained from the stem will not.

Tissue cultures do not always essentially behave exactly as the **intact plant**, as has been observed with *Catharanthus roseus*, wherein the cultures of either leaf or stem effectively carried out the synthesis of certain alkaloids found in the intact plant, but no **dimeric alkaloids** could either be observed or detected. On the contrary, tissue cultures of tobacco might convert thebaine to morphine; however, no **benzylisoquinoline alkaloids** have been noticed in *Nicotiana tabacum*.

7.4 ALKALOIDS IN CHEMOSYSTEMATICS

Broadly speaking the alkaloids invariably occur across the entire plant kingdom. In reality, alkaloids are usually found more abundantly in plants specifically belonging to the *Dictyolendones* than in either the *Monocotyledones* or the non-flowering plants. However, it has been observed that amongst the *Pteridophyta* and *Gymnospermae*, only the *Lycopodiaceae* family happens to synthesize these compounds to any reasonable extent. The *Lycopodium* alkaloids are essentially the **quinolizidine derivatives**; and usually have the ring structure as displayed by **Lycopodine** below:



Lycopodine

(15R)-15-Methyllycopodan-5-one; (C₁₆H₂₅NO).

One school of thought believes that the alkaloids are confined rarely in dicotyledon orders classified before the Centrospermae, thereby establishing linkage of these simpler flowering plants with the *Gymnospermae*. The distribution and occurrence of alkaloids is found to be quite uneven among the remaining orders. Thus, an order rich in such compounds could be preceded and followed by an order wherein alkaloids are not synthesized at all. Such an erratic distribution may be noticed even more distinctly and apparently in certain plant families also. Therefore, one may safely conclude that alkaloids by themselves are absolutely spinless in establishing phylogenetic relationships either between orders or between families within an order. Of course, there exists some exceptionally few cases, for instance: *Centrospermae* which has been duly illustrated below.

Such natural orders that are found to be rich in **alkaloids** are: Centrospermae, Gentianales, Magnoliales and Ranunculales belonging to the class of *Dicotylendones*; and Liliales, and Orchidales belonging to the category of *Monocotylendones*. It has been reported duly tht both *Papaveraceae* and the absolutely unrelated *Apocynaceae* contain the largest number of these secondary metabolites. Surprisingly, each and every species of *Papaveraceae* thoroughly studied till date comprises of *alkaloids*; whereas, the *Apocynaceae* contains a rather more prominent diversity of complex *indole alkaloids*. There are a host of other plant families in which **alkaloids** usually occur more predominantly. and frequently are, namely: *Amaryllidaceae*, *Compositae*, *Leguminosae*, *Liliaceae*, *Loganiaceae*, *Orchidaceae*, *Ranunculaceae*, *Rubiaceae*, *Rutaceae* and *Solanaceae*. However, the *Amaryllidaceae* alkaloids are found to be specific only to that family, that contains no other variety.

A broad survey of literature has adequately proved the fact that while applying the chemosystematics to the classification of plants, the biosynthetic pathway is certainly more vital and important than that of the end-product. It may be further exemplified by considering, **quinine**, a quinoline derivative obtained in *Rubiaceae*, is biosynthesized from tryptophan by a biosynthetic pathway very similar to that forming the complex **indole alkaloidal** characteristic features of the family. Besides, quinoline derivatives are also found in the *Rutaceae*, but in this particular instance they are biosynthesized from anthranilic acid; and also by a pathway that is very specific to this family.

Lastly, one may draw an inference that the strategic application of alkaloid biosynthesis to classification is made rather complicated by both convergence and divergence. Some typical examples of convergence and divergence are given below:

- (a) **Examples of Convergence:** The synthesis of the **tropane alkaloids** from hygrine is on account of convergence. It normally occurs in a plethora of unrelated plant families, such as: *Convolvulaceae* (*Convolvulus* species), *Cruciferae* (*Cochleavia arctica*); *Erythroxylaceae* (*Erythroxylum coca*); *Euphorbiaceae* (*Phyllanthus discoidens*); and *Solanaceae* (includes several genera).
- (b) **Examples of Divergence:** *Papilionoideae* exhibit several examples of divergence; because, members of this subfamily synthesise virtually a wide spectrum of secondary metabolites. The **Papilionoideae alkaloids** are found to exhibit no apparent relationship either in molecular structure or in their respective biosynthetic pathways. Hence, the **spiroamine alkaloids** found in *Erythrina* species have virtually little structural relationship to the **pyrrolizidine alkaloids** that are characteristic of *Crotolaria* species, and are biosynthesized from tyrosine; whereas, the **pyrrolizidine alkaloids** are normally found to originate from ornithine.

FURTHER READING REFERENCES

1. Antkowiak, R. and Autkowiak, W.Z. **Alkaloids from Mushrooms**, *The Alkaloids, Chemistry and Pharmacology*, (ed Brossi A) Vol. 40, Academic, San Diego, pp 189-340, 1991.
2. Atta-ur-Rahman and Choudhary MI: **Chemistry and Biology of Steroidal Alkaloids**. *The Alkaloids, Chemistry and Pharmacology*, (ed Cordell GA), Vol 50, Academic, San Diego, pp 61-108, 1998.
3. Amiya T and Bando H: **Aconitum Alkaloids**, *The Alkaloids Chemistry and Pharmacology* (ed Brossi A) Vol 34, Academic, San Diego, pp 95-179, 1988.
4. Brossi A and Manske RHF., eds: **The Alkaloids**, Vols. XXI-XXV, 26-40, New York, Academic Press Inc., 1983-1991.
5. Brossi A and Cordell G A., eds: **The Alkaloids**, Vol 41, San Diego, Academic Press Inc., 1992.
6. Bentley KW: **β -Phenylethylamines and the Isoquinoline Alkaloids**. *nat Prod Rep* **17**, 247-268, Earlier Reviews: 1999, **16**, 367-388, 1998, **15**, 341-362, 2000.
7. Bisset NG: **Curare Alkaloids, Chemical and Biological Perspectives** (ed Pelletier S.W.) Vol. 8. Wiley, New York, pp 1-150, 1992.
8. Bosch J, Bonjoch J and Amat M: **The Sytrychnos Alkaloids**. *The Alkaloids, Chemistry and Pharmacology*, (ed Cordell GA). Vol 48, Academic, San Diego, pp 75-189, 1996.
9. Brossi A, Pei XF and Greig NH (1996): **Phenserine, a Novel Cholinesterase Related to Physostigmine: Total Synthesis and Biological Properties**, *Aust J. Chem.*, **49**, 171-181, 1996.

10. Cordell G.A.: **Introduction to Alkaloids: A Biogenetic Approach**, John Wiley Sons. Inc., New York, 1981.
11. Chiara GD and North RA: **Neurobiology of Opiate Abuse**. *Trends Pharmacol Sci*, **13**, 185-193, 1992.
12. Cervoni P. Crandall DL and Chan PS: **Cardiovascular Agents-Antiarrhythmic Agents**, Kirk-Othmer Encyclopedia of Chemical Technology, 4th edn. Vol. 5, Wiley, New York, 207-238, 1993.
13. Dey, P.M. and Harborne JB, (eds): **Methods in Plant Biochemistry**, Vol. 8., *Alkaloids and Sulphur Compounds*, Academic Press Ltd., London (1993)
14. Dalton DR: **Alkaloids: Kirk Othmer Encyclopedia of Chemical Technology**, 4th edn., Vol. 1, Wiley, New York, 1039-1087, 1991.
15. Emboden, W.: **'Narcotic Plants'**, Mc Millan Publishing Co. Inc. New York (1979).
16. Evans WC: **Datura-a Commercial Source of Hyoscine**, *Pharm. J.*, **244**, 651-653, 1990.
17. Fujii T and Ohba M: **The Ipecac Alkaloids and Related Bases**. *The Alkaloids, Chemistry and Biology* (ed Cordell GA) Vol. 51, Academic San Diego, pp 271-321, 1998.
18. Glasby, J.S.: **Encyclopedia of the Alkaloids**, Vols. 1-4, Plenum Press, New York, 1975, 1977, and 1983.
19. Goodyer L: **Travel Medicine (4): Malaria**, *Pharm. J.*, **264**, 405-410, 2000.
20. Groger D and Floss H: **Biochemistry of Ergot Alkaloids: Achievements and Challenges**. *The Alkaloids Chemistry and Pharmacology* (ed Cordell CA) Vol. 50, Academic, San Diego, pp 171-218, 1998.
21. Herbert, R. B. **The Biosynthesis of Secondary Metabolites**, 2nd ed. Chapman and Hall. New York, (1989).
22. Hibino S and Choshi T: **Simple Indole Alkaloids and those with a Non rearranged Monoterpenoid Unit**. *Nat Prod Rep* **18**, 66-87, 2001. Earlier Review: Lounasmaa M and Tolvanen A **17**, 175-191, 2000.
23. Hartmann T. and Witte L: **Chemistry, Biology and Chemecology of the Pyrrolizidine Alkaloids**. *Alkaloids, Chemical and Biological Perspectives* (ed Pelletier SW) Vol 9, Wiley, New York, 155-233, 1955.
24. Johnson EL and Emcho SSD: **Variation in Alkaloid Content in *Erythroxylum coca* Leaves from Leaf-bud to Leaf-drop** *Ann. Bot* **73**, 645-650, 1994.
25. Kutchan TM: **Molecular Genetics of Plant Alkaloid Biosynthesis**. *The alkaloids, Chemistry and Pharmacology*. (ed Cordell GA) Vol. 50, Academic, San Diego, pp. 257-316, 1998.
26. Kutchan TM: **Strictosidine: From Alkaloid to Enzyme to Gene**. *Phytochemistry* **32**, 493-506, 1993.
27. Kutney JP: **Plant Cell Culture Combined with Chemistry: A Powerful Route to Complex Natural Products**, *Account Chem. Res.* **26**, 559-566, 1993.
28. Kren V: **Bioconversions of Ergot Alkaloids**. *Adv. Biochem. Eng. Biotech* **44**, 123-144, 1991.
29. Kalix P: **The Pharmacology of Psychoactive Alkaloids from Ephedra and Catha**. *J. Ethnopharmacol.* **32**, 201-208, 1991.
30. Lee M: **Malaria in Search of Solutions**. *Chem. Brit.* **32**, (8), 28-30, 1996.
31. Leonard J: **Recent Progress in the Chemistry of Monoterpenoid Alkaloids Derived from Secologanin** *Nat Prod Rep* **16**, 319-338, 1999.
32. Lewis JR: **Amaryllidaceae, *Sealeetium*, Imidazole, Oxazole, Thiazole, Peptide and Miscellaneous Alkaloids**. *Nat Prod Rep* **18**, 95-128, 2001.
33. Misra N, Luthra R, Singh KL and Kumar S: **Recent Advances in Biosynthesis of Alkaloids**. *Comprehensive Natural Products Chemistry*, Vol. 4, Elsevier, Amsterdam, pp. 25-59, 1999.
34. Michael JP: **Indolizidine and Quinolizidine Alkaloids**. *Nat Prod Rep* **17**, 579-602, 2000.
35. Noble RL: **The Discovery of the Vinca Alkaloids-Chemotherapeutic Agents against Cancer**. *Biochem Cell Biol* **68**, 1344-1351, 1990.
36. O'Hagan D: **Pyrrole, Pyrrolidine, Pyridine, Piperidine and Tropane Alkaloids**. *Nat Prod Rep* **17**, 435-446, 2000.

37. Pelletier SW (ed) **Alkaloids**, Vols 1-6, John Wiley & Sons. Inc., New York, 1993-1988; Vols 7-8, New York, Springer-Verlag New York Inc., 1991-1992.
38. Philipson JD., Roberts, M.F., Zenk, M.H. eds: **The Chemistry and Biology of Isoquinoline Alkaloids**, Springer-Verlag; Berlin, 1985.
39. Quetin-Leclereq J, Angenot L. and Bisset NG: **South American *Strychnos* species. Ethnototany (except Curare) and Alkaloid Screening**. *J. Ethnopharmacol.* **28**, 1-52, 1990.
40. Ripperger H: **Solanum Steroid Alkaloids: an Update. Alkaloids, Chemical and Biological Perspectives** (ed Pelletier SW) Vol. 12. Elsevier, Amsterdam, pp 103-185, 1998.
41. Robins DJ: **Biosynthesis of Pyrrolizidine and Quinolizidine Alkaloids. The Alkaloids, Chemistry, Pharmacology** (ed Cordell GA) Vol. 46, Academic, San Diego, pp 1-61, 1995.
42. Stockigt, J. and Ruppert M: **Strictosidine-the Biosynthetic Key to Monoterpenoid Indole Alkaloids** *Comprehensive Natural Products Chemistry*, Vol. 4. Elsevier, Amsterdam, pp 109-138, 1999.
43. Schneider MJ: **Pyridine and Piperidine Alkaloids. an Update: Alkaloids, Chemical and Biological Perspectives** (ed Pelletier SW) Vol 10, Elsevier, Amsterdam, pp 155-299, 1996.
44. Singh, S: **Chemistry, Design, and Structure-activity Relationships of Cocaine Antagonists**. *Chem. Rev.* **100**, 925-1024, 2000.
45. Saxton JE: **The Chemistry of Heterocyclic Compounds: The Monoterpenoid Indole Alkaloids**, Vol 25, Part 4., John Wiley & Sons Inc., New York, 1983.
46. Toyota M. and Ihara M: **Recent Progress in the Chemistry of Non-monoterpenoid Indole Alkaloids** *Nat Prod Rep* **15**, 325-340, 1998, Earlier reviews: Ihara M and Fukumoto K **14**, 413-429, 1997, **13**, 241-261, 1996.
47. Wang FP and Liang XT: **Chemistry of the Diterpenoid Alkaloids: The Alkaloids, Chemistry and Pharmacology** (ed Cordell GA) Vol 42: Academic San Diego, 151-247, 1992.
48. Weiss, R. D., Mirin, SM and Bartel RL: **Cocaine**, 2nd ed., American Psychiatric Press, Inc., Washington DC, 1994.
49. Wills S: **Drugs and Substance Misuse: Plants** *Pharm. J.* **251**, 227-229, 1993.
50. Verpoorte R., Van der Heiden R and Memelink J: **Plant Biotechnology and the Production of Alkaloids: Prospects of Metabolic Engineering**; *The Alkaloids Chemistry and Pharmacology* (ed Cordell GA) Vol 50, Academic San Diego, pp 453-508, 1998.

8

Bitter Principles

- Introduction
- Classification of Bitter Principles
- Further Reading References

8.1 INTRODUCTION

In general, the **bitter principles** are heterogenous vegetative compounds that neither belong to the class of *alkaloids* nor to the *glycosides*, but they do possess a characteristic bitter taste.

It is, however, pertinent to observe that bitter principles are invariably of vegetative origin and essentially comprise of C, H, and O, but are found to be free from N.

Interestingly, at one point in time the **bitter principles** were frequently and extensively utilized in liquid medicaments to augment and stimulate appetite. It has been established that the bitter constituent particularly stimulate the salivary glands (gustatory-nerves) present in the mouth and cause an enhancement in the psychic secretion of the gastric juice in the stomach. Since the past several decades the extract of the following drugs have been employed both extensively and intensively in various herbal systems of medicine, namely: **calumba, cinchona (or quinine) gentian, quassia, nux-vomica**, etc.

The '**bitter principles**' are mostly found in a number of plants, and are observed to be present abundantly in certain families, such as: *Compositae Labiatae*, and *Gentianaceae*.

Over the years, considerable research has accelerated the investigation of a number of these **bitter compounds** possibly for other meaningful applications, for instance: the **bitters** (*i.e.*, **bitter principles**) of the *Simaroubaceae* as *antitumour* and *antimalarial* agents.

8.2 CLASSIFICATION OF BITTER PRINCIPLES

Bitter principles have been judiciously classified into *six* categories based on the typical chemical structures present in them, namely:

- (a) Phenolic Bitter Principles,
- (b) Lactone Bitter Principles,
- (c) Chromone Bitter Principles,
- (d) Coumarin Bitter Principles,
- (e) Coumarone Bitter Principles, and
- (f) Miscellaneous Bitter Principles.

All these six different groups of **bitter principles** shall now be discussed at length along with certain important members from each class separately.

8.2.1 Phenolic Bitter Principles


The crystalline **acidic bitter principles** having a phenolic function are found in naturally occurring plant sources, such as: **humulon, lupulon**.

8.2.1.1 Humulon

Synonyms Humulone; α -Lupulic acid, α -Bitter acid.

Biological Source It is obtained as an antibiotic constituent from the strobiles of *Humulus lupulus* L. belonging to the natural order *Moraceae* (Hops).

Chemical Structure



Humulon

(R)-3, 5, 6-Trihydroxy-4, 6-bis (3-methyl-2-butenyl)-2-(3-methyl-1-oxobutyl)-2, 4-cyclohexadien-1-one; (C₂₁H₃₀O₅)

Isolation The various steps involved in the isolation of **humulon** from hops strobiles are:

1. Hops strobiles are extracted with ethanol for several hours and the alcoholic extract is filtered.
2. The filtrate is heated with animal charcoal and then cooled. In this manner, the charcoal adsorbs the bitter principles.
3. The charcoal is filtered and the adsorbed bitter principle is extracted with ethanol.
4. The alcoholic extract is evaporated over a water-electric bath and **humulon** is subsequently extracted from the resulting resinous residue by the help of boiling water repeatedly.
5. The combined aqueous fraction is cooled to 20°C and extracted with solvent ether successively.
6. The ethereal layer is filtered and evaporated in a Thin-Film Rotary Evaporator to obtain the bitter principle humulon as a residue.

Characteristic Features

1. The crystals obtained from ether have mp 65-66.5°C
2. It has a distinct bitter taste especially in alcoholic solution.
3. **Humulon** is observed to be more stable to air than lupulon.
4. It is a monobasic acid.
5. It has a specific optical rotation $[\alpha]_D^{20} - 212^\circ$ (1.0 g in 15.5 g 96% v/v ethanol).
6. It has uv_{max} (ethanol): 237, 282 nm (ϵ 13, 760; 8330).
7. It is found to be soluble in usual organic solvent.

8. It is slightly soluble in boiling water from which it normally separates out as a milky precipitate on cooling.
9. It readily forms a sodium salt which is rapidly soluble in water.
10. **Bacteriostatic Potency: Humulon** suffers no loss of bacteriostatic potency against *Staphylococcus aureus* upon autoclaving 40 ppm in phosphate buffer at pH 6.5 or 8.5. However, the addition of ascorbic acid in low concentrations extends the duration of bacteriostatic action.

Identification Tests These are as follows:

1. An ethanolic solution of **humulon** gives a reddish-violet colouration with a few-drops of FeCl_3 solution (0.5% w/v).
2. A few mg of **humulon** when dissolved in 0.5-1 ml NaOH solution (0.1 N) it produces a yellow colour.
3. **Humulon** reduces Tollen's Reagent (*i.e.*, ammoniac silver nitrate solution) *in cold* and gives a silver mirror.
4. **Humulon** on being heated with an alcoholic solution of NaOH (0.5 N) undergoes complete decomposition to yield: humulinic acid (an unsaturated acid), acetic acid, isobutyric aldehyde and an unsaturated liquid volatile acid.

Uses


1. **Humulon** exerts bacteriostatic action.
2. It contributes to the bitterness of hops extract used in making beer.

8.1.2.2 Lupulon

Synonyms β -Bitter Acid; β -Lupulic Acid.

Biological Sources The biological sources of **lupulon** are the same as for humulon given under sections 8.2.1.1.

Chemical Structure



Lupulon

3, 5-Dihydroxy-2, 6, 6-tris (3-methyl-2-butenyl)-4-(3-methyl-1-oxobutyl)-2, 4-cyclohexadien-1-one; ($\text{C}_{26}\text{H}_{38}\text{O}_4$).

Isolation **Lupulon** may be isolated from the commercial hops *Humulus lupulus* L. (*Moraceae*) by the method suggested by Lewis *et al.**

* Lewis *et al.*, *J. Clin. Invest.*, **28**, 916 (1949).

Characteristic Features

1. It is obtained as prisms from 90% (v/v) methanol having mp 92-94°C.
2. It possesses a distinct bitter taste especially in alcoholic solutions.
3. It behaves as a monobasic acid.
4. It is perfectly stable in *vacuo* even upto a temperature of 60°C.
5. It exhibits a slight acid reaction.
6. It is found to be optically inactive.
7. It is freely soluble in ethanol, methanol, hexane, petroleum ether, isooctane; and slightly soluble in either neutral or acidic aqueous solutions.
8. It readily forms a sodium salt which is rapidly soluble in water.
9. The addition of 0.1% solution of ascorbic acid affords a marked and pronounced protective action upon the bacteriostatic activity of **lupulon** steamed or autoclaved at a concentration of 4 ppm in phosphate buffer at pH 6.5 and 8.5.

Identification Tests

1. **Lupulon** turns yellow and amorphous in nature within a few days with the development of a characteristic odour.
2. **Lupulon** on being subjected to oxidation with potassium permanganate solution gives rise to the formation of *valerianic acid*.

Uses

1. **Lupulon** contributes exclusively to the bitterness of hops extract and is employed in the manufacture of beer.
2. It possesses the properties of an aromatic bitter and are said to have a sedative activity.

8.2.2 Lactone Bitter Principles


The **lactone bitter principles** essentially possess a five-membered lactone ring which may be exemplified by the help of *two* glaring potent compounds belonging to this category, namely: **α -santonin** and **picrotoxinin**. These compounds shall now be discussed individually in the sections that follow:

8.2.2.1 α -Santonin

Synonym *l*-Santonin.

Biological Sources It is obtained from the dried unexpanded flower heads of *Artemisia maritima* L., sens. lat. (*Compositae*) (**Levant Wormseed**); and other spices of *Artemisia* found mostly in Russia, China and Turkestan besides the Southern Ural Region.

Chemical Structure



α -Santonin

1, 2, 3, 4, 4 α , 7-Hexahydro-1-hydroxy- α , 4 α -8-trimethyl-7-oxo-2-naphthaleneacetic acid γ -lactone; (C₁₅H₁₈O₃).

Isolation The dried unexpanded flower heads of **levant wormseed** are treated with milk of lime so as to obtain calcium santoninate. The resulting product is subsequently converted into the corresponding soluble salt of **sodium santoninate** by the careful treatment with either sodium carbonate or sodium hydroxide. A stream of CO₂ is passed through the reaction mixture to get rid of calcium hydroxide as a precipitate of calcium carbonate which is filtered off conveniently. The filtrate is acidified with dilute sulphuric acid (6 N) when the crude **santonin** gets precipitated. The crude product thus obtained is made to dissolve in minimum quantity of ethanol (95%) and treated with activated charcoal powder to decolourize the solution. It is finally filtered off, ethanol is evaporated on an electric water bath and allowed to cool in a refrigerator overnight to obtain the pure **α -santonin**.

Characteristic Features The *three* different forms of **santonin** have the following characteristic features:

(a) **(–)-Form of Santonin:**

1. It may be obtained either as tabular crystals or as orthorhombic sphenoidal crystals having mp 170-173°C.
2. It is found to be practically tasteless with a positive bitter aftertaste.
3. Its specific optical rotation $[\alpha]_D^{25}$ ranges between – 170° to 175° (C = 2 in ethanol).
4. It turns yellow on being exposed to light.
5. It causes irritation to the mucous membranes.
6. It has specific gravity d 1.187.
7. **Solubility Profile:** One part dissolves in 5000 parts of cold water, in 250 parts of boiling water, in 280 parts of 55% ethanol at 17°C, in 10 parts of boiling 50%-ethanol, in 44 parts of cold 90% ethanol, in 3 parts of boiling 90% ethanol, in 125 parts of cold ether, in 72 parts of boiling ether, and in 4.3 parts of cold chloroform.

(b) **(+)-Form of Santonin:**

1. It is obtained as colourless plates from methanol having mp 172°C.
2. Its specific optical rotation $[\alpha]_D^{20} + 165.9^\circ$ (C = 1.92 in ethanol).

(c) **(±)-Form of Santonin:**

1. It is obtained as colourless plates from methanol having mp 181°C.
2. It has uv_{max} (ethanol): 241 nm (log ϵ 4.10).

Identification Tests The various identification tests for **α -santonin** are as stated below:

1. **Chromosantonin (Photosantonin): Santonin** is fairly stable in air, however, it turns yellow on exposure to *light* whereby it gets converted into its isomeric form chromosantonin, also known as photosantonin. The latter may be regenerated into santonin by simply crystallisation from ethanol.
2. **Santonin** when warmed with ethanolic solution of KOH or NaOH, it first and foremost produces a violet-red colouration, which gradually alters to reddish-yellow.

- Heat 0.01 g of **santonin** with 2 ml of a mixture of sulphuric acid and water (1 : 1) no colour is produced apparently; but on the addition of 2-3 drops of dilute FeCl₃ solution (0.1% w/v) to the hot liquid a violet colouration is produced instantly.
- Santonin** when dissolved in a few drops of ethanol containing furfural, 2 ml of concentrated H₂SO₄ and the resulting mixture is heated in a porcelain dish over water-bath it gives a purple-red colouration that gets changed to bluish-violet, to dull blue, and finally to almost black.
- Santonin** 0.1 g when dissolved in 5 ml of ethanol (95% v/v) gives a clear solution which being neutral to litmus paper and is levorotatory.


Uses

- It is mostly used as an anthelmintic (Nematodes).
- It is very efficient in its action on round worms; but shows less effect on the thread worms and none on taenia.

8.2.2.2 Picrotoxinin

Biological Sources It is the toxic component of **picrotoxin** obtained from the seed of *Anamirta cocculus* L. Wight & Arn. (*Menispermaceae*); and also found in *Tinomiscium philippinense* Diels.

Chemical Structure



Picrotoxinin

[1aR – (1a α , 2a β , 3 β , 6 β , 6a β , 8aS*, – 8b β , 9R*)]-Hexahydro-2a-hydroxy-8b-methyl-9-(1-methyl-ethenyl)-3, 6-methano-8H-1, 5, 7-trioxacyclopenta [ij] cycloprop [a] azulene-4, 8 (3H)-dione; (C₁₅H₁₆O₆).

Preparation **Picrotoxinin** may be prepared from picrotoxin by the method suggested by Horrmann.*

Characteristic Features

- It is obtained in two different forms: *first*—as large prisms, and *secondly*—as small crystals containing water having mp 209.5°C.
- Its specific optical rotations are: $[\alpha]_D^{17} + 4.4^\circ$ (C = 4.28 in absolute alcohol); and $+ 3.49^\circ$ (C = 7.57 in acetone).
- It is found to be soluble in hot common organic solvents; and also in cold chloroform and ethanol.
- Nevertheless it has a very bitter taste.


* Horrmann, *Ber.*, **45**, 2090 (1912).

Uses



1. It is employed as a CNS and respiratory stimulant.
2. It may also be used as an antidote to barbiturates.

8.2.3 Chromone Bitter Principles


The following heterocyclic moieties, such as: **chromone**, **coumarin** and **coumarone** are derived from **γ -pyrone**, **α -pyrone** and **furan** nucleus respectively in combination with a benzene nucleus.

Chromone γ -Pyrone

Benzene

Benzo- γ -pyrone or
Chromone**Coumarin** α -Pyrone


Benzene

Benzo- α -pyrone or
Coumarin**Coumarone:**

Furan



Benzene



Benzofuran or Coumarone


The important members belonging to the class of **chromone bitter principles** are, namely: **khellin**, **khellol glucoside** and **visnagin**. These *three* drug substances shall be described as under:

8.2.3.1 Khellin

Synonyms Kellin; Kelamin; Kelicor; Keloid; Kelicorin; Khelfren; Gyno khellan; Eskel; Norkel; Amicardine; Ammivisnagen; Viscardan; Visnagen; Visnagalin; Visokellina, Cardio-Khellin; Coronin; Ammivin; Ammipurin; and Ammicardine.

Biological Sources It is the major active chemical constituent obtained from the seeds of *Ammi visnaga* Lam. (*Umbelliferae*) (**Toothpick Ammi; Chellah; Khella**). It is present in the plant substance to the extent of 1%.

Chemical Structure



Khellin

4, 9-Dimethoxy-7-methyl-5H-furo [3, 2-g]-[1]-benzopyran 5-one; (C₁₄H₁₂O₅).

Khellin is a *Furanochromone* compound.

Isolation The various steps involved in the isolation of **khellin** from the seeds of *A. visnaga* are as stated below:

1. The seeds are dried, powdered, sieved and extracted in Soxhlet apparatus with solvent ether for several hours.
2. The ethereal extract is concentrated in a rotary thin-film evaporator and stored in a refrigerator for a few days.
3. The cold ethereal extract eventually comprise of *three* distinct layers: an upper green oily layer; a middle cream coloured fatty layer; and a lower green crystalline layer. The upper green oil is removed by filtration with gentle suction, the middle cream coloured fatty layer is removed by the help of petroleum ether, and the remaining lower solid residue is duly purified by repeated crystallization from methanol to obtain pure **khellin**.

Note: The methanol mother liquor is kept aside for the isolation of ‘visnagin’.

Characteristic Features

1. The crystals of **khellin** are obtained from methanol having mp 154–155°C.
2. It has a characteristic bitter taste.
3. It boils at bp_{0.05} 180-200°C.
4. It has uv_{max} (ethanol): 250, 338 nm (E_{1cm}^{1%} 1600, 200).
5. **Solubility Profile:** Its solubility in g/100 ml at 25°C are: water 0.025; acetone 3.0; methanol 2.6; isopropanol 1.25; ether 0.5; and skellysolve B 0.15.
However, it is found to be much more soluble in hot water and hot methanol.
6. **Khellin** is observed to be significantly stable when mixed with the normal tableting excipients.

Identification Tests These are as enumerated below:

1. **Khellin** decolourizes potassium permanganate solution.
2. When 5-8 mg **khellin** is mixed with a small piece of solid KOH or NaOH it produces a distinct rose-red colouration.

Note: This test is not positive when either K or Na carbonate/bicarbonate used.

3. **Wagner’s Reagent Test:** A saturated aqueous solution of khellin yields a precipitate with Wagner’s Reagent.*

* **Wagner’s Reagent:** It is a solution of iodine with KI.

4. Khellin gives a faint precipitate with tannic acid solution.

Uses


1. **Khellin** is used as a potent vasodilator (coronary).
2. It also finds its application as a potent selective bronchodilator.
3. It is used extensively in the treatment and control of coronary insufficiency, angina pectoris and in chronic bronchial asthma.

8.2.3.2 Khellol Glucoside

Synonyms Khellinin

Biological Sources It is obtained from the seeds of *Ammi visnaga* Lam., (*Umbelliferae*); and also from *Eranthis hyemalis* L. (*Ranunculaceae*) upto 0.3%.

Chemical Structure



Khellol Glucoside

7-[(β -D-glucopyranosyloxy)-methyl]-4-methoxy-5H-furo [3, 2-g] [1] benzopyran-5-one; ($C_{19}H_{20}O_{10}$).

Isolation It may be isolated from *Eranthis hyemalis* by the method put forward by Egger.*

Characteristic Features

1. It is obtained as crystals from ethanol having mp 179°C.
2. It has uv_{max} (ethanol): 250, 325 nm.
3. It is found to be soluble in acetic acid, hot ethanol slightly soluble in hot methanol; and almost insoluble in acetone, ethyl acetate, ether, chloroform, cold alkali.

Identification Test **Khellol glucoside** may be identified by making its following tetraacetate derivative due to the presence of four OH moieties.

Khellol Glucoside Tetraacetate It is obtained as flakes from ethanol having mp 153°C. It is freely soluble in acetone, ethanol, ethyl acetate; and almost insoluble in petroleum ether.

Uses It is used as a vasodilator.


8.2.3.3 Visnagin

Synonym Visnacorin.

Biological Sources It is obtained from *Ammi visnaga* Lam., (*Umbelliferae*).

* Egger. *Z. Naturforsch.*, **16B**, 697 (1962).

Chemical Structure



Visnagin

4-Methoxy-7-methyl-5H-furo [3, 2-g] [1]-benzopyran-5-one; (C₁₃H₁₀O₄).

Isolation Various sequential steps involved are as under:

1. The methanol mother-liquor remaining after the isolation of **khellin**, is evaporated to dryness under vacuo.
2. The resulting residue is taken up in benzene and treated subsequently with petroleum ether, until a distinct turbidity is accomplished.
3. The reaction mixture is cooled, and some small quantum of **khellin** shall separate out which is removed by filtration.
4. To the filtrate further addition of petroleum ether shall initiate the process of separation of visnagin as different crops that are removed, dried and subjected to distillation under vacuum carefully.
5. The fraction distilling between 150-155°C is collected and **visnagin** is finally recrystallized from methanol as prisms.

Characteristic Features These are as follows:

1. **Visnagin** is obtained as thread-like needles from water having mp 142-145°C.
2. It is found to be very slightly soluble in water; sparingly soluble in ethanol; and freely soluble in chloroform.


Identification Tests **Visnagin** when triturated with a piece of solid KOH or NaOH, it gives rise to a distinct rose-red colour which is certainly lighter in shade than that obtained with **khellin**.

8.2.4 Coumarin Bitter Principles

Coumarin nucleus is generated by the combination of benzene and α -pyrone as already mentioned under section 8.2.3. A good number of *coumarin bitter principles* have been isolated and characterized for their efficacious medicinal values, such as: **psoralen, methoxsalen, bergapten, imperatorin, angelicin and pimpinellin**. Out of these the first four drugs have already been discussed under the chapter on 'Phenylpropanoids' under Sections 6.2.3.3.1 through 6.2.3.3.4. The remaining *two* drugs shall now be described in the sections that follow:

8.2.4.1 Angelicin

Biological Sources It occurs in the fruit or root of *Angelica archangelica* L. (*A. officinalis* Moench) (*Umbelliferae*) (**Angelica; Garden Angelica; European Angelica**).

Chemical Structure


Angelicin

It is an angular furocoumarin.

Uses *Angelica* is useful for dyspepsia, enteritis, flatulence, gastritis, insomnia, neuralgia, rheumatism and ulcers.

8.2.4.2 Pimpinellin

Biological Sources It occurs in the fruits and rhizomes of *Pimpinellin saxifraga* L., *Heracleum spondylium* L.; *H. lanatum* Michx; and *H. panaces* belonging to the natural order *Umbelliferae*.

Chemical Structure


Pimpinellin

5, 6-Dimethoxy-2H-furo [2, 3-h]-1-benzopyran-2-one; (C₁₃H₁₀O₅)

Isolation **Pimpinellin** may be isolated by the methods suggested by Fujita and Furuya*, and by Svendsen *et al.***

Characteristic Features

1. It is obtained as off-white needles from methylene chloride/hexane having mp 119°C.
2. It is found to be practically insoluble in water; and soluble in ethanol.
3. It also undergoes isomerism to give rise to **isopimpinellin** as shown below:




Isopimpinellin

* Fujita, Furuya, *J. Pharm. Soc. Japan*, **74**, 795 (1954); **76**, 535 (1956).

** Svendsen *et al. Planta Med.*, **7**, 113 (1959).

Biosynthesis of Psoralen, Methoxsalen (Xanthotoxin), Bergapten, Angelicin and Isopimpinellin The various steps that are involved in the *biosynthesis of psoralen, xanthotoxin, bergapten, angelicin and isopimpinellin* are given below in a sequential manner:

1. **Umbelliferone** is first produced by the interaction of an isoprene unit with an appropriate alkylating agent *e.g.*, dimethylallyldiphosphate (DMAPP). Thus, the aromatic ring in the former gets duly activated at positions *ortho* to the hydroxyl group present in it.
2. The newly introduced dimethylallyl function present in *demethyl-suberosin* gets subsequently cyclized, having the phenolic moiety intact, to yield *marmesin*. However, this specific biotransformation is found to be catalyzed by a cytochrome P-450-dependent mono-oxygenase and also essentially requires cofactors such as NADPH and molecular oxygen.
3. It has been suggested appropriately that a *second* cytochrome P-450 dependent mono-oxygenase enzyme then cleaves off the hydroxyisopropyl portion (as a mole of acetone) from *marmesin*, thus producing the *linear furocoumarin psoralen*.
4. **Psoralen** is supposed to act as a precursor for the production of the subsequent series of further **substituted furocoumarins**, namely: **bergapten, xanthotoxin, and isopimpinellin** as shown below. Interestingly, such modifications are usually afforded due to steps taking place rather late in the biosynthetic pathway than occurring at the cinnamic-acid stage.
5. **Angelicin**—the so called *angular* furocoumarins, is the outcome of an identical sequence of reactions; however, these steps specifically involve dimethylallylation due to DMAPP at the alternative position *ortho* to the phenol.



Biosynthesis of Psoralen, Methoxsalen, Bergapten, Angelicin and Isoimpinellin

8.2.5 Coumarone Bitter Principles


The combination of the furan and benzene rings gives rise to the formation of benzofuran nucleus, otherwise termed as ‘**coumarone**’ as already depicted under Section (8.2.3). The most important coumarone based bitter principle is **rotenone** which shall now be discussed in an elaborated fashion in given below.

8.2.5.1 Rotenone

Synonym Canex

Biological Sources The principal insecticidal constituent of the dried **derris roots**, *Derris elliptical* Roxb. and *D. malaccensis* Prain, belonging to family *Leguminosae*; from **cube roots**, *Lonchocarpus utilis* and *L. urucu*, belonging to the natural order *Leguminosae*; from *Lonchocarpus nicou* (Aubl.) D.C. (*Leguminosae*); fruits and plant of *Piscidia piscipula* Sarg. (**Jamaica Dogwood**); and the roots of *Tephrosia virginiana* (L.) Pers (*Fabaceae*) (**Devil’s Shoe String**).

Chemical Structure



Rotenone

[2R-(2 α , 6 α , 12 α)]-1, 2, 12, 12a-Tetrahydro-8, 9-dimethoxy-2-(1-methylethenyl)-[1] benzopyrano [3, 4-6] furo [2, 3-h] [1] benzopyran-6 (6aH)-one; (C₂₃H₂₂O₆). It is a *rotenoid*.


Isolation Various steps involved in the isolation of **rotenone** are as follows:

1. The derris roots and rhizomes are dried, powdered, sieved and extracted with carbon tetrachloride in a Soxhlet apparatus for at least 24 hours.
2. The CCl₄ extract is filtered, concentrated under vacuo and allowed to cool at an ambient temperature for 24 hours, when crystals of **rotenone** separate out.
3. The resulting mixture is filtered through a gouche crucible under suction, and the crystals this collected are washed with a little CCl₄; and finally dried in the air.

Characteristic Features Its chemical features are:

1. It is an ‘**isoflavone compound**’ wherein the 2 : 3 double bond has undergone reduction.
2. Its heterocyclic portions are:
 - (a) A hydrobenzopyran moiety, and
 - (b) A hydrocoumarone (or 2 : 3-benzofuran) function.

- Rotenone** is a derivative of *tubic acid lactone* and is more commonly known as **6, 7-dimethoxy-2, 3-dihydro-benzopyran tubic acid lactone**.
- Decomposition of **rotenone** yields derric acid and tubic acid; and the latter further gives rise to a 'lactone' termed as the **tubic acid lactone** as shown below:



The physical parameters of **rotenone** are:

- It is usually obtained either as orthorhombic or as six-sided plates from trichloroethylene having mp 165–166°C; however, the dimorphic form has mp 185–186°C.
- Its specific optical rotation $[\alpha]_D^{20} - 228^\circ$ (C=2.22 in benzene).
- It gets decomposed upon exposure to air and light.
- Rotenone** is almost insoluble in water; and soluble in ethanol acetone, carbon tetrachloride, chloroform, ether, in addition to many other organic solvents.

Identification Tests These are as given below:

- Dissolve 2-3 mg of **rotenone** in 1 ml acetone and add to it 1 ml dilute HNO₃ (50% v/v), and allow it to stand for about one hour to cause the oxidation. Now, add to it a few drops of NaOH solution (10% w/v) when a distinct blue colour gets developed.
- Its colourless solutions in organic solvents normally oxidize upon exposure and become yellow, orange and then deep red finally. It may also deposit crystals of **dehydrorotenone** and **rotenonone** that are found to be toxic to insects.

Uses


- It is mostly used as a potent pesticide.
- It is widely employed as an acaricide and actoparasiticide in cattles.
- The action of **rotenone** closely resembles to that of *pyrethrin* in affecting a rapid *knock-down* of the flying insects (*e.g.*, house-flies, mosquitos etc.); and is found to be comparatively harmless to the warm-blooded animals.

4. As **rotenone** does not leave any harmful residue, it may be employed with enormous safety for most delicate and precious garden crops and garden plants.

Note Interestingly, though the derris roots do contain a natural insecticidal principle (**rotenone**) they are nevertheless prone to infestation by some specific types of insects obviously unaffected by rotenone.

Biosynthesis of Rotenone In general, many thousands of wide variety of **isoflavonoids** have been duly isolated, characterised and identified; and subsequently their structural complexity have been resolved logically and methodically by first carrying out the hydroxylation, and secondly by alkylation reactions. These reactions not only helped in varying the oxidation level of the heterocyclic ring, but also produced additional heterocyclic rings.

In the **biosynthesis of rotenone** a simple **isoflavone** called **daidzein** is the starting material which undergoes methylation in the *para* position of the phenyl ring attached to the pyran ring with a covalent bond thereby forming an isoflavone termed as **formononetin**. This undergoes further biotransformations as stated earlier to yield **rotenone**. It contains a C₅ isoprene unit, as could be observed in most of the natural **rotenoids**,* which is afforded *via* dimethyl-allylation of **demethylmunduserone**.



8.2.6 Miscellaneous Bitter Principles

There are some **bitter principles** which normally do not fall into the various categories already discussed from Sections 8.2.1 through 8.2.5. A few important potent drugs used as bitter principles that belong to this group are, namely: **picrotoxin**, **quassin**, **cantharidin**, which shall now be described separately as under:


8.2.6.1 Picrotoxin

Synonym Cocculin.

Biological Sources It is obtained as the **bitter principle** from the seed of *Anamirta cocculus* L. Wight & Arn. (*Menispermaceae*); and also found in *Tinomiscium philippinense* Diels.

* **Rotenoids:** The rotenoids take their name from the first known example *rotenone* and are usually generated by ring cyclization.

Chemical Structure **Picrotoxin** is a molecular compound of one mole **picrotoxinin** ($C_{15}H_{16}O_6$), q.v., and one mole **picrotin** ($C_{15}H_{18}O_7$), q.v., into which it is readily separated. Thus, **picrotoxin** may be resolved into the two components by boiling with 20 parts of benzene. In this manner, **picrotoxinin** remains dissolved in benzene whereas **picrotin** that is practically insoluble in benzene can be separated easily. Likewise, this cleavage may also be accomplished by chloroform more efficiently. Thus, we have:



Isolation Various steps involved in the isolation of **picrotoxin** are:

1. The seeds are dried, powdered coarsely, sieved and defatted with petroleum ether in a Soxhlet apparatus.
2. The defatted powder (marc) is subsequently extracted by boiling with ethanol or with water.
3. The filtrate thus obtained is treated with lead acetate solution (5% w/v), filtered and the excess of 'lead' is removed by passing freshly generated H_2S gas (*i.e.*, Pb is precipitated as Pb S).
4. The resulting solution is filtered, residue discarded and the filtrate is concentrated to a syrupy consistency in a Rotary Thin Film Evaporator. The syrupy liquid is kept in a refrigerator overnight.
5. **Picrotoxin** crystallizes out as a crude substance.
6. It may be further purified by treating with ethanol or boiling water and activated charcoal powder to obtain the pure substance.

Characteristic Features

1. It is obtained as shiny rhomboid leaflets mp $203^\circ C$.
2. It has an intense bitter taste and is *extremely poisonous*.
3. It has specific optical rotation $[\alpha]_D^{16} - 29.3^\circ$ ($C = 4$ in absolute ethanol).
4. **Solubility Profile:** 1 g dissolves in 150 ml cold water; 45 ml boiling water, in 13.5 ml 95% ethanol, in 3 ml boiling ethanol; sparingly soluble in ether, chloroform; and readily soluble in aqueous solution of NaOH and in strong NH_4OH .
5. It is highly toxic to fish.
6. It is stable in air, but is affected by light.
7. Picrotoxin is almost neutral to litmus.

Identification Tests These are as stated below:

1. **Sulphuric Acid Test:** Dissolve 2-3 mg of **picrotoxin** in a few drops of sulphuric acid, a golden-yellow-colour is produced that gets changed to reddish-brown gradually.

- Anisaldehyde Test:** Moisten a few crystals of **picrotoxin** with H_2SO_4 and just add 1-2 drops of a solution of anisaldehyde in dehydrated ethanol (1 : 5), a permanent blue colouration is produced.
- Potassium-Cupric Tartrate Test:** Add about 5-10 mg of **picrotoxin** to 2 ml of potassium-cupric tartrate solution (0.5%) with 10 ml of water, a red precipitate is formed gradually in the cold, but a little faster on warming.
- Vanillin HCl Test:** A few mg of picrotoxin when boiled with vanillin hydrochloride solution (0.1% w/v) it gives rise to a green colouration.
- Reduction Tests:** **Picrotoxin** reduces Fehling's solution to give a brick-red precipitate; and **Tollen's reagent** to give a silver mirror.
- Mix 0.2 g KNO_3 with four drops of H_2SO_4 in an evaporating dish. Sprinkle a few crystals of **picrotoxin** on the resulting mixture and add dropwise NaOH solution (2N), until it is present in a little excess quantity. The crystals of picrotoxin shall initially acquire a red colouration that fades out slowly.

Uses


- It is used as a CNS-stimulant. Therefore, it may be employed intravenously as an antidote in barbiturate poisoning and other narcotics also.
- It also finds its application as an effective respiratory stimulant.
- Very small quantities of the powdered seeds are sufficient to stupify fish.

8.2.6.2 Quassin

Synonym Nigakilactone D

Biological Sources It is obtained from the wood of *Quassia amara* L., (*Simaroubaceae*) commonly known in commerce as **Surinam quassia**. It is also obtained from the stem wood of *Picrosma excelsa* (Sw.) Planch. (*Aeschrion excelsa* or *Picroena excelsa*) known in commerce as **Jamaican quassia**. All these species belong to the natural order *Simaroubaceae*.

Chemical Structure



Quassin

2, 12-Dimethoxypicrosa-2, 12-diene-1, 11, 16-trione; (C₂₂H₂₈O₆).

Isolation The following steps may be adopted in a sequential manner for the isolation of **quassin**.

- The **quassia wood** is chopped into small pieces and subjected to aqueous decoction, which is filtered and concentrated to the original weight of the wood taken; and finally neutralized carefully with Na_2CO_3 .

2. Tannic acid solution (5% w/v) is added slowly until no more precipitate is obtained.
3. The precipitate thus obtained is filtered, collected and transferred to a pestle and mortar, triturated with solid lead carbonate (or with freshly prepared lead oxide), so as to liberate **quassin** and form lead tannate; and the resulting mass is dried on a water bath.
4. The dried mass is powdered and then subjected to extraction with 80% (v/v) ethanol successively.
5. The combined ethanolic extract is filtered and concentrated under vacuo and left for cooling overnight, when the crystals of **quassin** would separate out.

Quassin may also be obtained by the resolution of the mixture of bitter constituents of **quassia wood** by the method of London *et al.**

Characteristic Features

1. **Quassin** is obtained as rectangular plates from dilute methanol having mp 222°C.
2. Its specific optical rotation $[\alpha]_D^{20} + 34.5^\circ$ (C = 5.0 g in CHCl₃).
3. It has uv_{max} : ~ 255 nm ($\epsilon \sim 11,650$).
4. It is extremely bitter; and it has the bitterness threshold 1 : 60,000.
5. It is found to be freely soluble in benzene, acetone, ethanol, chloroform, pyridine, acetic acid, hot ethyl acetate; and sparingly soluble in ether and petroleum ether.

Identification Tests

1. Add to a few crystals of **quassin** 2-3 drops of concentrated, H₂SO₄ and sucrose when a red colouration is produced.
2. **Phloroglucin Test:** Dissolve 2-3 mg **quassin** in 1-2 ml ethanol, and add to it a few crystals of phloroglucin and a few drops of concentrated. HCl, when a crimson red colour is obtained.


Uses

1. It possesses insecticidal properties.
2. The *quassia wood extract* is used as a bitter tonic.
3. **Quassin** exhibits anthelmintic properties, and on being administered as enema *expels thread worms* specifically.

8.2.6.3 Lactucin

Biological Sources It is obtained from the dried milky juice of *Lactuca virosa* L. (*Asteraceae*) (**Bitter Lettuce; Wild Lettuce**); and from the plant of *Cichorium intybus* L. (*Compositae*).

Chemical Structure



Lactucin

* London *et al.*, *J. Chem. Soc.*, 3431, 1950.

[3aR – [3a α , 4 β , 9a α , 9b β)]-3,3a,4,5,9a,9b-Hexahydro 4-hydroxy-9-(hydroxymethyl)-6-methyl-3-methyl-eneazuleno [4, 5-b] furan-2, 7-dione; (C₁₅H₁₆O₅).

Isolation **Lectucin** may be isolated by the method suggested by Schenck *et al.**

Characteristic Features

1. It is obtained as crystal from methanol which sinters at 218°C and has mp 228-233°C.
2. It exhibits specific optical rotation $[\alpha]_D + 49^\circ$ (C = 0.90 in methanol); and $+77.9^\circ$ (C = 3.44 in pyridine).
3. It has uv_{max} : 257 nm (ϵ 14,000).
4. It is found to be soluble in water, ethanol, methanol, ethyl acetate, anisol and dioxane.


Identification Tests It may be identified from its derivative:

Lectucin *para*-hydroxyphenylacetate hydrate (C₂₃H₂₂O₇) (Intybin; Lactucopicrin); It is obtained as crystals from water which get decomposed at 148-151°C. It shows specific optical rotation $[a]_D^{17.5} + 67.3^\circ$ (pyridine).

8.2.6.4 Erythrocentaurin

Biological Sources It is obtained from the plant *Centaureum umbellatum* Gilib. (*Erythraea centaureum* Pers.), *Gentianaceae* or *Swertia japonica* (Maxim.) Makino *Gentianaceae*. It is also accomplished by carrying out the hydrolysis of **swertiamarin** and **erytaurin** with emulsin.

Chemical Structure



Erythrocentaurin

5-Formyl-3, 4-dihydroisocoumarin; (C₁₀H₈O₃).

Isolation **Erythrocentaurin** may be isolated from *C. umbellatum* by the method of Kariyone and Matsushima.**

Characteristic Features

1. It is obtained as long needles having mp 140-141°C.
2. It turns red on being exposed to sunlight.
3. It has uv_{max} : 223, 290 nm ($\log \epsilon$ 4.30, 3.13).

Uses It is mostly employed as a bitter tonic.

* Schenck *et al.*, *Arch. Pharm.*, **294**, 17 (1961).


** Kariyone and Matsushima J., *Pharm. Soc.*, Japan, **47**, 25 (1927).

8.2.6.5 Gentisin

Synonyms Gentianin; Gentiin; Gentianic Acid.

Biological Sources It is obtained from the roots of *Gentiana lutea* L. (*Gentianaceae*) (**Yellow Gentian**).

Chemical Structure



Gentisin

1, 7-Dihydroxy-3-methoxy-9H-xanthen-9-one; (C₁₄H₁₀O₅).

Characteristic Features

1. It is obtained as yellow needles from ethanol having mp 266-267°C.
2. It has uv_{max} (methanol): 260, 275, 315, 410 nm ($\log \epsilon$ 4.35, 4.30, 4.10, 3.70).
3. It is observed to be very slightly soluble in water or organic solvents.

Identification Test

Gentisin Diacetate (C₁₈H₁₄O₇) It is obtained as crystals from ethanol having mp 196-197°C. Its absorption max (metanol): 240, 270, 300 nm ($\log \epsilon$ 4.58, 4.05, 4.10).


Uses It may be used to stimulate gastric secretion, improve appetite and digestion, and alleviate debility.

8.2.6.6 Cantharidin

Synonym Cantharides Camphor.

Biological Sources It is the active vesicating principle of cantharides (q. v) and other insects, in notorious '**Spanish Fly**' aphrodisiac, which essentially comprise of the dried insects (Beetles) *Lytta* (*Cantharis*) *vesicatoria* belonging to the order *Coleoptera*; and family *Meloidae*. It has been found that the soft parts of the insect are the chief seat of **cantharidin**. Besides, cantharidis contain 0.5 to 0.95 of **cantharidin**.

Chemical Structure



Cantharidin

Exo-1, 2-*cis*-Dimethyl-3, 6-epoxy hexahydrophthalic anhydride; (C₁₀H₁₂O₄).

Isolation The various steps involved in the isolation of **cantharidin** are:

1. The dried insects are collected and powdered. It is now treated with an acid whereby the **cantharidin** gets liberated in the form of its corresponding salts.
2. The resulting product is subjected to extraction, of both **cantharidin** and fat, by the help of ethyl acetate in a Soxhlet apparatus.
3. The solvent is removed carefully under reduced pressure and the crude cantharidin crystallizes out.
4. The fat may be removed by the help of petroleum ether, in which **cantharidin** is only negligibly soluble.
5. Ultimately, the crude defatted **cantharidin** is dissolved in a minimum quantity of hot ethanol and allowed to cool when cantharidin crystallizes out in its purest form.

Characterstic Features These are as follows:

1. **Cantharidin** is obtained as orthorhombic plates or as scales having mp 218°C.
2. It sublimes at 110 °C (12 mm Hg, 3-5 mm distances).
3. It is practically insoluble in cold water and somewhat soluble in hot water. 1g dissolves in 40 ml acetone; 65 ml chloroform; 560 ml ether; 150 ml ethyl acetate; and soluble in oils.

Identification Tests

1. **Formaldehyde Test:** Add to a few crystals of **cantharidin** 1-2 drop of dilute formaldehyde solution mixed with H₂SO₄, the development of a brown to black colouration on warming identifies it.
2. A solution of **cantharidin** in olive oil is vesicant to the skin (*i.e.*, sensitive upto an extent of 0.14 mg).

Uses

1. It is mostly used as a vesicant.
2. It is also employed as a rubefacient and counterirritant in veterinary practice.

FURTHER READING REFERENCES

1. Barakat, Z., and Badran, N., 'Identification of Khellin, Visnagin and Khellol Glucoside', *J. Pharm. Pharmacol.*, **3**, 576, (1951).
2. Bisset N.G. (Ed.): Max Wichtl—**Herbal Drugs and Phytopharmaceuticals**, CRC Press, London, 1994.
3. Brown, S.A., **Recent Studies on the Formation of Natural Coumarins**, *Lloydia*, **26**, 211, 1963.
4. Dewick, P.M., 'Medicinal Natural Products—A Biosynthetic Approach', John Wiley & Sons, Ltd., England, 2nd. edn., 2002.
5. Duke, J.A., 'Handbook of Medicinal Herbs', CRC-Press, New York, 2001.
6. Evans, W.C., **Trease and Evan's Pharmacognosy**, W.B. Saunders Company Ltd., London, 14th, edn, 1996.
7. Hostettmann K. *et al.* (eds.), **Phytochemistry of Plants Used in Traditional Medicine**, Proceedings of Phytochemical Society Europe, 37, Oxford Science, New York, 1995.
8. Jisaka M. *et al.*, **Antitumoral and Antibacterial Activities of Bitter Sesquiterpene Lactones of *Vernonia amygdalina*: a possible Medicinal Plant used by Wild Chmpanzees**, *Biosci. Biotech. Biochem.*, **57**: 833-834, 1993.
9. Newall, C.A., Anderson, L.A., and Phillipson, J.D.,: **Herbal Meicines—A Guide for Health care Professionals**. The Pharmaceutical Press, London, 1996.
10. Ramstad, E.: **Modern Pharmaceognosy**, McGraw Hill Book, Co., London, 1959.

9

Antibiotics

- Introduction
- Antibiotic Development
- Classification of Antibiotics
- Further Reading References

9.1 INTRODUCTION

Antibiotics, in today's most up-to-date therapeutic armamentarium, occupy strategically the most coveted and key position during the span of past half-century across the globe. This conglomeration of drugs affords an effective management and critical control of a host of deadly human related pathogenic microorganisms which previously caused pathetic prolonged human sufferings or ultimately leading to death irrespective of the physical condition, age factor or economic status of an individual.

The word '**antibiotic**' has been coined from the term antibiosis that evidently means '**against life**' (*anti*—against and *bios*—life).

Over the years various versions of 'definitions' for an **antibiotic** have been postulated which are enumerated as under.

The most widely accepted definition of an **antibiotic** accepted by the scientific jargons is—'**a chemical substance produced by a microorganism, that has the capacity, in low concentration, to inhibit or kill, selectively, other microorganisms.**' This definition lays particular emphasis on the terminology '*selectivity*' or '*selective toxicity*' that explicitly suggests that the substance either checks the growth of pathogens or exerts a bactericidal action on the microbes without displaying a likewise action on the host organism *i.e.*, the human beings.

The above definition clearly excludes the compounds having the pure synthetic genesis (origin). However, in a rather broader perspective these '*synthetic substances*' are virtually treated at par with the natural compounds along with their corresponding derivatives under the terminology '**antimicrobials**' which may be further categorized into **antifungals** and **antibacterials** based on the particular type of **microbe** undergoing inhibition. Hence, in order to circumvent the practical aspects, both the terminologies *viz.*, '**antibiotic**' and '**antimicrobial**' may be employed interchangeably irrespective of the particular source of the compound.

Even in the ancient and primitive era, dating back to 2500 years, the anti-infective characteristic features of *fungi* and *moulds* usually observed in various food products like: mouldy bread, yoghurt, and soybean curds, and other similar materials to wounds and boils to curb their infection. This sort of age-old treatment one may regard as a **folk-medicine style of antibiotic therapy**.

It is, however, pertinent to mention here that the real impetus and legitimate recognition of the antibiotics in the so called 'modern drugs' was virtually accomplished by the famous french scientist Louis Pasteur. The epoch making introduction of *pyocyanase* interestingly extracted from *Pseudomonas aeruginosa* as a prominent therapeutic agent under the '**antibiotics**' is indeed one of the greatest achievements in the history of medicine. This event was immediately followed by another historic invention of Alexander Fleming for the drug penicillin; and the subsequent antimicrobial activity of *Penicillium notatum* discovered by **Chain Florey and his co-workers**.

In fact, the most effective and wonderful class of life saving antibiotics comprise of a plethora of active substances that are found to be effective on either Gram +ve or Gram -ve micro-organisms; besides the ones that are invariably known as the **broad-spectrum antibiotics**.

In general, the **antibiotics** are produced on a large scale by *three* known methods, namely: (a) fermentation process; (b) semi-synthetic process; (c) synthetic process. Recently, with the advent of a tremendous quantum jump and diversification in the specific field of '**biotechnology**', the first two processes stated above have not only gained an enormous increase in the rate of production but also improved their yield and purity. Nevertheless, the fermentation process is further categorized into *two* types: (a) surface method; (b) submerged method. It is worthwhile to mention here that the second method has a much greater efficiency limit and hence used commercially. Over the years, a vast number of altogether newer, purer, and high-yielding **microbial strains** have been developed, tried and tested for evaluating their **antibiotic** yielding strength besides the efficiency in their extraction.

9.2 ANTIBIOTIC DEVELOPMENT

The latest progressive trend in the logistic features of **antibiotic development** may be expatiated by the following sequence of objectives, namely:

- (a) To screen and evaluate different types of sources of microorganisms for detection of purposeful antagonism.
- (b) To identify and select modified versions of microbial mutants, establish optimal environmental and nutritional conditions, and to develop appropriate methods for recovering antibiotics from cultures.
- (c) To induce the production of particular desired metabolites.
- (d) To improve upon and modify the fermentative metabolites either by the help of chemical or biological manipulations to accomplish more useful antibiotic products (compounds).
- (e) To develop detailed methods for '**total synthesis**' of antibiotics from *ab initio* for a feasible economic advantage, and
- (f) To make use of an adjunct agent to distinctly enhance the impact or availability of an '**antibiotic**'.

9.2.1 Quest for New Antibiotics

In the quest for new **antibiotics**, rather simpler, standardized and quicker procedures have been developed and established for screening viable microorganisms having antibiotic-yielding capability. In actual practice, however, the soil samples are the choicest candidates towards an endeavour to identify the microbes for the simplest logical reason that they are considered as the richest source of

antibiotic-producing organisms. Interestingly, majority of these organisms happen to be the bonafide members of a specific class of branching, procaryotic microorganisms which essentially retain a coveted status in their morphologic characteristic features between bacteria and fungi. A survey of literature reveals that between early fifties to late seventies the microbial sources of antibiotics discovered in Japan and USA mainly comprise of actinomycetes (85%), fungi (11%), and bacteria (4%).

The following are the summary of the most prominent **genera** and their **taxonomic relations**.

Genus	Streptomyces	Order	Penicillium
Phylum	–	Eumycophyta (Fungi)	–
Class	–	Ascomycetes	Deuteromycetes (Fungi Imperfecti)
Order	Actinomycetales (Actinomycetes)	Aspergillales	Moniliales
Family	Streptomycetaceae	Aspergillaceae	Moniliaceae

In general, nowadays a great deal of emphasis is being focused upon the pathogens responsible for causing mostly incurable fungal and viral infections, besides the bacterial infections, such as: **methicillin-resistant** *Staphylococcus* and *Pseudomonas* species.

Following are the various steps involved in the so called ‘general method’ for the methodical screening of **newer antibiotics**, namely:

Step I: Treatment of the soil sample (or sample from other sources) by an antifungal chemical antibiotic, **cycloheximide** which specifically checks the growth of interfering bacteria and fungi but nevertheless affects the *actinomycetes*. Besides, a diluted solution of phenol (1 : 140) may also be used as an antibacterial agent.

Step II: The treated sample, in their varying known dilutions are subsequently streaked on agar plates containing medium (nutrients) which augments and accelerates the growth of actinomycetes.

Step III: The streaked agar plates are incubated for 3 to 7 days between 25-30°C; and examined carefully for their characteristic colonies of actinomycetes. After due physical identification these colonies are selectively transferred onto fresh medium aseptically.

Step IV: Well grown big cluster of colonies of the above selected organisms are cut in such a manner that the ‘*plugs*’ comprise of both the organisms and the underlying agar.

Note: In case, the isolated organisms produces an antibiotic, it must normally diffuse into the agar medium.

Step V: The ‘*plugs*’ are meticulously removed and placed on an agar plate which has already been seeded with a specific ‘**test organism**’ that clearly shows a positive indication of the potential effectiveness and usefulness of the antibiotic in question.

Step VI: All the ‘**test plates**’ are duly incubated for a stipulated temperature and duration required for the maximum (optimum) growth of the ‘**test organisms**’. In case, there exists a clear zone of inhibition around the ‘**plug**’ of the actinomycete, it may be inferred that an ‘**antibiotic component**’ is present in the ‘**plug**’ which obviously inhibited the growth of the ‘**test organisms**’.

Step VII: ‘*Dereplication*’ *i.e.*, to establish by appropriate means as to whether the chemical substance (**‘antibiotic component’**) which affected the inhibition is either an already known compound* or happens to be a ‘*new antibiotic*’. In short, if the newly discovered **‘antibiotic component’** is really promising and possesses remarkable marked and pronounced **antibiotic activity** only then it will be subjected to further thorough investigation.

9.2.2 Large-Scale Production

Always, the ultimate decision to carry out the large-scale production of a ‘*new antibiotic*’ is based on several cardinal qualifying factors, such as: (a) its chemical properties, (b) its physical characteristics, and (c) its detailed biological activities.

However, there are *two* extremely vital requirements for production, namely:

- (i) The organism should produce the **‘new antibiotic’** most preferably, in a submerged culture as opposed to a surface culture, and
- (ii) The organism should liberate and excrete the **‘new antibiotic’** right into the prevailing culture medium.

There are, of course, some other important considerations also for the large-scale production of a **‘new antibiotic’** that are of rather minor nature, such as:

- (i) A few **‘antibiotics’** are produced in the cells of the organisms and therefore, requires altogether special cost-involving extraction procedures for their final recovery.
- (ii) Some other minor but equally important related considerations are, namely: **minimum inhibitory concentration (MIC)** against the strains of pathogenic organisms, chemical stability, activity *in vivo*, and lastly the toxic manifestations in mammals.

The most intricate, diligent and marvellous exploitation of the wisdom of the man in the application of the in-depth knowledge of **microbiology, biotechnology, pharmaceutical chemistry, and engineering** has ultimately opened the flood gate towards the development of **‘newer antibiotics’** and their commercial production to curtail the existing human sufferings.

The various important sequential procedural steps that are essentially required for the large-scale production of antibiotics are stated as under:

- (i) Invariably requires growth of the producing organisms in aerated stainless steel tanks with a capacity to hold thousands of gallons of the respective nutrient medium.
- (ii) The fermentation process is duly initiated with the help of spores or occasionally, vegetative growth from a pure **stock culture**** of the organism.
- (iii) The inoculation of the huge fermentation tanks are normally accomplished by carrying out successively the transfer of the organism to increasingly greater volumes of nutrient. The major advantages of making use of a large standard inoculum are as stated below:

* Based on the chromatographic, physico-chemical properties, antibiotic spectrum and comparing the same to a database of previously identified compounds.

** **Stock Cultures:** These are maintained very carefully (*e.g.*, by lyophilization) that essentially require transfer as infrequently as possible, as repeated transfers may ultimately select only those cells of the organism which are rather poor generators of antibiotic.


- (a) Considerable reduction in the total incubation time required for the normal production of the antibiotic,
- (b) Reduces importantly the slightest possible chance for undesired costly contamination by foreign microorganisms, and
- (c) Caters for the best ever possible scope and opportunity for the entire control and management of subtle nutritional and environmental factors that vitally influence the ultimate yield of the *antibiotic*.

9.2.2.1 Phases in Fermentative Process

In fact, there are *two* important and distinct phases normally encountered in the fermentative process, namely:

- (a) **Growth Phase of the Organism:** It is also sometimes referred to as the ‘**trophophase**’; wherein the number of organisms per unit time increases progressively, and
- (b) **Idiophase of the Organism:** In the idiophase there is a substantial antibiotic production; and hence, invariably termed as the ‘**antibiotic production phase**’.

The above mentioned *two* phases in the fermentative process may be further explained with the help of the following diagram:



(Adapted from: Robbers, J.E. *et al.* 'Pharmacognosy and Pharmacobiotechnology', Williams & Wilkins, London, 1996)

In this particular instance, both the **growth phase** and the **idiophase** in the course of a typical '**penicillin fermentation**' performed in a culture-medium consisting of:

- (i) **Source of carbon nutrition:** – *e.g.*, lactose and glucose;
- (ii) **Nitrogen sources:** – *e.g.*, corn steep liquor; and
- (iii) **Phosphate buffer:** – to provide *P* in the medium and also to maintain the pH of the medium.

The observations from the above diagram are as follows:

- (a) The growth of microorganisms is shown in the above diagram by the curve indicating an enhancement of mycelial nitrogen (Mycelial N). This particular phenomenon continues right from the beginning (0 hours) of the culture period to nearly one day (24 hours).

Note: In the 'growth phase', the culture becomes thick by virtue of the formation of 'aggregates of fungal cells' usually known as mycelium.

- (b) Glucose is preferentially consumed as compared to lactose specifically in the 'growth phase', as it may be employed as a prime source of C directly.
- (c) Ammonia (NH₃) gets liberated also in the 'growth phase' which is caused due to the deamination of various amino acids present in the *corn-steep liquor* (medium).
- (d) Release of NH₃ evidently increases the pH of the medium from acidic to almost 7 (neutral). Thus, the ideal and optimum pH necessarily required for the stability of 'penicillin' is 7, which is maintained by adding adequate 'phosphate buffers' into the medium.
- (e) The 'penicillin production' happens to rise very progressively and rapidly between 24-48 hours.

Note: Just in the initial stage of 'penicillin production', glucose gets fully utilized, and subsequently the fungus makes use of 'lactose' as a source of C.

- (f) Interestingly, no additional growth takes place as the lactose cannot be used as such unless and until it gets converted to glucose and galactose *via* hydrolysis. Hence the prevailing decreased availability of C in the medium obviously offers a 'triggering mechanism' in the production of penicillin.

9.2.2.2 Enhancing Yield in Large-Scale Production

During the past half-a-century an enormous volume of intensive and extensive research has been duly carried out by different groups/individuals across the world to determine and establish the optimal nutritional and environmental parameters required necessarily for antibiotic production. In reality, these conditions are certainly not quite similar to those required for maximum vegetative growth.

The various factors that exert vital impact upon the qualitative and quantitative antibiotic production are enumerated below:

- Sources of nutritional C and N
- Ratio of C/N in nutrients
- Mineral composition of medium
- Temperature of incubation
- Initial pH, control and management of pH during the entire course of fermentation.
- Aeration mode and rate
- Time-phase for addition of special growth and antibiotic enhancing materials.

Empiric Observations The selection of optimal fermentation parameters is not only based on certain empiric observations but also serve as critical factors.

Examples

1. A few strains of microorganism *Bacillus subtilis* give rise to the maximum yields of **bacitracin*** at a C & N ratio of 1 : 15; but at a lower ratio 1 : 6 it forms **licheniformin**** which happens to be a structurally related but an undesired commercial antibiotic.

* **Bacitracin:** Its antibacterial actions are similar to those of penicillin, including Gram + ve cocci and bacilli and some Gram -ve organisms. Because of its toxicity when used parenterally, it is normally used topically in ointment form.

** **Licheniformins:** These are antibiotic substances usually produced by *Bacillus licheniformis*.

2. **Phenylacetamide** or related substances when added to the culture medium of penicillin production though exhibits a very negligible effect on the yield of penicillin compounds, yet shows a very significant improvement upon the ultimate composition of the penicillin mixture.
3. **Phenylacetic Acid Derivative's** inclusion as a part and parcel in the nutrient mixture composition is observed to influence favourably the production of **Penicillin G**; and this particular vital step has considerably minimised the tedious problems with regard to the use of either unknown or variable composition of mixtures; besides, the significant cost, time and energy involved unnecessarily in separating the individual antibiotic substances.
4. **Acyl Moieties:** The application of different acyl groups so as to achieve the fermentative production of certain other penicillins, for instance: phenoxy-methylpenicillin (or Penicillin V) could not achieve appreciable feasible success in large-scale production; but surprisingly, the various semisynthetic techniques evolved not only superseded this specific line of action but also greatly enhanced the production of specialized penicillins.
5. **Mercaptothiazole:** The incorporation of mercaptothiazole in cultures of *Streptomyces aureofaciens* certainly approves the doctrine that certain 'chemical additives' might be useful without necessarily being introduced into the antibiotic molecule partially or fully.
6. **Effect of Enzyme Induction:** It has been proved beyond any reasonable doubt there are certain 'chemical additives' that may enhance the antibiotic production by means of an enzyme induction effect.

Example: Methionine when added to a *cephalosporin C* fermentation process, during the growth phase of the organism (*i.e.*, '**trophophase**') there is an apparent stimulation observed in the actual production of the antibiotic. As methionine does not behave as a precursor to the antibiotic in its biosynthetic process, in comparison to the performance of phenylacetic acid in the biosynthesis of Penicillin G, one may conclude and infer with rather stress and emphasis that methionine stimulates the ultimate production of cephalosporin C biosynthetic enzymes.

7. **Inhibition of Antibiotic Production:** Lysine exhibits an inhibition of penicillin fermentation by its presence in the culture medium which ultimately retards the antibiotic production. This particular phenomenon may, however, be explained by the fact that both lysine and penicillin are the end products of a branched biosynthetic pathway wherein the alpha-amino adipic acid serves as a '**common precursor**'. The production of '**lysine**' is regulated and monitored by two processes, *viz.*, repression or inhibition of the requisite enzymes needed for the production of alpha-amino adipic acid. Hence, lysine puts a halt of alpha-amino adipic acid formation which finally causes a decrease in the production of penicillin.
8. **Mutation and Strain Selection: Mutation*** influenced and persuaded by virtue of exposure of the parent-strain to uv-light, X-rays, or a host of mutagenic chemical substances *e.g.*, analogues of purines and pyrimidines, nitrogen mustards (*viz.*, **mechlorethamine hydrochloride, mephalan, cyclophosphamide, chlorambucil**)** is widely recognized as the most virile and versatile means for the selection of **improved strains**.

* **Mutation:** A change in a gene potentially capable of being transmitted to offspring.

** Kar, A., **Medicinal Chemistry**, New Age International (Pvt) Ltd, New Delhi, 4th edn, 2006.

It is, however, pertinent to mention here that a constant search across the globe of natural sources for either newer wild-type(s) or various diversified species of organisms that afford to yield the ‘antibiotic’ in much higher percentage than the original one. In the particular instance of induced mutations, lethal levels of the mutagen are adjusted in such a manner so that nearly 90-99% of the cells of the organism are destroyed (killed). Thus, the high-antibiotic-yielding mutants are selected meticulously from the remaining surviving cells.

Example: Production of Penicillin: Initially, a penicillin antagonism was noticed from a culture of *Penicillium notatum* Westling, that yielded a meagre 4 mg L^{-1} of penicillin from its culture medium. In other words, no mutation of *Penicillium notatum* were ever observed in the early selection process which could have given a significant yield of penicillin in the submerged fermentation technique. In 1944, there was an unique breakthrough in research whereby through the natural selection, a strain of *Penicillium chrysogenum* Thom was invented that raised the yield of penicillin almost by 10 times *i.e.*, 40 mg L^{-1} . Later on, with the help of vigorous modification of mutation techniques amalgamated with strain-selection, the ultimate yield of penicillin has been successfully enhanced to $21,000 \text{ mg L}^{-1}$.

The recent quantum advancement in the field of **molecular biology** there has been a tremendous expansion with specific reference to the knowledge of molecular regulation related to antibiotic biosynthesis. In this manner perhaps one may accomplish greater heights in the antibiotic production through such measures as:

- Rational manipulation(s) of the antibiotic-producing organisms to enhance its yield significantly.
- Deregulating the particular rate-limiting biosynthetic enzymes.
- Introduction of additional ‘copies of genes’ matching the rate-limiting steps.
- Rational implementation of specific genes for parallel/alternate biosynthetic routes.
- Production of ‘hybrid-antibiotics’ through the fermentatively-generated structural analogues of the natural antibiotic molecules.

9.2.2.3 Separation and Isolation of Antibiotics

Generally, the large-scale-produced antibiotics are released rapidly right into their nearest environment *i.e.*, the nutrient medium, where they get accumulated. However, there are some other instances *e.g.*, the peptide antibiotics, wherein the specific antibiotic is stored endocellularly (within the cells); the fermentation is maintained unless and until the cells accomplish an advanced matured physiologic age, the process of fermentation is arrested (ceased) whereby majority of the cell membranes have either lost their selective retention characteristic property or have undergone lysis—thereby releasing the antibiotic into the surrounding medium. In other words, therefore, the isolation process of various antibiotic substances is nothing but purely a recovery from the culture broth. The various **standard operating procedures (SOPs)** essentially comprise of: selective precipitation, specific adsorption, or finally the chosen extraction with an immiscible solvent.

In fact, in an ideal situation the very first isolation process must be as crisp, selective and efficient as possible so as to achieve the maximum yield, besides to help in subsequent purification without any cumbersome method. However, the particular chemical characteristic feature of an antibiotic shall be the ultimate determining factor, and also their accompanying metabolites to guide and dictate the manipulative procedures which may be adopted effectively in any particular instance.

Obviously, a balanced compromise procedure that is economically viable and feasible shall be the **'ideal procedure'** for all practical purposes.

The various means of extraction and purification of **'antibiotic substances'** may be accomplished through a number laid-down, tested and tried techniques that shall now be discussed briefly as under:

- (a) **Liquid-Liquid Extraction:** Invariably the application of certain water-immiscible organic solvents *e.g.*, chloroform, solvent ether, carbon tetrachloride etc., are exercised for the extraction of most antibiotics. This particular process has evidently *two* major disadvantages, namely:
- (i) Lacks high-degree of selectivity because majority of solvents, which are fairly cheap and hence economical, tend to be employed on a large-scale production, and
 - (ii) Comparatively inefficient as most of the known *'antibiotic substances'* are generally highly polar molecules.

Interestingly, in most instances the above *two* serious drawbacks are easily circumvented by adopting a **chemical-engineered-flow process**; but even then the highly polar 'antibiotics' fail to separate in which the partition-coefficient obviously favours the aqueous phase.

- (b) **Recovery through Adsorption:** Extremely polar antibiotics *i.e.*, **the aminoglycoside antibiotics**, such as: **neomycin, streptomycin, paromomycin, kanamycin, amikacin, gentamycin, tobramycin, netilmicin**, are normally recovered from the culture medium through adsorption on certain appropriate adsorbent. It has been observed that—

- Most adsorbents remove highly polar antibiotics from culture media with varying degree of selectivity.
- Selecting a suitable adsorbent offers major limitations by virtue of the fact that while applying reversal of the adsorption process for recovering the antibiotic(s) very careful and moderate conditions be applied so as to avoid its possible denaturation/destruction.
- Ideally, the application of controlled-activity grade charcoal as an adsorbent, and subsequent elution with a dilute mineral acid (H_2SO_4) is normally employed as an universal method of choice.

- (c) **Chromatography-Recrystallization-Standard Manipulations:** It is, however, pertinent to state here that as soon as one is able to lay hands onto the **'crude antibiotic'** recovered from the culture medium (or nutrient broth), it becomes absolutely necessary to accomplish the said product in its purest form within the permissible attainable limits of purity. In order to achieve this the 'crude product' is subjected to various advanced techniques of chromatography, followed by meticulous recrystallization procedure, and ultimately subjected to the standard manipulative operations using specific skill and wisdom.

Salient Features Some of the salient features required to cause a suitable extent of purification are:

1. The attempt to achieve a very high degree of *'chemical purity'* is neither practicable nor necessary for therapeutic purposes.
2. Foreign proteins *i.e.*, extraneous metabolites, responsible for undesirable side-effects are excluded automatically through the process of purification.

3. Complete separation/elimination of closely structurally related antibiotic substances is invariably unfeasible.
4. Antibiotics derived from various fermentative procedures most frequently employed in therapy are, in true sense, admixtures of very intimately related chemical entities having one of the metabolites predominantly present in the mixture.
5. Reproducible therapeutic response is of prime importance, which must be attained through permissible practical limits due to the fact that a given antibiotic compound always constitute a major component of the mixture.
6. It also furnishes the economic viability of antibiotic substances in various drug formulative operations by virtue of the fact that the inefficiency and total expenses involved for complete separation of closely related chemical substances having unequal relative concentrations, may be avoided to a great extent.

Example: Chlortetracycline present upto 6% in the commercial *tetracycline* fairly represents an actual realistic and practical approach of such purification considerations.

Note: The overall accepted standards of purity for antibiotics and other antibiotic formulations (i.e., dosage forms) are strictly controlled and monitored by the pharmacopocia of various countries, such as: USP; B.P.; Eur. P.; Int. P.; Ind. P.; Japanese P., etc.

(d) **Purity of Antibiotic:** The highest attainable purity of an antibiotic is an absolute necessity so as to minimise its undesirable side effects.

Example: Vancomycin is a glycopeptide antibiotic particularly effective for the treatment of **endocarditis*** caused by Gram +ve bacteria. However, its wide application and usefulness was grossly restricted due to its **nephrotoxicity.**** Interestingly, upon much improved purity status of vancomycin not only reduced nephrotoxicity significantly but also raised its position in the therapeutic armamentarium.

(e) **Antibiotic Masking of Microbial Contaminants:** The **parenteral preparations** need to be guaranteed for their stringent sterility test(s) in the presence of an antibiotic. Therefore, it has become almost necessary to assess the masking of the very presence of the microbial contaminants by means of the bacteriostatic action exerted by the prevailing antibiotic.

There are, in fact, *three* basic approaches, which are not only vital but also fundamental in nature, that may be employed so as to eliminate as far as possible the '**antibiotic masking**' of microbial contaminants, such as:

1. All antibiotic formulations (dosage forms) which are essentially inactivated promptly either by chemical or biological methods must be suitably treated before carrying out the test for sterility.

Examples:

- (a) Inactivation of the enzyme *penicillinase* by **Penicillin G**, and
- (b) Inactivation of hydroxylamine hydrochloride by **Streptomycin**.

* **Endocarditis:** Inflammation of the lining membrane of the heart. It may be due to invasion of microorganisms or an abnormal immunological reaction.

** **Nephrotoxicity:** A toxic substance that damages specifically the kidney tissues.

2. Most parenteral antibiotic preparations, particularly those having the relatively more stable ones, may be evaluated conveniently by subjecting the preparations to such a level of dilution so that the '**antibiotic level**' is definitely below the **minimum threshold concentration** for its activity, and
3. Physically removing, at the very first instance, any possible microorganisms by the help of a sterile Millipore filter in such a manipulative manner such that the organisms (undesired) are evidently separated from the antibiotic.

9.2.2.4 *Sophisticated Skillful Antibiotic Preparations*

A lot of wisdom, skill and knowledge has been rightly incorporated in accomplishing fairly stable sophisticated antibiotic preparations. There are various ways and means that have been explored meticulously in order to achieve these objectives, namely:

- (a) Shielding of relatively less stable antibiotics in gastric juice (acidic) through various chemical and physical approaches,
- (b) '**Prodrug Approach**': Usage of rather insoluble corresponding antibiotic analogues so as to get rid of objectionable taste, and thus make it more patient-friendly especially in certain vital oral formulations.

Example: Chloramphenicol Succinate/Palmitate—The bitter taste of chloramphenicol is completely masked by preparing its corresponding esters for use in suitable parenteral preparations.

- (c) **Soluble/Insoluble Derivatives**: The preparation of various soluble or insoluble derivative of antibiotics are afforded so as to make it convenient for its desired delivery at a particular site *in vivo*.

Example: Gentamycin sulphate, Neomycin sulphate, Tetracycline Hydrochloride, Penicillin G sodium etc. These salts are more readily absorbed *in vivo* and hence enhance their therapeutic efficacy.

It is pertinent to cite here certain **classical examples** highlighting the sophisticated skillful antibiotic preparations, namely:

- (i) Use of '**buffers**' in oral penicillin G formulations significantly minimise its loss of potency due to gastric juice,
- (ii) Enteric coating of erythromycin tablets with synthetic polymers, definitely protect the **macrolactone** ring present in it, till it sails through the entire distinctly acidic environment of the stomach (*i.e.*, gastric juice) and subsequently makes it pass into the long small intestinal canal where it eventually gets absorbed.

Example: The two commonly used modified versions of erythromycin are, namely:

- (a) Erythromycin ethylsuccinate, and
- (b) Erythromycin estolate (*i.e.* the lauryl sulphate salt of the propionyl ester).

These two salts are very much insoluble than the parent macrolide antibiotic; and provide *dual usefulness* in oral parenteral suspensions *viz.*, *first*, to refrain of their very bitter taste due to poor solubility; and *secondly*, to protect their safe journey till the lower end of the intestine.

- (iii) Enhancing the solubility characteristics of erythromycin for allowing it to be given intravenously could be accomplished by making its **glucoheptonate** and **lactobionate** salts.
- (iv) Benzathine penicillin G possesses insoluble property, and this contributes heavily as a stability factor for its usage in oral suspensions.

- (v) Penicillins give rise to insoluble procaine and benzathine salts that are used extensively through IM route for prolonged and sustained effects.
- (vi) **Probenecid** is invariably employed as an adjunct substance to the *penicillins*; and this affords *two* vital classical plus points: *first*, it checks the tubular excretion of penicillins; and *secondly*, to accomplish significant sustained blood levels of these antibiotics.
- (vii) **Amoxicillin and Ticarcillin** supplemented with a **β -lactamase inhibitor**, clavulanic acid, in various preparations usually offers an expanded therapeutic spectrum.

In short, the classical examples enumerated above from (i) through (vii) paints a beautiful rosy picture which further testifies the reality that a constant research in the applications of different aspects of pharmaceutical technology with a very strong bearing on the basic fundamental knowledge of medicinal chemistry shall ever open the limitless boundaries of 'wonderful drug formulations' to save the mankind of its sufferings.

9.3 CLASSIFICATION OF ANTIBIOTICS


Antibiotics are broadly classified on the basis of their inherent chemical structures as stated below:

- (i) Aminoglycosides,
- (ii) Anthracyclines,
- (iii) Cephalosporins,
- (iv) β -Lactams,
- (v) Lincosamides,
- (vi) Macrolides,
- (vii) Penicillins,
- (viii) Polypeptide antibiotics,
- (ix) Tetracyclines, and
- (x) Miscellaneous antibiotics.

All these different categories of antibiotics shall now be described individually in the sections that follows:

9.3.1 Aminoglycosides

The **aminoglycosides** each contain one or more amino sugars, for instance: **neosamine** or **glucosamine**, bridged by glycoside linkages to a basic, either amino or guanidino, six-membered carbon ring, such as: streptomine or streptidine as given below:



Aminoglycosides occupy a coveted status in the domain of antibiotics exclusively for the control, management and treatment of infections caused by Gram-negative bacilli. However, the overall treatment of most nosocomial Gram negative basillary infections with the aid of third-generation cephalosporins, carbapenems and new fluoroquinolones have made the aminoglycosides more or less as the alternative drugs unless and until resistant strains are suspected invariably amongst the immuno suppressed patients.

It is worthwhile to mention here that the major spectrum of activity of the **aminoglycosides** essentially comprise of aerobic Gram-negative bacilli and *Staphylococcus aureus*.


A few important members of the **aminoglycoside antibiotics** are, namely: **amikacin**, **gentamycin**, **paramycin**, **kanamycin**, **netilmicin**, **streptomycin** and **tobramycin** which shall be discussed separately as under:

9.3.1.1 Amikacin

Synonyms Lukadin.

Biological Source It is a semisynthetic aminoglycoside antibiotic derived from **Kanamycin A**.

Chemical Structure



Amikacin

1-N-[L (-)-4-Amino-2-hydroxybutyryl] kanamycin A; (C₂₂H₄₃N₅O₁₃).

The presence of the 4-amino-2-hydroxybutyryl moiety protects the antibiotic against the enzymic deactivation at many locations, while the activity of the parent molecule is still maintained.

Preparation **Amikacin** is obtained by acylation of the C-1 amino function of the 2-deoxystreptamine group of **kanamycin** with L-(-)-4-amino-2-hydroxybutyric acid.

Characteristic Features

1. It is obtained as white crystalline powder from a mixture of methanol-isopropanol having mp 203-204°C (sesquihydrate).
2. Its specific optical rotation $[\alpha]_D^{23} + 99^\circ$ (C = 1.0 in water).

Identification Tests

Amikacin Sulphate ($C_{22}H_{43}N_5O_{13} \cdot 2H_2SO_4$) (**Amikin; Amiklin; Biklin; Amikavet; Fabianol; Kaminax; Mikavir; Novamin; Pierami**): It is obtained as an amorphous form which gets decomposed at 220-230°C. Its specific optical rotation $[\alpha]_D^{22} + 74.75^\circ$ (water).


Uses

1. **Amikacin** is observed to be fairly stable to most of the aminoglycoside inactivating enzymes, and is, therefore, considered to be valuable for the treatment of serious infections usually caused by Gram -ve bacteria that are resistant to gentamycin or tobramycin.
2. It is mostly employed in a wide range of infections, such as: septicemia, serious infections due to burns, urinary tract, respiratory tract and various soft tissues, meningitis, peritonitis, osteomyelitis, omphalitis in neonates, and other serious surgical infections.

9.3.1.2 Gentamicin

Synonym Gentamycin.

Biological Sources It is an antibiotic complex produced by the fermentation of *Micromonospora purpurea* and *M. echinospora*; and a number of variants thereof.

Chemical Structure

Preparation **Gentamicin** is normally recovered from a fermentation broth produced when submerged cultures of two subspecies of *Micromonospora purpurea* are grown in the yeast extract-celulose medium.

Characteristic Features

1. It is a white amorphous powder having mp 102-108°C
2. It has specific optical rotation $[\alpha]_D^{25} + 146^\circ$.
3. It is found to be freely soluble in water, pyridine, DMF, in acidic media with salt formation; moderately soluble in methanol, ethanol, acetone; and almost insoluble in benzene and halogenated hydrocarbons. Characteristic features of some of its congeners are as follows:

S. No.	Gentamicin Congeners	Molecular formula	mp (°C)	$[\alpha]_D^{25}$
1	Gentamicin C ₁	C ₂₁ H ₄₃ N ₅ O ₇	94-100	+ 158
2	Gentamicin C ₂	C ₂₀ H ₄₁ N ₅ O ₇	107-124	+ 160
3	Gentamicin C _{1a}	C ₁₉ H ₃₉ N ₅ O ₇	—	—

Identification Tests

1. **Gentamicin Complex Sulphate:** [Synonyms Alcomycin; Bristagen; Cidomycin; Duragentum; Garamycin; Garasol; Genoptic; Gentacin; Gentak; Gentalline; Gentalyne; Gentibiopthal; Genticin; Gentocin; Gentogram; Gent-Ophtal; Gentrasul; Lugacin; Nichogencin; Ophtagram; Pangram; Refobacin; Septopal; Sulmycin; and U-Gencin.]

It is obtained as a white, hygroscopic powder having mp 218-237°C. It has specific optical rotation $[\alpha]_D^{25} + 102^\circ$. It is soluble in formamide and in ethylene glycol.

2. **Gentamicin Hydrochloride:** It has mp 194-209°C; and specific optical rotation $[\alpha]_D^{25} + 113^\circ$. It is found to be freely soluble in water, methanol; slightly soluble in ether; and practically insoluble in other organic solvents.

Uses

1. It is currently the most important drug of choice for the treatment of infections caused by most aerobic Gram-negative bacteria, besides several strains of *Staphylococci*.
2. It essentially exhibits a broad-spectrum antibacterial activity.
3. It is found to be specifically effective against *Pseudomonas*, because species of this genus resistant to 'other antibiotics' have proved to be an important cause of **surgical infections**. In the same vein, gentamicin, is also very effective in severe burned-skin patients *i.e.*, third-degree burns; and severe UTI* infections, both caused by *Pseudomonas*.
4. It is employed topically in the treatment of impetigo, infected bed sores, burns and nasal staphylococcal carrier state, pyodermata and also in the infections of external-eye.

Note: Because of gentamicin's systemic toxicity, its present systemic usage is restricted and limited to life-threatening infections produced by *Citrobacter*, *Klebsiella-Enterobacter-Serratia*, *Proteus* and *Pseudomonas*. To cause an effective control it is invariably combined along with either penicillin or cephalosporin.


9.3.1.3 Kanamycin

Biological Sources Kanamycin is an 'antibiotic complex' produced by *Streptomyces kanamyceticus* Okami & Umezawa from the Japanese soil.** The antibiotic complex is comprised of three distinct components, namely: **kanamycin A**—representing the *major* component, and usually designated as **kanamycin**; besides two *minor* components (congeners more precisely) usually known as **kanamycins B and C**.

* UTI = Urinary tract infections.

** Umezawa *et al.*, *J. Antibiot.* 10A, 181 (1957); US patent 2, 931, 798 (1960).

Chemical Structure



Interestingly, these *three* antibiotics essentially comprise of *two* aminosugars (*i.e.*, 6-amino-6-deoxy-D-glucose) which are linked individually to one single 2-deoxystreptamine aglycone (*i.e.*, non-sugar) residue.

Characteristic Features The characteristic features of all the *three* kanamycins and their respective salts shall be discussed as under:

(a) **Kanamycin A:** [C₁₈H₃₆N₄O₁₁]; O-3-Amino-3-deoxy- α -D-glucopyranosyl-(1 \rightarrow 6)-O-[6-amino-6-deoxy- α -D-glucopyranosyl-(1 \rightarrow 4)-2-deoxy-D-streptamine.

It is obtained as crystals from a mixture of methanol and ethanol. It has specific optical rotation $[\alpha]_D^{24} + 146^\circ$ (0.1N. H₂SO₄).

Kanamycin A Sulphate [*Synonyms:* Cantrex; Crystalomicina; Enterokanacin; Kamycin; Kamynex; Kanabristol; Kanacedin; Kanamytrex; Kanasig; Kanatrol; Kanicin; Kannasyn; Kantres; Kantrox; Klebcil; Otokalixin; Resistomycin; Ophthalmokalixan; Kantrexil; Kano; Kanesein; Kanaqua.]

It is obtained as irregular prisms that decompose over a wide range above 250°C. It is freely soluble in water; and almost insoluble in nonpolar solvents and the common alcohols.

Note: USP-requires that Kanamycin A sulphate contains not less than 75% Kanamycin A on an anhydrous basis.

(b) **Kanamycin B:** [C₁₈H₃₇N₅O₁₀]; [*Synonyms:* Bekanamycin; Aminodeoxy-kanamycin; NK-1006].

It is obtained as crystals having mp 178-182°C (dec.).

It has specific optical rotations $[\alpha]_D^{18} + 130^\circ$ (C = 0.5 in H₂O); $[\alpha]_D^{21} + 114^\circ$ (C = 0.98 in H₂O). It is found to be soluble in water, formamide; slightly soluble in chloroform, isopropanol; and practically insoluble in the common alcohols and nonpolar solvents.

Kanamycin B Sulphate [*Synonyms:* Coltericin; Kanendomycin; Kanendos.]


(c) **Kanamycin C: [C₁₈H₃₆N₄O₁₁]**: It is obtained as crystals from methanol + ethanol which get decomposed above 270°C. It has specific optical rotation $[\alpha]_D^{20} + 126^\circ$ (H₂O). It is found to be soluble in water; slightly soluble in formamide; and practically insoluble in nonpolar solvents and the common alcohols.

9.3.1.4 Neomycin

Synonyms Fradiomycin; Mycifradin; Neomin; Neolate; Neomas; Pimavekort; Vonamycin Powder V.

Biological Source It is an 'antibiotic complex' comprised of neomycins A, B and C. It is obtained from *Streptomyces fradiae**.

Chemical Structure



Neomycin is usually obtained as a mixture of neomycin B (*Framycetin*) and its epimer **neomycin C**, the latter constitutes 5-15% of the mixture. Interestingly, in contrast to the other clinically useful aminoglycosides, neomycin is observed to comprise essentially of *three* sugar residues strategically attached to *2-deoxystreptamine* as shown above. One of the three sugars present is the *D-ribose* (a common sugar).

Characteristic Features

(a) **Neomycin Complex:** It is an amorphous base. It is soluble in water, methanol and acidified ethanol; and almost insoluble in common organic solvents.

* Waksman and Lechevalier, *Science*, **109**, 305 (1949).

(b) **Neomycin A: [Synonym: Neamine]: $C_{12}H_{26}N_4O_6$:** It is obtained as crystals either from water or aqueous ethanol that get decomposed at 225-226°C. It has specific optical rotation $[\alpha]_D^{25} + 112.8^\circ$ (C = 1)

Neomycin A Hydrochloride: ($C_{12}H_{26}N_4O_6 \cdot 4HCl$): It is obtained as an amorphous powder decomposing between 250-260°C. Its specific optical rotation is $[\alpha]_D^{25} + 83^\circ$ (C = 1).

Neomycin A, N-Acetyl Derivative: [$C_{12}H_{26}N_4O_6 \cdot (CH_3CO)_4$]: It is obtained as crystals from methanol having mp 334-336°C. It has specific optical rotation $[\alpha]_D^{25} + 87^\circ$ (C = 1).

(c) **Neomycin B: [Synonyms: Antibiotique EF 185; Framycetin; Enterfram; Framygen; Soframycin; Actilin;] ($C_{23}H_{46}N_6O_{13}$):** It yields on hydrolysis *neomycin A* and *neobiosamine B*.

Neomycin B Hydrochloride: It is an amorphous white powder having specific optical rotation $[\alpha]_D^{20} + 57^\circ$ (H_2O). Its solubility in $mg\ ml^{-1}$ at $\sim 28^\circ C$: water 15.0; methanol 5.7; ethanol 0.65; isopropanol 0.05; isoamyl alcohol 0.33; cyclohexane 0.06; benzene 0.03; and is almost insoluble in acetone, ether, other organic solvents.

Neomycin B Sulphate: [Synonyms: Biosol, Bykomycin; Endomixin; Fraquinol; Myacine; Neosulf; Neomix; Neobreltin; Nivemycin, Tuttomycin;]. It is an amorphous white powder which is almost tasteless. It has specific optical rotation $[\alpha]_D^{20} + 54^\circ$ (C = 2 in H_2O). Its solubility in $mg\ ml^{-1}$ $\sim 28^\circ C$: water 6.3; methanol 0.225; ethanol 0.095; isopropanol 0.05; and almost insoluble in acetone, ether, chloroform. The aqueous solutions are quite stable between a pH 2 to 9. The highly purified preparations are very stable to alkali, but unstable to acids. On being refluxed with $Ba(OH)_2$ for 18 hours it exhibited no loss of activity. On boiling with mineral acids it gives rise to furfural (an aldehyde), and also an organic base.

(d) **Neomycin C: [$C_{23}H_{46}N_6O_{13}$]:** It yields on hydrolysis *neomycin A* (*i.e.*, neamine) and *neobiosamine C*.

Uses

1. It has good activity against Gram-positive and Gram-negative bacteria, but is very *ototoxic*. Therefore, its usages has been severely restricted to the oral treatment of intestinal infections.

Note: It is poorly absorbed from the digestive tract.


2. It also finds its enormous use in topical applications, such as: eardrops, eyedrops, and ointments.

9.3.1.5 Netilmicin

Synonyms 1-N-Ethylsisomicin; Sch-20569.

Biological Sources **Sisomicin** is known to be the dehydro analogue of **Gentamicin C_{1a}** (see section 9.3.1.2), and is produced by cultures of *Micromonospora inyoensis*. Nevertheless, the semisynthetic N-ethyl derivative, **netilmicin**, is mostly used medicinally because it has an almost identical activity to **gentamicin**, but produces significantly much less ototoxicity.

Chemical Structure



Characteristic Features It has specific optical rotation $[\alpha]_D^{26} + 164^\circ$ (C = 3 in H₂O).

Netilmicin Sulphate [(C₂₁H₄₁N₅O₇)₂.5H₂SO₄] [**Synonyms: Certomycin; Nettlein; Netilyn; Netromicine; Netromycin; Nettarein; Vectacin; Zetamicin.**]

Preparation It is a semi-synthetic derivative of sisomicin and is skillfully prepared by ethylation of the amino group in the 1-position of the 2-deoxy-streptamine ring.

Characteristic Features It is an off-white powder; pH (1 in 25 solution) ranges between 3.5-5.5; and pKa 8.1. It is found to be very soluble in water.


Uses Its antibiotic profile is very similar to that of **Gentamicin**.

9.3.1.6 Streptomycin

Synonym Streptomycin A.

Biological Sources After the qualified success and the overwhelming recognition of the therapeutic potential of penicillin an extensive and intensive search for other antibiotic substances gathered a tremendous momentum and stimulation. A major target and goal was the discovery of such antibiotics that are antagonistic to the Gram-negative microorganisms. **Streptomycin** was obtained from a strain of *Streptomyces griseus* (Krainsky) Waksman et Henrici (*Actinomycetaceae*); and produced by the soil Actinomycete.

Chemical Structure



Streptomycin has essentially *two* sugar components, namely: L-streptose and 2-deoxy-2-methylamino-L-glucose, which are linked to a non-sugar moiety streptidine evidently through *two* ether-linkages.

Characteristic Features **Streptomycin** is normally available as the trichloride, trichloride-calcium chloride double salt, phosphate or sesquisulphate, which invariably occur as powder or granules. It is more or less odourless but possesses a slightly bitter taste. It has been observed that most of its salts are hygroscopic and deliquesce on exposure to air; however, they are not affected by air or light. Nevertheless, the salts are very soluble in water; and practically insoluble in ether, ethanol and chloroform. The solutions of its salts are levorotatory.

(a) **Streptomycin Trichloride:** [C₂₁H₃₉N₇O₁₂·3HCl] [**Synonym: Streptomycin Hydrochloride**]:

It has specific optical rotation $[\alpha]_D^{25} - 84^\circ$. Its solubilities in mg ml⁻¹ at ~ 28°C are: water > 20; methanol > 20; ethanol 0.90; isopropanol 0.12; isoamyl alcohol 0.117; petroleum ether 0.02; ether 0.01; and carbon tetrachloride 0.042.

(b) **Streptomycin Trihydrochloride—Calcium Chloride Double Salt:** [(C₂₁H₃₉N₇O₁₂·3HCl)₂·CaCl₂] [**Synonym: Streptomycin hydrochloride-Calcium chloride complex**]: It is prepared from the streptomycin trihydrochloride salt. It is highly hygroscopic in nature, and gets decomposed at ~ 200°C. Its specific optical rotation $[\alpha]_D^{25} - 76^\circ$.

(c) **Streptomycin Sesquisulphate:** [(C₂₁H₃₉N₇O₁₂)₂·3H₂SO₄] [**Synonyms: Streptomycin sulphate; Agristrep; Streptobrettin; Vetstrep**]: It is a white to light gray or pale buff powder having faint amine-like odour. Its solubilities in mg ml⁻¹ at ~ 28°C are: water > 20; methanol 0.85; ethanol 0.30; isopropanol 0.01; petroleum ether 0.015; ether 0.035; and carbon tetrachloride 0.035.

Uses

1. It is a potent antibacterial, and more so as a tuberculosstatic agent. The MIC of streptomycin for *M. tuberculosis* is nearly 0.5 mcg ml⁻¹; whereas many sensitive Gram-negative bacteria have MICs in the range of 2-4 mcg ml⁻¹.
2. **Streptomycin** exerts its action in the control and management of *Yersinia pestis* (*plague*) and *Francisella tularensis* (*tularemia*); and in such typical incidences, it is invariably combined with a **sulphonamide drug**. Mostly hyphenated therapeutic approaches are practised, such as: **streptomycin-penicillin** used for **endocarditis***; and **streptomycin-tetracyclin** employed for **brucellosis****

Note: The incidence of serious auditory impairment is now recognized and established to be far greater with dihydro-streptomycin than the parent drug streptomycin.

3. **Streptomycin** exerts bacteriostatic action in low concentrations and bactericidal in high concentrations to a good number of Gram-negative and Gram-positive microorganisms.
4. It is an alternate choice drug in the treatment of **chancroid*****, rat-bite fevers (*Spirillum* and *Streptobacillus*).

Biosynthesis of Components of Streptomycin The so called ‘**components of streptomycin**’ essentially comprise of two major portions, namely: *first*—the *streptidine* moiety; and *secondly* – the **streptobiosamine**, which is, in fact a *disaccharide*, that consists of the two sugar residues *viz.*, **streptose** plus *2-deoxy-2-methylamino-L-glucose*. It has been revealed through an elaborated biosynthetic studies that all the *three* aforesaid ‘**components of streptomycin**’ are derived exclusively from D-glucose. As on date no exact scientific evidence is available which may give an ample proof about the point of attachment of the three different components in the streptomycin molecule. Further, there is scanty and paucity of an elaborated explanation or information with regard to the manner whereby the individual moieties present in an aminoglycoside antibiotic. However, based on the ground realities derived from the metabolic relationships of glucose to the different moieties could be gathered and prevailed, upon directly from the various biosynthetic origins of the ‘**components of streptomycin**’ as shown under.


Sailent Features The salient features of the biosynthesis of **components of streptomycin** are, namely:

1. **Streptose: D-Glucose** is first converted to 4-hexosulose which on being subjected to transannular rearrangement gives rise to **streptose**.
2. **Streptidine:** D-Glucose upon demethylation yields myoinositol which upon amination produces **streptamine**. The resulting **streptamine** in the presence of L-arginine ultimately affords the formation of streptidine.

* **Endocarditis:** Inflammation of the lining membrane of the heart.

** **Brucellosis:** A widespread infection febrile disease affecting mostly cattle, goats, swine, and sometimes humans. In humans, it is called brucellosis or Malta fever and is caused by several *Brucella* species.

*** **Chancroid:** A highly infectious nonsyphilitic ulcer; and is caused by *Haemophilus ducreyi*—a Gram-negative bacillus.



Biosynthetic Pathway of Components of Streptomycin


3. **2-Deoxy-2-methylamino-L-glucose:** D-Glucose through deoxidation and methylation yields 2-deoxy-2-methylamino-L-glucose.

9.3.1.7 Tobramycin

Synonyms Nebramycin Factor 6; NF 6; Gernebcin; Tobracin; Tobradistin; Tobralax; Tobramaxin; Tobrex;

Biological Sources It is a single factor antibiotic comprising about 10% of nebramycin, previously known as **tenebrimycin**, **tenemycin**; and the **aminoglycosidic antibiotic complex produced by *Streptomyces tenebrarius***.


Chemical Structure



Tobramycin essentially contains two aminosugar residues namely: nebrosamine and 3-deoxy-3-amino-D-glucose, and a 2-deoxystreptamine moiety. It has been found to be structurally related to **kanamycin B** (section 9.3.1.3); but evidently differs only in the absence of the 3-hydroxyl group present in the **kanasamine residue**.

9.3.2 Anthracyclines

The **anthracyclines** *i.e.*, the **anthracycline antibiotics** essentially contain an anthraquinone moiety fused with a non-aromatic ring:



There are quite a few '**anthracycline antibiotics**' which have been isolated, characterized and evaluated for their therapeutic activities, namely: **doxorubicin**, **epirubicin**, **aclacinomycin A**, and **idarubicin**. However, **mitoxantrone (mitozantrone)**, a synthetic structural analogue of the **anthracyclines** wherein both the non-aromatic ring and the respective aminosugar have been duly replaced by aminoalkyl side-chains.


All the above mentioned potent compounds shall be described in the sections that follows:

9.3.2.1 Doxorubicin

Synonyms 14-Hydroxydaunomycin; NSC-123127; FI-106.

Biological Sources **Doxorubicin**, an anthracycline antibiotic is obtained from the cultures of *Streptomyces peucetius* var *caesius*.

Chemical Structure



(8*S-cis*)-10 [(3-Amino-2, 3, 6-trideoxy- α -L-lyxo-hexopyranosyl) oxy]-7, 8, 9, 10-tetrahydro-6, 8, 11-trihydroxy-8-(hydroxyacetyl)-1-methoxy-5, 12-naphthacenedione; [C₂₇H₂₉NO₁₁]

Characteristic Feature **Doxorubicin** has mp 229-231°C.

Uses

1. It has one of the broadest spectra of antitumour activity displayed by antitumour drugs.
2. It is extensively employed to treat acute leukemias, lymphomas, and a large number of solid tumours.
3. It has been found to inhibit the synthesis of RNA copies of DNA by virtue of the intercalation of the planar molecule between base pairs on the DNA helix.

Note: The inherent 'sugar moiety' affords an additional strength besides playing a critical role in the sequence-recognition required for the binding.

4. **Doxorubicin** also exerts a few of its cytotoxic effects on account of the inhibition of the enzyme **topoisomerase II**, which is solely responsible for both cleaving and resealing of the double-stranded DNA during the replication process.

Doxorubicin Hydrochloride [C₂₇H₂₉NO₁₁·HCl] [*Synonyms* **Adriacin; Adriblastina; Adriamycin**]

1. It is obtained as orange-red coloured thin needles having mp 204-205°C (decomposes).
2. It has specific optical rotation $[\alpha]_D^{20} + 248^\circ$ (C = 0.1 in methanol).
3. It exhibits uv_{max} (methanol): 233, 252, 288, 479, 496, 529 nm.
4. It is found to be soluble in water, methanol, aqueous alcohols; and almost insoluble in acetone, benzene, chloroform, ethyl ether and petroleum ether.
5. The aqueous solutions show different colours at different pH ranges, *e.g.*, at acidic pH yellow-orange; at neutral pH orange-red; and at pH > 9 violet-blue.
6. The aqueous solutions are unstable at higher temperatures or at either alkaline or acidic pHs.

Note: Doxorubicin may reasonably be anticipated to be a carcinogen*.


* *Seventh Annual Report on Carcinogen* (PB 95-10978, 1994), p. 86.

9.3.2.2 Epirubicin

Synonyms 4'-Epidoxorubicin; 4'-Epiadriamycin; Pidorubicin; 4'-EpiDX; IMI-28.

Biological Source It has the same biological source as that doxorubicin. It is the structural analogue of the **anthracycline antibiotic doxorubicin**, wherein the only point of difference is in the position of the C-4 hydroxy group of the sugar moiety.

Chemical Structure



4'-Epidoxorubicin; $C_{27}H_{29}NO_{11}$

Characteristic Features Epirubicin Hydrochloride [$C_{27}H_{29}NO_{11} \cdot HCl$] [Synonyms Farmorubicin; Pharmorubicin].

1. It is obtained as red-orange crystals having mp 185°C (decomposes).
2. It has specific optical rotation $[\alpha]_D^{20} + 274^\circ$ (C = 0.01 in methanol).

Caution: Its solution should be protected from sunlight.

Uses

1. It is broadly employed as an antineoplastic agent.
2. It is proved to be particularly effective in the treatment of breast cancer, producing much lower side effects than **doxorubicin** itself.


Biosynthesis of Doxorubicin and Epirubicin The various steps involved in the **biosynthesis of doxorubicin and epirubicin** are as follows:

1. The propionyl-CoA is used not only as a 'starter moiety' but also as a chain-extender *via* the methylmalonyl-CoA. The *Actinomycetes* (e.g., **Streptomyces**) has a tendency to use propionate *via* methylation using SAM followed by incorporation of propionate by methylmalonyl-CoA.

Note: It has been observed that the incorporation of propionate by the methyl malonate extender units may undergo unusual frequent interruption, which process may be combated by the addition of further malonate extenders. The said phenomenon usually gives rise to an irregular sequence of methyl side-chains.

2. The alkanonic acid in the presence of S-adenosylmethiorine (SAM) and through aldol condensation yields **aklaviketone**.

3. The resulting **aklaviketone** undergoes reduction of $\text{—}\overset{\text{O}}{\parallel}{\text{C}}\text{—}$ at C-7 with NADPH generating the **aklavinsonone**.
4. Further aklavinsonone with NADPH in the presence of oxygen causes hydroxylation at C-11 to produce **ϵ -rhodomycinone**.
5. At this juncture thiamine diphosphate-D-glucose (TDP) yielding thiamine diphosphate-L-daunosamine is introduced. Consequently, a series of *five* sequential reactions, such as: (i) glycosylation of 7-hydroxyl moiety; (ii) hydrolysis of ester; (iii) de-carboxylation 10-carboxylic acid group; (iv) oxidation to corresponding 13-ketone; and (v) methylation of 4-hydroxyl moiety, ultimately produces **daunorubicin (daunomycin)**.
6. The resulting daunorubicin undergoes hydroxylation at C-14 in the presence of oxygenated NADPH finally yields **doxorubicin (adriamycin)**.
7. Doxorubicin undergoes epimerization at C-4' (see inset) in the following biosynthetic pathway, to produce **epirubicin**.




Biosynthetic Pathway of Doxorubicin and Epirubicin
 [Adapted from: Dewick P.M. *Medicinal Natural Products* 2nd edn, 2001,
 John Wiley & Sons Ltd., U.K.]

9.3.2.3 Aclacinomycin A

Synonyms Aclarubicin; Antibiotic MA 144 A1; NSC-208734; Jaclacin.

Biological Source It is obtained from *Streptomyces galilacus*.

Chemical Structure It is a complex glycoside of **aklavinone** ($C_{42}H_{53}NO_{15}$).



Characteristic Features

1. It is obtained as a yellow microcrystalline powder from a mixture of chloroform and hexane having mp 151-153°C (decomposes).
2. It has specific optical rotation $[\alpha]_D^{24} -11.5^\circ$ (C = 1 in methylene chloride).
3. It has uv_{max} (methanol): 229.5; 259; 289.5; 431 nm ($E_{1cm}^{1\%}$ 550, 326, 135, 161); (0.1 N HCl) 229.5, 258.5, 290, 431 nm ($E_{1cm}^{1\%}$ 571, 338, 130, 161); (0.1 N NaOH) 239, 287, 523 nm ($E_{1cm}^{1\%}$ 450, 113, 127).
4. It is found to be soluble in chloroform, ethyl acetate; insoluble in ether, n-hexane, petroleum ether.

Identification Tests

1. To an aqueous solution of **aclacinomycin A** add a few-drops of NaOH solution when an intense reddish purple colour is obtained.
2. To a few mg of it add 0.5 ml of pure concentrated HCl when it gives a distinct yellow colouration.


Uses It shows an enhanced antineoplastic activity with much less cardiotoxicity.

9.3.2.4 Idarubicin

Synonyms 4-Demethoxy daunomycin; 4-Demethoxydaunorubicin; DMDR; IMI-30; NSC-256439.

Biological Source It is an orally active semi-synthetic structural analogue of **daunorubicin** (Section 9.3.2.2).

Chemical Structure

Idarubicin (C₂₆ H₂₇ NO₉)

Characteristic Features **Idarubicin Hydrochloride:** [C₂₆H₂₇NO₉·HCl] [Synonyms: **Idamycin, Zavedos**]: It is obtained as orange crystalline powder having mp 183-185°C. It has specific optical rotation $[\alpha]_D^{20} + 205^\circ$ (C = 0.1 in methanol).

Uses


1. It is mostly used as an antineoplastic agent.
2. It may show increased activity with comparatively much lesser cardiotoxicity.

Note: The major drawback of structurally related and modified semi-synthetic doxorubicin-type antibiotic is due to their significant cardiotoxicity that invariably comes into being by virtue of the distinct inhibition of cardiac Na⁺, K⁺-ATpase.

9.3.2.5 Mitoxantrone

Synonyms Mitozantrone; DHAQ; NSC-279836.

Chemical Structure **Mitoxantrone** is purely a ‘synthetic structural analogue’ of the ‘anthracyclines’ wherein *two* important components viz., the *aminosugar* and the *non-aromatic ring* have been strategically replaced with a pair of amino-alkyl side chains *i.e.*, amino-ethyl.



Mitoxantrone

1, 4-Dihydroxy-5, 8-bis [[2-[(2-hydroxyethyl) amino] ethyl] amino]-9, 10-anthracenedione; (C₂₂H₂₈N₄O₆).

Characteristic Features

1. It is obtained as crystals from a mixture of ethanol and hexane having mp 160-162°C.
2. It has uv_{\max} (ethanol): 244, 279, 525, 620, 660 nm (log ϵ 4.64, 4.31, 3.70, 4.37, 4.38).
3. It is sparingly soluble in water; slightly soluble in methanol; and almost insoluble in acetone, acetonitrile, chloroform.

Mitoxantrone Dihydrochloride [C₂₂H₂₈N₄O₆·2HCl] [Synonyms: Novantrone; DHAD; CL-232315; NSC-301739].

1. It is obtained as a hygroscopic blue-black solid from a mixture of water and ethanol having mp 203-205°C.
2. It has uv_{\max} (water): 241, 273, 608, 658 nm (ϵ 41000, 12000, 19200, 20900).
3. It is found to be sparingly soluble in water; slightly soluble in methanol; and practically insoluble in acetone, acetonitrile, chloroform.

Uses

1. **Mitoxantrone** has a marked and pronounced reduced toxicity as compared to **doxorubicin** (Section 9.3.2.1)
2. It is found to be extremely effective and useful in the treatment of leukemias and solid tumours exclusively.

9.3.3 Cephalosporins



Brotzu*, in 1948, was pioneer in isolating a novel microorganism from the sea water meticulously sampled very close to a sewage outpours off the coast of Sardinia. Interestingly, he noticed its marked and pronounced antagonism to both Gram-positive and Gram-negative microorganisms. Almost after seven years, Abraham** at Oxford first and foremost gave the scientific world the report on the isolation of *three* 'antibiotic substances' from the culture of this specific organism, namely: **cephalosporin P**, **penicillin N**, and **cephalosporin C**. Out of these three isolated antibiotics, the *first*: **cephalosporin P** has practically accomplished little therapeutic significance; the *second*: **penicillin N** (originally termed as **cephalosporin N**) obtained as the major component and differs significantly from the common penicillin by its antibacterial activity and hydrophilic character; and the *third*: **cephalosporin C** showed low toxicity and *in vitro* activity against the penicillin-resistant *Staphylococci*.

In view of the above statement of facts, it is quite evident that there exist an apparent contrast with regard to the typical features of **cephalosporin C** and the **penicillins** (*viz.*, **benzylpenicillin**) besides other possible structural modifications as given below:

* Brotzu, G., *Lav. Ist. Igiene Caligari*, (1948)

** Abraham, Newton, *Nature*, **175**, 548, (1955).

Points of Contrast Between Cephalosporin C and Penicillins

S. No.	Typical Features & Structural Modifications	Cephalosporin C	Penicillins (Benzyl Penicillin)
1	Structure		
2	Stability	Stable under acidic conditions	Unstable under acidic conditions
3	Effect of Enzyme Penicillinase (β -Lactamase)	Not attacked	Attacked
4	Antibacterial Profile	Low Normal	
5	Absorption after oral administration	Poor	Normal
6	Hydrolysis	Yield 7-ACA (7-Aminocephalosporanic acid) having a fused dihydrothiazine β -lactam ring.	Yields 6-APA (6-Aminopenicillanic acid) having a fused thiazolidine β -lactam ring.
7	Removal of side-chain at C-7 by enzymes or suitable microorganisms	Afforded fruitful and elusiveresults	—
8	Removal of ester side-chain at C-3	Possible either through (i) Enzymatic hydrolysis by fermentation with a yeast, or (ii) Displacement of acetoxy moiety by nucleophilic reagents.	Possible either through (i) Enzymic removal of side chain, or (ii) Chemical ring expansion
9	Scope for side-chain modifications at C-7.	Ample scope exist	Restricted scope only

In the broader perspective the semi-synthetic cephalosporins may be classified into *three* different manners, namely: (a) chemical structure; (b) β -lactamase resistance; and (c) antibacterial spectrum. However, in usual widely accepted prevailing practice the cephalosporins are logically and legitimately classified by a more arbitrary system, dividing them into '*generations*', such as: **First generation; Second generation; and Third generation cephalosporins.**

It is pertinent to mention *two* important points with regard to the '**cephalosporin antibiotics**', namely:

- All cephalosporins commence with the prefix *ceph-* or *cef-*; however, the latter spelling now being preferred over the former, though both spellings are usually encountered in certain branded drugs; and
- The basis for the classification into the said *three* generations depends primarily and solely upon the antibacterial spectrum shown by the drugs, besides the year they were first introduced.

Note: 1. There are several instances in which the drugs belonging to the 'second generation' may have been introduced after the 'third generation' of drugs had been accomplished.

- Categorically, there is no prevalent practice or demarkation to suggest that the drugs belonging to the '**third generation**' automatically supercede second and first genera-

tion ones. In fact, cephalosporins from all the three aforesaid categories are still being used across the globe.

Prodrugs (e.g., cefuroxime-axetil; cefpodoxime-proxetil) The prodrugs of some cephalosporin antibiotics, for instance: cefuroxime-axetile and **cefpodoxime-proxetil** have been duly developed having an additional ester moiety attached to the C-4 carboxyl function. However, these tailor-made 'prodrugs' get duly hydrolyzed to their respective active agents by the aid of esterases.


Cephameycins These represent another group of **cephalosporin antibiotics** that are characterized by a 7α -methoxy function, and are usually produced by two cosecutive reactions, namely: **hydroxylation and methylation**.

Example

Cephameycin C: In this particular instance, the introduction of a carbamate function derived from carbamoyl phosphate on the hydroxymethyl function.

A few important and typical examples of the '**Cephalosporin Antibiotics**' belonging to the various recognized groups, such as: first generation, second generation, third generation, prodrugs, and cephamycins have been duly summarized below along with their structural variants, names, synonyms and special remarks:













Cephalosporin Antibiotics: Typical Examples



Class	Sl. No.	R ₁	R ₂	Name (Synonyms)	Special Remarks
First Generation	1		{-CH ₃ }	Cefalexin [Cephalexin]	Orally Active
	2		{-CH ₃ }	Cefradine [Cephadrine]	Orally Active, superseded generally
	3		{-CH ₃ }	Cefadroxil	Orally Active
Second Generation	4		{-Cl}	Cefaclor	Orally Active
	5		{-CH=CH-CH ₃ }	Cefprozil	Orally Active,

(Contd.)

(Table contd.)

Third Generation	6			Cefamandole [Cepha- mandole]	High Resis- tance to β -lactamases
	7			Ceftazidime	Broad- Spectrum Gram- Negative Activity; good Activity against Pseudo- monas
	8			Ceftriaxone	Broad- Spectrum Gram- Negative Activity; Longer Half-Life Than Other Cepha- losporins
Prodrugs	9		$\{-\text{CH}=\text{CH}_2$	Cefixime	II/III Generation, Orally Active, Long Duration of Action
	10			Cefuroxime -Axetil	II Generation, Orally Active; Hydrolysed by Esterases to Liberate Cefuroxime
	11			Cefpodoxime -Proxetil	II/III Generation, Orally Active; Hydrolysed by Esterases to Liberate Cefpodoxime
Cepha mycins	12		$\{-\text{CH}_2-\text{O}-\text{C}(=\text{O})-\text{NH}_2$	Cefoxitin	Stable to β -Lacta- mases and Mammalian Esterases

A good number of **cephalosporins** belonging to the three categorized generations are available in the therapeutic armamentarium, besides the cephamycins, which are given as under:

- (i) **First generation Cephalosporins:** Cefalotin (Cephalothin); D-Cephalexin (D-Cefalexin); Cephapirin; Cefazolin; D-Cephadrine (D-Cefradine); D-Cefadroxil;
- (ii) **Second Generation Cephalosporins:** D-Cefactor; D-Cefamandole; Cefuroxime; D-Cefonicid; Ceforanide;
- (iii) **Third Generation Cephalosporins:** Cefotaxime; Ceftizoxime; D-Cefoperazone; Ceftazidime; Ceftriaxone; Cefmonoxime, Moxalactam;
- (iv) **Prodrugs:** Cefpodoxime proxetil; Cefuroxime axetil;
- (v) **Cephamycins:** Cephamycin C; Cefoxitin.

A few of these important compounds representing the above said categories shall now be discussed individually in the sections that follows:

9.3.3.1 First Generation Cephalosporins


The **first generation cephalosporin** antibiotics are found to be effective against a host of Gram-positive microorganisms, including penicillinase-producing *Staphylococcus*. Besides, being resistant to **penicillinase** they are found to be inactivated by another **cephalosporinase** termed as **β -lactamase**. The Gram-negative organisms that are observed to be highly sensitive to these compounds are, namely: *Escherichia coli*; *Proteus mirabilis*; and *Klebsiella pneumoniae*. These antibiotics are also found to be less active against *Haemophilus influenzae* as compared to the extended-spectrum penicillins, such as: **ampicillin**.

A. Cephalothin

Synonyms Cefalotin; 7-(Thiophene-2-acetamido) cephalosporanic acid).

Biological Source It is a **semi-synthetic cephalosporin antibiotic** derived from *Cephalosporium acremonium*.

Chemical Structure



Cephalothin

(6R-*trans*)-3-[(Acetyloxy) methyl]-8-oxo-7-[(2-thienylacetyl) amino]-5-thia-1-azobicyclo [4.2.0] act-2-ene-2-carboxylic acid; (C₁₆H₁₆N₂O₆S₂).

Preparation First of all the 7-aminocephalosporanic acid is N-acetylated with 2-thiopheneacetyl chloride in a dehydro-chlorinating environment. The starting acid may be prepared from the natural antibiotic, **cephalosporin C**, either by means of enzymatic hydrolysis or by proton-catalyzed hydrolysis. The **cephalothin** thus obtained may be purified from acetonitrile.

Characteristic Features

1. It is obtained as a white amorphous powder having mp 160-160.5°C.
2. It has specific optical rotation $[\alpha]_D^{20} + 50^\circ$ (C = 1.03 in acetonitrile).

Cephalothin Sodium $[C_{16}H_{15}N_2NaO_6S_2]$ [*Synonyms* Averon-1; Cefalotin; Cemastin; Cephation; Ceporacin; Cepovenin; Coaxin; Keflin; Lospoven; Microtin; Synclothin; Toricelocin].

1. It is obtained as a white to off-white, crystalline powder, almost odourless, moderately hygroscopic and has mp 204-205°C.
2. It has dissociation constant pKa 2.2.
3. It has specific optical rotation $[\alpha]_D + 135^\circ$ (C = 1.0 in water).
4. It has uv_{max} : 236, 260 nm (ϵ 12950, 9350).
5. **Solubility Profile:** It is freely soluble in water, normal saline or dextrose solution; slightly soluble in ethanol; and practically insoluble in most organic solvents.


Uses

1. It is a potent antibacterial agent.
2. It is a **first-generation cephalosporin** given IM and IV.
3. It is found to be a short-acting antibiotic and exhibits the weakest spectrum of its class.

B. Cephazolin

Synonym CEZ.

Biological Source It is also a semi-synthetic antibiotic derived from **7-aminocephalosporanic acid** obtained from *Cephalosporium acremonium*.

Chemical Structure

Cephazolin

7-(1-(1H)-Tetrazolyl acetamido)-3-[2-(5-methyl-1, 3, 4-thiadiazolyl) thiomethyl]- Δ^3 -cephem-4-carboxylic acid; ($C_{14}H_{14}N_8O_4S_3$).

Preparation The sodium salt of **7-aminocephalosporanic acid** is acylated with 1H-tetrazole-1-acetyl chloride. The acetoxy moiety present in the resulting product is then displaced by reaction with 5-methyl-1, 3- 4-thiadiazole-2-thiol to produce the desired product *i.e.*, **cephazolin**. It is then further purified from aqueous ethanol.

Characteristic Features

1. Cephazolin is obtained as needles from aqueous acetone having mp 198-200°C (decomposes).
2. It has uv_{max} (buffer pH 6.4): 272 nm (ϵ 13150).

3. It is found to be freely soluble in DMF, pyridine; soluble in aqueous acetone, aqueous dioxane, aqueous ethanol, slightly soluble in methanol; and practically insoluble in benzene, chloroform, ether.

Cephazolin Sodium [$C_{14}H_{13}N_8NaO_4S_3$] **Synonyms Acef; Ancef; Atirin; Biazolina; Bor-Cefazol; Cetacidal; Cefamedin; Cefamezin; Cefazil; Cefazina; Elzogram; Firmacef; Gramaxin; Kefzol; Lampocef; Liviclina; Totacef; Zolicef]:**

1. It is obtained as white to yellowish white, odourless crystalline powder having a bitter salty taste. It crystallizes out in α -, β -, and γ -forms.
2. It is found to be freely soluble in water; slightly soluble in methanol, ethanol; and almost insoluble in benzene, acetone, chloroform.

Uses


1. **Totacef** may be given IV or IM; however, its Gram-negative activity is essentially limited to *E. coli*; *Klebsiella*; and *Pr mirabilis*.
2. Some Gram-negative organisms and penicillinase-producing staphylococci which are resistant to both **Penicillin G** and **Ampicillin** are found to be sensitive to cefazolin.
3. It may be used to treat infections of the respiratory tract skin, soft tissues, tones, joints and urinary tract and *endocarditis* and septicemia caused by susceptible organisms. However, amongst the UTIs, **cystitis*** specifically responds much better than **pyelonephritis****.
4. It is one of the preferred cephalosporins for most surgical prophylaxis, by virtue of its inherent long half-life (*i.e.*, 1.5 to 2 hours in normal persons but 3 to 42 hours in renal failure).

C. Cefadroxil

Synonyms BL-S578; MJF-11567-3; Baxan; Bidocef; Cefa-drops; Cefamox; Ceforal; Cephos; Duracef; Duricef; Kefroxil; Oracefal; Sedral; Ultracef.

Biological Source It is also an orally active **semi-synthetic cephalosporin** antibiotic obtained from the species *Cephalosporium aeremonium*.

Chemical Structure



para-Hydroxycephalexine monohydrate; ($C_{16}H_{17}N_3O_5S \cdot H_2O$).

Characteristic Features It is obtained as white to yellow white crystals having mp 197°C (decomposes). It is found to be soluble in water, and fairly stable in acidic medium.

* **Cystitis**: Inflammation of the bladder usually occurring secondary to ascending urinary tract infections (UTIs).

** **Pyelonephritis**: Inflammation of kidney and renal pelvis.

Uses

1. It is intermediate acting and quite effective against *Staphylococcus* and certain enteric Gram-negative bacilli.
2. Because of its prolonged excretion criterion, it has an added advantage of catering for more sustained serum and urine concentrations than are usually obtained with other **oral cephalosporins**.
3. Clinical studies have revealed that **cefadroxil** administered, 1g twice daily, is as effective as cephalixin given 500 mg four times daily.


9.3.3.2 Second Generation Cephalosporins

Generally, the 'second generation cephalosporins' exhibit the same spectrum of antibacterial activity as that of the first generation cephalosporins. The glaring exceptions being that these are comparatively much more active against certain specific organisms, namely: *Haemophilus influenzae*, gonococcus, and some enteric Gram-negative bacilli. Interestingly, most of the second generation cephalosporins are adequately absorbed through the oral administration. Some typical examples of **second generation cephalosporins** shall now be described as under:

A. Cefamandole

Synonyms CMT; Compound 83405.

Biological Sources It is a broad-spectrum semi-synthetic cephalosporin antibiotic obtained from *Cephalosporium acremonium*.

Chemical Structure

Cefamandole

7-D-Mandelamido-3-[[[(1-methyl-1H-tetrazol-5yl) thio]-methyl]-3-cephem-4-carboxylic acid; (C₁₈H₁₈N₆O₅S₂).

Characteristic Features

Cefamandole Nafate [C₁₉H₁₇N₆NaO₆S₂] [**Synonyms Bergacef; Cedol, Cefam; Cefiran; Cemado; Cemandil; Fado; Kefadol; Kefandol; Lampomandol; Mandokef; Mandol; Mandolsan; Neocefal; Pavecef**]:


1. It is obtained or white odourless needles having mp 190°C (decomposes).
2. It has uv_{max} (H₂O): 269 nm (ϵ 10800).
3. Its dissociation constant pKa 2.6-3.0.
4. It is soluble in water, methanol; and almost insoluble in ether, chloroform, benzene, cyclohexane. It is also found to be soluble in saline TS or dextrose solutions.

Uses

1. It is administered effectively through IM and IV.
2. It is found to be short-acting.

B. Cefuroxime

Several antibiotics belonging to the category of ‘**second generation antibiotics**’ are absorbed orally. Interestingly, cefuroxime is available in two different versions; *first*—as its sodium salt and *secondly*—as its *prodrug* cefuroxime axetil that are hydrolyzed once they are absorbed and its absorption rate increased by the intake of food in-take.

Chemical Structure

Cefuroxime

(6R, 7R)-3-Carbamoyloxymethyl-7-[2-(2-furyl)-2-(methoxy-amino) acetamido] ceph-3-em-4-carboxylic acid; (C₁₆H₁₆N₄O₈S).

Characteristic Features

1. It is obtained as a white crystalline solid.
2. It has specific optical rotation $[\alpha]_D^{20} + 63.7^\circ$ (C = 1.0 in 0.2 M pH 7 phosphate buffer).
3. It has uv_{max} (pH6 phosphate buffer): 274 nm (ϵ 17600).

Cefuroxime Sodium [C₁₆H₁₅N₄NaO₈S]: [Synonyms Anaptivan; Biociclin; Biofurex; Bioxima; Cefamar; Ceffoprim; Cefumax; Cefurex; Cefurin; Curocef; Curoxim; Duxima; Gibicef; Ipaccef; Kefurox; Kesint; Lampsporin; Medoxim; Novocef; Spectrazole; Ultroxim; Zinacef].

It is a white solid; specific optical rotation $[\alpha]_D^{20} + 60^\circ$ (C = 0.91 in water); uv_{max} (water): 274 nm (ϵ 17400). It is found to be freely soluble in water and buffered solutions; soluble in methanol; very slightly soluble in ether, ethyl acetate, octanol, benzene and chloroform. Its solubility in water is 500 mg/2.5 ml. Its dissociation constant in water pKa 2.5; in DMF 5.1. The stability of cefuroxime sodium salt in water at room temperature stands valid upto 13 hours; and at 25°C for 48 hours nearly 10% decomposition takes place.

Uses


1. Its activity against *H. influenzae* and ability to penetrate into the CSF makes it specifically useful for the treatment, control and management of meningitis caused by this organism. However, it is also recommended to treat meningitis caused by *Strep. pneumoniae*, *N. meningitidis* and *Staph. aureus*.
2. It exhibits an excellent and super activity against all species of gonococci, hence it is recommended for the treatment of gonorrhoea.

3. It may also be employed to treat lower respiratory tract infections normally caused by *H. influenzae* and *parainfluenzae*, *Klebsiella* species, *E. coli*, *Strep pneumoniae* and *pyrogenes* and *Staph aureus*.
4. It is also approved for use against UTIs caused by *E. Coli* and *Klebsiella*; of course, a rather more restricted approval as compared to other second-generation drugs.
5. It is also recommended for use in bone infections, septicemias and surgical prophylaxis.

C. Cefonicid

Biological Source It is essentially an injectable **semi-synthetic cephalosporin** antibiotic related to **cefamandole** which is obtained from *Cephalosporium acremonium*.

Chemical Structure



Characteristic Features

Cefonicid Disodium [$C_{18}H_{16}N_6Na_2O_8S_3$] [*Synonyms* SKF-75073; Cefodie; Monocid; Monocidur; Praticef;] The pH of 5% (w/v) solution is between 3.5 to 6.5.

Uses

1. It is administered through IV and IM.
2. It is an **intermediate-acting second generation cephalosporin**.

9.3.3.3 Third Generation Cephalosporins


The '**third generation cephalosporins**' are logically differentiated from the first and second generation cephalosporins by virtue of their extended activity against a wide spectrum of **enteric Gram-negative bacilli**, along with the **β -lactamase-producing strains**.

A few potent and typical examples of drugs belonging to this category shall be discussed in the section that follows:

A. Cefotaxime

Biological Source It is a broad spectrum **third generation cephalosporin** antibiotic derived from *Cephalosporium acremonium*. The name cefotaxime applies to the isomer having a *syn*-methoxyamino moiety.

Chemical Structure It is the desacetyl active metabolite of **cefotaxime** *i.e.*, another **third generation cephalosporin**.



Cefotaxime

7-[2-(2-Amino-4-thiazolyl)-2-methoxyimino acetamido] cephalosporanic acid; ($C_{16}H_{17}N_5O_7S_2$).

Characteristic Features

Cefotaxime Sodium [*syn-Isomer*]: [$C_{16}H_{16}N_5NaO_7S_2$] [**Synonyms**: *Cefotax*; *Chemcef*; *Claforan*; *Pretor*; *Tolycar*; *HR-756*; *RU-24756*]:

1. It is a white to off-white solid.
2. The pH of a 10% (w/v) solution is approximately 5.5.
3. It has specific optical rotation $[\alpha]_D^{20} + 55^\circ \pm 2$ ($C = 0.8$ in water).
4. Its pKa (acid) is 3.75.
5. It is found to be freely soluble in water; and almost insoluble in most organic solvents.

Uses


1. It is found to be active against a good number of Gram-negative bacilli and its action is almost equivalent to the **amino glycosides**, except against *Ps aeruginosa*, *Acinetobacter* and a few *Enterobacter*.
2. It is highly resistant to the β -lactamases.
3. It is found to be less active than either the *first* or the **second generation cephalosporins**.
4. It is a recognized and preferred **third generation cephalosporin** for Gram-negative meningitis and other serious Gram-negative bacillary infections outside the CNS.
5. It is also recommended widely for surgical prophylaxis.
6. **Cefotaxime** has a slightly longer half-life which permits 8 to 12 hour dosing in comparison to 6 to 8 hours for **cefotaxime**.

B. Ceftriaxone

Synonym Cefatriaxone.

Biological Source It is a parenteral **third generation cephalosporin antibiotic** obtained from *Cephalosporium acremonium*.

Chemical Structure [$C_{18}H_{18}N_8O_7S_3$]



Ceftriaxone

Characteristic Features

Ceftriaxone Disodium Hemiheptahydrate [$C_{18}H_{18}N_8Na_2O_7S_3 \cdot 3\frac{1}{2} H_2O$]: [**Synonyms** *Rocefin*; *Rocephin(e)*]:

1. It is obtained as white crystalline powder having mp > 155°C. (decomposes).
2. It has specific optical rotation $[\alpha]_D^{25} -165^\circ$ (C = 1 in water) (calculated for anhydrous substance).
3. It has uv_{max} (water): 242, 272 nm (ϵ 32, 300; 29530).
4. It shows dissociation constant pKa : ~ 3 (COOH); 3.2 (NH₃⁺), 4.1 (enolic OH).
5. Its solubility in water at 25°C: ~ 40 g/100 ml.
6. The colour of its solution varies from light yellow to amber depending solely on the concentration (g. L⁻¹) and the duration of storage (hrs.)
7. The pH of a 1% (w/v) solution is nearly 6.7.
8. It is found to be sparingly soluble in methanol; and very slightly soluble in ethanol.

Uses


1. **Ceftriaxone sodium** is considered as the drug of choice for uncomplicated and disseminated gonococcal infections.
2. It is also an effective alternative for meningitis in infants essentially caused by *H. influenzae*, *N. meningitidis* and *Streptococcus pneumoniae*.
3. It is also recommended for Gram-negative bacillary meningitis and some other serious Gram-negative infections, including complications associated with **Lyme disease**.*
4. It may also be used for the treatment of bone and joint infections, intra-abdominal infections, lower respiratory tract infections, pelvic infections, skin and urinary tract infections (UTIs).
5. It is also indicated for preoperative prophylaxis, for which its efficiency is almost equivalent to that of **cefazolin**.

C. Ceftazidime

Synonym GR-20263; Fortaz; Tazicef; Tazidime.

Biological Source It belongs to the class of third generation cephalosporin antibiotic.

Chemical Structure The O-substituted oxime function certainly improves upon the potency of the ‘**third generation cephalosporin**’, and ultimately exerts resistance to **β-lactamases**. It is, however, pertinent to mention here that the oximes with *syn* stereochemistry are appreciably more potent and efficient than the corresponding *anti* isomers.



Ceftazidime

* **Lyme Disease:** A multisystem disease caused by the tick-transmitted spirochete *Borrelia burgdorferi*.

1-[[[(6R, 7R)-7-[2-(2-(2-amino-4-thiazolyl) glyoxylamido)-2-carboxy-8-oxo-5-thia-1-azabicyclo-[4.2.0] oct-2-en-3-yl] methyl] pyridinium hydroxide inner salt 7²-(Z)-[0-(1-carboxy-1-methylethyl) oxime]; (C₂₂H₂₂N₆O₇S₂).

Characteristic Features

1. It is obtained as an ivory-coloured powder.
2. Its pKa values are : 1.8, 2.7, 4.1.
3. It has uv_{max} (pH6) : 257 nm ($E_{1cm}^{1\%}$ 348).

Uses


1. **Ceftazidime** is a broad-spectrum antibiotic and is administered IV or IM.
2. It is specifically of great interest because of its distinct high activity against *Pseudomonas* and *Enterobacteriaceae*, but not enterococci.
3. It is an alternative drug for the treatment of hospital-acquired Gram-negative infections especially in immuno-compromised patients when *Ps aeruginosa* is a potential causative organism.
4. It is also recommended for use in the treatment, control, and management of bone and joint infections, CNS-infections, gynecological infections, lower respiratory tract infections, septicemia, skin and UTIs.
5. Its activity is fairly comparable to that of **cefotaxime** and **coftizoxime** *in vitro* but is much more active against *Pseudomonas aeruginosa* and fairly less active against staphylococci and *Bacteroides fragilis*.
6. It is an agent of choice for the ‘**empirical antibiotic therapy**’ when *pseudomonas* happens to be one of the suspected pathogens.

D. Moxalactam

Synonyms Lamoxactam; Latamoxef.

Biological Source It is an oxa-substituted **third generation cephalosporin** antibiotic (oxacephalosporin).

Chemical Structure [C₂₀H₂₀N₆O₉S]



Moxalactam

Characteristic Features

1. **Moxalactam** is obtained as a colourless powder having mp 117-122°C (decomposes).
2. Its specific optical rotation $[\alpha]_D^{25} -15.3 \pm 2.6^\circ$ (C = 0.216 in methanol).
3. It has uv_{max} (methanol): 276 nm (ϵ 10200).

Moxalactam Disodium [C₂₀H₁₈N₆Na₂O₉S] [**Synonyms** Festamoxin; Moxalactam; Moxam; Shiomarin; LY-12735; S-6059].

1. It has specific optical rotation $[\alpha]_D^{22} -45^\circ$ (water).
2. It has uv_{\max} (water): 270 nm (ϵ 12000).

Uses

1. **Moxalactam** exhibits a spectrum of activity which is almost identical to that of cefotaxime. However, its usage is very much restricted and limited on account of the occurrence of bleeding disorders of serious nature.

Note: The presence of methyltetrazolethiomethyl moiety may be responsible for causing hypoprothrombinemia; and the α -carboxyl group may attribute to the platelet dysfunction. Perhaps both these vital factors ultimately lead to bleeding disorders.

2. It has an extended Gram-negative spectrum, and are found to be most active against enteric Gram-negative bacilli, but may be less active against certain Gram-positive microorganisms, especially *Staphylococcus aureus*.

9.3.3.4 Prodrugs


A major noticeable chief disadvantage of plethora of the current **cephalosporins** is that they are not rapidly and effectively absorbed through oral route. This specific serious drawback is perhaps due to the nature of the side-chain present at C-3. An attempt has been made to design orally active prodrugs, namely: **cefuroxime-axetil** and **cefpodoxime-proxetil**, which have been developed meticulously by providing an additional ester function on the C-4 carboxyl moiety. Nevertheless, these prodrugs are designed in such a manner that they are easily hydrolysed to the active agents *i.e.*, drugs by the *esterases*.

A. Cefpodoxime Proxetil

Synonyms Banan; Cefodox; Orelox; Otreon; Vantin; CS-807; U-76252;

Biological Source It is a broad spectrum, orally absorbed third generation cephalosporin, tailor-made ester prodrug of the active-free-acid metabolite, **cefpodoxime**.

Chemical Structure The skill and wisdom of a pharmaceutical chemist has made it possible to design an ester of the metabolite, **cefpodoxime**, in which the free carboxyl function located at C-4 of the **thiazine ring** *i.e.*, the heterocyclic six-membered ring with one each of S and N atom studded at alternate position in the ring.



Cefpodoxime Proxetil

1-(Isopropoxy carbonyloxy) ethyl (6R, 7R)-7 [2-(2-amino-4-thiazolyl)-(Z)-2-(methoxyamino)acetamido]-3-methoxymethyl-3-cephem-4-carboxylate; (C₂₁H₂₇N₅O₉S₂).

Uses

1. It is a '**third generation cephalosporin**'-prodrug administered orally.
2. It is found to be an '**intermediate acting cephalosporin**'.
3. Its pharmacologic activity is almost similar to that of **cofixime**.


9.3.3.5 Cephameycins

Streptomyces clavuligerus gave rise to the isolation of the natural antibiotic known as *Cephameycin C*, which essentially has an α -methoxy function at C-7 present in the **basic cephalosporin ring system**. It has been observed that the **steric hindrance** caused due to the presence of the additional methoxy moieties, affording thereby a possible resistance to **β -lactamase** hydrolysis, might be solely responsible for the weak antibacterial activity of both **cephameycin C** and other natural **cephameycins**. Various semi-synthetic structural analogues have been designed either:

- (a) By chemical introduction of the α -methyl moiety at C-7 of the **basic cephalosporin ring system**,
or
- (b) By modification of the side-chains of the naturally occurring **cephameycins**.

A. Cefoxitin

Biological Source It is a semi-synthetic derived from the **Cephameycin C** through the method 'a' stated above. The synthesis of **Cefoxitin** has been successfully carried out by Karady *et al.**

Chemical Structure

Cefoxitin

3-Carbamoyloxymethyl-7 α -methoxy-7-[2-(2-thienyl)-acetamido]-3-cephem-4-carboxylic acid; (C₁₆H₁₇N₃O₇S₂).

Characteristic Features

1. It is obtained as crystals having mp 148-150°C (decomposes).
2. Its dissociation constant pKa is 2.2.
3. **Solubility Profile:** It is very soluble in acetone; soluble in aqueous NaHCO₃; very slightly soluble in water; and almost insoluble in ether and chloroform.

Cefoxitin Sodium [C₁₆H₁₆N₃NaO₇S₂] [**Synonyms** Betacef; Farmoxin; Mefoxin; Mefoxitin; Merxin; Cenomycin;]:

1. It is obtained as white crystals with a characteristic odour having mp 150°C.
2. It has specific optical rotation $[\alpha]_{589\text{nm}}^{25} + 210^\circ$ (C = 1 in methanol).

* Karady *et al.* *J. Am. Chem. Soc.*, **94**, 1410, (1972).

- It has dissociation constant pKa 2.2 (acid).
- Solubility Profile:** It is found to be very soluble in water; soluble in methanol; sparingly soluble in ethanol or acetone; and practically insoluble in aromatic and aliphatic hydrocarbons.






Uses

- It is mostly used as an alternative drug for intra-abdominal infections, colorectal surgery or appendectomy and ruptured viscus by virtue of the fact that it is active against most enteric anaerobes including *Bacteroides fragilis*.
- It is also recommended for use in the treatment of bone and joint infections usually caused by *S. aureus*, gynecological and intra abdominal infections by *Bacteriodes* species
- It is also approved for lower respiratory tract infections caused by *Bacteriodes* species; *E. coli*; *H. influenzae*; *Klebsiella* species; *S. aureus* etc.

9.3.4 β -Lactams







The **β -lactam antibiotics** (or **β -lactams**) essentially comprise of the **penicillins, cephalosporins, imipenem, nocardicin A, aztreonam, clavulanic acid, moxalactam, and thienamycin**. Interestingly, the β -lactam heterocyclic nucleus consists of a 4-membered cyclic ring with a N-atom. There exist a number of structural variants of β -lactam ring whereby the highly-strained β -lactam nucleus is strategically stabilized by means of the fusion of a variety of either 5-membered or 6-membered heterocyclic moieties to give rise to a wide spectrum of newer antibiotics as enumerated below.

β -Lactam Variants

S. No.	Nomenclature	Chemical Structure (Nucleus)	Notes
1	β-Lactam		Fused skeletons found in penicillins, cephalosporins, imipenem, and aztreonam.
2	Penam		β -Lactam ring fused with a 5-membered dihydrothiazine ring e.g., penicillins (discussed under section 9.3.7).
3	Cephem		β -Lactam ring fused with a 6-membered dihydrothiazine ring e.g., cephalosporins (discussed under section 9.3.3).
4	Penem		β -Lactam ring fused with a 5-membered thiazolidine ring having a double bond between C-2 and C-3.
5	Oxapenam		S-Heteroatom in penam nuclei replaced by an O-atom e.g.,  Clavulanic Acid

(Contd.)

(Table contd.)


6	Oxacepham		S-Heteroatom in <i>cepham</i> nuclei replaced by an O-atom eg., moxalactam (section 9.3.3.3.D)
7	Carbopenem		Variant of the penicillin-ring-system wherein the S-heteroatom is replaced by carbon eg.,
8	Monobactam		Monocyclic β -lactams e.g.,
			Thienamycin
			Aztreonam
			Nocardicin A

A few important compounds belonging to the category of so called '**other β -lactams**', such as: **thienamycin**, **aztreonam**, **nocardicin A**, **imipenem** and **meropenem** shall be treated separately as under:

9.3.4.1 Thienamycin

Biological Source It belongs to the first member of a family of **des-thia-carbapenam nucleus antibiotics** with a thioethylamine side-chain strategically positioned on the enamine portion of the fused 5-membered ring. It is obtained from cultures of *Streptomyces cattleya*.

Chemical Structure



Thienamycin

[5R-[5 α , 6 α (R*)]]-3 [(2-Aminoethyl) thio]-6-(1-hydroxyethyl)-7-oxo-1-azabicyclo [3, 2, 0]-hept-2-ene-2-carboxylic acid; (C₁₁H₁₆N₂O₄S).

Characteristic Features


1. It is obtained as a white hygroscopic solid powder.
2. Its specific optical rotation $[\alpha]_D^{27} + 82.7^\circ$ (C = 1.0 in water).
3. It has uv_{max} (water pH 4.8): 296.5 nm (ϵ 7900); (pH 2): 309 nm; and (pH 12): 300.5 nm.
4. It is found to be freely soluble in water; and sparingly soluble in methanol.
5. In dilute solution its stability is observed to be optimal between pH 6.7, declining with unusual rapidity above that range.
6. It is found to be susceptible to inactivation by dilute solutions of hydroxylamine and cysteine.

9.3.4.2 Aztreonam

Synonyms Azthreonam; Azactam; Azonam; Aztreon; Nebactam; Primbactam; SQ-26776.

Biological Sources Aztreonam enjoys the reputation of being the first totally synthetic monocyclic β -lactam (monobactam) antibiotic.

Chemical Structure



Aztreonam

[2S-[2 α , 3 β (Z)]]-2-[[[1-(2-Amino-4-thiazolyl)-2-2-methyl-4-oxo-1-sulfo-3-azetidiny] amino]-2-oxoethylidene] amino]oxy]-2-methylpropanoic acid; (C₁₃H₁₇N₅O₈S₂).

Characteristic Features

1. It is obtained as white crystalline odourless powder which decomposes at 227°C.
2. **Solubility Profile:** It is found to be very slightly soluble in ethanol; slightly soluble in methanol; soluble in DMF, DMSO; and almost insoluble in toluene, chloroform, ethyl acetate.


Uses It offers a significantly high degree of resistance to β -lactamases and displays specific activity *Vs* aerobic Gram-negative rods.

9.3.4.3 Imipenem

Synonyms Imipemide; N-fomimidoylthienamycin monohydrate; MK-787.

Biological Source **Imipenem** is an extremely broad-spectrum semi-synthetic antibiotic produced by *S. cuttleya*. It is being recognized as the first stable derivative of thienamycin.

Chemical Structure



Imipenem

[5R-[5 α , 6 α (R*)]]-6-(1-Hydroxyethyl)-3[[2-[(iminomethyl) amino] ethyl] thio]-7-oxo-1-azabicyclo-[3, 2, 0] hept-2-ene-2-carboxylic acid monohydrate; (C₁₂H₁₇N₃O₄S·H₂O).

Preparation It is obtained as a crystalline derivative of **thienamycin** by the method suggested by Leanza *et al.**

Characteristic Features

1. It is obtained as crystals from a mixture of water and ethanol.
2. It has specific optical rotation $[\alpha]_D^{25} + 86.8^\circ$ (C = 0.05 in 0.1 M phosphate pH 7).
3. It has dissociation constants pK_{a1} ~ 3.2, pK_{a2} ~ 9.9.
4. It shows uv_{max} (water): 299 nm (ϵ 9670, 98% NH₂ OH ext.)
5. It has a solubility profile (mg . ml⁻¹): water 10, methanol 5, ethanol 0.2, acetone < 0.1, dimethyl formamide < 0.1, and dimethyl sulphoxide 0.3.

Note: It is available in combination with cilastatin sodium as Imipem, Primaxin, Tan Acid, Tienam, Tracix, Zienam.

Uses

1. It exhibits a broader antibacterial spectrum than any other **β -lactams**.
2. It happens to surpass cephalosporins against staphylococci, equals penicillin G against streptococci, equals **third generation cephalosporins** against most aerobic Gram-negative bacilli and is found to be fairly comparable to **ceftazidime** against *Ps aeruginosa*.
3. It is also equally comparable to both **metronidazole** and **clindamycin** against the anaerobes.
4. It is specifically recommended for the treatment, control and management of mixed bacterial infections.


9.3.4.4 Meropenem

Synonyms Merrem; Meronem; ICI-194660; SM-7338.

Biological Source It is also another semi-synthetic structural analogue of thienamycin, produced by *S. cuttleya* by the method proposed by Sunagawa *et al.***

* Leanza W.J. *et al.*, J. Med. Chem., **22**, 1435 (1979).

** M. Sunagawa *et al.*, *Eur. pat. Appl.* 126, 587; M. Sunagawa, *U.S. Pat.* 4, 943, 569 (1984, 1990 both to Sumitomo).

Chemical Structure

Meropenem

[4R-[3 (3S*, 5S*) 4 α , 5 β , 6 β (R*)]]-3-[[5-[(Dimethylamino) carbonyl]-3-pyrrolidinyl] thio]-6-(1-hydroxyethyl)-4-methyl-7-oxo-1-azabicyclo [3, 2, 0] hept-2-ene-2-carboxylic acid trihydrate; (C₁₇H₂₅N₃O₅S·3H₂O).

Characteristic Features


1. It is obtained as white to pale yellow crystalline powder.
2. The colour of its solutions vary from colourless to yellow depending on the concentration.
3. **Solubility Profile:** It is found to be sparingly soluble in water; soluble in 5% monobasic sodium phosphate [H₂NaO₄P] solution; very slightly soluble in ethanol; and practically insoluble in acetone or ether.

Uses

1. The resistance of meropenem to most β -lactamases is fairly good.
2. It has a similar distribution as imipenem.
3. It is not degraded by renal dehydropeptidases.
4. It possesses slightly different affinity for specific PBPs* (primary target includes PBPs 2 and 3) depending on the strain of Gram-negative microorganisms.

9.3.4.5 Nocardicin A

Biological Sources It is a monocyclic β -lactam (monobactam) antibiotic with antimicrobial activity that specifically inhibits bacterial cell wall biosynthesis. In short, **nocardicins A, B, C, D, E, F, G** have been isolated and identified duly. All are produced by *Nocardia uniformis* subspecies *tsuyamensis*, A being the most important component. However, nocardicin A has also been produced by *Actinosynnema mirum*.**

Chemical Structure

Nocardicin A

* Penicillin Binding Proteins.

** K. Watanabe *et al.*, *J. Antibiot.*, **36**, 321 (1983).

[3S-[1(S*), 3R* [Z(S*)]]]-3-[[[4-(3-Amino-3-carboxypoxy) phenyl] (hydroxy imino) acetyl]-amino]- α -(4-hydroxyphenyl)-2-oxo-1-azetidineacetic acid; [C₂₃H₂₄N₄O₉].

Isolation Nocardicin A has been isolated and characterized by Aoki *et al.**

Characteristic Features

1. It is obtained as colourless needles from acidic water having mp 214–216°C (decomposes).
2. It has specific optical rotation $[\alpha]_D^{25}$ -135° (for the sodium salt).
3. Its uv_{max} (1/15 M phosphate buffer); 272 nm ($E_{1cm}^{1\%}$ 310); (0.1 N. NaOH): 244, 283 nm ($E_{1cm}^{1\%}$ 460, 270).
4. It is found to be soluble in alkaline solutions; slightly soluble in methanol; and practically insoluble in chloroform, ethyl acetate, solvent ether.

9.3.5 Lincosamides

Lincosamides essentially comprise of two clinically useful antibiotics, namely: **Lincomycin** and **Clindamycin**; however, their overall general usage has been restricted and limited on account of their inherent potentially fatal side effect called **pseudomembranous colitis****


These two aforesaid antibiotics shall now be treated individually in the sections that follows:

9.3.5.1 Lincomycin

Synonyms Lincolnensin; Lincolcina; U-10149; NSC-70731;

Biological Source It is produced by *Streptomyces lincolensis* var. *lincolensis*.

Chemical Structure Lincomycin has an amide function in its molecule which may have been contributed essentially by a unique strategic combination of amino acid and carbohydrate metabolites. It is also obtained through **stereoselective synthesis** by the method of Knapp and Kukkola***



Lincomycin

(2S-*trans*)-Methyl 6, 8-dideoxy-6-[[[(1-methyl-4-propyl-2-pyrrodinyl) carbonyl] amino]-1-*thio-p*-erythro- α -D-galacto-octopyranoside; (C₁₈H₃₄N₂O₆S).

* H. Aoki *et al.*, *J. Antibiot.*, **29**, 492 (1976)

** **Pseudomembranous Colitis**: Colitis associated with antibiotic therapy. It is indicated by formation of a pseudomembrane on the mucosa of the colon. The symptoms are: diarrhoea, abdominal cramps, fever and leukocytosis normally after 4 to 10 days after the start of antibiotic therapy.

*** S. Knapp and P.J. Kukkola, *J. Org. Chem.* **55**, 1637 (1990).

Characteristic Features

1. It is obtained as a free-base invariably.
2. Its dissociation constant is $pK_a' 7.6$.
3. It is found to be more stable in salt form.
4. It is soluble in methanol, lower alcohols, acetone, ethyl acetate, chloroform; and slightly soluble in water.

Lincomycin Hydrochloride Hemihydrate ($C_{18}H_{34}N_2 \cdot O_6S \cdot HCl \cdot \frac{1}{2}H_2O$): [**Synonyms** **Frademicina; Lincocin; Mycivin; Wayenecomycin;**]

1. It was obtained formerly as needle-like crystals of low specific gravity from aqueous solution by rapid addition of acetone at low temperatures; but nowadays it is mostly obtained as crystals of relatively higher specific gravity, having distinct cubic crystal structure with greater solubility in HCl, by the slow addition of acetone. It has mp $145-147^\circ C$.
2. It has specific optical rotation $[\alpha]_D^{25} + 137^\circ$ (water).
3. It is found to be freely soluble in water, methanol, ethanol; and sparingly soluble in most organic solvents other than the hydrocarbons.

Uses

1. It is active against most Gram-positive bacteria, including pneumococci, staphylococci, and streptococci, with the exception of *Enterococcus faecalis*.
2. Its anaerobic spectra (both Gram-positive and Gram-negative) are also recognized as significant and distinctive.
3. Its use is restricted due to its serious side-effects which essentially include: diarrhoea, occasionally serious *pseudomembranous colitis* (caused by overgrowth of resistant strains of *Clostridium difficile*), which may cause serious fatalities especially in aged patients.


Note: Lincomycin inhibits protein biosynthesis due to the blockade of peptidyltransferase site on the 50S subunit of the bacterial ribosomes 70S.

9.3.5.2 Clindamycin

Synonyms Antirobe; Cleocin; Dalacin C; Klimicin; Sobelin; Clinimycin (rescinded); 7-Deoxy-7(S)-chloro-lincomycin.

Biological Source **Clindamycin** (7-chloro-7-deoxy-lincomycin) is synthetically derived from **lincomycin**, which is obtained from the cultures of *Streptomyces lincolensis* var *lincolensis*.

Chemical Structure The semi-synthetic derivative is obtained by the chlorination of the *lincomycin* with resultant inversion of stereochemistry.



Clindamycin

(2*S-trans*)-Methyl 7-chloro-6,7,8-trideoxy-6-[[[(1-methyl-4-propyl-2-pyrrolidiny) carbonyl] amino]-1-thio-L-threo- α -galacto-octo-pyranoside; (C₁₈H₃₃Cl N₂O₅S).

Characteristic Features

1. It is obtained as yellow amorphous solid.
2. It has specific optical rotation $[\alpha]_D + 214^\circ$ (chloroform).

Clindamycin Hydrochloride Monohydrate [C₁₈H₃₃Cl N₂O₅S.HCl.H₂O]: [*Synonym Dalacin*]:

1. It is obtained as white crystals obtained from a mixture of ethanol-ethyl acetate having mp 141-143°C.
2. It has specific optical rotation $[\alpha]_D + 144^\circ$ (H₂O).
3. Its dissociation constant is pKa 7.6.
4. It is found to be soluble in water, ethanol, DMF, and pyridine.


Uses

1. It appears slightly more effective quantitatively than its structural analogue lincomycin.
2. It is more readily eliminated from the body. The normal 300 mg dose of lincomycin gives a peak serum level of 2.6 to 3.6 mcg. ml⁻¹ in 1-2 hours, and the normal half-life ranges between 2-4 hours.
3. **Clindamycin** is reported to be one of the most effective antibiotics against strains of *Bacteroides fragilis*, the Gram-negative anaerobe that is responsible for a number of abdominal infections.
4. The antianaerobic property of **clindamycin** renders it useful in pneumonias caused by anaerobes.
5. **Clindamycin phosphate** is employed topically for the treatment, control and management of serious acne and vaginally for bacterial vaginosis.


9.3.6 Macrolides

Macrolide antibiotics are typically characterized by macrocyclic lactones having a ring-size ranging between 12-16 atoms and also possess inherent extensive branching through the methyl substituents. However, the macrolactone ring essentially bears a glycosidal linkage to either one or several sugar functions. Exhaustive biosynthetic studies have revealed that the genesis and formation of macrocyclic lactone due to the condensation of either acetate and/or propionate units, evidently *via* malonyl-CoA and 2-methylmalonyl-CoA. Interestingly, the methyl substituents present on the macrolactone ring seem to be contributed exclusively as the residual from incorporation of **propionate units** instead of the so called **terminal biological methylation**. It has been established beyond any reasonable doubt that two or even more '**sugar units**' are attached through the glycoside linkages. Further, these sugars are found to be somewhat unusual 6-deoxy structures normally restricted to this particular class of compounds *i.e.*, **macrolides**.


The four '**sugar components**' present often in macrolides are, namely: **L-cladinose**; **L-mycarose**; **D-mycinose**; and **L-oleandrose**, whose structures are given below:




L-Cladinose



D-Mycinose




L-Mycarose



L-Oleandrose

Out of all the ‘**sugar components**’ present, at least one sugar is an **amino-sugar**, such as: **D-desosamine**; **D-forosamine**; and **D-mycaminose**, whose structures are as depicted below.



D-Desosamine

D-Forosamine

D-Mycaminose

The various important members of ‘**macrolide antibiotics**’ are, namely: **Erythromycin**, **Clarithromycin**, **Arithromycin**, **Oleandomycin**; **Troleandomycin**; and **Spiramycin**.

In general, these antibiotics essentially exhibit a narrow spectrum of antibacterial activity, most importantly against the Gram-positive microorganisms. It is, however, pertinent to mention here that the antibacterial spectrum of the aforesaid antibiotics resembles, but is not very much identical to, that of the **penicillins**; hence, they cater for an extremely valuable alternative or substitute for such patients who are found to be allergic to the penicillins. It is worthwhile mentioning at this point in time that **Erythromycin** is one of the most important and principal **macrolide antibiotics** presently employed in medicine.

Some of these antibiotics shall now be treated individually in the sections that follows:


9.3.6.1 Erythromycin

Synonyms Erythromycin A; Abomacetin; Ak-Mycin; Aknin; E-Base; EMU; E-Mycin; Eritrocina; Ery Derm; Erymax; Ery Tab; Erythromast 36; Erythromid; ERYC, Erycen; Erycin; Erycinum; Ermysin; Ilotycin; Inderm, Retcin; Staticin; Stiemycin, Torlamicina.

Biological Sources It is produced by cultures of *Saccharopolyspora erythraea* (formerly known as *Streptomyces erythreus*). Waksman and Henrici were the pioneer in finding this antibiotic in a soil sample collected from the Philippine Archipelago.

Erythromycin is, in fact, a mixture containing principally *Erythromycin A*. Together with small quantum of **Erythromycins B and C**.

Chemical Structure



Erythromycin

E-Mycin (3R*, 4S*, 5S*, 6R*, 7R*, 9R*, 11R*, 12R*, 13S*, 14R*)-4-(2, 6-Dideoxy-3-C-methyl-3-O-methyl- α -L-ribo-herapyranosyl) oxy]-14-ethyl-7, 12, 13-trihydrox-3, 5, 7, 9, 11, 13-hexamethyl-6-[[3, 4, 6-trideoxy-3 (Dimethyl-amino)- β -D-xylo-hexopyranosyl] oxy] oxacyclotetradecane-2, 10-dione; (C₃₇H₆₇NO₁₃).

Characteristic Features

1. It is obtained as white or slightly yellow-crystals or powder, odourless or practically odourless, slightly hygroscopic in nature, having mp 135-140°C.
2. It is found to get resolidified with second mp 190-193°C.
3. It has specific optical rotation $[\alpha]_D^{25} -78^\circ$ (C = 1.99 in ethanol).
4. It has uv_{max} (pH 6.3): 280 nm (ϵ 50).
5. Its dissociation constant is pKa₁ 8.8.
6. It usually shows basic reaction and readily forms salts with acids e.g., acetate, estolate, glucoheptanoate, lactobionate, propionate, stearate and the like.
7. Its solubility in water is nearly 2 mg . ml⁻¹.
8. It is found to be freely soluble in alcohols, acetone, chloroform, acetonitrile, ethyl acetate; and moderately soluble in solvent ether, ethylene dichloride and amyl acetate.

Uses

1. It exhibits a relatively broad spectrum of activity which usually overlaps the activity of penicillin.
2. It is found to be *most effective* against a host of Gram-positive cocci, namely: *Enterococci*, Group A hemolytic streptococci, pneumococci, and *Staphylococcus aureus*, *N. meningitidis* and *gonorrhoeae*, *Listeria*, *Corynebacterium diphtheria*, acnes and certain specific strains of *H. influenzae* are also reported to be sensitive.
3. A low concentration of **erythromycin** also inhibit *mycoplasma* and the agent of Legionnaire's disease*.

* **Legionnaire's Disease:** A severe, often total disease characterized by pneumonia, dry cough, myalgia and sometimes gastrointestinal symptoms. It may occur in epidemic or sporadically and has become an important cause of nosocomial pneumonia. An organism, *Legionella pneumophila*, cause the disease when it is inhaled from aerosols produced by air-conditioning units, shower heads etc.,

6-O-Methylerythromycin; (C₃₈H₆₉NO₁₃).

Characteristic Features

1. It is obtained as colourless powder from a (1 : 2) mixture of chloroform and diisopropyl ether having mp 217-220°C (decomposes).
2. It is also obtained as crystals from ethanol with mp 222-225°C (Morimoto).
3. It has uv_{max} (CHCl₃) : 288 nm (ϵ 27.9).
4. It has specific optical rotation $[\alpha]_D^{24}$ -90.4° (C = 1 in CHCl₃).
5. It is found to be stable at acidic pH.

Uses

1. It is invariably used as an alternative to **erythromycin** for treating streptococcal pharyngitis, community-acquired respiratory tract infections, skin and soft tissue infections and an acute attack of **sinusitis**.*
2. It is found to be two-to-four times more active than **erythronycin** itself against a host of *streptococci* and *staphylococci* species; however, certain organisms that are resistant to **erythromycin** are also observed to be resistant to clarithromycin.
3. It exhibits a moderate activity against *H. influenzae* and *N. gonorrhoea*.
4. Clarithromycin is found to be influenced by *Branhamella catarrhalis*, *Legionella pneumophilia*, *Mycoplasma pneumoniae*, *Chlamydia trachomatis* and *pneumoniae* and *Borrelia burgdorferi* (agent of **Lyme's disease****).
5. It also shows activity against *Mycobacterium avium* and *Mycobacterium intracellulare*; and is mostly employed as primary agent for the treatment of disseminated mycobacterial infections.

9.3.6.3 Azithromycin


Synonyms Azitrocin; Sumamed; Trozocina; Zithromax; Zitromax; CP-62993; XZ-450.

Biological Source It is a semi-synthetic **macrolide antibiotic** related to **erythromycin A** which is obtained from *Saccharopolyspora erythraea*.

Chemical Structure **Azithromycin** is a tailor-made ring-expanded aza-macrolide wherein the carbonyl moiety at C-6 has been subjected to reduction; and this sort of minor alternation *vis-a-vis* the complex structure has significantly increased the activity when compared to the parent compound **erythromycin A**.

* **Sinusitis**: Inflammation of a sinus, especially a paranasal sinus. It may be caused by various agents, including viruses, bacteria or allergy:

** **Lyme's Disease**: A multisystem disorder caused by the tick-transmitted spirochete *Borhelia burgdorferi*.



Azithromycin

9-Deoxo-9 α -methyl-9 α -aza-9 α -homoerythromycin A; (C₃₈H₇₂N₂O₁₂).

Characteristic Features

1. It is obtained as white crystals having mp 113-115°C .
2. It has specific optical rotation $[\alpha]_{20}^D -37^\circ$ (C = 1 in CHCl₃).

Use


It is found to be active against staphylococci and streptococci but is more active than **erythromycin** against *H. influenzae* and some aerobic Gram-negative bacilli.

9.3.6.4 Oleandomycin

Synonyms Amymicin; Landomycin; Romicil.

Biological Source It is an **antibiotic** substance produced by fermentation cultures of *Streptomyces antibioticus* no. ATCC 11891.

Chemical Structure

Oleandomycin [C₃₅ H₆₁ NO₁₂]

Characteristic Features

1. It is obtained as white amorphous powder.
2. It has uv_{max} (methanol): 286-289 nm.
3. It is found to be freely soluble in methanol, ethanol, butanol, acetone; and almost insoluble in hexane, carbon tetrachloride, dibutyl ether.

Oleandomycin Hydrochloride ($C_{35}H_{61}NO_{12} \cdot HCl$):

1. It is obtained as long needles from ethyl acetate having mp 134-135°C.
2. It has specific optical rotation $[\alpha]_D^{25} -54^\circ$ (methanol).
3. It is freely soluble in water; and forms various crystalline hydrates.

Oleandomycin Triacetate Ester [*Synonym* Troleandomycin; Cyclamycin; Wytrion; Evramycin; Triocetin; TAO; NSC-108166.]

It is a **semi-synthetic macrolide antibiotic** prepared from oleandomycin wherein the three hydroxyl functions each at C-11, and the two sugar moieties replaced by the acetate groups.

1. It is obtained as crystals from isopropanol that yet decomposed at 176°C.
2. It is practically tasteless.
3. It has specific optical rotation $[\alpha]_D^{25} -23^\circ$ (methanol).
4. It shows a dissociation constant pKa 6.6.
5. It is found to be soluble in water < 0.1 g per 100 ml.


Uses It is useful against a number of Gram-positive bacterial infections.

9.3.6.5 Spiramycins

Synonyms Selectomycin; Revamicina; Rovamycin; RP-5337.

Biological Sources **Spiramycins** are **macrolides** produced by cultures of *Streptomyces ambofaciens* from the soil of northern France.

Chemical Structure The mixture of **spiramycins** have been successfully separated into *three* different components termed as: **Spiramycin I, II and III.**



Spiramycin I : R = H
Spiramycin II : R = COCH₃
Spiramycin III : R = COCH₂CH₃

Characteristic Features

1. It is obtained as an amorphous base.
2. It has specific optical rotation $[\alpha]_D^{20}$ -80° (methanol).
3. It has uv_{\max} (ethanol): 231 nm.
4. It is found to be slightly soluble in water; and soluble in most organic solvents.

Uses

1. It exhibits activity on Gram-positive organisms and *rickettsiae*.
2. It also shows cross resistance between microorganisms resistant to **erythromycin** and **carbomycin**.

Spiramycin I [$C_{43}H_{74}N_2O_{14}$] [**Synonym Foromacidin A**]: It is obtained as crystals having mp ranging between 134-137°C, and specific optical rotation $[\alpha]_D^{20}$ -96°.

Spiramycin II [$C_{45}H_{76}N_2O_{15}$] [**Synonym Foromacidin B**]: It is obtained as crystals having mp 130-133°C, and $[\alpha]_D^{20}$ -86°.

Spiramycin III [$C_{46}H_{78}N_2O_{15}$] [**Synonym Foromacidin C**]: It is also obtained as crystals having mp ranging between 140-142°C, and $[\alpha]_D^{20}$ -98.4°.

Spetamycin III Diacetate: It is obtained as crystals from cyclohexane having mp 140-142°C; and specific optical rotation $[\alpha]_D^{20}$ -90.4°.


Use It is used for the treatment of toxoplasmosis, and also the infections caused by the protozoan *Toxoplasma gondii*.

9.3.7 Penicillins

Alexander Fleming observed in 1928 during the course of examination of certain culture plates in the laboratory of St. Mary's Hospital, London, that the lysis of the staphylococcus organisms takes place by a contaminating mold. Subsequently, the mold was subcultured in a sterile broth under aseptic condition; and it was revealed that it resulted into a powerful, nontoxic antibacterial product. Fleming baptized this substance as '**penicillin**' based on its parent organism *Penicillium notatum* that eventually paved the way for the creation of the so called '**generation of the antibiotic**'—a historic remarkable landmark in the field of medicine derived from the natural products.

'**Penicillin**' categorically symbolizes a host of vital and significantly prominent antibiotic substances produced by the growth of different *Penicillium* species or by various semi-synthetic or synthetic means.

In general, the penicillins are nomenclatured in the literature invariably as the derivatives of: (a) (2*S*-*cis*) -4-thia-1-azabicyclo-(3, 2, 0) heptane-2-carboxylic acid [I]; (b) 3, 3-dimethyl-7-oxo-derivative of [I] and also known by its trivial name **penicillanic acid** [II]; and (c) α -carboxamido derivative of it [III], here only the '*R*' of the α -carboxamido moiety is identified ultimately, as depicted below:



In actual practice, one comes across *three* different types of **penicillins**, such as:

(a) **Biosynthetic Penicillins:** These are usually accomplished by the introduction of various acids, amines or amides directly incorporated into the medium in which the mold is being developed thereby leading to the ultimate production of a spectrum of biosynthetic penicillins which essentially differ only in 'R' in III. By adopting this unique well-developed and articulated process dozens of **biosynthetic penicillins** have been prepared with a view to obtain newer molecules that have an edge over penicillin G with regard to various physical parameters, microbiological or pharmacological characteristics.

It is pertinent to mention here that in 1958 an altogether new dimension was added to update and boost up the on-going development of penicillins. Methods were devised for modifying the very '**penicillin nucleus**' thereby making it feasible to biosynthesize penicillins which earlier could not be accomplished in a just normal medium. The concerted efforts made by the researchers resulted into the formation of plethora of altogether new series of medicinally potent substances that were often found to be more acid-stable, more-penicillinase resistant or had a wider antibacterial spectrum.

(b) **Commercial Penicillins:** In fact, a major portion of the commercial penicillin is pure crystalline penicillin G. It is invariably obtained in the fermentation liquors along with variable quantities of penicillin K and F and relatively smaller amount of others. However, penicillin G is eventually separated from the other congeners during the process of purification. Nevertheless, the commercial process of producing penicillins is observed to suppress, to some extent, the inherent natural tendency of the mold to give rise to penicillins other than the desired **penicillin G** by the introduction of a precursor of G, such as: **phenylacetic acid, phenylacetamide, phenylethylamine** or such other chemical entities containing the '**phenylacetyl**' radical, that is incorporated directly into the **penicillin G** molecule. It is worth while to state here that **penicillin G** enjoys the additional advantage of being crystallized out more easily than K or F.

(c) **Salts of Penicillins:** From the figures I, and II and III it is quite evident that penicillin are acids. The potassium salt is more prevalent and hence predominates in actual usage, with the sodium salt next. The inherent acidic moiety may be exploited skillfully and judiciously to combine **penicillins** with various bases, namely: **procaine, benzathine**, to design and evolve rather insoluble salts, for repository application, or for the objective of minimising solubility so as to render the substance more resistant to gastric acid in the stomach.

9.3.7.1 Classification and Spectrum

Initially, **penicillins** were classified either on the basis of pseudohistorical categories, or according to various numbered "**generation**", very much identical to the classification of the '**cephalosporins**' (section 9.3.3). In fact, it is rather more convenient and beneficial as well to classify them according to a well-defined chemical and antimicrobial designations, namely:

- (i) **Natural Penicillins** (best streptococcal and narrow spectrum)
- (ii) **Penicillinase-resistant Penicillins** (antistaphylococcal)
- (iii) **Aminopenicillins** (improved Gram-negative: *H. influenzae*, *Enterococcus*, *Shigella*, *Salmonella*)
- (iv) **Extended-spectrum (antipseudomonal) penicillins**
- (v) **Beta Lactamase Combinations** (expand spectrum to staph, beta-lactamase producers)

It will be worthwhile to treat a few **important penicillin drugs** individually from each category in the sections that follow:

9.3.7.1.1 Natural Penicillins

A. Penicillin G Potassium

Synonyms Crystapen; Cosmopen; Eskacillin; Forpen; Hylenta; Hyasorb; Monopen; Notaral; Pentid.

Preparation It is prepared by the interaction of 6-amino-penicillanic acid and phenyl acetyl chloride in an inert organic solvent.

Chemical Structure Please refer to section 9.3.7.1.

Characteristic Features

1. It is obtained as colourless or white crystals, or a white crystalline powder, odourless or practically so; moderately hygroscopic and gets decomposed between 214-217°C.
2. Humidity and moisture accelerates decomposition.
3. It has specific optical rotation $[\alpha]_D^{22} + 285-310^\circ$ (C = 0.7).
4. It is not appreciably affected either by air or by light.
5. The solutions usually deteriorate at room temperature, but solutions stored lower than 15°C remain stable for several days.
6. It gets rapidly inactivated by acids and alkalies, and also by oxidizing agents.
7. The pH (aqueous solution 30 mg · mL⁻¹) 5 and 7.5.
8. The dissociation constant pKa (acid) 2.8.
9. **Solubility Profile:** It is found to be very soluble in water, saline TS or dextrose solutions; soluble in ethanol (*but is inactivated by this solvent*), glycerol and several other alcohols.
10. **Penicillin G Potassium** 1 mg \equiv 1595 U.S.P. Penicillin Unit or International Unit (IU).








Uses

1. It is still recommended as an important and useful drug for the treatment of many Gram-positive organisms, such as streptococci, pneumococci, gonococci, and meningococci infections.
2. It is mostly destroyed by gastric juice and is, therefore, not given by oral route, and is best administered as IM or IV injection.
3. The K-salt as such has no advantage over the corresponding Na-salt except when high doses are used in patients on sodium restriction *e.g.*, blood-pressure patients.
4. The K-salt also avoids the incidence of *hypokalemic alkalosis* which occasionally takes place during prolonged treatment with high doses of penicillins.
5. The half-life ranges between 0.5 to 0.7 hour; except 2.5 to 10 hour in renal failure or after probenecid.




Penicillins

Class	S.No.	Name (Synonyms)	Chemical Structure	Remarks
I. Natural Penicillins (best streptococcal and narrow spectrum)	1	Penicillins G sodium or Penicillin G potassium [Benzylpenicillin sodium or Benzylpenicillin potassium]		Best narrow spectrum (streptococci), IV, IM.
	2	Penicillin V [Phenoxymethylpenicillin; Penicillin phenoxymethyl]		Same spectrum profile as Penicillin G; Oral only
II. Penicillinase-Resistant Penicillins (antistaphylococcal)	3	Cloxacillin		Oral
	4	Dicloxacillin [Maclicine]	R = Cl; in 3 above.	Preferred oral
	5	Methicillin [Dimethoxyphenecillin]		IV; interstitial nephritis may take place.
	6	Nafcillin [6-(2-Ethoxy-1-naphthamido) penicillin]		Preferred IV drug for staphylococcus
	7	Oxacillin [Oxazocilline]		Oral

(Contd.)

Class	S.No.	Name (Synonyms)	Chemical Structure	Remarks
III. Aminopenicillins (improved Gram-negative <i>H. influenzae</i> , <i>Enterococcus</i> , <i>Shigella</i> , <i>Salmonella</i> ;))	8	Amoxicillin [<i>para</i> -hydroxyampicillin; Amoxycillin; Delacillin; Sawacillin; Widecillin]		Good oral absorption
	9	Ampicillin [Bonapicillin; Doktacillin; Domicillin; Tokiocillin]		Preferred IV drug; incomplete oral absorption; diarrhoea; rash
	10	Bacampicillin		Oral prodrug converted to ampicillin
IV. Extended-Spectrum (Antipseudomonal) Penicillins	11	Carbenicillin [α -Phenyl (carboxymethyl penicillin)]		IV, high sodium, oral prodrug (Indanyl sodium) available
	12	Tiacarcillin [α -Carboxy-3-thienylmethylpenicillin]		IV, similar to carbenicillin but less sodium
	13	Mezlocillin		IV; similar to piperacillin
	14	Piperacillin [4-Ethyl-2, 3-dioxo piperazine carbonyl ampicillin]		Preferred IV; best Gram-negative spectrum

(Contd.)


Class	S.No.	Name (Synonyms)	Chemical Structure	Remarks
V. β-Lactamase Combination (expand spectrum to staph., β -lactamase producers)	15	Clavulanate Amoxicillin	 Clavulanic Acid	+ Amoxicillin Oral; more diarrhoea than amoxicillin
	16	Sulbactam-Ampicillin	 Sulbactam	+ Ampicillin IV; active vs. staph. and β -lactamase-producing <i>H. influenzae</i> <i>Strep pneum</i>
	17	Clavulanate-Ticarcillin	Clavulanic Acid + Ticarcillin	IV; active vs more Gram-negative bacilli
	18	Tazobactam-piperacillin	 Tazobactam	+ Piperacillin IV; active vs. more Gram-negative bacilli

(Abstracted from: Remington-The Science and Practice of Pharmacy Vol. II., 20th edn., 2000)

B. Penicillin V

Synonyms Acipen-V; Distaquaine V; Fenospin; Meropenin; Oracilline; Oratren; V-Cillin.

Biological Source It is obtained by the addition of phenoxyacetic acid to the *Penicillium chrysogenum* culture using yeast autolyzate as a source of nitrogen, as shown below:



Chemical Structure Please refer to Section 9.3.7.1.

Characteristic Features

1. It is obtained as crystals that get decomposed between 120-128°.
2. It is found to be fairly stable in air upto 37°C.
3. It is relatively stable to acid.
4. It has uv_{max} : 268, 274 nm (ϵ 1330, 1100).
5. **Solubility Profile** It is found to be soluble in water at pH 1.8 (acidified with HCl) = 25 mg/100 ml; soluble in polar organic solvents; and almost insoluble in vegetable oils and in liquid petrolatum.

Uses

1. **Phenoxymethylpenicillin (Penicillin V)** enjoys the greatest advantage of being recognized as ‘acid-resistant’, which is solely due to the introduction of an electron-withdrawing heteroatom (*i.e.*, O-atom of phenoxy-moiety) into the side-chain.
2. It is, therefore, suitable for oral administration.
3. It is specifically recommended for respiratory tract infections and tonsillitis.

Penicillin V Potassium Salt [C₁₆H₁₇KN₂O₅S] [*Synonyms* Antibiocin; Apsin VK; Arcacin; Beromycin; Betapen VK; Calciopen; Cliacil; Compocillin VK; Distakaps V-K; Dowpen VK; Fenoxypen; Ledercillin VK; Penavlon V; Pen-Oral; Stabicilline; Uticillin VK; Vepem; Suspen]

It is soluble in water; and has specific optical rotation $[\alpha]_D^{25} + 223^\circ$ (C = 0.2).

Uses It exhibits an antibacterial spectrum very identical to that of **penicillin G** against Gram-positive bacteria but this is less potent and effective against Gram-negative bacteria. Its biological half-life is about 0.5 to 1 hour.

9.3.7.1.2 Penicillinase-resistant Penicillins

A. Cloxacillin

Biological Source Cloxacillin is a semi-synthetic antibiotic related to penicillin; and is the chlorinated derivative of oxacilline which contains an isoxazole group.

Preparation 6-APA is acylated with 3-(o-chlorophenyl)-5-methyl-4-isoxazolecarboxylic acid. The resulting **cloxacillin** is subsequently purified by recrystallization.

Chemical Structure Please refer to section 9.3.7.1.

Characteristic Features

Cloxacillin Sodium Monohydrate: $[C_{19}H_{17}Cl N_3NaO_5S \cdot H_2O]$ [*Synonyms* Bactopen; Cloxapen; Cloxypen; Gelstaph; Orbenin; Methocillin-S; Prostaphlin-A; Staphybiotic; Tegopen]

1. It is obtained as a white, odourless crystalline powder having a bitter taste and decompose at 170°C.
2. It is stable in light; and is slightly hygroscopic in nature.
3. It has specific optical rotation $[\alpha]_D^{20} + 163^\circ$ (C = 1 in water).
4. The pH of 1% aqueous solution is 6.0-7.5.
5. Its dissociation constant pKa (COOH) is 2.7.
6. It is found to be soluble in water, methanol, ethanol, pyridine and ethylene glycol; and slightly soluble in chloroform.

Uses

1. It is **penicillinase-resistant penicillin** (antistaphylococcal) which is administered orally.
2. It is a first-choice agent against penicillin-resistant *Staphylococcus aureus*.

B. Nafcillin

Biological Source It is a semi-synthetic antibiotic related to penicillin bearing essentially a maphthamido moiety.

Preparation 6-APA is first acylated by treatment with 2-ethoxy-1-naphthoyl chloride in an anhydrous organic solvent containing triethylamine. An aqueous extract of this product is admixed with a water-immiscible solvent and nafcillin is precipitated by the addition of H_2SO_4 . The crude product may be recrystallized from chloroform.

Chemical Structure Please refer to Section 9.3.7.1.

Characteristic Features

Nafcillin Sodium $[C_{21}H_{21}N_2NaO_5S]$: [*Synonyms* Nafcil; Naftopen; Unipen]

1. It is obtained as white to yellowish white powder having not more than a slight characteristic odour.
2. It is freely soluble in water or chloroform; and soluble in alcohol.

Uses

1. It is considered as a preferred drug given through IV for staphylococci.
2. It is a **penicillinase-resistant penicillin**, the use of which is restricted to the treatment of infections caused by penicillinase-producing cocci (mostly staphylococci).

Note: After oral administration serum levels are low and invariably unpredictable, hence the oral route is not recommended.

9.3.7.1.3 Aminopenicillins

A. Amoxicillin

Synonyms Amoxycillin; Amocilline; Amolin; Amopenixin; Amoram; Amoxipen; Anemolin; Aspenil; Betamox; Cabermox; Delacillin; Efpenix; Grinsil; Helvamox; Optium; Ospamox; Pasetocin; Penamox; Penimox; Piramox; Sawacillin; Sumox.

Biological Source It is a **semi-synthetic antibiotic** related to **penicillin** with side-chain containing a basic amino moiety.

Preparation It may be prepared by carrying out the acylation of 6-aminopenicillanic acid with D-(-)-2-(p-hydroxyphenyl) glycine.

Chemical Structure Please refer to Section 9.3.7.1.

It is usually obtained as its trihydrate product.

Characteristic Features

Amoxicillin Trihydrate [$C_{16}H_{19}N_3O_5S \cdot 3H_2O$]: [**Synonyms** Alfamox; Almodan; Amoxidin; Amoxyphen; Clamoxyl; Cuxacillin; Flemoxin; Ibiamax; Moxaline; Polymox; Robamax; Sigamopen; Silamax; Trimox; Utimox; Zamocillin].

1. It is obtained as fine, white to off-white, crystalline powder, bitter taste.
2. Exposure to high humidity and temperature beyond 37°C adversely affect the stability of amoxicillin.
3. It has specific optical rotation $[\alpha]_D^{20} + 246^\circ$ (C = 0.1).
4. It has uv_{max} (ethanol): 230, 274 nm (ϵ 10850, 1400); (0.1N HCl) : 229, 272 nm (ϵ 9500, 1080); (0.1N KOH): 248, 291 (ϵ 2200, 3000).
5. **Solubility Profile:** In mg . ml⁻¹: water 4.0; methanol 7.5; absolute ethanol 3.4. It is found to be insoluble in hexane, benzene, ethyl acetate and acetonitrile.

Uses

1. Its antibacterial spectrum is very much similar to that of **Ampicillin**, except that its activity is less against *Streptococcus*; *N. meningitidis*; *Clostridium*; *Salmonella*; and *Shigella*.
2. It is found to be more acid stable than **ampicillin** and absorption is not affected appreciably by food intake.
3. It is the drug of choice for various infections caused by *Enterococcus faecalis* (*enterococcus*) *Branhamella catarrhalis* or *Bacteroides fragilis* (mild to moderate infections).
4. It is an alternate drug for infections by penicillinase-producing *Staphylococcus* (combined with clavulanic acid), *N. gonorrhoeae* (with probenecid), *E. coli* (with clavulanic acid) or *Pasteurella multocida* (with clavulanic acid).

Note: It cannot be given parenterally in conditions with severe infections.

B. Ampicillin

Synonyms Albipen; Amfipen; Ampipenin; Bonapicillin; Britacil; Doktacillin; Domicillin, Dumopen; Nuvapen; Omnipen; Penicline; Tokiocillin.

Biological Source It is orally active, semi-synthetic antibiotic which is structurally related to penicilline *i.e.*, penicillin with side-chain having a basic amino function.

Preparation The outline of the synthesis is that 6-APA is appropriately acylated with D-glycine under specific experimental parameters. Kajfez *et al.*, in 1976 put forward an alternate method of synthesis.*

Chemical Structure Please see Section 9.3.7.1. It is mostly obtained as its trihydrate product.

Characteristic Features

1. It is obtained as crystals that get decomposed between 199–202°C.
2. It has specific optical rotation $[\alpha]_D^{23} + 287.9^\circ$ (C = 1 in water).
3. It is found to be sparingly soluble in water.

Uses

1. It is the **first aminopenicillin antibiotic** which exhibits its *in vitro* spectrum against Gram-positive cocci very much similar to but usually somewhat less effective than that of penicillin G, with an exception that it is somewhat effective against *Enterococcus faecalis* (enterococcus).
2. It is $1/20$ as effective against *Streptococcus pyogenes*.
3. It is the drug of choice for treatment of infections due to sensitive strains of Strep Group B, *Enterococcus faecalis* (combined with **gentamycin**); *Listeria monocytogenes* (with or without **gentamycin**); *E. coli* (with or without **gentamycin**); and *Prot mirabilis*, and *Salmonella* (not typhi).
4. It is employed invariably as an alternative drug against *Kl pneumoniae* (with **sulbactam**), indole-positive *Proteus* (*M. morgani*, *Pr vulgaris* and *Providencia rettgeri*; with **sulbactam**), *Salmonella typhi*, *Shigella*, *Gardnerella vaginalis*, *H. influenzae* (serious infections; initially combined with **chloramphenicol**) or *Nocardia*.

Note: 1. A good number of these organisms rapidly acquire resistance by elaboration of penicillinase, hence it is invariably administered in combination with sublactam.

2. It causes allergic reactions typical of other penicillins and is found to be five-times as allergenic as penicillin G.

Ampicillin Sodium [C₁₆H₁₈Na₃O₄S] [*Synonyms* Alpen-N; Amcill-S; Ampicin; Cilleral; Omnipen-N; Penbritin-S; Pentrex; Polycillin-N; Synpenin; Viccillin.]

Preparation It is prepared by *first* dissolving ampicillin in a suitable organic solvent, and *secondly* by precipitating it as its sodium salt by the addition of sodium acetate.

Characteristic Features

1. It is obtained as white to off-white, crystalline powder, hygroscopic in nature; the L(+) form decomposes at about 205°C.
2. Its dissociation constants are pK_{a1} 2.66; pK_{a2} 7.24.
3. L(+) form has specific optical rotation $[\alpha]_D^{20} + 209^\circ$ (C = 0.2 in water).
4. It is found to be very soluble in water, isotonic NaCl or dextrose solutions.

* F. Kajfez *et al.*, *J. Heterocycl. Chem.*, **13**, 561, (1976)

5. It has been observed that L(+) form is less active as an antibiotic than the corresponding D(-) isomer.

Uses It is employed for IM or IV administration; and its actions and uses are similar to *Ampicillin*.

C. Bacampicillin

Biological Source It is a **semi-synthetic antibiotic** related to penicillin. It is an acyloxymethyl ester through the **thiazolidine** carboxyl moiety (*i.e.*, a 'prodrug'), and is duly hydrolyzed to **ampicillin** by esterases in the gut.

Chemical Structure Please refer to Section 9.3.7.1.

Characteristic Features

Bacampicillin Hydrochloride [C₂₁H₂₇N₃O₇S.HCl]: [*Synonyms* Ambacamp; Ambaxin; Bacacil; Bacampicine; Spectrabid.]

1. It is obtained as white crystals from a mixture of acetone and petroleum ether having mp 171-176°C. (decomposes).
2. It has specific optical rotation $[\alpha]_D^{20} + 161.5^\circ$; and also reported as + 173° (Bodin).
3. The pH of a 2% (w/v) aqueous solution ranges between 3 to 4.5.
4. **Solubility Profile:** It is found to be soluble 1g in 15ml. water, 7ml. alcohol and 10 ml. chloroform.

Uses

1. It is an oral prodrug converted to **ampicillin** *in vivo*.
2. It is an improved version of **aminopenicillin** with modified and enhanced Gram-negative activity against *H. influenzae*, *Enterococcus*, *Shigella* and *Salmonella*.

9.3.7.1.4 Extended-Spectrum Penicillins The various typical examples of **extended-spectrum (antipseudomonal) penicillins** are **carbenicillin** and **ticarcillin** wherein the penicillins contain an *additional-COOH moiety in the side-chain* and their overall activity is certainly broad-spectrum. Another type of **extended-spectrum penicillins** essentially include the **acylureido penicillins**, namely: **Mezlocillin** and **Piperacillin**, which are found to be much more active against *Pseudomonas aeruginosa* together with other Gram-negative organisms, such as: *Klebsiella pneumoniae* and *Haemophilus influenzae*.

These aforesaid antibiotics shall now be discussed in the sections that follows:

A. Carbenicillin

Biological Source It is a **semi-synthetic antibiotic** related to **penicillin** that essentially has an additional carboxylic function present in the side-chain.

Chemical Structure Please refer to Section 9.3.7.1.

Preparation First of all the starting esters may be prepared by acylating 6-aminopenicillanic acid (*i.e.*, 6-APA) with monoesters of phenylmalonic acid. The resulting esters are subsequently hydrolyzed with the help of an appropriate *esterase*, for instance: **α -chymotrypsin** or **pancreatin**, and extracting the liberated acid with a suitable organic solvent.

Characteristic Features

Carbenicillin Disodium [$C_{17}H_{16}Na_2O_6S$]: [*Synonyms* Anabactyl; Carbapen; Carbecin; Geopen; Hyoper; Microcillin; Pyocianil; Pyopen.]

1. It is obtained as white to off-white, crystalline powder having a bitter taste, odourless; and hygroscopic in nature.
2. The pH of a 1% (w/v) aqueous solution is 8.0.
3. It gives rise to two distinct values for dissociation constant viz., pK_{a1} 2.76 and pK_{a2} 3.50.
4. **Solubility Profile:** 1 g in 1.2 ml water; 2.5 ml ethanol; and almost insoluble in chloroform and ether.

Uses

1. It is a **carboxy benzyl penicillin** with enhanced antibacterial profile against non- β -lactamase producing Gram-negative bacilli, precisely *Pseudomonas aeruginosa*.
2. It has been observed that the D-and L-isomers actually show very slight differences in their biologic activity, besides they undergo rapid interconversion when in solution; hence, most logically the racemic mixture is employed invariably.
3. It may be safely administered to a maximum extent of 4 g per day so as to obtain serum concentration exceeding 50-60 mcg ml^{-1} , which concentrations normally inhibit most *Pseudomonas aeruginosa* strains.
4. It has been observed that the clinical efficacy may be increased appreciably by the combination therapy of carbenicillin disodium either with *tobramycin* or *gentamycin* in their respective full therapeutic dosages.

Note: There is an obvious possibility of a chemical interaction between aminoglycosides and β -lactam antibiotics, whereby the amino moieties of the aminoglycoside molecules afford to attack the β -lactam ring, ultimately result into the formation of a covalent adduct and finally the inactivation of the antibiotics. This serious drawback is easily overcome by their administration through different routes.

5. It is particularly effective in UTIs by virtue of its attainment of very high urine levels through IM.

B. Ticarcillin

Biological Source It is a broad-spectrum, semi-synthetic antibiotic related to penicillin; and it also essentially has an additional carboxylic function at *alpha* position in the side-chain.

Chemical Structure Please refer to section 9.3.7.1.3.

Preparation It may be prepared by the conversion of 2-(3-thienyl) malonic acid monobenzyl ester to the corresponding acid chloride which is subsequently condensed with 6-APA, followed by hydrogenation to convert the ester to the free acid.*

* Belgian, Pat 646,991.

Characteristic Features

Ticarcillin Disodium [C₁₅H₁₄N₂Na₂O₆S₂] [*Synonyms* Aerugipen, Monapen; Ticar; Ticarpen; Ticillin]

1. It is obtained as creamy-white, hygroscopic non-crystalline powder.
2. It is found to be quite unstable in an acidic medium.
3. It has dissociation constant pK_a (acid form) 2.44 and 3.64.
4. The pH of a concentrated solution (> 100 g. mL⁻¹) is approximately 7.0.
5. Its aqueous solutions are relatively stable; and the acidic solutions comparatively unstable.

Uses

1. Its antibacterial profile very much resembles to that of *Carbenicillin* (Section 9.3.7.1.4.A).
2. It is found to be twice as active against *Ps aeruginosa*.
3. Though it has an inherent tendency to develop resistance readily; however, with many infections the resistance is obviated by inclusion of clavulanic acid.
4. For the treatment, control and management of Gram-negative infections, it is invariably combined with either **gentamycin** (Section 9.3.1.2) or **tobramycin** (see section 9.3.1.7) so as to enhance activity and delay resistance to an appreciable extent.

C. Mezlocillin

Biological Source It is a **semisynthetic, broad-spectrum antibiotic** related to **penicillin** and **azlocillin**; and belong to the class of acylureido penicillins.

Chemical Structure Please refer to Section 9.3.7.1.

Characteristic Features

Mezlocillin Sodium Monohydrate [C₂₁H₂₄N₅NaO₅S₂.H₂O] [*Synonyms* Baycipen; Baypen; Mezlin]:

1. It is obtained as either yellowish-white powder or as pale yellow crystalline substance.
2. It has dissociation constant pK_a 2.7.
3. It is found to be soluble in water, methanol and DMF; and insoluble in acetone and ethanol.

Uses

1. It is one of the most active penicillins against *Ps aeruginosa*, with a potency almost at par with **gentamycin**.
2. It is found to be more potent against *Klebsiella* and a host of other enteric bacilli than is **carbenicillin** and **ticarcellin** (A and B above).
3. It is employed frequently as an '**alternative drug**' against infections caused by *Acinetobacter*, *Bacteroidis fragilis* (G.I. strains), *Enterobacter*, *E. coli*, *Kl pneumoniae*, *Morganella morganii*, *Pr vulgaris*, *Providencia rettgeri*, *Ps aeruginosa* (UTIs) or *Serratia*.

D. Piperacillin

Biological Source It is a **broad spectrum semi-synthetic antibiotic** related to **penicillin** bearing essentially an acylureido function.

Chemical Structure Please refer to Section 9.3.7.1.

Characteristic Features

Piperacillin Sodium [$C_{23}H_{26}N_5NaO_7S$] [*Synonyms* *Isipen; Pipril Pentcillin; Pipracil*]: It is obtained as white crystals having mp 183-185°C (decomposes). 1g gets dissolved in approximately 1.5 ml water or methanol, and 5 ml of ethyl alcohol.

Uses

1. It is an **extended-spectrum (antipseudomonal) penicillin**.
2. It is usually administered IV.
3. Its activities are very much similar to **mezlocillin sodium** (sec C above).

9.3.7.1.5 Beta-Lactamase Combinations **β -Lactamases** are the enzymes that help in opening up the β -lactam rings of penicillins, **cephalosporins** and also the related compounds exclusively at the **β -lactam bond**. Generally, the **β -lactamases** may be classified into *three* major categories, namely:

- (a) Substrate selectivity and inhibition,
- (b) Acidity/basicity of the enzyme protein, and
- (c) Intra-and extracellular location of enzyme.

Penicillinases are the enzymes which get excreted exclusively from the *bacterium* and the *genes* located on plasmids. These are broadly regarded as **Type II β -lactamases**; and are essentially responsible for the penicillin-resistant Gram-positive organisms, Gram-negative cocci, besides a host of Gram-negative bacilli.

It has been observed that the penicillinase-resistant penicillins usually get bound to the penicillinases; however the actual dissociation of the '**drug**'-enzyme complex is rather quite rapid. In actual practice, they have been successfully supplanted by three substances, namely: **clavulanic acid, sulbactam** and **tazobactam**. All these are regarded as newer breeds of β -lactamase inhibitors that specifically acylate the enzymes by creation of a '**double-bond**' (**greater electronic bondage**) and consequently afford dissociation very slowly, thereby significantly enhancing the potency of the penicillins against certain organisms and ultimately increase their therapeutic efficacy.

The combination of **β -lactamase inhibitors** with other antibiotics helps to expand the spectrum of the **antibiotic** to a significant extent which may be observed evidently by carrying out the *in vitro* studies.

There are *three important β -lactamase inhibitors* duly recognized, namely: **clavulanic acid, sulbactam, and tazobactam**, which shall now be discussed individually and also the combinations with antibiotics which are available commercially in the sections that follow:

A. Clavulanic Acid

Synonyms MM 14151

Biological Source It is a β -Lactamase inhibitor, and an antibiotic obtained as a fermentation product of *Streptomyces clavuligerus*, structurally related to the penicillins. **Clavulanic acid** enjoys the status of being the first ever reported naturally occurring **fused β -lactam containing oxygen**.*

* *J. Antibiot.*, **29**, 668, 1976

Chemical Structure Please refer to Section 9.3.7.1.

Characteristic Features

Clavulanate Potassium [C₈H₈KNO₅]: It is obtained as a white powder having a bitter taste. 1g is soluble in 2.5 ml of ethanol or in less than 1 ml of water.

Uses

1. The sulphur at position 1 of the β -lactam ring has been strategically replaced by oxygen (less electro negative); and also there is an ethylidene function present at position 2, that significantly increases reactivity with the typical exopenicillinases of *Staphylococcus aureus* and *Epidermatitis* and the Gram-negative β -lactamases of the Richmond Types II and III (*Haemophilus*, *Niesseria*, *E. coli*, *Salmonella* and *Shigella*), IV (*Bacteroides*, *Klebsiella* and *Legionella*) and V. Interestingly, these are all plasmid-mediated enzymes; and the chromosomally mediated enzymes are not inhibited at all.
2. It is absorbed well orally, but is also suitable for parenteral administration. The half-life is about 1 hour.

Clavulanate Amoxicillin Trihydrate (*i.e.*, combination of potassium salt with amoxicillin trihydrate).

Synonyms Augmentin; Amoksiklav; Co-Amoxiclav; Ciblor; Klavocin; NeO-Duplamox.

Uses

1. It is a **β -lactam antibiotic** with a β -lactamase inhibitor.
2. It extends the *in vitro* activity of **amoxicillin** to include **β -lactamase** producing strains of *H. influenzae*, *E. coli*; *Pr. Mirabilis*; and *S. aureus*.

Note:

(a) **It is pertinent to mention here that it may not extend the spectrum to various bacteria not usually killed by amoxicillin (such as: Pseudomonas aeruginosa) in the absence of β -lactamase resistance.**

(b) **Clavulanate Ticarcillin Disodium** (*i.e.*, combination of potassium salt with ticarcillin disodium):

Synonyms Betabactyl; Timentin.

Uses

1. It is employed for parenteral treatment of UTIs, skin and soft tissue, and lower respiratory tract infections, and sepsis caused due to susceptible bacteria.
2. The combination exerts an appreciable increase in activity that takes place against particularly the **β -lactamase**-producing strains of *S. aureus*, *H. influenzae*, *gonococcus*, *E. coli*, and *Klebsiella*.

Note: It fails to inhibit the β -lactamases generated by majority of strains of pseudomonas, Enterobacter and certain other Gram-negative bacilli; besides, β -lactamase-producing strains of those bacteria which eventually remain resistant to ticarcillin.

B. Sulbactam

Synonyms Penicillanic acid sulfone; Penicillanic acid 1, 1-dioxide; CP-45899.

Biological Source It is also a **semi-synthetic β -lactamase** inhibitor; and is structurally related to the **penicillins**.

Preparation 6-APA is diazotized to result into the formation of the unstable diazo derivative, which is subsequently and rapidly converted to the corresponding 6, 6-dibromo compound by carrying out the reaction in the presence of bromine. Finally, the resulting product is subjected to catalytic hydrogenolysis of the bromine atoms from the product.*

Chemical Structure Please refer to Section 9.3.7.1.

Characteristic Features

1. It is obtained as white crystalline solid having mp 148-151°C.
2. It has specific optical rotation $[\alpha]_D^{20} + 251^\circ$ (C = 0.01 in pH 5.0 buffer).
3. It is found to be soluble in water.

Sulbactam Sodium [$C_8H_{10}NNaO_5S$] [*Synonyms* Betamaze; Unasyn; CP-45899-2.]

Uses

1. It shows greater activity against **Type-I β -lactamases** than clavulanic acid, but fails to penetrate the cell walls of Gram-negative organisms.
2. It also exerts its own feeble antibacterial activity.
3. It is absorbed by the oral route but is also suitable for parenteral administration.

Sulbactam Ampicillin [*i.e.*, mixture of sodium salt with ampicillin sodium]: [*Synonyms* Bethacil (inj.); Loricin; Unacid; Unacin (inj.)]

Uses It extends the antibacterial profile of ampicillin to include **β -lactamase**-producing strains of *Acinetobacter*, *Bacteroides*, besides other anaerobes, such as: *Branhamella*, *Enterobacter*, *E. coli*, *Klebsiella*, *Neisseria*, *Proteus*, and *Staphylococcus*.

C. Tazobactam

Synonyms CL-298741; YTR-830H.

Biological Source It is a **β -lactamase inhibitor** and structurally related to the **penicillins**. It also supplants the general approach of expanding the antibacterial spectrum of certain antibiotic(s) (*e.g.*, **piperacillin sodium**) to include some **β -lactamase-producing strains**.

Chemical Structure Please see section 9.3.7.1.

Characteristic Features

Tazobactam Sodium [$C_{10}H_{11}N_4NaO_5S$] [*Synonyms* YTR-830; CL-307579]: It is an amorphous solid having mp > 170°C (decomposes).

Tazobactam Piperacillin [*i.e.*, mixture of tazobactam sodium with piperacillin sodium] [*Synonyms* Tazocilline; Tazocin; Zosyn;]: The combination product is administered by IV-infusion over 30 minutes duration. The usual total daily dose is usually 12 grammes of piperacillin and 1.5 grammes of tazobactam gives as 3.375 grammes every 6 hours. It is found to be active *Vs* more Gram-negative bacilli.

* *J. Org. Chem.*, **47**, 3344, 1982.

9.3.8 Polypeptide Antibiotics

Interestingly, a plethora of **polypeptides** of bacterial origin that are found to comprise of D- and L-amino acids, do exert a marked and pronounced antibiotic activity. It is, however, pertinent to mention here that these specific antibiotics have two inherent major anomalies, namely: *first*, very poor absorption from the intestinal tract; and *secondly*, possess high degree of nephrotoxicity* when used systemically. Generally, the **polypeptide antibiotics** exert a predominantly Gram-positive spectrum; however, there are a few-exceptions that are solely active against Gram-negative organisms, such as: the strongly basic **polymyxins**.

It has been observed that these **polypeptide antibiotics** have a tendency to occur as mixtures of very close structurally related compounds. Nevertheless, the exact composition of commercial mixtures depend to a great extent upon the skilful usage of selected strains of producing organisms. Besides, a precise and reliable strength of therapeutic response against certain susceptible organisms is exclusively based on the quantitative microbial assay.

The various important members of '**polypeptide antibiotics**' are, namely: **cycloserine**; **polymyxin-B**; **colistin (polymixin-E)**, **bacitrasin**; **vanomycin**; and **teichoplanin**, which shall now be treated separately as under:


9.3.8.1 Cycloserine

Synonyms Closina; Farmiserina; Micoserina; Orientomycin; Oxamycin; Seromycin; PA-94.

Biological Sources It is a **polypeptide antibiotic** substance produced by *Streptomyces garyphalus* sive *orchidaceus*.**

Preparation It may also be synthesized by the method of Stammer *et al.****

Chemical Structure



Cycloserine

D-4-Amino-3-isoxazolidinone; C₃H₆N₂O₂.

Characteristic Features

1. It is obtained as crystals that decompose at 155-156°C.
2. Its specific optical rotations are: $[\alpha]_D^{23} + 116^\circ$ (C = 1.17); $[\alpha]_{546}^{25} + 137^\circ$ (C = 5 in 2N NaOH).
3. It has uv_{max} : 226 nm ($E_{1cm}^{1\%}$ 402).
4. Its aqueous solutions have a pH 6.
5. It is fairly soluble in water; and slightly soluble in methanol and propylene glycol.

* A toxic substance that causes damage to kidney tissues.

** Kuehl, Jr. *et al. J. Am. Chem. Soc.*, 77, 2344, (1955).

*** Stammer *et al. J. Am. Chem. Soc.*, 77, 2346, (1955).


- It is found to form salts readily with acids and bases.
- Its aqueous solutions buffered to pH 10 with Na_2CO_3 may be stored without any loss of activity upto a duration of one week between 0-10°C (*i.e.*, at refrigerated temperatures).

Uses

- It exhibits a fairly broad spectrum of activity; however, its therapeutic efficacy is exclusively associated with its inherent inhibitory effect on *Mycobacterium tuberculosis*.
- It precisely inhibits alanine racemase, which action precludes the incorporation of D-alanine strategically into the pentapeptide side-chain of the specific **murein*** component of bacterial cell walls. Perhaps this unique features is solely responsible for its antibiotic activity.
- It is invariably regarded as an '**antibiotic of second choice**'; and is frequently used in conjunction with **isoniazid** in the control, management and treatment of tuberculosis who usually fail to respond to the **first-line agents.****
- It is readily absorbed orally and is subsequently exerted quickly through the kidneys (*i.e.*, newly 50% without any metabolic alteration whatsoever).

9.3.8.2 Polymixin B

Polymyxins represent a group of cycle polypeptide antibiotics produced by various species of *Bacillus*. However, **polymyxins A to E** were primarily isolated from *Bacillus polymyxa*. Subsequently, it was shown that both **polymyxin B** and **polymycin E** (or **colistin**) were mixtures of two components each. The structures of **polymyxin B₁** and **polymyxin B₂**; and **polymyxin E₁** (**colistin A**) and **polymyxin E₂** (**colistin B**) are as given below:



Actually, these molecules essentially contain ten amino acids, of which six happen to be **L-α-, γ-diaminobutyric acid (L-Dab)**, having a fatty acid*** strategically bonded to the N-terminus; besides, a cyclic peptide portion meticulously designed *via* an amide bond located in between the **γ-amino** of one of the **Dab-residues** and the **carboxyl terminus**. Interestingly, the **γ-amino** functions

* Murien: Chloride (Cl⁻).

** Rifampin and Rifabutin.


*** 6-Methyloctanoic acid; or 6-Methylheptanoic acid.

of the remaining Dab residues distinctly attribute a rather strong basic property to the various antibiotics. This particular characteristic feature confers detergent-like properties and perhaps permits them to either get bound or cause damage to bacterial membranes.

Characteristic Features

Polymyxin Hydrochloride

1. It is obtained as nearly colourless powder that gets decomposed at 228-230°.
2. It has specific optical rotation $[\alpha]_D^{23} -40^\circ$ (C = 1.05).
3. It is very soluble (> 40%) in water and methanol; the solubility decreases considerably in higher alcohols; and almost insoluble in ethers, esters, ketones, hydrocarbons and the chlorinated solvents.
4. It usually gives rise to water insoluble salts with the help of a host of precipitants, such as: **helianthic acid** (C₇H₉O₄) **picric acid**; and **Reincke salt**.



Picric Acid



Reinecke Salt

Polymyxin B: It is a mixture of **Polymyxins B₁ and B₂**. The mixture also contains minimal amounts of the more toxic **polymyxins A, C and D**. Both **polymyxins B₁ and B₂** essentially possess a cyclopeptidic structure and comprise of six residues of **α, γ -diaminodutyric acid (DABs)**. However, the latter characteristic feature affords an exceptionally strong basic property to the **polymyxin antibiotics**.

It has specific optical rotation $[\alpha]_{5461} -106.3^\circ$ (1N. HCl).

Uses

1. It is used topically in ointments (usually 5000 or 10,000 Units/g) and ophthalmic solutions (10,000 Units/ml).
2. It was employed formerly for control, management and treatment of infections of the intestinal tract caused by *Shigella*, *Pseudomonas aeruginosa*, and *E. coli*.

Polymyxin B Sulphate [Synonyms Aerosporin; Mastimyxin;]: It is the sulphate salt of a substance produced by the growth of *Bacillus polymyxa* (Prazmowski) Mignla belonging to the natural order *Bacillaceae*. It has a potency of not less than 600 Units of **polymyxin B**. mg⁻¹, calculated on the anhydrous basis.

Preparation The filtered broth obtained from the fermentation process (section 9.3.7) is eventually treated with a 'certified dye', and the resulting **polymyxin B-dye salt complex** thus precipitated is collected by means of filtration, washed with water and finally treated with an alcoholic solution of a lower aliphatic amine sulphate. The **polymyxin B sulphate** thus produced is filtered off and subsequently purified and lyophilized. Polymyxin B is a mixture of **polymyxin B₁** (C₅₆H₉₈N₁₆O₁₃),

and **polymyxin B₂** (C₅₅H₉₆N₁₆O₁₃) the only vital point of difference is nothing but the composition of the N-acyl moiety (see Section 9.3.8.2).

Characteristic Features

1. It is a white to buff-coloured powder; either odourless or having a very faint odour.
2. It has dissociation constant pKa 8 to 9.
3. Its solutions are either slightly acidic or are neutral to litmus (pH 5 to 7.5).
4. It is found to be freely soluble in water; and slightly soluble in alcohol.

Uses

1. The antimicrobial spectrum of activity of **polymyxin B** sulphate for its *in vitro* and *in vivo* profile is solely restricted to Gram-negative organisms, namely: *Aerobacter*, *Escherichia*, *Haemophilus*, *Klebsiella*, *Pasteurella*, *Pseudomonas*, *Salmonella*, *Shigella*, most *Vibrio* and *Yersinia*; all strains of *Pr. providencia* and most of *Serratia marcescens* are found to be unaffected by this antibiotic.
2. It is used topically either for the treatment or the prevention and treatment of external ocular infections caused by susceptible microorganisms, especially *Ps aeruginosa*.
3. In topical therapy, it is invariably combined with **neomycin**, **gramicidin** and **bacitracin**.
4. It also forms an integral component in **glucocorticoid ophthalmological topical preparations**.

Note: Substances like soap, which is a triglyceride of fatty, acids, and hence specifically antagonize cationic surface-active agents, is found to impair the activity of the antibiotic.

Polymyxin B Sulphate mixture with Trimethoprim [Synonyms Polytrim]: The combination of polymyxin B sulphate with trimethoprim enhances the overall antibacterial profile rather than each one used alone.

Polymyxin B₁ [C₅₆H₉₈N₁₆O₁₃]


Polymyxin B₁ Pentahydrochloride [C₅₆H₉₈N₁₆O₁₃·5HCl]: It is obtained as a white powder. It has specific optical rotation $[\alpha]_D^{25} -85.11^\circ$ (C = 2.33 in 75% ethanol).

Polymyxin B₂ [C₅₅H₉₆N₁₆O₁₃]: It has specific optical rotation $[\alpha]_{5461}^{22} -112.4^\circ$ (2% acetic acid).

9.3.8.3 Colistin

Synonyms Polymyxin E; Colimycin; Coly-Mycin; Colisticina; Totazina.

Biological Source It is a cyclopolypeptide antibiotic produced by *Bacillus colistinus* (*Aerobacillus colistinus*) first isolated from Japanese soil). It is comprised of **colistins A, B and C**.*



* Suzuki *et al.*, *J. Biodyem.* (Tokyo), **54**, 25 (1963).

DAB = α, γ -diamobutyric acid

Polymyxin E₁ (Colistin A) : R = (+)-6-Methyloctanoyl

Polymyxin E₂ : R = 6-Methylheptanoyl

Uses This antibiotic has more or less the same spectrum and therapeutic application as that of **polymyxin B**.

Colistin Sodium Methanesulphonate [C₅₈H₁₀₅N₁₆Na₅O₂₈S₅] [**Synonyms Colistimethate sodium; Alficetin; Methacolimycin**]: It is the injectable form of colistin. It is soluble in water and fairly stable in the dry form. It is inactive in itself but releases active polymyxin in the body.

Colistin Sulphate [**Synonyms Malimyxin; Multimycine**]: It is mostly used either orally or topically.

Colistin Formaldehyde-Sodium Bisulphite*: It is obtained as crystals that decompose between 290-295°. It is found to be soluble in water; and slightly soluble in methanol, ethanol, acetone and ether.

Polymyxin E₁ [C₅₃H₁₀₀N₁₆O₁₃] [**Synonym Colistin A**]: It has specific optical rotation $[\alpha]_{5461}^{22} -93.3^\circ$ (2% acetic acid).

Polymyxin E₂ [C₅₂H₉₈N₁₆O₁₃]: It has specific optical rotation is $[\alpha]_{5461}^{22} -94.5^\circ$ (2% acetic acid).


Note: The use of methacolimycin nowadays is rarely justified on account of the availability of less toxic alternative antibiotics.

9.3.8.4 Bacitracin

Synonyms Altracin; Ayfivin; Fortracin; Penitracin; Topitracin; Zutracin.

Biological Source It is a polypeptide antibiotic complex produced by *Bacillus subtilis* and *licheniformis* (family: *Bacillaceae*).** The commercial **bacitracin** is found to be a mixture of at least nine bacitracins. The purification of **bacitracin** may be affected by 'carrier displacement method'.

Chemical Structure The major component of the mixture is '**Bacitracin A**', which is essentially a dodecylpeptide having five of its **amino-acid-residues** arranged strategically in a cyclic structure as shown below:



* Koyama *et al. Japan. pat.* 57, 4898 (1957).

** Anker *et al. J. Bacteriol.* 55, 249 (1948).

Characteristic Features

1. It is obtained as a Grayish-white powder having a very bitter taste, odourless and hygroscopic in nature.
2. It is found to be soluble in water and ethanol; and almost insoluble in ether, chloroform, and acetone.
3. It is fairly stable in acid solution and unstable in alkaline solutions.
4. It affords a loss in potency most probably on account of the transformation of **bacitracin A** to **bacitracin F**, and the latter does not have any antimicrobial activity.
5. Its solutions undergo rapid deterioration at room temperature, and ultimately affords precipitation.
6. Its activity is significantly negated by salts of many of the heavy metals.
7. Its aqueous solutions invariably retain their potency for several weeks when stored in a refrigerator.

Uses

1. It is found to be effective exclusively against Gram-negative organisms.
2. Its applications are more or less limited to such infections only which may be treated either by topical application or by local infiltration.
3. It is significantly effective topically in the control, management and treatment of the following cutaneous bacterial infections where the pathogenic organism is specifically **bacitracin-sensitive**, such as: impetigo-contagiosa; folliculitis; pyoderma; ecthyma; furunculosis; decubitus, ulcer; infectious eczematoid dermatitis; scabies and dermatophytosis.
4. **Bacitracin** also finds its applications in the treatment of various ophthalmological conditions.
5. Its zinc salt invariably is preferred for topical therapy; and is the form most often incorporated into combinations.
6. It is mostly combined with **neomycin** and **polymyxin B sulphate**.

Note: 1. **Due to the relatively high incidence of nephrotoxicity (albuminuria, cylindruria, azotemia, accumulation of drug) which essentially follows its parenteral administration precludes systemic usage except in life-endangering staphylococcal infections, such as: pneumonia, empyema, particularly in infants wherein other antibiotics have proved to be either ineffective or in the treatment of antibiotic-associated (pseudomembranous)-enterocolitis caused by *Cl difficile*.**

2. **Development of bacterial resistance is much less frequent and slower for bacitracin as compared to penicillin, and for most organisms it is found to be almost nil.**


9.3.8.5 Vancomycin

Synonyms Vancocin; Vncoled; Lyphocin (Lyphomed).

Biological Source It is an **amphoteric glycopeptide antibiotic** substance produced by *Streptomyces orientalis* (Family: *Streptomycetaceae*) from *Indonesian* and *Indian* soil that essentially inhibits bacterial nucleotide biosynthesis by formation of complexes.

Chemical Structure The structure of the primary component of the mixture has been established beyond any reasonable doubt to be a **complex tricyclic aglycone** linked **glycosidically** to *glucose* and **vincosamine** functions. The vancomycin molecule essentially contains one free carboxylic acid moiety, two chloro substituted aromatic residues, and seven amide bonds, one of which is a prominent **primary-amide**.

The apparently novel feature of **vancomycin** is the tricyclic structure exclusively generated by *three* phenolic oxidative coupling reactions.



Preparation It is produced by the submerged fermentation process as described earlier under penicillins.

Characteristic Features

Vancomycin Monohydrochloride [$C_{66}H_{75}Cl_2N_9O_{24} \cdot HCl$] [*Synonyms* Lyphocin; Vancor;]:

1. It is obtained as white solid, free-flowing powder, odourless and having a bitter-taste.
2. It has uv_{max} (H_2O): 282 nm ($E_{1cm}^{1\%}$ 40).
3. Its solubility in water is more than 100 mg. ml^{-1} .
4. It is found to be moderately soluble in dilute methanol; and insoluble in the higher alcohols, acetone and ether.
5. Its solubility in neutral aqueous solutions is enhanced by low concentrations of urea.
6. The acidic solutions precipitate out the antibiotic on addition of either NaCl or $(NH_4)_2 SO_4$.

Uses

1. It has a Gram-positive antibacterial spectrum.
2. It specifically acts on bacterial cell walls by inhibiting murein biosynthesis by virtue of its complexation with the D-alanyl-D-alanine precursor and hence is **bactericidal**, which eventually renders it particularly useful in serious infections besides in the immunocompromised patients.
3. It also exerts to a certain extent the '**secondary modes of action**' *i.e.*, enhancing cytoplasmic membrane permeability and impairing RNA synthesis.
4. **Vancomycin hydrochloride** is widely recommended for the control, management and treatment of serious infections, such as: septicemia, endocarditis, wound infections caused by Gram-positive bacteria, specifically in those patients who are allergic to **β -lactam antibiotics**.

5. **Vancomycin HCl** is also found to be effective in *Enterococcus faecalis* strains that are inadequately controlled and managed by **β -lactam antibiotics**.
6. **Vancomycin** is not absorbed orally; however, oral administration is usually recommended for the treatment of staphylococcal-enterocolitis and antibiotic-associated pseudomembranous colitis produced by *Clostridium difficile*.
7. IM administration is rather painful and very often associated with local necrosis; therefore, systemic therapy with **vancomycin** makes use of IV-infusion extended over a span of 20 to 30 minutes.

Note: (i) It is irritating to tissue and hence may cause thrombophlebitis, or pain at the site of injection and neurosis takes place if extravasted; also produces chills, fever, occasional urticaria and maculopapular rashes with hypotension (Red Man's Syndrome), nephrotoxicity and ototoxicity and, rarely, thrombocytopenia and neuropathy.

(ii) Recently, the plasmid-mediated resistant strains of enterococcus have virtually clamped restriction for the use of vancomycin in hospitals with a view to control the spread of resistance.

9.3.8.6 Teicoplanin

Synonyms Tiecoplanin A₂; Teichomycin A₂; Targocid; Targosid; MDL-507.

Biological Source It is a **glycopeptide antibiotic** complex produced by *Actinoplanes teichomyceticus* nov. sp.; structurally related to **vancomycin** and comprised of a mixture of five **teicoplanins**, which essentially differ only in the nature and length of the fatty acid-chain attached to the sugar residue.

Characteristic Features

1. It is obtained as an amorphous powder having mp 260°C (decomposes).
2. It has uv_{max} in 0.1 N HCl 278 ($E_{1cm}^{1\%}$ 53); and in 0.1 N NaOH: 297 ($E_{1cm}^{1\%}$ 74).
3. It is found to be soluble in aqueous solution at pH 7.0; partially soluble in methanol, ethanol; and insoluble in dilute mineral acids, and also in non-polar organic solvents.

The various physical parameters of the five major components of teicoplanin are as follows:

- (i) **Teicoplanin A₂-1: (C₈₈H₉₅Cl₂N₉O₃₃):** It is a white amorphous powder which darkens at 220°C and decomposes at 255°C.
- (ii) **Teicoplanin A₂-2: (C₈₈H₉₇Cl₂N₉O₃₃):** It is a white amorphous powder, darkens at 210°C and gets decomposed at 250°C.
- (iii) **Teicoplanin A₂-3: (C₈₈H₉₇Cl₂N₉O₃₃):** It is a white amorphous powder, darkens at 210°C and decomposed at 250°C.
- (iv) **Teicoplanin A₂-4: (C₈₈H₉₉Cl₂N₉O₃₃):** It is a white amorphous powder, darkens at 210°C and gets decomposed at 250°C.
- (v) **Teicoplanin A₂-5: (C₈₉H₉₉Cl₂N₉O₃₃):** It is a white amorphous powder, darkens at 210°C and gets decomposed at 250°C.










Uses

1. **Teicoplanin** has almost similar antibacterial profile to **vancomycin** (Section 9.3.8.5), but possesses a longer duration of action, and may be administered by IM as well as IV injection.

- It is also employed against Gram-positive pathogens that are resistant to established antibiotics.
- The 'Red-Neck Syndrome' as observed upon rapid administration of **vancomycin** is rarely seen, besides the incidence of autotoxicity also seems to be reduced considerably.

9.3.9 Tetracyclines

The **tetracyclines** are a conglomerate of broad spectrum orally active **actinomycete antibiotics** produced by cultures of *Streptomyces* species, and possessing appreciable therapeutic value. **Chlortetracycline** was the first bonafide member of this group isolated from *Streptomyces aureofaciens* and discovered by Duggar in 1948. It was immediately followed by **oxytetracycline** in 1950 from the cultures of *Streptomyces rimosus*; and in 1953 **tetracycline** was eventually discovered in the antibiotic mixture from *S. aureofaciens* as a minor antibiotic.



Consequently, the intensive and extensive research and development in the selection of **mutant strains**, and specifically in the manipulations and manifestations to monitor and control both '**methylation**' and '**chlorination**' procedures have resulted in the fermentative production of a good number of **tetracycline variants**, namely: **demeclocycline**, **methacycline**, **doxycycline**, **minocycline**, **lymecycline** as given below:

These compounds shall now be discussed individually as under:

9.3.9.1 Tetracycline

Synonyms Deschlorobiomycin; Tsiklomitsin; Abricycline; Ambramycin; Bio-Tetra; Cyclomycin; Dumocyclin; Tetradeclin;

Biological Source It is obtained from a *Streptomyces* species cultured in an appropriate nutrient medium.

Preparation It may be prepared by removal of chlorine from chlortetracycline and subjecting it to hydrogenation.

Chemical Structure Please see Section 9.3.9.

Characteristic Features

1. It is obtained as a yellow crystalline powder; odourless, and stable in air.
2. It usually darkens on exposure to strong sunlight.
3. Its potency is seriously affected in solutions of $\text{pH} < 2$.
4. It is destroyed rapidly by alkali hydroxide solutions.
5. It is found to be more soluble than chlortetracycline.
6. It is rather more stable within the physiological and moderately alkaline spectrum of pH.
7. The solutions of **tetracycline** gets darkened more rapidly than **chlortetracycline** but less than oxytetracycline.
8. The pH of an aqueous suspension ($1 \text{ mg} \cdot \text{mL}^{-1}$) ranges between 3.0 to 7.0.
9. Its dissociation constant pK_a are: 3.3; 7.7; 7.9.
10. **Solubility Profile:** 1g is soluble in $\sim 2500 \text{ mL}$ water; $\sim 50 \text{ mL}$ ethanol; freely soluble in dilute HCl or alkali hydroxide solutions; and almost insoluble in ether or chloroform.

Uses

1. It is found to be useful in the treatment of toxoplasmosis.
2. The GI side effects are comparatively less than those from chlortetracycline and **oxytetracycline** but more than from **demeclocycline**.
3. The plasma half-life ranges between 6 to 11 hours in patients with normal renal function.

Tetracycline Hydrochloride [$\text{C}_{22}\text{H}_{24}\text{N}_2\text{O}_8 \cdot \text{HCl}$] [**Synonyms** Achro; Achromycin; Ala Tet; Ambracyn; Ambramicina; Bristaciclina; Cefracycline; Cyclopar; Diocyclin; Hostacyclin; Mephacyclin; Panmycin; Polycycline; Quadracyclin; Remicyclin; Sanclomycine; Supramycin; Tetramycin; Topicycline; Totomycin; Unicin.

Characteristic Features

1. The crystals of **tetracycline hydrochloride** are obtained from butanol + HCl which decomposes at 214°C .

2. Its specific optical rotation $[\alpha]_D^{25}$ -257.9° (C = 0.5 in 0.1 N HCl).
3. It is freely soluble in water; soluble in methanol, ethanol; and insoluble in ether and hydrocarbons.
4. The pH of a 2% (w/v) aqueous solution ranges between 2.1-2.3.

9.3.9.2 Chlortetracycline

Synonyms 7-Chlorotetracycline; Acronize; Aureocina; Aureomycin; Biomitsin; Centraureo; Chrysoomykine; Orospray.

Biological Source It is obtained from the substrate of *Streptomyces aureofaciens*.

Chemical Structure Please refer to Section 9.3.9.

Characteristic Features

1. It is obtained as golden-yellow crystals having mp 168-169°C.
2. It has specific optical rotation $[\alpha]_D^{23}$ -275.0° (methanol).
3. It has uv_{max} (0.1 N HCl): 230, 262.5, 267.5 nm, and (0.1N NaOH): 255, 285, 345 nm.
4. Its solubility in water ranges between 0.5-0.6 mg mL⁻¹, very soluble in aqueous solutions above pH 8.5; freely soluble in the cellosolves, dioxane and carbitol; slightly soluble in methanol, ethanol, butanol, acetone, benzene, ethyl acetate; and almost insoluble in ether and petroleum ether.

Uses

1. It exerts antiamebic activity.
2. It is the first tetracycline antibiotic available for topical application, including ophthalmic purposes.
3. Though its general use has been replaced by other tetracycline antibiotics in human beings, but it is still employed in veterinary medicine.

9.3.9.3 Oxytetracycline

Synonyms Glomycin; Riomistin; Hydroxytetracycline.

Biological Sources It is an antibiotic substance obtained from the elaboration products of the actinomycete, *Streptomyces rimosus*, grown on a suitable medium. **Oxytetracycline** may also be obtained from *Streptomyces xanthophaeus*.

Chemical Structure Refert to Section 9.3.9.

Characteristic Features

1. It is obtained as pale yellow to tan, odourless, crystalline powder.
2. It is fairly stable in air, but an exposure to strong sunlight gets darkened.
3. Like tetracycline it also gets deteriorated in solution of pH less than 2, and is quickly destroyed by alkali hydroxide solutions.
4. Its saturated solution is almost neutral to litmus and shows a pH ~ 6.5.
5. **Solubility Profile:** 1g in 4150 mL water; 100 ml ethanol; > 10,000 ml chloroform; 6250 mL ether; and freely soluble in diluted HCl or alkaline solutions.
6. **Stability:** Its crystals exhibit no loss in potency on heating for a duration of 4 days at 100°C; whereas the hydrochloride crystals show < 5% inactivation after 4 months at 56°C. It has been

observed that the aqueous solutions of the hydrochloride at pH 1.0 to 2.5 are quite stable for at least 30 days at 25°C. It has been observed that the aqueous solutions of the hydrochloride at pH 1.0 to 2.5 are quite stable for at least 30 days at 25°C. However, its solutions at pH 3.0 to 9.0 show no detectable loss in potency on storage at + 5°C for at least 30 days. Half life in hours at aqueous oxytetracycline solutions at 37°C: pH 1.0 = 114; pH 2.5 = 134; pH 4.6 = 45; pH 5.5 = 45; pH 7.0 = 26; pH 8.5 = 33; and pH 10.0 = 14.

Oxytetracycline Hydrochloride [C₂₂H₂₄N₂O₉·HCl] [*Synonyms Alamycin, Duphacycline; Engemycin; Geomycin; Oxlopar; Oxybiocycline; Oxycyclin Oxy-Dumocyclin; Oxytetracid; Oxytetrin; Tetran; Vendarcin*]

It is obtained as yellow platelets from water and is found to be extremely soluble in water (1g/mL). It is also soluble in absolute ethanol: 12,000 g/mL; and in 95% (v/v) ethanol: 33,000 g/mL.

Note: Its concentrated aqueous solutions at neutral pH hydrolyze on standing and consequently deposit crystals of oxytetracycline.

Oxytetracycline Dihydrate [C₂₂H₂₄N₂O₉·2H₂O] [*Synonyms Abbocin; Clinimycin; Oxymycin; Stevacin; Terramycin; Unimycin*]

It is obtained as needles from water or methanol which decompose at 181-182°C. Its specific optical rotations are: $[\alpha]_D^{25} -2.1^\circ$ (0.1N NaOH); and $[\alpha]_D^{25} -196.6$ (0.1N HCl). It has uv_{max} (pH 4.5 phosphate buffer 0.1 M): 249, 276, 353 nm ($E_{1cm}^{1\%}$ 240, 322, 301). It is found to be soluble in water at 23°C at various pH's: pH 1.2 = 31, 400 g/mL; pH 2.0 = 4600 γ /mL; pH 3.0 = 1400 γ /mL; and pH 9.0 = 38,600 γ /mL. It is soluble in absolute ethanol 12,000 γ /mL and in 95% (v/v) ethanol 200 g/mL.

9.3.9.4 Demeclocycline

Synonyms Biotercilin; Declomycin; Deganol; Ledermycin; Periciclina; Demethylchlortetracycline (obsolete).

Biological Source Demeclocycline is related to tetracycline and produced by *Streptomyces aureofaciens*.


Preparation A suitable strain of *S. aureofaciens* is grown in an appropriate liquid nutrient medium under controlled experimental parameters of pH, temperature, and extent of aeration. Subsequently, the duly harvested broth is acidified carefully and filtered. The demeclocycline is isolated from the resulting filtrate, either by solvent extraction or by chemical precipitation.

Chemical Structure Refer to section 9.3.9.

Characteristic Features

Demeclocycline Hydrochloride [C₂₁H₂₁ClN₂O₈·HCl] [*Synonyms: Clortetrin; Demetraciclina; Detravis; Meciclin; Mexocine*]

1. It is obtained as yellow, crystalline powder, odourless and having a bitter taste.
2. The pH of 1 in 100 solution is ~ 2.5.
3. It essentially has three distinct dissociation constants, namely: pKa_{1,2,3}: 3.3, 7.2, 9.3 attributed by three separate zones in its complex molecule as shown below:



4. **Solubility Profile:** 1g soluble in ~ 60 mL water; 200 mL ethanol or 50 mL methanol; sparingly soluble in alkali hydroxides or carbonates; and almost insoluble in chloroform.

Uses

1. It is an intermediate-acting tetracycline and causes comparatively a greater extent of **phytotoxicity** than other members of its class.
2. Its better absorption and slower excretion by the body render blood levels that distinctly afford certain minor therapeutic advantages than other members of its class.

Demeclocycline Sesquihydrate It has mp 174-178°C (decomposes); and specific optical rotation $[\alpha]_D^{25}$ -258° (C = 0.5 in 0.1 N H₂SO₄).

9.3.9.5 Methacycline

Synonyms Metacycline, Bialatan; 6-Methylene-5-hydroxytetracycline.

Biological Source It is broad spectrum, semi-synthetic antibiotic related to **tetracycline**, which is obtained from *Streptomyces rimosus*.

Preparation It may be prepared by a chemical dehydration reaction from **oxytetracycline**; besides, it has a methylene function at C-6 position.

Chemical Structure Please refer to Section 9.3.9.

Characteristic Features

Methacycline Hydrochloride [C₂₂H₂₂N₂O₈.HCl] [*Synonyms* *Adriamicina*; *Ciclobiotic*; *Germiciclin*; *Metadomns*; *Metilenbiotic*; *Londomycin*; *Optimycin*; *Physiomycline*; *Rindex*; *Randomycin*]:

1. It is invariably obtained as crystals containing 0.5 mole water and 0.5 mole methanol; and also from a mixture of methanol + acetone + concentrated HCl + ether. It is a yellow crystalline powder which decompose at ~ 205°C and has a bitter taste.
2. It has uv_{max} (methanol + 0.1 N HCl): 253, 345 nm (log ε 4.37, 4.19).
3. It is found to be soluble in water; sparingly soluble in ethanol; and practically insoluble in chloroform and ether.

Uses

1. The utility of methacycline is particularly associated with good oral absorption.
2. It has a prolonged serum half-life.

9.3.9.6 Doxycycline

Synonyms Jenacylin; Supracyclin; Vibramycin.

Preparation **Methacycline** (*i.e.*, 6-deoxy-6-demethyl-6-methylene-5-oxytetracycline) is either dissolved or suspended usually in an inert organic solvent, for instance: *methanol* and subjected to hydrogenation under the influence of catalytic quantities of noble metals, namely: *Rhodium* or *Palladium* to yield a mixture of the 6 α - and 6 β -methyl epimers. The desired epimer *i.e.*, **α -6-deoxy-5-hydroxytetracycline**, is subsequently isolated by specific chromatographic methods (US Pat 3,200,149).

Chemical Structure Refer to Section 9.3.9.

Characteristic Features

Doxycycline Hydrochloride Hemihydrate Hemihydrate [C₂₂H₂₅Cl N₂O₈.1/2C₂H₆O.1/2H₂O]

[**Synonyms** *Doxycycline hyclate*; *Azudoxat*; *Diocimex*; *Doxatet*; *Doxychel hyclate*; *Duradoxal*; *Hydramycin*; *Paldomycin*; *Sigadoxin*; *Tetradox*; *Unacil*; *Vibramycin hyclate*; *Vibra-Tabs*; *Zadorin*];

1. It is obtained as light yellow, crystalline powder from ethanol + HCl; and gets charred without melting at ~ 201°C.
2. It has specific optical rotation $[\alpha]_D^{25}$ -110°C (C = 1 in 0.01 N methanolic HCl).
3. It has uv_{max} (0.01N methanolic HCl): 267, 351 nm (log ϵ 4.24, 4.12).
4. It is found to be soluble in water.
5. Both the inherent ethanol and water of crystallization (1/2 mol of each) are usually lost by subject to drying at 100°C under reduced pressure.
6. Its dissociation constant has three values, namely: pKa 3.4, 7.7, and 9.7 (see **demeclocycline**).
7. **Solubility Profile:** It is very slightly soluble in water; freely soluble in dilute acid or alkali hydroxide solution; sparingly soluble in ethanol; and practically insoluble in ether or chloroform.

Uses

1. The 6 α -isomer of **doxycycline** is found to be more active biologically than the corresponding 6 β -epimer hydrochloride.
2. It is active against Gram-positive organisms wherein it is almost *twice* as potent as **tetracycline**; and having an exception that it is virtually 10 times as potent against *Streptomyces viridans*.
3. Interestingly, strains of *Enterococcus faecalis* that are observed to be more resistant to other **tetracyclines** may prove to be sensitive to this drug.
4. Against Gram-negative organisms it is found to be twice as potent as tetracycline.
5. It is considered to be the drug of first choice for the prophylaxis of *traveler's diarrhea*, commonly caused by enterotoxigenic *E. coli*.
6. It is found to be the best amongst the '*tetracyclines*' against anaerobes.
7. It is absorbed almost completely *i.e.*, 90 to 100% through oral administration than the rest of tetracyclines, and its absorption does not seem to be retarded by intake of foods.
8. Its plasma-protein binding is almost 93%.
9. Its volume of distribution stands at 0.75 mL g⁻¹.
10. It is found to penetrate rapidly body fluids, cavities and cells.
11. It is invariably eliminated upto 65% through hepatic metabolism, and the balance 35% through biliary/renal excretion.

12. The rate of exertion is rather slow and the half-life is the longest among the 'tetracyclines', namely, 12 to 22 hr.

Note: 1. Photosensitization usually takes place more frequently as compared to other shorter-acting tetracyclines.

2. Complexation with Ca^{2+} is to a lesser extent than other tetracyclines; besides, it is not affected by either dairy products or foods.

9.3.9.7 Minocycline

Synonyms Minocyn.

Biological Source It is a semi-synthetic antibiotic obtained from 6-demethyl tetracycline.

Preparation 6-Demethyl tetracycline is first dissolved in tetrahydrofuran (solvent) containing aliquot quantity of methanesulphonic acid, and is subsequently reacted with dibenzyl azodicarboxylate to form 7-[1, 2-bis (carbobenzoxy) hydrazino]-6-demethyl-tetracycline. The resulting-product is subjected to Pd-catalyzed hydrogenation in the presence of formaldehyde to yield the desired product minocycline.

Chemical Structure Refer to section 9.3.9.

Characteristic Features

1. It is obtained as bright yellow-orange amorphous solid.
2. Its specific optical rotation $[\alpha]_D^{25} -116^\circ$ ($C = 0.524$).
3. It has uv_{max} (0.1N HCl): 352, 263nm ($\log \epsilon$ 4.16, 4.23); (01 N NaOH) : 380, 243 nm ($\log \epsilon$ 4.30; 4.38).

Uses

1. It is readily absorbed from the intestinal tract.
2. It has a slow renal clearance to afford prolonged blood levels; and is normally characterized by relatively lower MICs as compared to other tetracycline antibiotics for certain pathogenic organisms.

Minocycline Hydrochloride [$\text{C}_{23}\text{H}_{27}\text{N}_2\text{O}_7 \cdot \text{HCl}$] [*Synonyms* *Klinomycin* *Minomycin*; *Vettrin*]

Characteristic Features

1. It is obtained as yellow, crystalline powder, odourless, slightly bitter taste and slightly hygroscopic in nature.
2. It is fairly stable in air when protected from light and moisture; however, strong uv-light and/or moist air causes it to darken rather rapidly.
3. Its **potency*** in solution is primarily affected on account of *epimerization*.
4. The pH of 1 in 100 solution ranges between 3.5 to 4.5.
5. It distinctly gives rise to *four* dissociation constant values, namely: $\text{pK}_{\text{a}1}$ 2.8; $\text{pK}_{\text{a}2}$ 5; $\text{pK}_{\text{a}3}$ 7.8; and $\text{pK}_{\text{a}4}$ 9.3; mainly due to an additional dimethylamino moiety at e-7 position (compare with demeclocycline, Section 3.9.4).

* **Potency:** It is equivalent to not less than 785 mcg of minocycline mg^{-1} .

6. **Solubility Profile:** 1 g in nearly 60 mL water and ~ 70 mL alcohol; soluble in solutions of alkali hydroxides or carbonates; and almost insoluble in chloroform and ether.

Uses

1. Generally, it is found to be 2-4 times as potent as tetracycline against majority of **Gram-positive bacteria**.
2. It is found to exhibit an equally low-potency against *Enterococcus faecalis*.
3. It is almost 8 times as potent as **tetracycline** against *Streptococcus viridans*.
4. It is 2 to 4 times as potent as **tetracycline** against Gram-negative organisms.
5. It is now the drug of choice for treating infections caused by *Mycobacterium marinum*.

Note: It particularly differs from other tetracyclines wherein the bacterial resistance to the drug is not only of low incidence but also of a lower order; which is especially true to staphylococci, in which cross-resistance is observed to be as low as 4%.

6. It is absorbed by the oral route to the extent of 90-100%.
7. Diminution in absorption is caused exclusively by food and milk and substantially by iron preparations and nonsystemic antacids.
8. It is normally protein-bound in plasma between 70-75%.
9. Its '**volume of distribution**'* ranges between 0.14 to 0.7 mLg⁻¹.
10. Its half-life varies between 11 to 17 hours.

9.3.9.8 Lymecycline

Synonyms Armyl; Ciclolysol; Mucomycin; Tetralisal; Tetramyl; Tetralysal; N-Lysinomethyl tetracycline.

Biological Source It is a **semi-synthetic antibiotic** related to **tetracycline**. It is a classic example of an antibiotic developed by qualified chemical modification of the primary amide function at C-2.

Preparation It may be prepared by the method suggested by Tubaro and Raffaldoni.**

Chemical Structure Refer to Section 9.3.9.

Characteristic Features

Lymecycline Sodium [C₂₉H₃₇N₄NaO₁₀]: It has uv_{max} (CH₃ OH): 376 nm. It is used as a potent antibacterial agent.

9.3.9.9 Biosynthesis of Chlortetracycline

The various steps involved in the **biosynthesis of chlortetracycline** are as stated below:


1. A malonamyl-CoA residue probably caters for as a 'primer'; and eight such malonate entities undergo stepwise condensations with the addition of C₂ units and followed by decarboxylation to yield a linear C₁₉ *polyketide*.
2. Subsequently, the carbonyl-methylene condensations give rise to the **tetracyclic pretetramide nucleus**.

* **Volume of Distribution** It is pharmacokinetic parameter representing a proportionality constant that relates drug concentration in a reference fluid, typically plasma, to the amount of drug distributed throughout the body.

** Tubaro and Raffaldoni, *Bull. Chim. Farm.*, **100**, 9 (1961).

3. Importantly, **methylation** at the *C-6 position* of the pretetramide is normally regarded an initial step in the biosynthesis of most tetracyclines; however, this particular step is usually left out in the creation of the naturally occurring dimethyl tetracyclines.
4. Hydroxylation at the C-4 position followed by dearomatization to produce a **4-keto intermediate** appears to precede 7-chlorination.
5. It is necessary that halogenation should precede introduction of the 4-amino group, which is methylated in a stepwise manner.
6. Terminal reactions in the biosynthetic sequence are carried out in *two* stages: *first*, hydroxylation at C-6 position; and *secondly*, reduction of double bond in ring B.
7. It is, however, interesting to absence that the presence of a 7-halogen substituent evidently blocks 5-hydroxylation.

The various steps involved in the **biosynthesis of chlortetracycline** may be summarized as given below:



Biosynthesis of Chlortetracycline

9.3.10 Miscellaneous Antibiotics

In general, the ‘**antibiotics**’ are such a divergent group of medicinal compounds that usually fall into *Four* major categories, namely:

- (a) Antibiotics based on Mechanism of Action,
- (b) Synthetic Antimicrobial Agents,
- (c) Antifungal Agents, and
- (d) Anticancer Antibiotics.

A few typical and potent antibiotic substances shall be discussed under the so called ‘**miscellaneous antibiotics**’ because of the simple reason that these compounds could not be accommodated under various Sections from 9.3.1 through 9.3.9 discussed earlier in this chapter on ‘**Antibiotics**’.

9.3.10.1 Antibiotics Based on Mechanism of Action

An intensive and elaborated research findings, with regard to the various established mechanism of action of certain antibiotics, may further sub-divide this group into the following heads, namely:

- (i) Disruption of DNA-metabolism,
- (ii) Inhibition of Protein Synthesis,
- (iii) Inhibition of Cell wall formation, and
- (iv) Alteration in Cellular Membrane function.

These *four* sub-groups shall now be discussed separately with the help of *one important example* in the sections that follows:

9.3.10.1.1 Disruption of DNA-Metabolism The antibiotics belong to this specific category help to disrupt the DNA metabolism of the pathogenic organisms thereby inhibiting their growth effectively.

A. Rifampin


Synonyms Rifampicin; Rifaldazine; Rifamycin; Abrifam; Eremfat; Rifa; Rifadin(e); Rifaldin; Rifabrodin; Rifoldin; Rimactan(e); R/AMP.

Biological Sources The rifamycins are '**ansamycin antibiotics**'* produced by cultures of *Amycolatopsis mediterranei* (previously known as *Nocardia mediterranei* or *Streptomyces mediterranei*). The crude antibiotic mixture was found to consist of *five* closely related substances *i.e.*, **rifamycins A-E**.

The most clinically useful **rifamycin** is **rifampin**, a semi-synthetic derivative produced from **rifamycin SV** *via* a **Maanich reaction** by making use of formaldehyde and N-amino-N' methylpiperazine.

Preparation **Rifamycin SV**, which may be prepared by the method of Maggi *et al.*,** is converted to the 8-carboxaldehyde derivative, known also as **3-formylrifamycin SV**. It is finally condensed with 1-amino-4-methylpiperazine to give rise to a Schiff base, which is **Rifampin**.

Chemical Structure



* **Ansamycin Antibiotics (or Ansamycins)** These are a class of macrocyclic compounds wherein the non-adjacent positions on an aromatic ring system are usually spanned by the long aliphatic bridge (**Latin**: ansa = handle). The aromatic portion may comprise of either a *substituted benzene* ring or a *substituted naphthalene* or *naphthaquinone* moiety. The macrocycle present in the **ansamycins** is normally closed by an *amide* rather an *ester* linkage, *i.e.*, ansamycins are '**Lactams**'.

** Maggi *et al. Chemotherapy*, **11**, 285, (1966).

3-[[[(4-Methyl-1-piperazinyl) imino]-methyl]rifamycin; [C₄₃H₅₈N₄O₁₂].

Characteristic Features

1. It is obtained as red to orange platelets from acetone that decompose at 183-188°C.
2. It has absorption max. (pH 7.38): 237, 255, 334, 475 nm (ϵ 33200, 32100, 27000, 15400).
3. **Rifampin** has two values for its dissociation constant, namely:
 - (a) pK_{a1} 1.7 related to the 4-hydroxy function, and
 - (b) pK_{a2} 7.9 related to the 3-piperazine nitrogen.

Rifampin behaves as a 'zwitterion' *i.e.*, it behaves both as an acid and as a base.
4. It is odourless; and fairly unstable in light, heat, air and moisture.
5. It is found to be very stable in DMSO; rather stable in water; freely soluble in CH₂Cl₂, DMSO; soluble in ethyl acetate, methanol, tetrahydrofuran; and slightly soluble in water (pH < 6), acetone and carbon tetrachloride.

Uses

1. It acts as a broad-spectrum antibiotic effective against most **Gram-positive organism**, especially *Staph pyogenes*, *Strep pyogenes*, *viridans* and *pneumoniae*.
2. It shows variable activity against **Gram-negative organisms**, particularly *H.influenzae*, *meningococci*, and *gonococci*.
3. Importantly, both *Mycobacterium tuberculosis* and *Mycobacterium leprae* are very susceptible to **rifampin**.
4. Its clinical usage is mainly in the treatment of tuberculosis. However, it is invariably employed in combination with other antitubercular drugs, such as: isoniazid and PAS.
5. It is considered to be an *excellent drug* for prophylaxis of *Meningococcal meningitis* and *pneumonia* caused due to *H.influenzae* Type B; and also in the treatment of *meningococcal* carrier state.
6. It is remarkably absorbed 100% after oral administration; however, the food in the stomach seems to delay absorption of the drug.
7. In plasma it is protein-bound upto 98%.
8. The volume of distribution is 0.9 mL. g⁻¹.
9. Almost 85% of the drug gets eliminated through biotransformation in the liver.
10. One of its '**active metabolite**' gets secreted directly into the bile, where it is therapeutically effective.

Note: The most serious untoward adverse reactions involve hepatotoxicity, and the apparent enhanced risk of toxicity in persons having incidence of liver-damage e.g., chronic alcoholics suffering from cirrhosis*, may obviously preclude the use of this antibiotic.

9.3.10.1.2 Inhibition of Protein Synthesis There are certain antibiotics that specifically retard the anabolism of proteins *i.e.*, protein synthesis, at the ribosome level. For instance, **chloramphenicol** that binds preferentially to the 50S subunit of microbial 70S ribosomes and disrupts **peptidyl transferase** which catalyzes peptide bond formation exclusively. It is also found to specifically inhibit mitochondrial 70S ribosomes that ultimately results in the dose-related bone marrow suppression.

* A chronic disease of the liver marked by formation of dense perilobular connective tissue, degenerative changes in the parenchymal cells, structural alternation of the cords of liver lobules, fatty and cellular infiltration.

A. Chloramphenicol

Synonyms Ak-Chlor; Amphicol; Anacetin; Aquamycetin; Chlovamex; Chloromycetin; Chloramfen; Intramycetin; Leukomycin; Novomycetin; Pantovernil; Ronphenil; Synthomycetin; Tifomycine; Veticol.

Biological Sources It is a broad-spectrum antibiotic originally obtained by Burkholder from a culture of *Streptomyces venezuelae* that was initially isolated in 1947 from soil samples collected in the vicinity of Caracas (Venezuela). As the organism incidentally was not reported previously, Burkholder thought it proper to baptize it as *venezuelae*. **Chloramphenicol** was also isolated from the moon snail, *Lunatia heros*.

Chloramphenicol attracted surmountable fame and glory for being the first truly discovered broad-spectrum antibiotic having its span from Gram-negative, Gram-positive organisms, a variety of rickettsial pathogens to a few-specific viruses.

Preparation **Chloramphenicol** enjoys the reputation of being the first even naturally occurring antibiotic known to contain a **nitro function**, besides being a derivative of dehydroacetic acid. Nevertheless, its stereochemical configuration is very much identical to that of **(-)-norpseudoephedrine**; and is the only one of the four related stereoisomers that essentially possess the antibiotic activity stated earlier.

It may be obtained either from the natural source or by the synthetic route as given under:

- (a) **Natural Source:** It may be obtained from the filtrate of a *Streptomyces venezuelae* culture by extraction with ethyl acetate. In case, the charcoal extract is rich in **chloramphenicol**, the latter may be crystallized from the ethyl acetate by affecting dilution with several volumes of deodorized kerosene oil.
- (b) **Synthetic Route:** **Chloramphenicol** may be synthesized by many different routes of preparation, but one of the better known starts with *para*-nitroacetophenone and, after due conversion it into *para*-nitro-2-amino-acetophenone, proceeds through the following steps, namely:

Step I: Acetylation of the primary $-NH_2$ moiety,


Step II: Interaction with formaldehyde (HCHO) to induct the terminal primary $-CH_2OH$ function,

Step III: Reduction with **aluminium isopropoxide** $Al [OCH (CH_3)_2]_3$ to yield a mixture of the racemates of the *threo* and *erythro* forms of *para*- NO_2 Ph $CH(OH) CH(NH_2) CH_2OH$,

Step IV: Isolation of the *threo* racemate and its resolution using **camphorsulphonic acid**, and

Step V: Condensation of the (-)enantiomorph with methyl dichloroacetate.

Chemical Structure



Chloramphenicol

[R-(R*, R*)]-2, 2-Dichloro-N-[2-hydroxy-1-(hydroxymethyl)-2-(4-nitrophenyl) ethyl]acetamide; [C₁H₁₂Cl₂N₂O₅].

Characteristic Features

1. It is obtained either as needles or as elongated plates from water or ethylene dichloride having mp 150.5-151.5°C.
2. It usually sublimes under in the vacuum.
3. It appears as white to grayish white or yellowish white crystals.
4. It is practically odourless and possesses an intense bitter taste.
5. It has specific optical rotation $[\alpha]_D^{27} + 18.6^\circ$ (C = 4.86 in ethanol); $[\alpha]_D^{25} - 25.5^\circ$ (ethyl acetate).
6. The pH of a saturated solution ranges between 4.5 and 7.5.
7. It is found to be reasonably stable either in neutral or moderately acidic medium; but rapidly gets destroyed in an alkaline medium.
8. It has dissociation constant pKa 5.5.
9. **Solubility Profile:** It is found to be very soluble in methanol ethanol, butanol, ethyl acetate, acetone; fairly soluble in ether; insoluble in benzene, petroleum ether, vegetable oils; and soluble in 50% (w/v) acetamide solution to an extent of 5%.

Uses

1. The drug is found to be effective in **Rickettsial diseases*** including epidemic, murine and scrub typhus, Rocky Mountain spotted fever, rickettsial pox and **Q fever****.
2. It is also effective in chlamydial diseases including the **psittacosis*****-**lymphogranuloma****** group.
3. **Chloramphenicol** is useful in a good number of Gram-negative and Gram-positive bacterial infections including the anaerobes (especially *Bacteroides fragilis*).
4. It is extensively used topically for superficial conjunctival **infections******* and **blepharitis******* caused by *E. coli*, *H. influenzae*, *Moraxella Lacunata*, *Staphylococcus aureus*, and *Streptococcus hemolyticus*.
5. **Chloramphenicol** is very well absorbed through the oral route and crosses rapidly into the cerebrospinal fluid.
6. It is absorbed readily from the GI tract, having a bioavailability of 90%.
7. The drug in blood is bound to serum albumin to the extent of 60%.

* **Rickettsial Disease** A disease caused by an organism of the genus *Rickettsia*. The most common types are: spotted-fever group (Rocky Mountain spotted fever and rickettsial PoX), epidemic typhus, endemic typhus.

** **Q Fever** It is caused by the rickettsial organism, *Coxiella burnetii* and is contracted by inhaling infected dusts, drinking unpasteurized milk from infected animals.

*** **Psittacosis** An infections disease caused by *Chlamydia psittacoci* of parrots and other birds that may be transmitted in humans.

**** **Lymphogranuloma** Infections granuloma of the lymphatics, Hodgkin's disease.

***** **Conjunctival Infections** Infections in the mucous membrane that lines the eyelids and is reflected into the eyeball.

***** **Blepharitis** Ulcerative or nonulcerative inflammation of the hair follicles and glands along the edges of the eyelids.

8. Its volume of distribution stands at 0.7 mL g^{-1} .
9. Biotransformation in liver ranges between 85-95%.
10. Its half-life is normally between 1.5 to 5 hours; except over 24 hours in neonates 1 to 2 day old; and 10 hours in infants 10 to 16 days old.

Note: 1. The plasma levels should be monitored very closely due to the significant variability with half-life.

2. Bone-marrow injury is the major toxic effect.

3. It causes serious *hematopoietic disturbances, such as: *thrombocytopenia*** , *granulocytopenia**** and *aplastic anemia*.******

Chloramphenicol variants are as follows, namely:

(a) Chloramphenicol Palmitate [Synonyms Chlorambon; Chloropal; Chlorolifarina]:

1. It is obtained as crystals from benzene having mp 90°C .
2. It is practically tasteless and hence may be used as suspensions in pediatric formulations.
3. It has specific optical rotation $[\alpha]_{\text{D}}^{26} + 24.6^{\circ}$ ($C = 5$ in ethanol).
4. It has uv_{max} (ethanol): 27 nm ($E_{1\text{cm}}^{1\%} 179$).
5. It is very slightly soluble in water (1.05 mg mL^{-1} at 28°C); and petroleum ether (0.225 mg mL^{-1})
6. It is found to be freely soluble in methanol, ethanol, chloroform, ether and benzene.


(b) Chloramphenicol Monosuccinate Sodium Salt [$\text{C}_{15}\text{H}_{15}\text{Cl}_2\text{N}_2\text{NaO}_8$] [Synonyms Protophenicol]: It is freely soluble in water upto 50% (w/w).

Biosynthesis of Chloramphenicol The various steps involved in the biosynthesis of chloramphenicol are as follows:

1. Experimental studies with radioactive precursors have established a **Shikimic acid-phenylpropanoid pathway**.
2. Pathway evidently branches from normal phenylpropanoid metabolism prior to the formation of phenylalanine or tyrosine.
3. *para*-Aminophenylpyruvic acid (I) seems to be an early metabolite in the biosynthetic pathway.
4. Subsequent steps involved are, namely: transamination, hydroxylation, acylation, reduction of carboxyl moiety and terminal oxidation of the amino function.

The above steps (1) through (4) are summarized below:

* **Hematopoietic** A substance that assists in or stimulates the production of blood cells.
 ** **Thrombocytopenia** An abnormal disease in number of the blood platelets.
 *** **Granulocytopenia** An abnormal reduction of granulocytes in the blood.
 **** **Aplastic Anemia** A serious complication of infection with human parvovirus B-19 infection in patients with chronic hemolytic anemia, such as: sickle-cell disease.



Biosynthesis of Chloramphenicol

9.10.1.3 Inhibition of Cell-Wall Formation

A number of naturally occurring substances have been isolated and characterized that are found to exert their action by inhibiting specifically the cell-wall formation *in vivo*.


Examples

A. Novobiocin

Synonyms Biotexin; Cardelmycin; Cathocin; Cathomycin; Inamycin; Speromycin; Vulcamycin; Vulkamycin; Albamycin; Streptonivicin.

Biological Sources Novobiocin is produced by *Streptomyces spheroides* and *Streptomyces niveus*.

Chemical Structure



The structure of **novobiocin** implies a not so common biosynthetic origin; and it evidently seems to involve various essential groups derived from amino acid, acetate and above all carbohydrate metabolic pathways.

Characteristic Features

1. It is obtained as pale yellow orthorhombic crystals obtained from ethanol that get decomposed at 152-156°C (a rarer modification get decomposed at 174-178°C).
2. **Novobiocin** is sensitive to light.
3. It has a density 1.3448.
4. It is found to be acidic in reaction and has two dissociation constant values, namely: pK_{a1} 4.3; pK_{a2} 9.1.
5. It has specific optical rotation $[\alpha]_D^{24}$ -63.0° (C = 1 in ethanol).
6. It shows uv_{max} (0.1 N NaOH; 0.1N methanolic HCl; Phosphate Buffer pH 7): 307; 324; 390 nm ($E_{1cm}^{1\%}$ 600, 390, 350 respectively).
7. It is found to be soluble in aqueous solution > pH 7.5.
8. **Solubility Profile:** It is almost insoluble in more acidic solutions; soluble in acetone, ethyl acetate, amyl acetate, lower alcohols, pyridine.

Novobiocin Monosodium Salt $[C_{31}H_{35}N_2NaO_{11}]$ [Synonyms: Robiocina]

1. It is obtained as minute crystals that get decomposed at 220°C.
2. Its specific optical rotation $[\alpha]_D^{24}$ -38° (C = 2.5 in 95% ethanol); and $[\alpha]_D^{24}$ -33° (C = 2.5 in water).
3. The pH of a 100 mg mL^{-1} solution is 7.5.
4. It is found to be freely soluble in water.
5. It has a half-life of approximately 30 days at 25°C and of several months at 4°C.

Uses

1. The activity profile of **novobiocin** is predominantly Gram-positive.
2. It has been observed that it is unusually sensitive to *Staphylococci* (MIC* ranges between 0.1 to 2.0 mcg. mL^{-1}); however, resistance gets developed rather readily.
3. It finds its therapeutic application as an '**alternate drug**' for controlling **penicillin-resistant staphylococci**.
4. It also exerts antiviral activity.
5. It exhibits efficacy in canine respiratory infections.

Note: Based on its relatively high incidence of adverse reactions, such as: blood dyscrasia**, hepatic dysfunction and hypersensitivity, the therapeutic usage is no longer justified.

9.3.10.1.4 Alteration in Cellular Membrane Function There are a few antibiotic substances which essentially cause alteration in the cellular membrane function of various organisms.

A. Candicidin


Synonyms Levorin; Candivon; Vanobid.

* MIC = Minimum Inhibitory Concentration;

** **Dyscrasia** An old term meaning abnormal mixture of the four humors.

Biological Source It is a **heptane macrolide antifungal antibiotic complex** composed of candidicidins A, B, C and D (major component). It is produced by a strain of *Streptomyces griseus*.

Chemical Structure



Characteristic Features

1. It is obtained as small yellow needles or rosettes from aqueous tetrahydrofuran or a mixture of pyridine/acetic acid/water.
2. It has uv_{max} : 403, 380 ($E_{1cm}^{1\%}$ 1150), 360 nm.
3. **Solubility Profile:** It is almost insoluble in water, alcohols, ketones, esters, ethers, hydrocarbons and other lipophilic solvents; soluble in DMSO, DMF, and lower aliphatic acids (formic acid, acetic acid); very soluble in 80% aqueous tetrahydrofuran solution; and addition of 5-25% of water to alcohols greatly enhances its solubility.
4. It readily forms soluble salts in alkaline solutions.

Candidicin D [$C_{59}H_{84}N_2O_{18}$] [**Levorin A₂**]: It is found to be identical with **levorin A₂**.*

Uses

1. It is mostly used as a topical antifungal agent.
2. It is also found to exert anticholesteremic activity.


9.3.10.1.5 Synthetic Antimicrobial Agents There are a host of purely synthesized tailor-made medicinal compounds that have been used as antimicrobial agents in the ever increasing therapeutic armamentarium to combat human diseases across the globe. The various synthetic antimicrobial agents may be categorized as stated below:

- (i) Sulphonamides and Trimethoprim
- (ii) Nitroimidazoles
- (iii) Quinolones and Fluoroquinolones
- (iv) Agents for systemic UTIs
- (v) Agents for *Mycobacterium tuberculosis* Infections.
- (vi) Antifungal agents, and
- (vii) Anticancer antibiotics.


The aforesaid categories of synthetic antimicrobial agents shall now be treated individually with some typical potent examples in the sections that follows:

* Bosshardt, Bickel, *Experientia*, **24**, 422, (1968).

9.3.10.1.5.1 Sulphonamides and Trimethoprim **Sulphonamides**—the first and foremost antimicrobial agents, since discovered in 1930s, still hold the glory and fame of the modern antibiotic era. In general, **sulfanilamide** (*i.e.*, *para*-aminobenzene sulphonamide), obtained as a structural analogue of ***para*-aminobenzoic acid (PABA)**, is basically the core compound from which hundreds of congeners were synthesized over the years by suitable modifications at N₁ (amide) or N₄ (*p*-amino function) so as to alter the pharmacological characteristics of the parent compound (sulphanilamide). The following table summarizes some of the approved and clinically useful widespread sulphonamides (see page 667).


S. No.	Classification	Drug(s)	R1	R2	Brand Name	Therapeutic Uses
I	Sulphonamides for general Infections	1. Sulphanilamide	H	H	Rhinamid	Obsolete
		2. Sulphapyridine		H	M2B 693	Pneumonia; <i>Dermatitis herpetiformis</i> .
		3. Sulphathiazole		H	Cibazol	Bubonic plague staph. infections
		4. Sulphadiazine		H	Diazyl	Rheumatic fever; Chancroid due to <i>Haemophilus ducreyi</i> .
		5. Sulphamerazine		H	Solumedine	General infection
		6. Sulphadimidine (sulphamethazine)		H	Pirmazin	Meningal infections
II	Sulphonamides for Urinary Infections	1. Sulphacetamide		H	Albucid	UTIs; topical for eye and skin infections.
		2. Sulphafurazole		H	Sulfalar	UTIs; topically for vaginitis.
		3. Sulfacytine		H	Renoquid	Acute UTIs.
III	Sulphonamides for Intestinal Infections	1. Sulphaguanidine		H	Shigatox	Bacillary dysentery;
		2. Phthalylsulphathiazole		H	Thalazole	Bacteriostatic effect in GIT.
		3. Succinylsulphathiazole			Sulfauxidine	Cholera; Bacillary dysentery.
IV	Sulphonamides for Local Infection	Mafenide	H	X = CH ₂ NH ₂	Sulfonyl	<i>Pseudomonas aeruginosa</i> ; severe burns.

[Adapted From: Kar, Ashutosh, 'Medicinal Chemistry', 4th ed 2006, New Age International Publishers, New Delhi.]



Sulphonamide


Sulphonamides are regarded as '**antimetabolites**' which are found to be the structural analogues of PABA; the latter being a substrate in the biosynthesis of folic acid and also an absolutely necessary metabolite for microorganism which in turn utilizes it as a vital source of *single-carbon-units* required for the biosynthesis of *amino acids*, *purines* and *pyrimidines*.

p-Amino
Benzoic acid
(PABA)

Sulphanilamide

[Sulphonamides Structural
Analogues of (PABA)]



It has been observed that **sulphonamides** inhibit competitively the enzyme **tetrahydropteroyl synthetase** that makes use of PABA as a *substrate* ultimately in generating the '*pteroyl moiety*' in the **Folic Acid** as given below:



Pteroyl Moiety


Folic Acid

Sulphonamides are of *two* types, namely: *unionized form* e.g., **sulphamethoxazole**; and *ionized form* e.g., **sodium sulfisoxazole**, as shown below:

Sulphamethoxazole
[Unionized Form]Sodium Sulfisoxazole
[Ionized Form]

Sulphamethoxazole enjoys the status of a **medium-acting sulphonamide** which is comparatively less soluble than **sulfisoxazole**; and, therefore, accomplishes higher blood-levels. Interestingly, **sulfisoxazole** is a **short-acting sulphonamide** predominantly useful for UTIs caused by susceptible pathogenic bacteria.

The unique combination of **trimethoprim (TMP) and sulphamethoxazole (SMZ)*** affords a broad-spectrum '*antibiotic product*' which is found to be active (invariably bactericidal) against a plethora of Gram-positive *cocci* and Gram-negative *rods*. However, certain microorganisms are evidently resistant to this combination, namely: *Mycobacterium tuberculosis* and species of *Pseudomonas*, *Mycoplasma* and *Bacteroides*.




Trimethoprim

The therapeutic advantages of this combination are as follows:

1. It particularly inhibits the sequential steps in the formation of tetrahydrofolic acid. The resulting inhibition is magnified by the independent actions at two consecutive metabolic steps; and thus the **bacteriostasis** phenomenon** gets converted into the **bactericidal***** one. The double blockade broadens appreciably the antibacterial spectrum from that of each drug used alone.
2. The most significant usage of **TMP-SMZ** combination is in the control, management and treatment of UTIs, specifically recurrent, chronic or complicated infections not easily manaeuvrable and controllable by individual drugs; such as: UTIs caused by *E. coli*, *Klebsiella-Enterobacter* and *Proteus species*.
3. TMP-SMZ combination provides the treatment and serves as *prophylaxis of choice* particularly in immunocompromised patients for *pneumonia* caused by *Pneumocystis carinii* and *enterocolitis* caused by *Isospora*.
4. By virtue of the fact that **sulphamethoxazole**, exhibits poor tissue distribution, and also the pharmacokinetics of **TMP-SMZ** mixture is not optimal for the treatment of systemic infections.

9.3.10.1.5.2 Nitroimidazoles: **Nitroimidazole**, is a 5-membered heterocyclic nucleus having two N-atoms at positions 1 and 2, with the nitro group at position 5. Several structural variants of nitroimidazole have been duly synthesized and screened for their antibacterial activities.



5-Nitroimidazole

* **Trimethoprim-Sulfamethoxazole (Co-trimoxazole, TMP-SMZ, Bactrium, Spectra).** This combination is generally available in a fixed ratio of 1:5 for IV and oral administration.

** **Bacteriostasis:** The arrest of bacterial growth.

*** **Bactericidal:** Destructive to or destroying bacteria.


Two such potent compounds, namely: **Metronidazole** and **Tinidazole**, shall now be discussed as under.

A. Metronidazole

Synonyms Bayer 5630; RP-8823; Arilin; Clont; Deflamon; Elyzol; Flagyl; Fossypol; Klion; MetroGel; Metrolag; Metrolyl; Orvagil; Rathimed; Trichazol; Tricocet; Vagilen; Zadstat.

Preparation **Metronidazole** is prepared by the condensation of 2-methyl-5-nitroimidazole with ethylene chlorohydrin by subjecting it to heating in the presence of a large excess of the **chlorohydrin**. The surplus chlorohydrin is removed, the residue is extracted with water and the resulting extract is alkalized and extracted with chloroform. Chloroform is removed under vacuo to obtain crude **metronidazole** which is finally recrystallized from ethyl acetate.

Chemical Structure



Metronidazole

2-Methyl-5-nitroimidazole-1-ethanol; (C₆H₉N₃O₃).

Characteristic Features

1. It is obtained as cream-coloured crystals having mp 158-160°C, odourless, stable in air and darkens on exposure to light.
2. Its solubility profile at 20°C (g/100 ml): water 1.0; ethanol 0.5; ether < 0.05; chloroform < 0.05; and sparingly soluble in DMF.
3. The pH of a saturated aqueous solution is 5.8.
4. It has dissociation constant pK_a 2.62.

Uses

1. It is found to be bactericidal to **anaerobic** and **microaerophilic*** microorganisms, including: *Bacteroides*; *Clostridium sp.*; *Endolimax nana*; *Entameba histolytica*; *Fusobacterium vincentii*; *Gardnerella vaginalis*; *Giardia lamblia*; *Peptococcus*; *Peptostreptococcus*; and *Trichomonas vaginalis*.

Interestingly, all these organisms critically reduce the nitro moiety, and subsequently produce the metabolites that specifically inhibit DNA synthesis.

2. It is the drug of choice since long for the treatment of **trichomoniasis**** , and more recently in combination with **iodoquinol** for the treatment of symptomatic amebiasis (except in brain).
3. As it is absorbed so effectively from the oral route, hence its adequate concentrations in the

* **Microaerophilic:** Growing at low amounts of oxygen.

** **Trichomoniasis** Infestation with a parasite of the genus *Trichomonas i.e.*, Genus of flagellate parasitic protozoa e.g., *T. hominis*—a benign trichomonas found in the large intestine; *T. tenax*—a benign trichomonas that may be present in the mouth; and *T. vaginalis*—a species found in the vagina that produces discharge.

lower bowel invariably are not sufficient enough to cause eradication of amebas, so that it is combined with iodoquinol to render it a first-choice combination.


4. It is also recommended as the drug of first-choice for the treatment of *Dracunculus* (guinea worm) infestations.
5. It has been employed successfully in the treatment of antibiotic-associated **pseudomembranous colitis*** either through IV or orally.
6. It has been reported to be of value in **Crohn's disease** *i.e.*, inflammatory bowel.
7. Its half life is about 6-12 hour.

Note: Metronidazole may reasonably be anticipated to be a carcinogen. (Ref: Seventh Annual Report on Carcinogens (PB95-109781, 1994) p 257.

B. Tinidazole

Synonyms Fasigin; Fasigyn; Pletil; Simplotan; Sorquetan; Tricolam; Trimonase.

Chemical Structure



Tinidazole

1-[2-(Ethylsulfonyl) ethyl]-2-methyl-5-nitro-1H-imidazole; (C₈H₁₃N₃O₄S).

Characteristic Feature It is obtained as colourless crystals from benzene having mp 127–128°C.

Uses


1. It is used as an effective antiprotozoal agent for *Trichomonas* and *Giardia*.
2. It is also employed widely for the treatment of amebic dysentery.
3. It is broadly used as an antibacterial agent.
4. A combination with **norfloxacin** [**Nortec**, -Z^(R)] (German Remedies Ltd.) containing **norfloxacin** 400 mg + **tinidazole** 600 mg] is invariably employed for the treatment of serious types of dysentery caused due to food poisoning, loose motion etc.

9.3.10.1.5.3 Quinolones and Fluoroquinolones **Quinolone antibacterial drugs** have gained entry into the therapeutic armamentarium since 1964 with the evolution of **nalidixic acid**-a purely synthetic molecule, used exclusively for UTIs. Within a very short span (1968-1970) *two* more structurally related analogues, namely: **oxolinic acid** and **cinnoxacin** were introduced. However, all these drugs has shown a serious disadvantage by virtue of two major short-comings: (a) exhibited limited antibacterial spectra; and (b) rapid development of resistance to these bacteria.


In the usual constant, routine, untiring, sincere efforts towards designing newer and safer drug molecules an attempt was made whereby *two* new chemical entities were introduced into the parent

* Inflammation associated with antibiotic therapy. It is caused due to a toxin produced by *Clostridium difficile* and is marked by formation of a pseudomembrane on the mucosa of the colon; symptoms diarrhoea with gross blood and mucus, abdominal cramps, fever and leukocytosis.


structure, namely: (a) 6-fluoro; and (b) 7-(1-piperazinyl). These structural modifications properly expanded not only the spectrum but also enhanced the potency besides preventing the development of plasmid-mediated resistance. This significantly made the wonderful discovery of the so called '**fluoroquinolones**' that showed **bacteriostatic** action *at low concentrations* and **bactericidal** at **high concentrations**.



Nalidixic Acid



Oxolinic Acid



Cinoxacin

It is, however, pertinent to state here that the mechanism of action of **fluoroquinolones** predominantly involves inhibition of DNA gyrase, a member of the topoisomerase group of enzymes that essentially regulate the superhelicity of DNA within the cells. Evidently, inhibition of DNA gyrase leads to inhibition of DNA replication phenomenon, and ultimately cell death is accomplished. The latest marketed fluoroquinolones exert merely minimal activity against the mammalian topoisomerases. Another important feature with regard to the apparently enhanced activity of the **fluoroquinolones** *vis-a-vis* **nalidixic acid** is solely due to the much greater affinity of the former for getting bound to the DNA gyrase. Interestingly, clinical studies have revealed that certain extent of acquired resistance has been observed not so commonly with the newer designed fluoroquinolones and another most vital and important revelation is that till date absolutely not even a single incidence of **plasmid-mediated resistance** has been reported in any clinical isolates.

Thus, **fluoroquinolone** based synthetic compounds occupy a very strategically-useful status in combatting dreadful microbial infections in human beings. A few potent compounds belonging to this category are, namely: **ciprofloxacin**, **lomefloxacin**, **norfloxacin**, **ofloxacin**, **sparfloxacin** and **trovafloxacin**, which shall now be elaborated squarely in the sections that follows:


A. Ciprofloxacin

Synonyms Bay q 3939.

Preparation It is a **fluorinated quinolone antibiotic** prepared by the condensation of 3-chloro-4-fluoroaniline with diethyl ethoxymethylenemalonate to yield the *imine* that is thermally cyclized to form ethyl 7-chloro-6-fluoro-4-hydroxy quinoline-3-carboxylate. Subsequent N-alkylation with cyclopropyl iodide followed by nucleophilic displacement of the 7-chloro function by N-methyl piperazine; and finally the hydrolysis of the ester affords the desired product.*

* *J. Med. Chem.*, **19**, 1138, 1976.

Chemical Structure



Ciprofloxacin

1-Cyclopropyl-6-fluoro-1, 4-dihydro-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid; (C₁₇H₁₈FN₃O₃).

Characteristic Features

1. It is mostly obtained as pale-yellow crystals that get decomposed at 255-257°C.
2. It is found to be amphoteric in nature.
3. Its dissociation constant pKa is 6 and 8.8.
4. 1g of it is soluble in 25 ml of water.

Ciprofloxacin Monohydrochloride Monohydrate [C₁₇H₁₈FN₃O₃.HCl.H₂O] [Synonyms Baycip; Ciflox; Ciloxan; Ciprinol; Cipro; Ciprobay; Ciproxan; Ciproxin; Flociprin; Septicid; Velmonit]: It is obtained as light-yellow crystalline powder having mp 318-320°C.

Uses


1. It is recommended for use in the treatment of bone and joint infections, infectious diarrhoea caused by *Shigella* or *Compylobacter*), lower respiratory tract infections, skin infections and UTIs.
2. It is found to be the drug of choice for the treatment of infections caused by **Compylobacter jejuni***
3. Besides, it is an unlabeled but authoritatively alternative drug for the treatment of gonorrhoea, salmonella, and **yersinia**** infections.
4. The oral bioavailability is about 70-80%.
5. Its half life is about 4 hour.
6. Ciprofloxacin has improved activity against Gram-positive bacteria, particularly certain *Staphylococci* and *Streptococci* species.

B. Lomefloxacin

Preparations The active 2-chloro group of 2, 6-dichloro-3-nitropyridine is nucleophilically displaced by 3-methyl-N-carbetoxypiperazine; then the 6-chloro is displaced with ammonia, and the resulting amine is acylated to the acetamide. The nitro group is reduced, diazotized and subsequently treated with HBF₄ to yield the fluoro derivative. The balance of the synthesis is analogous to that for **ciprofloxacin**.

* **Compylobacter jejuni** A subspecies of *C. fetus* formerly called *Vibrio fetus*. It can cause an acute enteric disease characterized by diarrhea, abdominal pain, fever, nausea, and vomiting.

** **Yersinia** A genus of **Gram-negative organism**.

Chemical Structure

Lomefloxacin

1-Ethyl-6, 8-difluoro-1, 4-dihydro-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinoline carboxylic acid; (C₁₇H₁₉F₂N₃O₃).

Characteristic Features It is obtained as colourless needles from ethanol having mp 239-240.5°C.

Lomefloxacin Monohydrochloride [C₁₇H₁₉F₂N₃O₃.HCl] [Synonyms NY-198; Bareon; Chimono; Lomebact; Maxaquin; Uniquin]: It is obtained as colourless needles having mp 295°C with decomposition and is found to be soluble in water.


Uses

1. It is invariably recommended only for treatment of UTIs and bronchitis caused by *H. influenzae* or *Branhamella catarrhalis*.
2. It is found to cover Gram-negative bacteria frequently responsible for UTIs but does not necessarily possess the activity to cover the same bacterial infections that respond to either **ciprofloxacin** or **ofloxacin**.

C. Norfloxacin

Synonyms Baccidal; Barazan; Chibroxin(e); Chibroxol; Floxacin; Fulgram; Gonorcin; Lexinor; Noflo; Nolicin; Noracin; Noraxin; Norocin; Noroxin(e); Norxacin; Sebercim; Uroxacin; Utinor; Zoroxin;

Preparation Similar to **Ciprofloxacin**.*

Chemical Structure

Norfloxacin

1-Ethyl-6-fluoro-1, 4-dihydro-4-oxo-7-(1-piperazinyl)-3-quinoline-carboxylic acid; (C₁₆H₁₈FN₃O₃).

Characteristic Features

1. It is obtained as white to light-yellow crystalline powder having mp 220-221°C.

* H. Koga *et al.*, *J. Med. Chem.*, **23**, 1358, (1980).

- It has uv_{\max} (0.1N NaOH): $\sim 274, 325, 336 \text{ nm}$ ($A_{1\text{cm}}^{1\%} \sim 1109, 437, 425$).
- It has *two* distinct values of dissociation constant: $pK_{a1} 6.34$; $pK_{a2} 8.75$.
- It shows partition coefficient (octanol/water): 0.46.
- It is hygroscopic in air and forms a hemihydrate.
- Its solubility in water is pH dependent, increasing sharply at $pH < 5$ or at $pH > 10$.
- Solubility Profile:** Solubility at 25°C ($\text{mg} \cdot \text{mL}^{-1}$): water 0.28; methanol 0.98; ethanol 1.9; acetone 5.1; chloroform 5.5; diethyl ether 0.01; benzene 0.15; ethyl acetate 0.94; octanol 5.1; glacial acetic acid 340.

Uses


- It is used in the treatment of lower urinary tract infections.
- It is also employed in penicillin-resistant gonorrhoea.
- It is a limited-spectrum fluoroquinolone.
- It has incomplete oral absorption.

D. Ofloxacin

Synonyms **Ofloxacin**; DL-8280; HOE-280; Excin; Flobacin; Floxil; Floxin; Oflocet; Oflocin; Oxaldin; Tarivid; Visiren.

Preparation It is a broad-spectrum **fluorinated quinolone antibacterial**; and may be prepared by a method analogous to that for **Ciprofloxacin**.*

Chemical Structure



Ofloxacin

(\pm)-9-Fluoro-2, 3-dihydro-3-methyl-10(4-methyl-1-piperazinyl)-7-oxo-7H-pyrido [1, 2, 3-de], 4-benzoxazine-6-carboxylic acid; ($\text{C}_{18}\text{H}_{20}\text{FN}_3\text{O}_4$).

The C-atom to which the $-\text{CH}_3$ moiety is attached, in the oxazine ring, is *chiral* and the therapeutically used substance is a racemic mixture; whereas the (+)-form has twice the activity of the (–)-form.

Preparation It may be prepared by a method analogous to that for *Ciprofloxacin*.

Characteristic Features

- It is obtained as colourless needles having mp $250\text{-}257^\circ\text{C}$, with decomposition.
- Its dissociation constant pK_a is 7.9.
- It is found to be poorly soluble in water or ethanol.

* H. Egawa *et al.* *Chem. Pharm. Bull.* **34**, 4098 (1986).

Uses In general, it is found to be an **intermediate-spectrum fluoroquinolone**.


E. Sparfloxacin

Synonyms Spara; Zagam; AT-4140; CI-978; PD-131501.

Biological Sources It is a protein-synthesis inhibitor with antibiotic and antineoplastic activity. It is obtained from the fermentation both of *Streptomyces sparsogenes* var *sparsogenes*.* It may also be obtained from *Streptomyces cuspidosporus***

Preparation US Pat 4,795, 751 (1989); *J. Med. Chem.* **33**, 1645, 1990.

Chemical Structure



Sparfloxacin

(*cis*)-5-Amino-1-cyclopropyl-7-(3, 5-dimethyl-1-piperazinyl)-6, 8-difluoro-1, 4-dihydro-4-oxo-3-quinolinecarboxylic acid; (C₁₉H₂₂F₂N₄O₃).

Characteristic Features

1. It is obtained as crystals from a mixture of chloroform and ethanol having mp 266–269°C with decomposition.
2. It has two distinct values for dissociation constants. pK_{a1} 6.25; pK_{a2} 9.30.
3. **Solubility Profile:** It is sparingly soluble in glacial acetic acid or chloroform; very slightly soluble in ethanol; almost insoluble in water or ether; and soluble in dilute mineral acids or fixed bases (ca 0.1N).

Uses

1. It is a newer **fluoroquinolone** with much improved activity against *Streptococcus pneumoniae*, besides other lower respiratory pathogens covered by **grepafloxacin**.
2. It is found to be more active against *Mycoplasma* than other fluoroquinolones.
3. It exhibits an excellent oral bioavailability profile (92%) and gets metabolized chiefly by **hepatic glucuronidation** rather than **cytochrome P450-mediated pathways**. Evidently, it does not affect the clearance of other drugs (*e.g.*, **cimetidine**, **cyclosporine**, **digoxin**, **theophylline** and **warfarine**) that usually takes place with certain **fluoroquinolones**.
4. Its half-life is 20 hours.

* S.P. Owen *et al.* *Antimicrob Ag. Chemother.* 772 (1962).


** E. Higashide *et al.* *Takeda Kykusho Nempo.* **25**, 1 (1966).

F. Trovafloxacin

Synonyms Trovan; CP-9921g.

Preparation It is also a **fluorinated quinolone antibacterial** and may be prepared as per US Pat 5,164,402 (1992).

Chemical Structure



Trovafloxacin

(1 α , 5 α , 6 α)-7-(6-Amino-3-azabicyclo [3.1.0]hex-3-yl)-1-(2, 4-difluorophenyl)-6-fluoro-1, 4-dihydro-4-oxo-1, 8-naphthyridine-3-carboxylic acid; (C₂₀H₁₅F₃N₄O₃).

Characteristic Features It is usually obtained as white to off white powder.

- (a) **Trovafloxacin Hydrochloride** [C₂₀H₁₅F₃N₄O₃·HCl]: It is mostly obtained as pale-yellow crystals from a mixture of acetonitrile and methanol having mp 246°C (decomposes).
- (b) **Trovafloxacin Monomethanesulfonate** [C₂₀H₁₅F₃N₄O₃·CH₄O₃S] [**Synonyms: Trovafloxacin Mesylate**]

Uses

1. It is an altogether **newer fluoroquinolone** that exhibits a better activity against certain respiratory pathogens as compared to the **older fluoroquinolones**, for instance: **ciprofloxacin**.
2. It is found to be more active against *Streptococcus pneumoniae* (including **penicillin-resistant strains**), *Staphylococcus aureus* (including **methicillin-resistant strains**), *Enterococcus faecalis*.
3. It is observed to be highly active against *chlamydia*, *Mycoplasma*, and *Ureaplasma*; besides, it also covers important anaerobes, for instance: *Bacterioides fragilis*, and the Gram-negative *Enterobacteriaceae*, including *Ps aeruginosa*.
4. **Trovafloxacin** is recommended for a very broad range of infections including oral and IV treatment of nosocomial and community-acquired pneumonia, acute exacerbations of chronic bronchitis, acute sinusitis, complicated intra-abdominal and pelvic infections, diabetic-foot infection, uncomplicated UTIs, prostatitis, cervicitis, and uncomplicated gonorrhoea.
5. It has an excellent oral bioavailability (88%).
6. It has a half-life of 10 hours.
7. It hardly shows any untoward photosensitivity as other **fluoroquinolones**.

9.3.10.1.5.4 Agents for Systemic UTIs There are a few purely synthetic drug molecules which are being used exclusively for various systemic urinary tract infections (UTIs). The *two* typical and


potent drugs belonging to this category are, namely: **methenamine** and **nitrofurantoin**, which shall now be discussed as under:

A. Methenamine

Synonyms Hexamine; Hexamethylamine; Aminoform; Ammoform; Crystamin, Cystogen; Formin; Uritone; Urotropin.

Preparation **Methenamine** may be prepared by the interaction of a moderate excess of ammonia water to formaldehyde solution, and evaporating the mixture to absolute dryness.

Chemical Structure



Methenamine

1, 3, 5, 7-Tetraazatricyclo [3.3.1.-1^{3,7}] decane; (C₆H₁₂N₄).

Characteristic Features

1. It is obtained as colourless, lustrous crystals or a white crystalline powder, or granules almost odourless which sublimes at about 263°C without melting and with partial decomposition.
2. It is found to be somewhat volatile at low temperature.
3. Its aqueous solution is alkaline to litmus.
4. One subjecting it to ignition it burns with a smokeless flame.
5. **Solubility Profile:** 1g dissolves in 1.5 mL of water; 12.5mL ethanol; 320 mL ether; and 10 mL chloroform.
6. The pH of a 0.2 molar aqueous solution is 8.4.

Uses

1. It is found to be an extremely effective urinary tract anti-infective agent provided it is made to act in an acidic medium.
2. As it is exerted quite rapidly, hence it attains effective antiseptic concentrations in the urine.

Note: Methenamine depends on its action solely on the liberation of free formaldehyde.

Thus, it attains 20% of theoretical value at pH 5.0; 6% at pH 6.0; and nil at pH 7.6.

3. It is of particular value in the treatment of *E. coli* infections of the urinary tract.
4. It is also useful specifically in patients with *renal insufficiency*. By virtue of its low systemic toxicity, failure to excrete the drug produces absolutely very little harmful consequences; of course, unless and until the renal insufficiency is of a severe nature.


Note: The drug should not be used in patients having hepatic insufficiency.

B. Nitrofurantoin

Synonyms Berkfurin; Chemiofuran; Cyantin; Cystit; FuaMed; Furachel; Furalan; Furadantin; Furadantine MC; Furadoin; Furantoina; Furobactina; Furophen T-Caps; Ituran; Macrofantin; Parfuran; Trantoin; Urantoin; Urizept; Urocin; Urolong; Uro-Tablinen; Welfurin.

Preparation It is prepared by the interaction of 1-aminohydantoin sulphate and 5-nitro-2-furaldehyde diacetate.*

Chemical Structure



Nitrofurantoin

1-(5-Nitro-2-furfurylideneamino) hydantoin; (C₈H₆N₄O₅).

Characteristic Features

1. It is obtained as orange-yellow needles, crystals or fine powder from dilute acetic acid that get decomposed at 270-272°C. It is found to be odourless and gives a bitter aftertaste.
2. It has dissociation constant pK_a 7.2.
3. It shows uv_{\max} : 370 nm ($E_{1\text{cm}}^{1\%}$ 776).
4. **Solubility Profile:** It has solubility (mg/100 mL): water (neutral pH 7) 19.0; 95% (v/v) ethanol 51.0; acetone 510; DMF 8000; peanut oil 2.1; glycerol 60; and polyethylene glycol 1500.

Nitrofurantoin Monohydrate [C₈H₆N₄O₅·H₂O] [*Synonyms Macrobid*]

Uses

1. It is found to be effective against a majority of urinary tract pathogens, including certain strains of *E. coli*; *Enterobacter*; *Klebsiella*, *Proteus Spp.*; *Staphylococcus aureus*; and *Streptococcus faecalis*.
2. It is also effective against many *staphylococci*, *clostridia*, and *Bacillus subtilis*.
3. It is invariably indicated for the treatment of infections of the urinary tract caused by the aforementioned organisms: pyelonephritis; cystitis and pyelitis.
4. An acidic urine favours activity; and in chronic bacteriuria, it is a second- or third-choice agent.
5. Interestingly, as a prophylactic in the prevention of recurrences it is found to be effective, being slightly superior to methenamine mandelate but inferior to sulfamethiazole.
6. It has a half life for only 0.3 hour.

9.3.10.1.5.5 Agents for Mycobacterium Tuberculosis Infections One of the most dreadful diseases across the world is tuberculosis which is caused by the pathogenic organisms *Mycobacterium tuberculosis*. The **antimycobacterial drugs** existing **generally are usually divided into two main categories. namely: first-line drugs**, which essentially comprise of the natural products, such as: **streptomycin** (see section 9.3.1.6), **rifampin** (Section 9.3.10.1.1.A) and **rifabutin**, plus a good number of purely synthetic compounds including **ethambutol**, **isoniazid (INH)** and **pyrazinamide**; **second-line drugs**, are invariably employed in *three* specific events, for instance: (a) organisms resistant to first line drugs, (b) patient idiosyncrasy prohibit use of the first-line agents, and (c) serious adverse and untoward effects, and these mostly include **cycloserine** (Section 9.3.8.1) and

* Swirska *et al.* *Przem. Chem.*, **11**, (34), 306, (1955).

caperomycin belonging to the class of natural products, whereas **para-aminosalicylic acid (PAS)** and **ethionamide** belong to the synthetic origin.

It is, however, pertinent to mention here that, in general, *tuberculosis* is always treated with a combination **antimycobacterial agents** to minimise the emergence of resistance organisms and also to broaden its spectrum of activity.


It is thought worthwhile to discuss one synthetic drug each from the **first-line** and **second-line agents** in this section:

A. Pyrazinamide

Synonyms Pyrazinecarboxamide; Prezetamid; Pyrafat; Pirilene; Piraldina; Tebrazid; Unipyranamide; Zinamide.

Preparation **Pyrazinamide** may be prepared by the thermal decarboxylation of 2, 3-pyrazinedicarboxylic acid to form the monocarboxylic acid, which is esterified with methanol and finally subjected to controlled ammonolysis.*

Chemical Structure



Pyrazinamide

Pyrazine carboxamide; (C₅H₅N₃O).

Characteristic Features

1. It is obtained as white to practically white crystals from either ethanol or water having mp 189-191°C. It is found to sublime at 60°C.
2. Its dissociation constant pK_a is 0.5.
3. It has uv_{\max} : 269 nm ($E_{1\text{cm}}^{1\%}$ 660).
4. **Solubility Profile** (mg. mL⁻¹): water 15; methanol 13.8, absolute ethanol 5.7; isopropanol 3.8; ether 1.0; isooctane 0.01; chloroform 7.4.
5. Its aqueous solutions are found to be neutral.

Uses

1. It is used as an antituberculosis drug used for initial treatment in combination with **isoniazid** and **rifampin**.
2. It is invariably administered with **isoniazid**, which it potentiates.

Note: All patients intended to be treated with the drug must have thorough prior liver-function tests, which tests also should be repeated periodically during therapy.


* *J. Am. Chem. Soc.* **74**, 3617 (1952).

B. Ethionamide

Synonyms Amidazine; Ethioniamide; Bayer 5312; Nisotin; Actina; Trescatyl; Ethimide; Iridocin; Tio-Mid.

Preparation It may be prepared by the method suggested by Libermann *et al.**

Chemical Structure



Ethionamide

2-Ethyl-4-pyridinecarbothioamide; (C₈H₁₀N₂S).

Characteristic Features

1. It is obtained as minute yellow crystals from ethanol that get decomposed at 164-166°C.
2. It is found to be very sparingly soluble in ether and water; sparingly soluble in methanol, ethanol, propylene glycol; soluble in hot acetone, dichloroethane; and freely soluble in pyridine.

Uses

1. It is mostly used as a potent tuberculostatic agent.

9.3.10.1.5.6 Antifungal Agents: Immunocompromised hosts are usually pretty gullible and prone to dreadful pathogenic fungi solely responsible for causing a large number of infections. Generally, all fungi are eucaryotic organisms and by virtue of this fact they behave more like human cells than are bacteria. It is, however, pertinent to mention here that the antibacterial compounds may not necessarily either inhibit the growth of fungi or able to kill them completely; and, therefore, it has been virtually a difficult task to identify rather selective and specific targets within the fungal cells. The latest range of systemic antifungal agents invariably and largely target cell membrane synthesis or their integrity. In the recent past, significant emphasis has been attached towards the development of antifungal drugs based on the cardinal fact that the fungal membranes essentially contain a sterol, **ergosterol**, which is not present in any human cell membranes.

Broadly speaking fungi that cause painful human infections are of *four* major categories, namely: (a) **yeast-i.e.**, *Candida albicans*, usually found in the normal flora that may conveniently overgrow in various vital organs of the body, such as: bladder, intestine, vagina, mouth, skin and even infect the blood stream; besides another organism known as *Cryptococcus neoformans* which specifically causes meningitis or pneumonia; (b) **filamentous dermatophytes i.e.**, long, interwoven, irregularly placed fungal parasite that grows in or on the skin, which normally cause skin infections, for instance: ringworm (tinea corporis), athlete's foot (tinea pedis) and nail infections (onychomycosis); (c) **dimorphic fungi-i.e.**, which essentially have both filamentous and yeast-like structures, and are

* Libermann *et al. Compt. Rend.* **242**, 2409 (1956).

responsible for causing either exclusively primary pulmonary or secondary disseminated infections by such organisms as: *Blastomyces*, *Coccidioides*, *Histoplasma*, *Paracoccidioides*; and (d) **filamentous fungi**-*i.e.*, fungi having long, interwoven and irregularly shaped structures *e.g.*, *Aspergillus* that may duly attack on a variety of sites in the body.

In actual practice, there are *three* well-known **natural products** that are invariably used as the antifungal agents, namely: **griseofulvin**, **polyenes (amphotericin B)**, and **undecylenic acid**, that are employed to treat *dermatophytic infections*. Besides, there are some synthetic antifungal agents that are exclusively used to combat the dermatophytic infections, such as: *allylamines (naftifine, terbinafine)*; *5-fluorocytosine*; and the “*azoles*” (**fluconazole**, **miconazole**).

First, of all the antifungal agents derived from the natural products shall be discussed, followed by a brief discussion of the purely synthetic products in the sections that follows:

Natural Products


A. Griseofulvin

Synonyms Amudane; Curling Factor; Fulcin; Fulvicin; Grifulvin; Grisactin; Grisefuline; Grisovin; Gris-PEG; Grysio, Lamoryl; Likuden; Neo-Fulcin; Polygris; Poncyl-FP; Spirofulvin; Sporostatin.

Biological Sources **Griseofulvin** is an antibiotic substance produced by *Penicillium griseofulvum* Diercks and also by *Penicillium janzewskii* Zal. [Same as *P. nigricans* (Banier) Thom]. It is produced by *Penicillium patulum*.

Preparation It is produced commercially by employing the submerged process using *P. patulum*.

Chemical Structure



Griseofulvin

(1'S-*trans*)-7-Chloro-2', 4, 6-trimethoxy-6'-methylspiro[benzofuran 2(3H), 1'-[2] cyclohexene]-3, 4'-dione; (C₁₇H₁₇ClO₆).

Characteristic Features

1. It is obtained as white to creamy white, powder, wherein particles of the order of 4 μm in diameter usually predominate. It may also be obtained as octahedra or rhombs from benzene having mp 220°C; and is odourless.
2. It has specific optical rotation $[\alpha]_D^{17} + 370^\circ$ (in saturated chloroform solution).
3. It has uv_{max} : 286, 325 nm.
4. Its solubility in DMF at 25°C is 12-14 g per 100 mL; slightly soluble in ethanol, methanol, acetone, benzene, chloroform, ethyl cetate, acetic acid; almost insoluble in water, petroleum ether.

Uses

1. **Griseofulvin** is **fungistatic** and not **fungicidal**, and serves as an effective agent in the treatment of superficial fungal infections.
2. When administered systemically it is found to be highly effective in the control and management of *tinea capitis*, *tinea corporis*, *tinea unguium* (onychomycosis) and the chronic form of *tinea pedis* normally caused by the dermatophytes, namely: *Microsporon*, *Trichophyton*, and *Epidermophyton*.
3. As it evidently exerts a fungistatic activity thereby only arresting reproduction of the organism, it is absolutely necessary to continue medication long enough so that the entire epidermis undergoes shedding and replaced so as to get rid of reinfecting organisms.


Mechanism of Action: Griseofulvin gets deposited in the **keratin precursor cells** and is carried outwards into the epidermis as normal skin-growth proceeds. It also obviates for a long latency from the time medication is commenced until evidence of improvement takes place.

4. Its half-life is 24-36 hours.
5. It is the drug of choice for mucocutaneous infections.

Biosynthesis of Griseofulvin The various steps involved in the **biosynthesis of griseofulvin** are as follows:

1. **Griseofulvin** is produced biosynthetically from the head-to-tail condensation of 7-acetate units to yield **polyketide**—as the basic precursor.
2. **Griseophenone C** is obtained as an early intermediate in the biosynthetic pathway.
3. Subsequent methylation and chlorination are assumed to precede the oxidative coupling of the benzophenone to give rise to the formation of **dehydrogriseofulvin-a spiran**.
4. Obviously, the last and ultimate step is the production of **griseofulvin** *via* reduction.

The aforementioned intermediates involved in various biotransformations from step (1) through (4) may be summarized as given below:



Biosynthesis of Griseofulvin


B. Amphotericin B

Synonyms Ambisome; Amphozone; Fungizone; Fungilin; Ampho-Moronal.

Biological Sources It is a polyene antibiotic produced by M 4575 obtained from the soil of the Orinoco river region of Venezuela.*

Preparation **Amphotericin B** may be prepared by the growth of selected strains of *Streptomyces nodosus* in a suitable culture medium under specific controlled parameters of temperature, pH and aeration. Subsequently, after due extraction from the medium, the crude product is purified by treatment with different solvents at a controlled acidity (pH).

Chemical Structure ($C_{47}H_{73}NO_{17}$)




Amphotericin B

Characteristic Features

1. It is obtained as yellow to orange powder or as deep yellow-prisms or needles from DMF that gradually get decomposed above 170°C; and is odourless.
2. It has uv_{max} (methanol): 406, 382, 363, 345 nm.
3. It shows specific optical rotation $[\alpha]_D^{24} + 333^\circ$ (acidic MDF); -33.6° (0.1N methanolic HCl).
4. It has dissociation constant pKa (acid) 5.7, (amine) 10.0.
5. Solids and solutions are fairly stable for long durations between pH 4 and 10 when stored at moderate temperatures out of direct contact with light and air.
6. **Solubility Profile:** It is found to be soluble at pH 2 or pH 11 in water: about 0.1 mg mL⁻¹; soluble in DMF 2-4 mg mL⁻¹; in DMF + HCl: 60–80 mg mL⁻¹; in DMSO: 30-40 mg mL⁻¹. However, its aqueous solubility may be enhanced to nearly 50 mg mL⁻¹, by complexation with **sodium desoxycholate**.

* Walters *et al.* *J. Am. Chem. Soc.*, **79**, 5076, (1957).



Sodium Desoxycholate

Uses

1. It exhibits the widest spectrum of antifungal activity of any systemic antifungal drug.
2. It is proclaimed to be an extremely useful drug when given by IV route for the management and therapy of a host of systemic fungus diseases, especially coccidioidomycosis, cryptococcosis, systemic moniliasis, histoplasmosis, aspergillosis, rhodotorulosis, sporotrichosis, phycomycosis (mucormycosis), and North American blastomycosis.
3. It also is used topically in the treatment of superficial monilial infections.
4. It also finds its application by nasal spray in the prophylaxis of aspergillosis in immunocompromised patients.
5. It has an initial half-life of 24-hours, which is followed by a terminal half-life of about 15 days.

Note: It is highly bound predominantly to β -lipoproteins and is excreted gradually by the kidneys but neither renal failure nor hemodialysis has a consistent effect on the plasma levels.

C. Undecylenic Acid

Synonyms 10-Undecenoic acid; Declid; Renselin; Sevinon.

Preparation It is obtained either by pyrolysis of ricinoleic acid or by vacuum distillation of castor oil. It is also found to occur in sweat.

Chemical Structure

Undecylenic Acid

Characteristic Features

1. It is obtained as liquid or crystals having mp 24.5°C.
2. It has a distinct odour suggestive of perspiration.
3. It shows various densities as: $d_4^{24}(\text{vac.})$ 0.9072; d_{25}^{25} 0.9102; d_{45}^{45} 0.8993; $d_4^{79.9}(\text{vac.})$ 0.8653.
4. It has n_D^{25} 1.4486.
5. It shows various bp: bp₇₆₀ 275°C (decomposes); bp₁₃₀ 230-235°C; bp₉₀ 198-200°C; and bp_{1.0} 131°C.
6. It has neutralization value 304.5; and iodine value 137.8.
7. It is found to be insoluble in water, but soluble in chloroform, ethanol and ether.

Undecylenic Acid Zinc Salt [$C_{22}H_{38}O_4Zn$] [*Synonyms* Zinc undecylenate]: It is obtained as an amorphous white powder having mp 115–116°C. It resembles zinc stearate in appearance and physical properties. It may be prepared by dissolving zinc oxide in dilute undecylenic acid and concentrating the solution


Synthetic Products: The various important synthetic antifungal products are:

A. Allylamines Recently, two new allylamines have been synthesized that are found to be potent antifungal agents:

(a) Naftifine

Synonyms Naftifungin.

Chemical Structure It is an antimycolytic allylamine.



Naftifine

(E)-N-Methyl-N-(3-phenyl-2-propenyl)-1-naphthalene methanamine; ($C_{21}H_{21}N$).

Characteristic Features It is a colourless viscous oil bp_{0.015 torr} 162-167°C.

Naftifine Hydrochloride [$C_{21}H_{21}N.HCl$] [*Synonyms* Exoderil; Naftin; AW-105-843; SN-105-843]: It is obtained as colourless crystals from propanol having mp 177°C.


Uses It is mostly employed as a topical antifungal agent in the treatment of various skin infections.

(b) Terbinafine

Synonyms Lamisil; SF-86-327.

Preparation It is an antimycotic allylamine related to naftifine and may be prepared by the method provided by Lednicer *et al.**

Chemical Structure



Terbinafine

* D. Lednicer *et al.*, *Org. Chem. of Drug Syn.* Vol. 4, Wiley, NY, p-55 (1990).

trans-N-Methyl-N (1-naphthylmethyl)-6, 6-dimethylhept-2-en-4-ynyl-1-amine; (C₂₁H₂₅N).

Characteristic Features


Terbinafine Hydrochloride [H₂₁H₂₅N.HCl]

1. It is obtained as white to off-white crystalline powder or as crystals from a mixture of 2-propanol + diethyl ether having mp 195-198°C (alteration in crystal structure commences nearly at 150°C).
2. **Solubility Profile:** It is freely soluble in methanol and methylene chloride; soluble in ethanol; and slightly soluble in water.


Uses

1. It is the *first allylamine* known so far as a systemic drug recommended in the treatment of all dermatophytes, namely: *Trichophyton*, *Epidermophyton*, and *Microspora*.
2. It is also used for topical therapy of dermatophytes including various types of *tinea* infections.

Mechanism of Action **Terbinafine hydrochloride** selectively inhibits the *fungal squalene epoxidase* responsible for causing a *fungicidal action* by virtue of the intracellular accumulation of the toxic sterol, *squalene*; it is also found to exert a *fungistatic action* by depletion of *ergosterol*.



Squalene



Ergosterol


B. Fluorocytosine It essentially has basic pyrimidine nucleus that has been duly substituted with fluoro-, amino-, and oxo-functions. This relatively not-so-complicated molecule was introduced in early seventies as a potent drug against yeasts, such as: *Candida* and *Cryptococcus*.

(a) Flucytosine

Synonyms 5-Fluorocytosine; 5-FC; Alcobon; Ancobon; Ancotil; Cytosine, 5-fluoro-.

Preparation **Flucytosine** may be prepared by the interaction of **5-fluorouracil** with POCl₃ to yield **2,4-dichloro-5-fluoropyrimidine** which is subsequently reacted with ammonia (NH₃) to give rise to **2-chloro-4-amino-5-fluoropyrimidine**. Heating the latter in a medium of concentrated HCl forms the desired product.

Chemical Structure



Flucytosine


4-Amino-5-fluoro-2(H)-pyrimidinone; (C₄H₄FN₃O).

Characteristic Features

1. It is obtained as a odourless, white off-white crystalline powder with a mp 295-297°C (decompose).
2. It is found to be fairly stable in light, non hygroscopic; stable for at least 3 months at 45°C.
3. It has dissociation constants as pKa 2.9 and 10.7.
4. Its uv_{max} (0.1 N HCl): 285 nm (ϵ 8900).
5. **Solubility Profile:** Solubility in water: 1.5g/100 ml at 25°C; 1g in 12 mL 0.1N HCl; slightly soluble in ethanol; and almost insoluble in chloroform, ether.


Uses

1. **Flucytosine** gets converted in the fungus to **5-fluorouracil** in the presence of fungal deaminase, which is subsequently incorporated into RNA, that interferes with normal protein synthesis. It has been observed that certain fungal organisms are relatively more sensitive to interference from the drug than are the human cells, and hence it is definitely, useful in the radical treatment of some fungal infections.




2. It has been found that majority of clinical isolates of *Cryptococcus*; and 40-92% of *Candida* are sensitive to the drug.
3. It is the *drug of choice* to treat *chromomycosis*; and of *second choice* to treat systemic *candidiasis*.
4. It may be logically combined with amphotericin B [section 9.3.10.1.5.6.(B)] for the *first-choice treatment* of *aspergillosis* or *cryptococcosis*, especially associated with *meningitis*.
5. It has a normal half-life 0.5-1 hour.

C. Azoles In general, the chemical compounds containing either 'imidazole' or 'triazole' moieties are invariably clubbed together and baptized as "azoles".




Imidazole




Triazole


The **imidazole** bearing antifungal agents are: **miconazole**, **clotrimazole**, **econazole**, and **ketoconazole** as given below:




Miconazole



Clotrimazole




Econazole




Ketoconazole

The **triazole** containing antifungal agents are, namely: **fluconazole**, and **itraconazole** as shown under:



Fluconazole



Itraconazole

It is thought worthwhile to discuss at least one representative drug from each of the aforesaid *two* class of compounds:

(a) Miconazole

Synonyms Micatin; Monistat.

Preparation It is a topical drug for mucocutaneous infections exclusively. **Miconazole** may be prepared* by alkylation of imidazole with 2, 4-dichlorophenacyl bromide followed by reduction of the ketone function to a corresponding secondary alcohol which is subsequently converted to the alkoxide. Finally, the Williamson alkylation with α , *p*,-dichlorotoluene gives rise to the desired product.

Characteristic Features

Miconazole Nitrate [C₃₄H₃₂N₂O₅] [**Synonyms** Aflorix; Albistat; Andergin; Brentan; Conoderm; Conofite; Daktarin; Deralbine; Dermonistat; Florid; Fungiderm; Micotef; Prilagin; Vodol]: It is obtained as crystals having mp 170.5°C.

(+)-Form Nitrate It has mp 135.3°C; and $[\alpha]_D^{20} + 59^\circ$ (methanol).

(-)-Form Nitrate It has mp 135°C; and $[\alpha]_D^{20} - 58^\circ$ (methanol).

Uses

1. It exerts significant fungicidal action to various species of *Aspergillus*, *Blastomyces*, *Candida*, *Cladosporium*, *Epidermophyton*, *Histoplasma*, *Microsporon*, *Paracoccidioides*, and *Trichophyton*.
2. In *tinea pedis* (*i.e.*, **Athlete's Foot**) a mycological cure rate of 96% has been duly accomplished with the nitrate salt, which appreciably exceeds that of any other drugs except **clotrimazole** and **econazole**.
3. It is recommended for topical usage in *vulvovaginal candidiasis*, successful cure rate varies from 80-95% which is significantly superior to that accomplished with **nystatin** (65%), and **amphotericin B** (75%).
4. It has been observed that pruritus is cured even after a single application.
5. It is also found to be very effective against certain vaginal infections caused by *Trichophyton glabratum*.
6. **Miconazole** base is very useful in the topical treatment of various ophthalmic mycoses.
7. **Miconazole** base has been recommended very successfully for the systemic cure and treatment of a number of deep and systemic mycoses, such as: *Candidiasis* and *Cryptococcosis*.

Mechanism of Action It specifically inhibits ergosterol synthesis, thereby disrupting the fungal cell membranes. Thus, the drug rapidly penetrates deep into the *stratum corneum* where it remains in relatively high concentration upto 96 hours, which situation perhaps contributes its remarkable efficacy against the *dermatophytoses*.

* *J. Med. Chem.*, **12**, 784, (1969).

(b) Fluconazole

Synonyms Biozolene; Diflucan; Elazor; Triflucan.

Characteristic Features It is an orally active **bistriazole antifungal agent** which is obtained as white crystals having mp 139°C. It is also obtained as crystals from a mixture of ethyl acetate and hexane that melts about 138-140°C.

Uses

1. It is found to be a highly selective inhibitor especially of *fungal cytochrome P-450* together with *sterol C-14 α -demethylation* that is responsible for the inhibition of ergosterol synthesis.
2. **Fluconazole** is a broad-spectrum bistriazole antifungal substance which is observed to be primarily fungistatic with appreciable activity against *Cryptococcus neoformans* and *Candida spp.*
3. It is duly recommended for *systemic candidiasis, oropharyngeal and esophageal candidiasis; and cryptococcal meningitis.*
4. The bioavailability of oral fluconazole is more than 90% compared with IV administration.
5. The volume of distribution is 0.8gL⁻¹.
6. The plasma half-life is about 30 hours.

Note: Fluconazole may particularly alter cytochrome P-450 pathways of metabolism of a good number of therapeutic agents, such as: cyclosporine, phenytoin, sulphonyl ureas, and warfarin.

9.3.10.1.5.7 Anticancer Antibiotics A plethora of **anticancer antibiotics** have been derived exclusively from natural origin that are used either independently or in combination with other chemotherapeutic agents. Of course, it may not be within the scope of this text to deal with the entire compounds used in anticancer chemotherapy, but an attempt is made to discuss a few most potent drugs in the sections that follow, namely: **doxorubicin, daunorubicin, dactinomycin and mitomycin C.**

A. Doxorubicin It has already been discussed in this chapter under Section 9.3.2.1.


B. Daunorubicin

Synonyms Daunomycin; Leukaemomycin C; Rubidomycin; Cerubidin; RP-13057.

Preparation It is an anthracyclinic antibiotic related to the rhodomycins. It is produced from the fermentation broths of *Streptomyces peucetins* or *S. coeruleorubidus*. **Daunorubicin** is a glycoside formed by a tetracyclic aglycone, **daunomycinone** (C₂₁H₁₈O₈) and an aminosugar, **daunosamine** (C₆H₁₃NO₃), 3-amino-2, 3,6-trideoxy-L-lyxo-hexose. However, it may also be synthesized by the method of Acton *et al.**

* E.M. Acton *et al.*, *J. Med. Chem.* **17**, 659 (1974).

Chemical Structure



(8*S*-*cis*)-8-Acetyl-10-[(3-amino-2, 3, 6-trideoxy- α -L-lyxo-hexopyranosyl) oxy]-7, 8, 9, 10-tetrahydro-6, 8, 11-trihydroxy-1-methoxy-5, 12-naphthacenedione; (C₂₇H₂₉NO₁₀).

Characteristic Features It is obtained as crystals having mp 208–209°C.

Daunorubicin Hydrochloride [C₂₇H₂₉NO₁₀·HCl] [Synonyms Cerubidine; Daunoblastina; Ondena.]

1. It is obtained as thin red needles that decompose at 188-199°C.
2. The pH of an aqueous solution containing 5 mg mL⁻¹ ranges between 4.5 to 6.5.
3. The specific optical rotation $[\alpha]_D^{20} + 248 \pm 5^\circ$ (C = 0.05-0.1 in CH₃ OH).
4. The colour of aqueous solution changes from pink at acid pH to blue at alkaline pH.
5. It has uv_{max} (methanol): 234, 252, 290, 480, 495, and 532 nm ($E_{1cm}^{1\%}$ 665, 462, 153, 214, 218, and 112).
6. **Solubility Profile:** It is soluble in water, methanol, aqueous alcohols; and almost insoluble in chloroform, ether and benzene.

Uses


1. It is invariably employed for the treatment of severe (acute) lymphocytic and nonlymphocytic leukemias, usually as a component of combination chemotherapeutic regimens. It has been found to undergo very quick reductive metabolism specifically in the liver to give rise to the active **dauno-rubicinol**.
2. It inhibits DNA synthesis by means of a series of bio-chemical reactions, such as: intercalation into DNA, inhibition of topoisomerase II, and production of oxygen radicals. It may prevent cell division in such doses that do not interfere with nucleic acid synthesis.
3. In combination with other drugs it is included in the **first-choice-chemotherapy** of acute *myelocytic leukemia* in adults (for induction of remission) and also in the acute phase of *chronic myelocytic leukemia*.
4. The half-life of distribution is 45 minutes. The half-life of its active metabolite, **daunorubicinol**, is nearly 27 hours.

C. Dactinomycin

Synonyms Actinomycin D; Meractinomycin; Actinomycin A_{IV}; Actinomycin C₁; Actinomycin I₁; Actinomycin X₁; Cosmegen.

Preparation It is an antibiotic substance belonging to the **actinomycin** complex, elaborated during the culture of *Streptomyces parvulus*. After extracting from the fermentation broth by suitable solvents, it is subsequently subjected to purification through various chromatographic and crystallization processes (*US Patent: 2, 378, 876*).

Chemical Structure



Dactinomycin [C₆₂ H₈₆ N₁₂ O₁₆]

Characteristic Features

1. It is obtained as a bright-red crystalline powder, light sensitive and should be protected appropriately. It melts at 246°C with the decomposition.
2. It should be protected from excessive heat and moisture.
3. It essentially contains in one mg an amount of antibiotic activity of not less than 900 mcg of dactinomycin.
4. **Solubility Profile:** It is found to be soluble 1g in 8 mL ethanol; 25 mL water (at 10°C); 1 L water (at 37°C) and approximately 1666 mL ether.

Uses

1. It is an antineoplastic substance that specifically inhibits DNA-dependent RNA-polymerase, recommended for use in **Wilm's tumour***; **rhabdomyosarcoma****, carcinoma of the testis and the uterus.
2. It is an essential component of the first-choice combinations for the control, management and treatment of **choriocarcinoma,*** embryonal rhabdomyosarcoma**, and **Wilm's tumour**.
3. It is still recommended for use against **Ewing's sarcoma,****** testicular carcinoma, and sarcoma botyroides (*i.e.*, resembling a bunch of grapes).
4. Dactinomycin appreciably potentiates radiotherapy (*i.e.*, *radiation recall*).

* **Wilm's tumour** A rapidly developing tumour of the kidney that usually occurs in children.

** **Rhabdomyosarcoma** An extremely malignant neoplasm, originating in skeletal muscle.

*** **Choriocarcinoma** An extremely rare, very malignant neoplasm, usually of the uterus but sometimes at the site of the ectopic pregnancy.

**** **Ewing's sarcoma** A diffuse endothelial mycloma forming a fusiform swelling on a long bone.

5. It is found to be a secondary (efferent) immunosuppressive agents.
6. It has half-life approximately upto 36 hours.


Note: It must be handled with exceptional care, to prevent inhaling particles of it and exposing the skin to it.

D. Mitomycin C

Synonyms MMC; Ametycine; Mitocin-C; Mutamycin.

Preparation Mitomycin C is one of the three very intimately related entities isolated from the antibiotic complex produced by *Streptomyces caespitosus*, an organism from the Japanese soil.

Chemical Structure



Mitomycin C [C₁₅H₁₈N₄O₅]

Characteristic Features

1. It is obtained as blue-violet crystals or crystalline powder, that does not melt below 360°C.
2. It has uv_{\max} (methanol): 216, 360, 560 nm ($E_{1\text{cm}}^{1\%}$ 742, 742, 0.06).
3. **Solubility Profile:** It is found to be soluble in water, methanol, acetone, butyl acetate and cyclohexanone; slightly soluble in benzene, carbon tetrachloride, ether; and almost insoluble in petroleum ether.

Uses

1. It predominantly inhibits DNA synthesis by virtue of cross-linking double-stranded DNA through guanine and cytosine.
2. It is specifically recommended for the palliative treatment of **disseminated adenocarcinoma* of the stomach and the pancreas that have failed other treatment.**
3. It is an integral component of *second--line combinations* for the control, management and treatment of *cervical, gastric, pancreatic carcinomas*, and above all *non-small-cell bronchogenic carcinoma*.
4. It is usually introduced into the bladder in **papilloma.****
5. It is mostly employed as an *alternative drug* in the therapy of *head and neck squamous*** cell carcinoma, bladder carcinoma, and osteogenic sarcoma.*

Note: The active form of nitromycin C is produced metabolically *in situ* and evidently caters as an alkylating agent to suppress DNA synthesis categorically.

* **Adenocarcinoma** A malignant adenoma (*i.e.*, a neoplasm of glandular epithelium) arising from a glandular organ.
 ** **Papilloma** A benign epithelial tumour.
 *** **Squamous** Scalelike.

FURTHER READING REFERENCES

1. Bambeke FV *et al.*, **Glycopeptide Antibiotics from Conventional Molecules to New Derivatives**, *Drugs*, **64**: 913-36, 2004.
2. Berdy, J. *et al.* (eds): **Handbook of Antibiotic Compounds**, Vols. 1-8, Boca Taton, FL, USA : CRC-Press, 1982.
3. Brakhage, A: *Biosynthesis of β -Lactam Compounds in Microorganisms*. **Comprehensive Natural Products Chemistry**, Vol. 4, Elsevier, Amsterdam, 1999.
4. Calam, C.T.: **Process Development in Antibiotic Fermentation**. Cambridge, UK: Cambridge University Press, 1987.
5. Cavalleri, B. and Parenti F: Antibiotics [glycopeptides (Dalbaheptides)]. **Kirk-Othmer Encyclopedia of Chemical Technology**, 4th edn., Vol. 2. Wiley, New York, 1992.
6. Coute JE: **Manual of Antibiotics and Infections Diseases**, 8th ed., Williams & Wilkins, Baltimore, 1995.
7. Croom KF and Goa KL: **Levofloxacin—A Review of its use in the Treatment of Bacterial Infections in the United States**, *Drugs*, **63**: 2769-802, 2003.
8. Davies, J.E.: Aminoglycoside-aminocyclitol Antibiotics and Their Modifying Enzymes. In: Loraian, V., ed. **Antibiotics in Laboratory Medicine**, 3rd ed. Baltimore, Williams &-Wilkins, 1991.
9. Fenton C, Keating CM, and Curran MP: **Daptomycin**, *Drugs*, **64**: 445-55, 2004.
10. Hughes, M.A.: **Biosynthesis and Degradation of Cyanogenic Glycosides**. **Comprehensive Natural Products Chemistry**, Vol. 1, Elsevier, Amsterdam, 1999.
11. Jawetz *et al.*, **Medical Microbiology**, 20th ed., Appleton & Lange, Norwalk CT, 1995.
12. Kwon-Chung, KJ and JE Benett: **Medical Mycology**, Lea & Fabiger, Philadelphia, 1992.
13. Kucers A., Bennett, N. McK, and Kemp R.J. Eds.: **The Use of Antibiotics**, J.B. Lippincott, Philadelphia, 1987.
14. Lancini *et al.*, **Antibiotics: an Interdisciplinary Approach**, Plenum Press, New York, 3rd edn., 1995.
15. Luengo JM: Enzymatic Synthesis of Penicillins, **Comprehensive Natural Products Chemistry**, Vol. 4. Elsevier, Amsterdam, 1999.
16. Mandell GL, Bennett JE, and Dolin R, **Mandell, Douglas, and Benett's Principles and Practice of Infectious Diseases**, 4th ed, Churchill Livingstone, Inc., New York, 1995.
17. Page MI (ed.): **The Chemistry of β -Lactams**, Chapman & Hall, New York, 1992.
18. Sutcliffe J and NH Georgopapdakou, Eds.: **Emerging Targets in Antibacterial and Antifungal Chemotherapy**, Chapman and Hall, New York, 1992.
19. Vanden HB, DWR Mackenzie, G. Cauwenbergh, JV Custem, E Drouket, and B Dupont Eds.: **Mycoses in AIDS Patients**, Plenum Press, New York, 1990.
20. Wellington K and Noble S: **Telithromycin**, *Drugs*, **64**: 1683-94, 2004.
21. Wise EM: Antibiotics (Peptides). **Kirk-Othmer Encyclopedia of Chemical Technology**, 4th edn. Vol. 3., Wiley, New York, 1992.

10

Drug Molecules of Marine Organisms

- Introduction
- Classification of Drug Molecules
- Marine Natural Products: An Upgradation Profile
- Summary
- Further Reading, References

10.1 INTRODUCTION

The phrase '**drug molecules**' has been appropriately and judiciously employed in line with the classical concept, that is a specific chemical entity which essentially possesses a marked and pronounced pharmacological activity emphatically on the mammalian organism. In fact, it is the dedicated and concerted efforts of expert researchers from various specialized scientific fields that a new drug molecule is isolated, characterized and subsequently subjected to a rigorous preclinical and successful clinical studies and ultimately baptized as a therapeutically effective and potent '**drug**'.

Interestingly, innumerable products derived from the marine organisms in several '**crude forms**' have been widely used across the globe by the traditional practitioners for thousands of years. However, scientific as well as logical approach to such marine organisms could materialize only during the last five decades or so. Krebs* and Faulkner** gave the most comprehensive and excellent reviews on the chemistry of thousands of compounds obtained exclusively from the marine organisms during the eighties.

Kaul and Daftari*** presented the latest review on pharmacology of chemically well-defined molecules belonging to marine organisms. In the light of the ever mounting evidences pieced together over the recent past it has more or less strengthened the belief and hypothesis that there exists not only an ample hidden treasure but also a tremendous scope for the emergence of newer drug molecules from the marine environment to combat the human sufferings.

Marine toxins were reported to possess an extremely high potency with regard to their pharmacological actions, and, therefore, sometimes collectively referred to as '**toxins**'. The doctrine and philosophy of *Paracelsus* advocating that 'the clinical efficacy and usefulness of an active principle is nothing but a matter of the right dose administered by the right route' carries a lot of

* Krebs, M.C., *Fortschr. Chem. Organ. Naturstoffe*, **48**, 151 (1986).

** Faulkner, D.J., *Nat. Prod. Rep.* 1. 251 and 551 (1984): 3.1 (1986): **4**, 539 (1986).

*** Kaul, P.N., and Daftari, P., *Ann. Rev. Pharmacol. Toxicol.* **26**, 117 (1986).

value and impact. Nevertheless, the joint efforts by phytochemists and medical scientists may pave the way to assess the plethora of the currently labeled 'marine toxins' for their usefulness as drugs and or as physiological tools that may ultimately decode the mechanisms responsible for various cellular processes solely characteristic of life but unfortunately not revealed as on date.

It is indeed a very critical as well as a crucial turning point when phytochemists, medicinal chemists and pharmacologists will put their knowledge, wisdom, expertise, skill and resources together to unravel the wealth of drugs beneath the sea as they already had accomplished toward the terrestrial plants almost a century ago; and towards soil, samples nearly half-a-century ago for 'antibiotics' when penicillin was discovered by Alexander Fleming.

The present chapter exclusively focuses upon a good complication of certain vital and important 'marine biological' (marine organisms) having established chemical structures and proven pharmacological activities. So far, more than five lakh species of marine organisms have been well-documented both from the ocean and sea across the globe.

Interestingly, quite a sizable number of such marine organisms do possess a wide-spectrum of biological activities, namely: antibiotics, antiviral, antineoplastic, cytotoxic, antimicrobial, antiinflammatory, enzyme inhibitors, prostaglandins, neurophysiological and cardiovascular agents.

10.2 CLASSIFICATION OF DRUG MOLECULES OF MARINE ORGANISMS

The enormous quantum of newer and potent drug molecules derived from the wide spectrum of marine organisms across the world may be judiciously and logically classified based on their specific pharmacologic actions as stated below:

- (i) Cytotoxic/Antineoplastic Agents
- (ii) Cardiovascular active drugs
- (iii) Marine Toxins
- (iv) Antimicrobial drugs
- (v) Antibiotic substances.
- (vi) Antiinflammatory and Antispasmodic Agents
- (vii) Miscellaneous pharmacologically active substances.

The above mentioned broad classification of marine organisms shall be dealt with at length individually in the sections that follows:

10.2.1 Cytotoxic/Antineoplastic Agents

The two prominent US-based research institutes, namely: US **National Cancer Institute (NCI)** and US **National Sea Grant Office (NSGO)** have discovered thousands of pure and semi-pure compounds derived from the marine origin that distinctly exhibited antineoplastic/cytotoxic activities in a good number of cell lines; besides, *in vivo* actions against both malignant tumours and leukemias in various animal models. In the present context certain duly characterized drug molecules displaying known and potent activities shall be discussed, namely:

- (a) Cembranes,
- (b) Macrolides,

- (c) Depipeptides, and
 (d) Miscellaneous compounds.


10.2.1.1 Cembranes

Cembranoids, are the 14-membered cyclic diterpenes obtained from a wide variety of soft corals. A good number of cembranoids have been isolated and characterized. It has been observed that most of these compounds do contain an *exocyclic lactone* as their integral part.

A few typical examples of naturally occurring *cembranes* are, namely: sinularin, crassin acetate, cytarabine (Ara-C), fludarabine and aplysiastatin. These compounds shall now be discussed individually in the sections that follows:

A. Sinularin

Chemical Structure




Sinularin

Biological Source Sinularin and its dihydro congener are obtained from *Sinularia flexibilis*.

Uses It possesses anticancer activities.

B. Crassin Acetate

Chemical Structure



Crassin Acetate

Biological Source It is obtained from the **Caribbean gorgonian** *Pseudoplexaura porosa*.

Uses **Crassin acetate** was observed to be comparatively inert to the mammalian systems but on the contrary found to be extremely cytotoxic to human leukemic as well as Hella cells *in vitro* and also to the mouse fibroblasts.


Note: (a) The aforesaid activity is almost lost as soon as the exocyclic lactone portion of the compound undergoes cessation (*i.e.*, cleavage) by titration with NaOH.

(b) Moreover, the conversion of 'acetate' to 'hydroxyl' via hydrolysis may render this compound more water soluble and useful only if the 'lactone moiety' is kept intact.

C. Cytarabine

Synonyms Ara-C; Alexan; Arabitin; Aracytine; Cytarbel; Cytosar; Erpalfa; Iretin; Udicil; U-19920; CHX-3311; Aracytidine; β -Cytosine Arabinoside.

Chemical Structure




Cytarabine

4-Amino-1- β -D-arabinofuranosyl-2(1H)-pyrimidinone; (C₉H₁₃N₃O₅).

Cytarabine is a synthetic compound exclusively based on the knowledge of naturally occurring moieties found to be present in the '**Caribbean Sponges**' e.g., **spongosine**, and **spongouridine**.

It may be synthesised* by the acetylation of **uracil arabinoside** followed by treatment with phosphorus pentasulphide (P₂S₅) and subsequent heating with ammonia as given below:



Characteristic Features

1. It is obtained as prisms from ethanol (50% v/v) having mp 212-213°C.
2. Its specific optical rotation $[\alpha]_D^{23} + 158^\circ$; $[\alpha]_D^{24} + 153^\circ$ (C = 0.5 in water).
3. It has uv_{max} at pH 2 : 281.0, 212.5 nm (ϵ 13171, 10230); and at pH 12 : 272.5 nm (ϵ 9259).

Uses

1. It is indicated in both adult and childhood leukemia.


* Kar, A.: **Medicinal Chemistry**, New Aage International, New Delhi, 4th, edn., 491, 2006.

2. It is found to be specifically useful in acute granulocytic leukemia, and is more effective when combined with thioguanine and daunorubicine.
3. It is a potent antineoplastic and antiviral agent.
4. It is also employed in the treatment of acute myelogenous leukemia and human acute leukemia.

D. Fludarabine

Synonyms 2-Fluorovidarabine; 2F-Ara A.

Chemical Structure



Fludarabine

2-Fluorovidarabine; (C₁₀H₁₂FN₅O₄).


Characteristic Features

1. It is obtained as crystals from ethanol and water having mp 260°C.
2. Its specific optical rotation is $[\alpha]_D^{25} + 17 \pm 2.5^\circ$ (C = 0.1 in ethanol).
3. It has uv_{max} (pH1, pH7, pH13): 262, 261, 262 nm ($\epsilon \times 10^{-3}$ 13.2, 14.8, 15.0).
4. It is sparingly soluble in water and organic solvents.

Uses It is used as a antineoplastic agent.

E. Aplysistatin

Chemical Structure



Aplysistatin

Biological Source **Aplysistatin** is obtained from *Aplysia angasi* (Sea Hare).

Uses It is used as an antineoplastic agent.

Interestingly, certain **cembranoids** were observed to deciliate protozoa even at a 5 ppm level*

* Rerkins, D.L., and Ciereszko, L.S.: *Hydrobiologia*, **42**, 77 (1973).


thereby implying that perhaps quite a few compounds belonging to this category or their feasible structural analogues might also exhibit either spermicidal and/or parasiticidal properties. It is, however, pertinent to state here that these drug molecules derived from the marine origin obviously require both intensive and extensive preclinical studies by virtue of the fact that most of them are virtually inert to the mammalian systems.*

Non-Lactonic Cembranoid There are certain compounds which do not contain a lactone moiety but they do possess cytotoxic actions. A few such examples of **non-lactonic cembranoid** are, namely: geranylhydroquinone and asperidol. These *two* drugs are described as under:

A. Geranylhydroquinone

Synonyms Geroquinol; Geranyl-1, 4-benzenediol.

Chemical Structure



Geranylhydroquinone

trans-(3, 7-Dimethyl-2, 6-octadien-yl)-hydroquinone; (C₁₆H₂₂O₂).

Biological Sources Geranylhydroquinone is obtained from the chloroform extract of *Aplidium* species.


Characteristic Features It is usually obtained as colourless needles from n-hexane/ethyl acetate having mp 61–62°C.

Uses

1. It is found to be cytotoxic to leukemia and mammary carcinoma.
2. It is employed as a radioprotective agent.

B. Asperidol

Chemical Structure



Asperidol

* Kobayashi, Y., and Osabe, K.: *Chem. Pharm. Bull.* **37**, 631, (1989).

Biological Source It is obtained from the *gorgonian* coral.

Use It exhibits cytotoxic activities.

10.2.2 Cardiovascular Active Drugs

During the past three decades a huge number of extracts, fractions and pure isolates from thousands of **marine organisms** were subjected to thorough cardiovascular screening in various research laboratories around the world. Interestingly, most of these compounds did exhibit **cardiovascular activities** perhaps as frequently as observed antibiotic and antineoplastic activities. Unfortunately, as on date hardly any compound could surface out and obtain the FDA approval as a potential drug.

The cardiovascular active drugs may be broadly classified under the following *two* categories, namely:

- (a) Cardiotonics, and
- (b) Hypotensive compounds.

These *two* categories shall now be treated separately as under:

10.2.2.1 Cardiotonics

The **cardiotonic compounds** (or **cardiotonics**) showing positive response in either *in vivo* and/or *in vitro* inotropic activities on whole or part of the heart are included in this section.

In general, the **cardiotonics** may be further sub-divided into *two* groups, namely:

- (i) Marine peptides, and
- (ii) Marine glycosides

10.2.2.1.1 Marine Peptides The age-old belief and classical concept that the steroidal nucleus present in the aglycone residues of either **digitalis** or **strophanthus** could only exhibit cardiotonicity was virtually turned down when ‘**marine peptides**’, obtained from *coelenterates*, such as: *Anthopleura xanthogrammica* producing **anthopleurins A, B and C*** (also known as **AP-A, B, C**); and *Anthopleura elegantissima* giving **AP-C**.

Out of these *three* anthopleurins, only **AP-A** has been reported to be showing the most promising cardiotonic activity both on the isolated and *in situ* heart of different species. **AP-A** has a positive edge over the known **cardiac glycosides** because of its unique ability to afford a sustained significant inotropic effect under the ischemic conditions. Gross *et al.*** advocated strongly that perhaps AP-A could prove to be extremely beneficial especially in patients having a concomitant β -adrenergic blockade.

Furthermore, the **anemonia toxin II** (also termed as: **Cardiotoxin-II, ATX-II**) isolated from *Anemonia sulcata* (**Sea Anemone**) comprising of at least 47 amino acids demonstrated a very close resemblance to **AP-A***** Besides, ATX-II was shown to exhibit a significant dose-dependent cardiotonic activities in different mammalian heart experiments.

* Norton *et al.* *J. Pharm. Sci.*, **65**, 1368, (1976).

** Gross *et al.* *Eur. J. Pharmacol.* **110**, 271 (1985).

*** Kaul, P.N. and Dabarti, P.; *Ann. Rev. Pharmacol. Toxicol.* **26**, 117, (1986).

The cardinal points of difference between **AP-A** and **ATX-II** peptides are as follows:

S. No.	Anthopleurin-A (AP-A)	Anemonia Toxin-II (ATX-II)
1	Alanine is present at the residue 38	Lysine is present at the residue 38.
2	No similar studies as for ATX-II were carried out.	Modification of the acidic-COOH function at the 38 residue leads to absolute inactivation of cardiotoxic potency.*
3	If administered systemically gives an immune response; and when given orally loses its activity due to gastric juice (acidic) in stomach.	No such effect has been observed.


Likewise, the two compounds *viz.*, **AP-A** and **ATX-II**, display a number of similarities, namely: (a) conformation of the two peptides are almost identical; (b) amino-acid sequence similar; (c) disulphide bridges are alike; and (d) they have the same basicity. Based on these striking points of similarities one may rightly justify the very identical profiles of **cardiotoxic activity** of the said two peptides.

A host of other compounds showing **cardiotoxic activities** belonging to the class of '**marine peptides**' are, namely: **laminin**, **octopamine**, **saxitoxin** and **autonomium chloride**.

These compounds shall now be treated individually in the sections that follow:

A. Laminin

Chemical Structure



Laminin

Biological Source **Laminin** is obtained from a marine algae *Laminaria angustata*.

Characteristic Features

1. It is the abundant structural component of the basal lamina.
2. It is critical to the stability of the extracellular matrix and to the adhesion of cells to the basement membrane.
3. It belongs to the family of **heterotrimeric glycoproteins** composed of a heavy chain, designated as α (also known as A) and 2 light chains, designated as β (B1) and γ (B2), which are linked by disulphide bonds to form an asymmetrical cross-shaped structure.
4. Eight genetically distinct laminin subunits have been identified, namely: α_1 , α_2 , α_3 , β_1 , β_2 , β_3 , γ_1 and γ_2 .


Uses

1. It shows hypotensive effect.
2. It also exhibits diverse biological activities.

* Barhanin *et. el. Toxicol.* **20**, 59 (1982).

B. Octopamine

Chemical Structure



Octopamine

Biological Sources **Octopamine** is found in the salivary glands of *Octopus vulgaris*, *Octopus macropus* and of *Eledone moschata*.

Characteristic Features D(-)-Form:

1. It is obtained as crystals from hot water that gets changed at about 160° to a compound which melts above 250° (decomposes).
2. Its specific optical rotation $[\alpha]_D^{25} - 56.0^\circ$ (0.1 N.HCl); and $- 37.4^\circ$ (water).

DL (±)-Octopamine Hydrochloride (C₈H₁₁NO₂.HCl) (Epirenor, Norden, Norfen): It is obtained as crystals which gets decomposed at 170°C. It is freely soluble in water.


Uses

1. The natural D(-) form is almost 3 times more potent than the L(+) form in producing cardiovascular adrenergic responses in anaesthetized dogs and cats.
2. It gives distinct adrenergic responses.
3. In invertebrate nervous systems **octopamine** may function as a neurotransmitter.

C. Saxitoxin

Synonyms Mussel poison; Clam poison; Paralytic shellfish poison; Gonyoulax toxin; STX.

Chemical Structure [C₁₀H₁₇N₇O₄]²⁺



Saxitoxin

Biological Sources **Saxitoxin** is the powerful neurotoxin produced by the dinoflagellates *Gonyaulax catenella* or *G. tamarensis*, the consumption of which causes the California sea mussel *Mytilus californianus*, the Alaskan butterclam *Saxidomus giganteus* and the scallop to become poisonous.*

* Ghazarossian *et al. Biochem. Biophys. Res. Commun.*, **59**, 1219 (1974) Sehantz *et al., Can. J. Chem.* **39**, 2117 (1961)

Isolation Saxitoxin may be isolated by the method suggested by Schantz et al.*

Characteristic Features

Saxitoxin Dihydrochloride [(C₁₀H₁₇N₇O₄)²⁺ · 2HCl]:


1. It is obtained as a white hygroscopic solid.
2. It has pKa in water: 8.24, 11.60.
3. Its specific optical rotation $[\alpha]_D^{25} + 130^\circ$.
4. It is extremely soluble in water, methanol; sparingly soluble in ethanol, glacial acetic acid; and practically insoluble in lipid solvents.
5. It is found to be fairly stable in acid solutions; and decomposes rapidly in an alkaline media.
6. On boiling for 3 to 4 hours at pH 3 usually causes loss of activity.

Uses

1. It exhibits a hypotensive effect.
2. It is invariably employed as a tool in the neurochemical research.

D. Autonomium Chloride


Chemical Structure



Autonomium Chloride

Biological Source It is found in *Verongia fistularis*.

Characteristic Features Autonomium chloride possesses an isosteric structure of **adrenaline** and **acetylcholine** as given below:



Adrenaline

Acetylcholine

Uses

1. It exerts both α - and β -adrenergic effects.
2. **Autonomium chloride** also exhibits cholinergic action.
3. It distinctly shows CNS stimulant activity in *mice* which action is evidently exhibited by an apparent substantial increase in the spontaneous motor activity (SMA).
4. By virtue of the unique dual effects of adrenergic and cholinergic it may prove to be an asset in the control, management and regulation of the behaviour of heart *i.e.*, **cardiotonic effect**.

* Schantz et al. *J. Am. Chem. Soc.*, **79**, 5230 (1957).

In addition to the above cited examples of the marine peptides, there are quite a few polypeptides which have been isolated and characterized from a wide spectrum of sea anemones. A few such typical examples are given as under:

S. No.	Biological Source	Compound	Uses
1	Actinia equina	A polypeptide with 147 amino acids	Exhibits bradycardia, rapid hypotension, and respiratory arrest in the rat.
2	Condylactis gigantea	A polypeptide with 195 amino acid residue	Exerts a haemolytic action in rabbits.
3	Parasicyonis action stoloides	A polypeptide with very less number of amino acids	Shows a neurotoxic action.

10.2.2.1.2 Marine Glycosides The **marine glycosides**, in general, are of two types, viz., nonsulphated and sulphated ones. However, **holothurins** and **astrosaponins** are two typical examples of **marine glycosides** that generally display cardiotoxic activity having an unusual narrow margin between the effective dose (*i.e.*, ED₅₀) and the lethal dose (*i.e.*, LD₅₀).

Holothurins: These are the aglyconic residues obtained from the family *Holothuroidea* of phylum *Echinodermata* and essentially possess a **steroidal moiety** that very much resemble to the aglycones of the **digitalis glycosides**. In the recent past a large number of the '**hydroxylated steroidal glycosides**' have been isolated and characterized from the **holothuroids**, also known as the sea cucumbers.

Astrosaponins: These are the **marine glycosides** obtained from the star fishes belonging to the family *Asteroidae*.

In fact, both these **marine glycosides** (*i.e.*, sulphated and non-sulphated) inhibit Na⁺, K⁺, Mg²⁺ and ATPases, but the holothurins are reported to be far more active on the Na⁺, K⁺ and ATPase.*

Interestingly, the **astrosaponins** are found to exert an altogether different type of pharmacological actions, such as: antiinflammatory, analgesic, haemolytic, hypotensive; besides having the cytolytic activity on account of its interference in neuromuscular blocking effects and the protein metabolism.

However, the **holothurins** are found to exert both **cardiotonic** and **ichthyotoxic actions**. Besides, they also exhibit haemolytic activity.

Eledoisin: [C₅₄H₈₅N₁₃O₁₅S]

Chemical Structure 5-oxoPro-Pro-Ser-Lys-Asp-Ala-Phe-Ile-Gly-Leu-MethNH₂.

Biological Source **Eldoisin** is obtained from the posterior salivary glands of eledone spp. (small octopus spp.) *Eledone moschata*.**

Characteristic Features

1. **Eledoisin** is obtained as a sesquihydrate powder that gets decomposed at 230°C.
2. Its specific optical rotation $[\alpha]_D^{22} - 44^\circ$ (C = 1 in 95% acetic acid).
3. It is found to lose its activity gradually when incubated in blood.

Uses

1. Its physiologic action resembles that of the other **tachykinins**, **substance P** and **physalaemin**.
2. It is found to stimulate extravascular smooth muscle.

* Gorshkov *et al. Toxicon*, **20**, 655 (1982).

** Anastasi, Erspamer, *Brit. J. Pharmacol.*, **19**, 326, (1962); *idem. Experientia*, **18**, 58, (1962).

3. Eledoisin acts as a potent vasodilator and hypotensive agent.
4. It causes salivation, and enhances capillary permeability in certain specific species.
5. It also stimulates lacrimal secretion.

10.2.2.2 Hypotensive Compounds

There are quite a few potent hypotensive compounds that have been derived from a variety of **marine organisms**. These newer range of medicinally active chemical entities may be categorized into *two* groups, namely:

- (i) Marine nucleosides, and
- (ii) Hypotensive peptides and other compounds.

The various '**marine biomedicines**' belonging to each of these *two* categories shall now be discussed as under:


10.2.2.2.1 Marine Nucleosides Nucleosides are formed by the combination of a purine or pyrimidine base in glycosidic linkage with a sugar moiety, such as: adenosine, thymidine etc.; whereas, the '*phosphate ester*' of a nucleoside is known as *nucleotide* e.g., 5'-guanylic acid; 3'-cytidylic acid.

In the past two-and-a-half decade a plethora of **marine nucleosides** have been isolated and characterized, and also evaluated for their therapeutic efficacy.

Cryptotethia crypta, the well-known **Caribbean sponge** made a spectacular success in the history of marine biomedicine that indeed produced a very rare and unique arabinosynucleoside which resulted on slightest structural manifestation the wonderful drug of choice used for antileukemic treatment cytarabine or Ara-C (see section 10.2.1.1)*. A good number other nucleosides have been derived from *C. crypta* that showed varying interests. However, *spongosine*, being one such a drug that gained meaningful legitimate recognition because of its highly potential hypotensive activities. *Doridosine*, is another most promising and potent hypotensive nucleoside reported.** These *two* compounds shall now be discussed as under:

A. Spongosine It is a **nucleoside** and the methoxy derivative of **adenosine**.

Chemical Structure



* Cohen, S.S.: *Cancer*, **40**, 509 (1977).

** F.A. Fuhrman *et al. Science*, **207**, 193, (1980).


Biological Source It is obtained from the **Caribbean sponge** *Cryptotethia crypta* and with a minor structural modification of the parent isolated nucleoside known as **arabinosylnucleoside**.

Uses

1. It exhibits various coronary effects resembling to those of **adenosine**, for instance: coronary vasodilation and negative inotropy.
2. It is found to exert more marked and pronounced long-acting effects.
3. It acts as a hypotensive at such as dose level at which **adenosine** is observed to be absolutely inactive.
4. It reduces the rate as well as the force of contraction of heart.

B. Doridosine

Chemical Structure



Biological Source It is obtained from the nudibranch *Anisodoris nobilis*:

Characteristic Features

1. It has been revealed on the basis of the kinetics of the enzymatic degradation of these nucleosides that **doridosine** is not only the most active but also the long-lasting nucleoside.
2. **Doridosine** does not undergo oxidative deamination *in vitro* on being subjected to incubation in the presence of **adenosine deaminase**, while **adenosine** disappears very fast. Perhaps the intensity and the duration of the cardiovascular effects of doridosine is directly related to its half-life *in vivo*.

Note: The intermediate acting, spongiosine happens to disappear in a gradual manner.

Uses

1. It is the most potent hypotensive marine nucleoside known so far.
2. It also exerts hypothermic effect *i.e.*, it lowers the normal temperature of the body.

Hypothermic Effect Interestingly, both **spongiosine** and **doridosine**, when administered *intracerebroventricularly* to guinea pigs, they lower the body temperature by several degrees for a

duration ranging between 4-7 hours.* However, this hypothermic effect seems to have no bearing to their prevailing systemic actions.

5'-Deoxy-5-iodobericidin: It is also a **nucleoside** which has been observed to lower the body temperature of mice**. It has been isolated from *Hypnea valentiae* (Red Algae). Perhaps this is the first and foremost compound which possesses the following *two* specific criteria, namely:

- (a) First iodinated nucleoside discovered, and
- (b) First 5'-deoxyribosyl nucleoside discovered in nature.

Uses


1. It may prove to be a vital biochemical research tool of immense interest.
2. It is found to be a potent inhibitor of **adenosine kinase**.

10.2.2.2 Hypotensive Peptides and Other Compounds A number of peptides have been isolated from the **marine organisms** that exclusively showed distinct and significant hypotensive activities in experimental laboratory animals.

There are certain typical examples, such as: **aaptamine**, **hymenin** and **uro-tensins I and II**, which shall now be discussed briefly in the sections that follows:

A. Aaptamine

Chemical Structure



Aaptamine

Biological Source Aaptamine is obtained from *Aaptos aaptos*.

Characteristic Features It has an inherent interesting heterocyclic nucleus besides its structural characteristic and molecular size which may render this molecule a good candidate drug for future intensive as well as extensive pharmacological studies provided it should exhibit a bare minimum level of toxicity.

Uses


1. It has an α -adrenergic blocking effect.
2. In case, it does not undergo a rapid metabolism in mammalian species *in situ*, it may prove to cause a hypotensive effect.

* Kaul, P.N.,: *Pure Apl. Chem.*, **54**, 1963 (1982).

** L.P. Davis *et al.*: *Biochem. Pharmacol.* **33**, 347, (1984).

B. Hymenin

Chemical Structure



Biological Source Hymenin is obtained from *Hymeniacidon aldis*.

Its characteristic features and uses are very much similar to those of aaptamine.

C. Urotensins I and II (U I, U II)

Biological Sources Urotensins I and II are obtained from the specific caudal neurosecretory system of *Giltichthys miralilis* (**Telecost**); and also from *Catostomus commersoni*—a fish.

Characteristic Features Both U I and U II are naturally occurring **poly-peptides** which distinctly possess appreciable hypotensive activity as evidently shown in rat, dog, sheep and squirrel monkey.

Uses

1. These **polypeptides** exhibit hypotensive effect that seems to be on account of their vasodilatory action.
2. Nevertheless, the discoverers of these two vasoactive peptides (Laderis and McCannell) proposed that they may possibly prove to be clinically potential drugs, but till date no evidence for their clinical efficacy has yet been reported.

10.2.3 Marine Toxins

A host of **marine biotoxins** have been obtained from a wide variety of **marine organisms** in their crude, semipure and other forms between 1960-1970. However, during the eighties a good number of them possessing venomous and toxic properties and having most complex chemical structures have been isolated, characterized with the advent of rather exceedingly sophisticated analytical instruments and a concerted effort of dedicated marine chemists and pharmacologists across the globe.

The **marine toxins** are caused due to either the **external metabolites (ectocrine)** or the **endotoxins**.* The endotoxins are found to be the most potent substances.

The **marine toxins** may be classified appropriately under the following categories, namely:

- (a) Palytoxin,
- (b) Red-Tide toxins
- (c) Ciguatera toxins.

* **Endotoxin**—A lipopolysaccharide that is part of the cell wall of Gram Negative bacteria.


The aforesaid varieties of toxins shall now be discussed in the sections that follows:

10.2.3.1 Palytoxin

Palytoxin (PTX) was initially reported from the Pacific ocean by Moore and Scheuer* in 1971; and in 1974 by Attaway and Ciereszko** from the Caribbean island; incidentally poles apart by two research teams almost at the same-time-span.

Biological Source It is obtained from **zanthid coral** of the genus *Polythoa* found abundantly both in the Pacific and the Caribbean oceans. **It is the most poisonous non-proteinaceous compound known.**

Chemical Structure



PALYTOXIN
[Polytoxin (C51-55) Hemiactal]

Polytoxin is regarded as one of the most interesting chemical entity isolated and characterized from a marine organism which is found to be linear, polycyclic, and polyether molecule having molecular formula C₁₂₉ H₂₂₃ N₃ O₅₄. Perhaps PTX is the largest **non-peptide marine biotoxin** known till date.

* Moore, R.E., and P.J. Schever: *Science*, **172**, 495 (1971).

** Attaway, D.H. and Ciereszko, L.S: *Proc. Soc. Int. Coral Reef Sym.* **7**, 497, (1974).

Characteristic Features

1. **Palytoxin** is a white amorphous hygroscopic solid powder.
2. It does not have any definite mp, but gets charred when heated to 330°C.
3. It has specific optical rotation $[\alpha]_D^{25} + 26^\circ$ (water).
4. It is insoluble in ether, chloroform, acetone; sparingly soluble in methanol, ethanol; and fairly soluble in DMSO, water and pyridine.

Uses

1. It is the most potent coronary vasoconstrictor.
2. It is a versatile physiological tool to evaluate anti-anginal chemotherapeutic agents.
3. Based on the evidence that $\text{Na}^+ - \text{K}^+$ exchange is almost instantaneous and massive in cells following exposure to PTX, it may be inferred beyond any reasonable doubt that it exclusively acts by affecting the Na^+, K^+ -AT Pase.*

10.2.3.2 Red Tide Toxins

Starr** (1958) was pioneer in reporting the red tide toxin from the Gulf of Mexico. In fact, the terminology '**red tide**' became known from the very existence of the red coloured *peridinin*, a carotenoid natural pigment, obtained from the periodical blooming of the **dinoflagellates**, thereby imparting to the water a brown to red colouration.

Interestingly, there are only *three* well-known biological species which exclusively produce toxins, namely: (a) *Ptychodiscus brevis* [earlier called *Gymnodinium breve*]; (b) *Protogonyaulax catenella* [earlier known as *Gaunyalax catenella*]; and *Protogonyaulax tamarensis*. Because, these organisms possess a mutually beneficial association with the **shell fishes**, the toxins generated by the former get duly accumulated in the latter; and upon subsequent consumption by human beings usually response to a toxic epidemic frequently known as the —'**paralytic shell fish poisoning**' (PSP).

In 1975, Japan (in Owase Bay) first had the incidence of PSP; followed, by Philippines and Thailand in 1983; and in 1986 both at Korea and Taiwan.

The various **red tide organisms** (*i.e.*, **dinoflagellate species**) containing specific *toxins*, *geographical distributions*, and their *physiological effects* are as stated below:

S. No.	Red Tide Organisms	Toxins Present	Geographical Distribution	Physiological Effects
1	<i>Protogonyaulax catenella</i>	GTXs and STXs	Japan; South America; Pacific USA;	Block membrane Na^+ conductance
2	<i>Alexandrium tamarensis</i> (<i>Protogonyaulax tamarensis</i>)	GTXs and STXs	North Atlantic	-do- coastal regions
3	<i>Ptychodiscus brevis</i>	Brevetoxin	Florida; Gulf of Mexico.	Positive inotropic and arrhythmogenic.
4	<i>Protogonyaulax cohorticula</i>	GTXs	Gulf of Thailand	Paralytic syndrome.
5	<i>Gymnodinium catenatum</i>	GTXs and PX	Japan, Tasmania	-do-
6	<i>Pyrodinium bahamense</i>	STXs and GTXs	Palau Island	-do-
7	<i>Prorocentrum lima</i>	Pectenotoxins	Gulf of Lima	Diarrhetic action.

* Habermann, E.: *Toxicon*, **27**, 1171 (1989).

** Starr. T.J.: *Texas Reb. Biol. Med.*, **16**, 813 (1958).

GTX = Gonyautoxins;
STXs = Saxitoxins;
PX = Protogonyautoxin.


The various important red tide toxins, are, namely: **saxitoxin**, **tetrodotoxin**, and **brevetoxin**. These toxins shall be discussed as under:

10.2.3.2.1 Saxitoxin It has already been described under Section 10.2.2.1.1.

10.2.3.2.2 Tetrodotoxin

Synonyms Fugu poison; Maculotoxin; Spheroidine; Tarichatoxin; Tetrodontoxin; TTX.

Chemical Structure



Tetrodotoxin

Octahydro-12-(hydroxymethyl)-2-imino-5, 9:7, 10a-dimethano-10aH-[1, 3] dioxocino [6, 5-d] pyrimidine-4, 7, 10, 11, 12-pentol; (C₁₁H₁₇N₃O₈).

Biological Sources **Tetrodotoxin** is obtained from the ovaries and liver of a large number of species of *Tetraodontidae*, especially the *Spheroides rubripes* (**Globe Fish**). It is also obtained from the **puffer fish** (*Tetraodon* species) which begin a topmost delicacy in Japanese cuisine.


Characteristic Features

1. It usually darkens above 220°C without any decomposition.
2. It has specific optical rotation $[\alpha]_D^{25} - 8.64^\circ$ (C = 8.55 in dilute acetic acid).
3. It is slightly soluble in water, absolute ethanol, ether; and almost insoluble in other organic solvents.
4. It has dissociation constant pKa = 8.76 (water); 9.4 in 50% (v/v) ethanol.
5. The toxin gets destroyed completely both in strong alkali and acid solutions.
6. It is a **marine neurotoxin essentially** containing a polar guanidino moiety.

Uses


1. TTX gets bound particularly to the Na⁺ channels on the outside of excitable membranes, thereby inducing Na⁺ influx in exchange for K⁺ influx within a few milliseconds of the accompanying membrane depolarization.
2. It is invariably employed as a valuable pharmacological tool.

Biosynthesis of Tetrodotoxin **Arginine** is a precursor for **tetrodotoxin**. It has been adequately proved that the remainder of the C-skeleton in it is a C₅ *isoprene unit*, perhaps provided as **isopentenyl diphosphate (IPP)** as given below:



10.2.3.2.3 Brevetoxin

Chemical Structure



Brevetoxin [Pb Tx-1]

Biological Sources A plethora of **polycyclic polyether metabolites** have been obtained from the **dinoflagellate** *Ptychodiscus brevis*; and now the **brevetoxins** are commonly known as **PbTx** derived from the generic nomenclature *i.e.*, *P. brevis* toxins.


Characteristic Features

1. These are lipid soluble toxins.
2. The **PbTxs** do not contain any N-atom, which is quite contrary to the STXs and GTXs.
3. The **PbTxs** may be classified into *two* categories, such as: (a) *Hemolytic*, and (b) *Neurotoxic*. However, the latter category are larger in number.
4. The Structure of Pb TX-1, as given above, is the most potent one; and its structure essentially differs from the remaining 9 PbTxs.
5. The toxin Pb Tx-2, is found to be a *cardiotonic* and an *arrythmogenic* in experimental laboratory animal heart studies.

Uses

1. It exerts an excitatory effect on the isolated neuromuscular and other cells. Besides, this effect is further mediated through an increased Na^+ influx subsequent to their strategic binding to *site-5* located upon the voltage-sensitive Na-Channel.
2. Both **Pb Txs** and the **polyether ionophores** (*e.g.*, *norhalichondrins* obtained from *Halichondria okadae* and *okadaic acid*) are found to exhibits marked and pronounced biological actions.
3. It causes both *neurological* and *gastrointestinal* disorders.
4. Pb Txs broadly are splendid activators of voltage-sensitive Na^+ channels, with an unique ability to get bounded to very specific receptor sites located on the rat brain synaptosomes. Hence, it may be useful as an important pharmacological tool.

Biosynthesis of Brevetoxin [Pb T_{x-1}] Brevetoxin (Pb T_{x-1}) is postulated to be generated from a **polyunsaturated fatty acid (PUFA)** through epoxidation of the various double bonds, and subsequently a concerted sequential opening of the epoxide rings ultimately give rise to the extended polyether structure. It has also been proved through biosynthetic studies that the C-skeleton does not conform to a simple polyketide chain, that fragments from the citric acid cycle together with a 4-carbon starter unit derived from mevalonate are also involved, and that quite a few of the methyls actually obtained from the amino acid *methionine*, as depicted below.




10.2.3.3 Ciguatera Toxins

It is pertinent to state here that the **ciguatera toxins**, though not directly linked to the ‘**paralytic shell fish poisoning**’ (PSP) of the red tide, are found to be occurring in the toxic *dinoflagellates* that subsequently penetrate the human-food-chain precisely through the *reef fishes* and *not* through the shellfishes.

In fact, *two* prominent members of this particular category are, namely: **Ciguatoxin (CTX)**, and **Maitotoxin (MTX)** which shall now be discussed as under:

10.2.3.3.1 Ciguatoxin The terminology ‘**ciguatoxin**’ was first and foremost employed to enumerate a disease primarily caused due to the ingestion of *marine snails*. Ciguatera poisoning was thought to be due to the ingestion of blue green algae.

Ciguatoxin [C₅₃H₇₈NO₂₄]

Biological Sources Ciguatoxin (CTX) is found in *Gymnothorax javanicus* (Moray Eel), besides in a variety of coral reef fish, for instance: *Lutjanus bohar* (Red Snapper).

Characteristic Features

1. It is one of the most complex examples of a natural polyether structural chemical substance.
2. A **dinoflagellate** *Gambierdiscus toxicus* definitely causes the production of this polyether, thereby synthesizing a comparatively less toxic analogue, that is eventually passed *via* the food chain; and ultimately modified structurally into the extremely toxic **ciguatoxin** within the system of the fish.

Uses

1. It causes neurological, cardiovascular and gastro-intestinal problems.
2. It is found to be exceedingly toxic even at mcg levels, thereby affecting widespread food poisoning (**ciguatera**) both in subtropical and tropical regions that is evidently characterized by diarrhoea, vomiting and sometimes leading to acute neurological problems.
3. CTX at low IV doses displays bradycardia and respiratory stimulation in anaesthetized cats and dogs; whereas at higher dose levels there is an apparent depression in both heart rate and respiration followed by hypertension (probably on account of reflex compensation of bradycardia).
4. Interestingly, CTX acts as a cardiotoxic at a very low concentrations *i.e.*, picogram levels.
5. CTX—is found to exert its action at the nerve ends specifically.

10.2.3.3.2 Maitotoxin

Synonym MTX.

Biological Source It is obtained from the toxic **dinoflagellate** *Gambierdiscus toxicus*.

Characteristic Features MTX possesses several characteristic pharmacological features, such as:

1. MTX is a Ca²⁺ channel activator.
2. It specifically enhances Ca²⁺ uptake in cultured NG 108-15 neuroblastoma X glioma cells by changing the voltage-dependence of calcium channel activation.
3. MTX is found to enhance the rate as well as the force of contraction in rat myocardial cells immediately followed by arrhythmias.

4. **MTX** affords to induce a dose-dependent release of **GABA*** in the reaggregate clusters of striated neurons, that may be blocked by the calcium blocker D-600.

MTX also has a number of characteristic chemical features, namely:

1. Chemical structure of **MTX** not yet known.
2. **MTX** is a non-peptide having a large molecular weight of 3,425 as estimated by the help of fast atom bombardment mass spectrum.
3. It is found to react with **Dragendorff's Reagent** but fails to give any response with **Ninhydrin Reagent**, suggesting thereby that is basically alkaloidal in nature.
4. The large non-peptide molecule of **MTX** possesses a good number of –OH moieties, ethereal oxygen atoms, and *two* sulphate ester functions.

5. It is found to contain *no* repeatable carbohydrate or amino acid units, no $\text{—}\overset{\text{O}}{\parallel}{\text{C}}\text{—}$ (carbonyl) moieties, *no* carbocycles, and *no* side chains except either a vinyl or a methyl function.

Note: Excepting the 'sulphated ester moieties' all other chemical features very closely resemble to those of PTX (i.e.; Palytoxin).


Uses The overall toxicity of **MTX** almost comes at par to that of **PTX**; however, the latter is obviously much more toxic than the former.

10.2.4 Antimicrobial Drugs

A plethora of important **antimicrobial drug** substances have been isolated, characterized and studied extensively over the past three decades particularly from the vast domain of marine organisms. A few typical examples are, namely: **brown algae**, **red algae** (viz., *three* different species), **gorgonian corals**; and **sponge** (viz., *two* variant species). The chemical substances from these species shall now be discussed briefly as under:

10.2.4.1 Zonarol


Chemical Structure



Zonarol


Biological Source Zonarol and iso-zonarol are both obtained from *Dictyopteris zonaroides* (**Brown Algae**).

* **GABA:** γ -Aminobutyric acid.

10.2.4.2 Prepacifenol**Chemical Structure**

Prepacifenol


Biological Sources It is obtained from *Laurencia pacifica* and *Laurencia filiformis*, the two different species of **Red Algae**.

10.2.4.3 Polyhalo-3-butene-2-one**Chemical Structure**

Polyhalo-3-Butene-2-One


[R₁, R₂, R₃, R₄ and R₅ are Haloatoms]

Biological Source In all four isomers of polyhalo-3-butene-2-one and nearly seven isomer of polyhalo-acetones are obtained from *Asparogopsis taxiformis*, another species of **Red Algae**.

10.2.4.4 Tetrabromo-2-Heptanone**Chemical Structure**

Tetrabromo-2-heptanone

Biological Source It is obtained from yet another species of **Red Algae** *Bonnemaisonia hemifera*.

10.2.4.5 2-Cyano-4, 5-dibromopyrrole**Chemical Structure**


2-Cyano-4, 5-dibromopyrrole

It is perhaps one of the rarest examples of a chemical entity isolated from a **marine organism** which contains a cyano (-CN) function group.


Biological Source It is obtained from *Agelas oroides*, a specific type of **sponge** found in marine sources.

10.2.4.6 *Aerophysinin-1 (+) and Aerophysinin-1 (-)*

Chemical Structures



Aerophysinin-1 (+)




Aerophysinin-1 (-)

Biological Sources The *two* isomers, *viz.*, **aerophysinin-1(+)** and **aerophysinin-1(-)** are obtained from *Verongia aerophoba* another species of **sponge**.

10.2.4.7 *Eunicin*

Chemical Structure



Eunicin

Biological Source **Eunicin** is obtained from *Eunicia mammosa* the well known **Gorgonian Corals**.

10.2.5 Antibiotic Substances


Interestingly, between a span of almost twenty years (1969-1988) thousands of marine-based extracts, fractions and pure isolates were evaluated for their antibiotic activity. However, the success rate was not only miserable but absolutely nonsignificant. As on date, not even a single **marine-derived antibiotic substance** has been able to either supersede or gained enormous recognition with regard to their broad spectrum of activity and superb quality of the known and available **antibiotics** obtained from the innumerable **terrestrial organisms, semi-synthetic products**, and/or purely synthetic ones.

Nevertheless, the earnest efforts contributed by the **marine-chemists** across the globe have been able to evolve a few antibiotics from the various marine organisms, namely: **okadaic acid, acanthifolicin** and **norhalichondrin A**. These naturally occurring marine antibiotics shall now be discussed in the sections that follows:

10.2.5.1 Okadaic Acid

Synonym Halochondrine A;

Chemical Structure



Okadaic Acid

9,10-Deepithio-9, 10-didehydro-acanthifolicin; (C₄₄H₆₈O₁₃).

It is the first **ionophoric polyether** identified in marine organisms.

Biological Sources It is obtained from *Halichondria (okadai or melanodocia)* (**Marine Black Sponges**).

Characteristic Features The characteristic features of **okadaic acid** are as follows:


1. It is obtained from dichloromethane/hexane having mp 171-175°C.
2. It has specific optical rotation $[\alpha]_D^{20} + 21^\circ$ (C = 0.33 in chloroform).
3. It is also reported as crystals from benzene-chloroform mixture having mp 164-166°C; and $[\alpha]_D^{25} + 25.4^\circ$ (C = 0.24 in chloroform).

Uses

1. It is an important biochemical tool as tumour promoter and probe of cellular regulation.
2. It was found to be far more cytotoxic to KB-cells and to mice than compared to another **novel ionophoric marine substance acanthifolicin**.
3. It is able to transport divalent cations *e.g.*, Ca²⁺ across the lipoidal membranes conveniently.
4. **Okadaic acid** uniquely causes a prolonged contraction of the human umbilical artery and rabbit aorta without the presence of extracellular Ca²⁺; and it does not affect the Na⁺ and K⁺ AT Pase.

10.2.5.2 Acanthifolicin

Chemical Structure




Acanthifolicin

Biological Source **Acanthifolicin** is obtained from *Pandaros acanthifolium* (**Sponge**).

Characteristic Features

1. It possesses an antibacterial activity.
2. It also exerts cytotoxic actions.
3. It is found to be lethal to mice at low dose level of 0.14 mg kg^{-1} i.v.

Uses The uses are almost the same as given under **okadaic acid**.

10.2.5.3 Norhalichondrin A**Chemical Structure**

Norhalichondrin A

Biological Source **Norhalichondrin A** and several other **halichondrin** structural analogues have been obtained from *Halichondria okadae* (Sponge).

Characteristic Features

1. It is a **polyether macrolide**.
2. The structure-activity correlational studies with regard to their derivatives; and also their structural fragments are of significant biological interest.

Uses It is found to exert antitumour activity.

10.2.6 Antiinflammatory and Antispasmodic Agents

A plethora of chemical substances have been isolated from the broad spectrum of **marine organisms** which attribute either **antiinflammatory** or **antispasmodic** activities. A few typical examples are summarized below:

S. No.	Chemical Substance	Marine Organism	Common Name	Biological Activities
1	Dendalone-3-hydroxybutyrate	<i>Phyllospongia dendyi</i>	Sponge	Antiinflammatory
2	Flustramine A and B	<i>Flustra foliaceae</i>	Swedish Marine Moss	Antispasmodic
3	Tetradotoxin	<i>Spheroides rubripes</i>	Globe Fish	Strong antispasmodic.
4	6-n-Tridecyl salicylic acid	<i>Caulocystis cephalornithos</i>	Brown Algae	Antiinflammatory
5	Flexibilide	<i>Sinularia flexibilis</i>	Soft coral	Antiinflammatory
6	Monalide	<i>Luffariella variabilis</i>	Sponge	Antiinflammatory (non-steroidal compound).

10.2.7 Miscellaneous Pharmacologically Active Substances

A good number of **pharmacologically active substances** have been isolated and characterized from marine organisms that invariably exhibit a variety of interesting actions. A few important and typical examples have been grouped together under this section, such as: **Latrunculins; Kainic Acid; Domoic Acid; Vidarabine; Aplysinopsin; 28-Deoxyzoanthenammine; Baretin; Nereistotoxin; and Conotoxins.**


The above stated **marine-based chemical substances** shall now be treated individually in the sections that follows:

10.2.7.1 Latrunculins


Latrunculins (LAT) A through D, M and 6, 7-epoxy-LAT A have been isolated and characterized. Interestingly, these are the first and foremost family of natural marine based products essentially having the **2-thiazolidinone moiety**. All, except latrunculin M, are 16- or 14-membered macrocyclic diterpene alkaloids (macrolides) that have been duly isolated and characterized from various corals and sponges, and majority of them do possess certain definite biological activity.

It is, however, pertinent to mention here that out of the *four* toxic **latrunculins A-D**, the first two *i.e.*, **latrunculin A** and **latrunculin B** shall be discussed as under:

Chemical Structures



Latrunculin A
(C₂₂H₃₁NO₅S]



Latrunculin B
(C₂₀H₂₉NO₅S]

Biological Sources Latrunculins are obtained from *Latrunculia magnifica* Keller (**Red Sea Sponge**). They are also found in *Chromodoris elisabethina* (**Pacific Nudibranch**); and the *Spongia mycofjiensis* (**Fijian Sponge**).

Note: When *L. magnifica* is gently squeezed into an aquarium, the toxins are exuded into it spontaneously. The living fish experience sequentially agitation, hemorrhage, loss of balance and ultimately succumbs to death within a span of 4-6 minute.

Isolation Latrunculins have been isolated from *L. magnifica* by Neeman *et al.**

* Neeman *et al.*: *Marine Biol.* **30**, 293, (1975).

Characteristic Features

Latrunculin A (LAT-A)

1. It is obtained as a foam.
2. It has specific optical rotation $[\alpha]_D^{24} + 152^\circ$ (C = 1.2 in chloroform).
3. It has uv_{max} (methanol): 218 nm (ϵ 23500).

Latrunculin B (LAT-B)

1. It is obtained as crystals.
2. It has specific optical rotation $[\alpha]_D^{24} + 112^\circ$ (C = 0.48 in chloroform).
3. It has uv_{max} (methanol): 212 nm (ϵ 17200).


Uses

1. They are used exclusively in establishing the elucidation of molecular mechanisms of motile processes.
2. At nanomolar concentrations in cultured mouse neuroblastoma and fibroblasts, the **LAT-A** and **LAT-B** distinctly cause significant disruption of microfilament organization without affecting the microtubules, analogous to what the less active **cytochalsins** do.*
3. Unlike cytochalasins, the **LAT-A** and **LAT-B** do not afford any change in the polymerization rate of active filaments.

10.2.7.2 Kainic Acid

Synonyms α -Kainic Acid; Digenic Acid; L_S-Xylo-kainic acid; Digenin; Helminal.

Chemical Structure



2-Carboxy-3-carboxymethyl-4-isopropenylpyrrolidine; (C₁₀H₁₅NO₄).

Biological Source It is obtained from dried *Digenea simplex* (Wulf.) Ag., (**Red Algae**) (*Rhodomelaceae*).

Characteristic Features

1. It is obtained as needles which get decomposed at 251°C.
2. It has specific optical rotation $[\alpha]_D^{24} - 14.8^\circ$ (C=1.01).
3. It shows an intense absorption at 6.05 and 112 μ .
4. It is found to be soluble in water; insoluble in ethanol; and fairly stable in boiling aqueous solutions.


* Spector *et al.*: *Science*, **219**, 493 (1983).

Uses

1. It is used as a potent convulsant.
2. It is invariably employed as a vital neurobiological tool.
3. It is mostly employed as an **anthelmintic (Nematodes)**.

10.2.7.3 Domoic Acid

Chemical Structure It is a structural analogue of **kainic acid**.



[2S-[2 α , 3 β , 4 β (1Z, 3E, 5S*)]]-2-Carboxy-4-(5-Carboxy-1-methyl-1, 3-hexadienyl)-3-pyrrolidineacetic acid; (C₁₅H₂₁NO₆).


Biological Source It is obtained from *Chondria armata* Okamura (*Rhodomelaceae*) (**Red Algae**), also known in Japanese as '**DOMOI**'; and hence the name domoic acid.

Characteristic Features**Domoic Acid Dihydrate**

1. It has mp 217°C (decomposes).
2. It has specific optical rotation $[\alpha]_D^{12} - 109.6^\circ$ (C = 1.314 in water).
3. It has uv_{max} : 242 nm (log ϵ 4.24).
4. Its dissociation constant in water pKa: 2.10; 3.72; 4.93; 9.82.
5. It is found to be soluble in water, acetic acid; and insoluble in methanol ethanol, chloroform, acetone and benzene.

10.2.7.4 Vidarabine

Synonyms Ara-A; Vira-A; Arasena-A; Adenine arabinoside; Spongoadenosine; CI-673.

Chemical Structure

9- β -D-Arabinofuranosyl-9H-purine-6-amine monohydrate; (C₁₀H₁₃N₅O₄·H₂O).

It is a purine nucleoside first ever synthesized as a potential antineoplastic agent.

Biological Source It is obtained by culturing a strain of *Streptomyces antibioticus*.


Characteristic Features

1. It is obtained as crystals from water having mp 257.0 – 257.5°C (0.4 H₂O).
2. It has specific optical rotation $[\alpha]_D^{27} - 5^\circ$ (C = 0.25).
3. It has uv_{max} (pH1): 257.5 nm (ϵ 12700); pH 7: 259 nm (ϵ 13400); pH 13 : 259 nm (ϵ 14000).

Uses It is mostly used as an antiviral agent.

10.2.7.5 Aplysinopsin

Chemical Structure It is a tryptophan derivative.



Aplysinopsin


Biological Sources It is originally obtained from *Verongia spengelii* (**Yellow-Sponge**); and also from *Astroides calicularis* (Anthozoan).

Uses

1. It has cytotoxic activity against **KB.P 388 and L1210 leukemia cell lines**.
2. It has been found to exhibit antidepressant activity very much similar to that of **imipramine**.

10.2.7.6 28-Deoxyzoanthamine

Chemical Structure It is an **alkaloid** from the **marine organism**.




28-Deoxyzoanthamine

Biological Source It is obtained from a *Zonathus* sp. from the Bay of Bengal (India).

Uses It is found to possess strong analgesic as well as antiinflammatory activities.

10.2.7.7 Baretin

Chemical Structure It is structurally an **indole-related alkaloid**.




Baretin

Biological Source It is obtained from *Geodia beretti* (**Cold Water Sponge**).

Uses It acts as a smooth muscle stimulant.

10.2.7.8 Nereistotoxin

Chemical Structure ($C_5H_{11}NS_2$)



Nereistotoxin

Biological Source It is obtained from *Lumbriconeris heteropoda*, a marine annelid.

Uses It is mostly used as an effective *insecticide* by virtue of its inherent ganglion blocking effects.

Note: Another semisynthetic structural analogue of nereistotoxin, known as cartap is invariably employed as an useful insecticide.

10.2.7.9 Conotoxins

Biological Sources A plethora of toxins have been obtained and isolated from *Conus geographus*, specifically from the toxic venom of a fish-eating snail. Based on a logical and scientific thought these toxins may be classified into *two* categories, namely:

- (a) **μ -Conotoxins** *i.e.*, a congregation of *seven* homologous peptides having 22 residues each*, and
- (b) **ω -Conotoxins** *i.e.*, a subgroup in a class of peptides which are exclusively neurotoxic in nature.

Characteristic Features

10.2.7.9.1 μ -Conotoxins (I) Kobayashi *et al.*** have reported **geographutoxins I and II** that seem to be almost similar to conotoxins **GIIIA** and **BIIB*****.

10.2.7.9.2 ω -Conotoxins The inhibition of a depolarization-evoked ATP release from synaptosomes of ray electric organ, evidently by affording the blockade of the Ca^{2+} flux through the

* Cruz *et al.*: *J. Biol. Chem.*, **260**, 9280 (1985).

** Kobayashi *et al.*: *Pflugers Arch.*, **407**, 241 (1986).

*** Sato *et al.*, *FEBS lett* **155**, 177 (1983).

voltage-gated calcium channels were demonstrated by ω -CgTX and the related ω -Cm TX obtained from *Conus magnus*.*

Uses [m-Conotoxins]

1. Very similar to the actions shown by the *guanidium toxins viz.*, **tetrodotoxin (TTX)** and **saxitoxin (STX)**, these toxins also effectively block the action potential of skeletal muscle cells; however, unlike **TTX** they fail to affect the channels of the nerves.
2. It has been proved experimentally that the **conotoxin CIII** usually get bound to the same site on the Na⁺ channel where **STX** and **TTX** are found to bind.

ω -Conotoxins It has been duly demonstrated that **w-CgTXVIA** shows a very specific high affinity binding to the synaptosomal and membrane preparations from chick brain. Furthermore, it has given a fairly good evidence that this particular toxin acts onto a *new site* on the Ca²⁺ channel, because it has shown an altogether different sites *i.e.*, from both **dihydropyridine** and **verapamil binding sites**.**

10.3 MARINE NATURAL PRODUCTS: AN UPGRADATION PROFILE

In the last two decades an overwhelming thrust in the research towards accomplishment of an upgradation profile of **marine natural products** have taken place across the globe.

However, these wonderful achievements could only be possible either through various microbial transformations or by means of different semisynthetic structural analogues of puupephenone. Interestingly, **marine natural products** were subjected to rigorous bioconversion studies and different specific organic reactions, such as: acetylation, addition of halogen acids (HX), Grignardization, conjugate additions, and other addition reactions.

The important aspects of such modifications of **marine natural products** shall now be discussed briefly with the help of certain typical examples in the sections that follow:

10.3.1 Microbial Transformations

Microbial transformation may be defined as—‘the phenomenon by which certain organism (bacteria) incorporate DNA from related strains into their genetic make-up’.

The two typical examples of microbial transformations *vis-a-vis* the upgradation profile of **marine natural products** are as stated under:

10.3.1.1 *Sarcophine: Bioconversion Studies*

In general, it is almost essential to carry out the **metabolism studies** so as to obtain the legitimate approval of any clinically useful drug. It has already been duly established that ‘**microorganisms**’ may be employed successfully in carrying out the *in vitro* studies so as to predict the drug metabolism

* Yeager *et al.*: *J. Neurosci.*, 7, 2390, (1987).

** L.J. Cruz and B.M. Olivera: *J. Biol. Chem.*, 261; 6230 (1986).

in the mammals by virtue of the fact that there exists a very close resemblance of some microbial enzyme systems, particularly fungi, with the **mammalian liver enzyme systems**.*

Sarcophine is the **furanochembranoid diterpene** obtained from *Sarcophyton glaucum* having an appreciable yield of 3% dry weight basis. *S. glaucum* is a common soft coral of the Red Sea Pacific, besides other coral reefs. Bernstein *et al.*** advocated that **sarcophine** represents the *major-chemical-defence* entity against the natural predators.

Preparation The specimen of *S. glaucum* (Red Sea Soft Coral) was identified and collected from Hurghada (Egypt) in 1994, by snorkelling and SCUBA (-3m). Approximately 1.1kg of frozen coral was subjected to lyophilization and subsequently extracted with 95% (v/v) ethanol (3 × 21).

Microbial Metabolism Studies Lee's method*** was adopted in carrying out the elaborated microbial metabolism studies. In all, twenty-five authentic **microbial cultures****** were employed for screening. The method of San-Martin *et al.****** was followed for the exclusive microbial bioconversion studies of sarcophine, which evidently proved that it can be duly metabolized by many fungal species. Further, preparative-scale fermentation was carefully performed by using *Absidia glauca* (ATCC*****-22752), *Rhizopus arrhizus* (ATCC-11145), and *Rhizopus stolonifer* (ATCC-24795), ultimately gave rise to the isolation of ten (5-14) altogether new metabolites; besides, the known 7 β , 8 α -dihydroxydeepoxy-sarcophine (4).

It is, however, pertinent to mention here that the structure elucidation of these compounds was entirely based on **2D-NMR spectroscopic studies**. The relative stereochemistry and confirmation of the probable structure of metabolite was ascertained by **X-ray crystallographic studies**.

The various salient features of the **isolated metabolites** are as follows:

1. **Sarcophytols A (2) and B (3)** are simple **cembranoids** isolated from the **Okinawan soft coral** *S. glaucum*. These possess potent inhibitory activities against a wide spectrum of tumour promoters.***** Interestingly, compound (2) helped to mediate dose-dependent diminution of 12-O-tetradecanoyl-phorbol 13-acetate (TPA)-induced transformation of JB6 cells.*****
2. The ability and potential to inhibit TPA-induced TB6 cell transformation, it has been duly observed that plethora of the **sarcophine metabolites** (4 to 14) helped to mediate inhibitory responses much higher than exhibited by **sarcophytol A (2)** or **sarcophine (1)**; and most prominently 7 α -hydroxy- $\Delta^{8(19)}$ **deepoxysarcophine (6)**, which was fairly comparable to the 13-*cis*-retinoic acid.

Note: A good number of novel furanochembranoids as antineoplastic agents may be further developed along these lines.

* Clark, A.M. *et al.*, *Med. Res. Rev.* **5**, 231, (1985).

** Bernstein, J., *et al.* *Tetrahedron*, **30**, 2817, (1974).

*** Lee, I.S., *et al.* *Pharm. Res.* **7**, 199, (1990).


**** University of Mississippi, Dept: of Pharmacognosy Culture Collection Centre.

***** San-Martin *et al.* *J. Phytochem.*, **30**, 2165, (1991).



***** ATCC = American-Type-Culture-Collection.

***** Kobayashi *et al.* *Chem. Pharm. Bull.*, **27**, 2382, (1979).

***** El Sayed *et al.* *J. Org. Chem.*, **63**, 7449, (1998).



No.	R ₁	R ₂	R ₃	R ₄
1	H	CH ₃	H	H
7	H	CH ₃	α-OH	H
10	H	CH ₂ OH	H	H
11	H	CH ₃	β-OH	H
12	β-OH	CH ₃	H	H
13	H	CH ₃	H	β-OH
14	H	CH ₃	H	α-OH



10.3.1.2 Manzamine A and ent-8-Hydroxymanzamine A: Bioconversion Studies

Higa *et al.** in 1986 first and foremost reported the **manzamines**, a group of **novel polycyclic β-carboline-alkaloids**, from the *Haliclona* (**Okinawan Marine Sponge**). Interestingly, the **manzamines** demonstrated an unique and diverse spectrum of bioactivity, such as: antimicrobial, insecticidal and cytotoxic activities.**

Higa *et al.* isolated **manzamine A (15)** to the extent of ~0.85% on anhydrous basis; and elucidated its probable structure with the aid of ¹⁵N-NMR spectroscopy exclusively. Further, the same researchers reported the new **ent-8-hydroxymanzamine A (16)** from an unidentified Indo-Pacific, (a sponge belonging to Family *Petrosidae* and Order *Heplosclerida*). Importantly, both the aforesaid compounds exhibited marked and pronounced **cytotoxic activity** against several tumour *cell-lines*; and also displayed appreciable and significant **antimalarial activity** against *Plasmodium falciparum* exclusively (**D6 and W2 clones**).

However, the two compounds (15 and 16) were selected as candidate '**new molecules**' for the intensive microbial bioconversion studies. El Sayed *et al.**** demonstrated that the said two compounds (**15 & 16**) may be metabolized by a number of microbial species.

Preparative-scale fermentation of compounds 15 and 16 under specific experimental parameters have given the following outstanding findings.


* Sakai *et al. J. Am. Cham. Soc.*, **108**, 6404, (1986).

** Edrada *et al. J. Nat. Prod.* **59**, 1056, (1996).


*** El Sayed *et al. Abstract O30*, The 39th Annual Meeting of the American Society of Pharmacognosy, Orlando, FL, July, 1998.

Compounds	Organism (ATCC No.)	Major Metabolite	Remarks
Manzamine-A; (15) ent-8-Hydroxymanzamine A; (16)	<i>Fusarium oxysporium</i> f. <i>gladioli</i> (ATCC 11137) <i>Nocardia</i> sp. (ATCC 11925) and <i>Fusarium oxysporium</i> (ATCC 7601)	Ircinal A (Compound 17) 12, 34-Oxamanzamine F (Compound 18)	— No cytotoxicity against different cell-line (> 10 mcg mL ⁻¹)


The chemical structures of compounds 15 to 18 are given below:




Manzamine A
[15]



ent-8-Hydroxymanzamine A [16]



Ircinala
[17]



12,34-Oxamanzamine F
[18]

10.3.2 Puupehenone: Semisynthetic Analogues

Puupehenone (19) is a sesquiterpene linked strategically to a C₆-Shikimate function, duly exemplified by the **quinol** (*viz.*, **avarol**)-**quinone** (*viz.*, **avarone**) pair; and interestingly, belongs to a prominent and distinctive family of **sponge metabolites**. Loya *et al.* (1990)* observed that the diversified biological activities proclaimed for this relatively rare group of compounds is solely due to the characteristic feature of *ilimaquinone* (18A) to prevent categorically and check the replication


* Loya, S. *et al.*, *Antimicrob. Agents Chemother.* **34**, 2009, 1990.

phenomenon of the **HIV virus**. It has been established on the basis of the preliminary screening of puupehenone against the pathogenic organism *Mycobacterium tuberculosis* that it caused inhibition to the extent of 99%. Therefore, it has evidently paved the way towards modifying the chemical structures of **puupehenone** and to evaluate their biological activity.

These structural modifications may be accomplished by the following *four* different ways, which shall be treated separately in the sections that follows:

10.3.2.1 Acetylation and Addition HX


Acetylation of puupehenone (19) should have normally yielded the corresponding **monoacetyl structural analogue (20)**. However, it exclusively gives rise to the **triacetyl derivative (21)** through a mechanism whereby the addition of the acetyl moieties takes place specifically at the conjugated double-bond system of the parent compound *i.e.*, puupehenone. It is worthwhile to observe here that the monoacetyl derivative of **puupehenone (20)** is produced as a side product due to the sequence of addition-elimination reaction with HBr/HCl and puupehenone, immediately followed by acetylation which ultimately gave rise to the corresponding **diacetyl derivative (22)** as shown below.




10.3.2.2 Grignardization

Grignardization of puupehenone (19) with methyl-magnesium iodide (H₃CMgI) in diethyl ether as a reaction medium gave rise to *two* products of addition that solely based upon the stoichiometric proportions of the **Grignard Reagent**.


- (a) The addition of *three* times molar excess of H₃CMgI yielded a product having α -orientation of the CH₃ moiety (23). This assignment was, however, based on the extrapolation of NMR features for the 15 α -cyanopuupehenol (X) to (23). Exactly in a similar fashion the interaction of puupehenone with ethyl-magnesium bromide produced 15 α -ethylpuupehenol (24).



- 23 : R = α -CH₃;
 24 : R = α -C₂H₅;
 25 : R = β -CH₃;
 26 : R = β -C₂H₅;




15 α -Cyanopuupehenol
X



Ilimaquinone
18A

(b) **Puupehenone (19)** on being treated with a large excess of H₃C MgI in an ethereal medium, significant decomposition of (19) took place, another **stereochemically different isomer of (23)** was isolated, which has been named as **15 β -methyl puupehenol (25)**. In an identical treatment with ethyl magnesium iodide (H₅ C₂ MgI), gave rise to the corresponding **15 β -ethylpuupehenol (26)**. However, till date no logical explanation could be forwarded for the formation of the two isomeric species *viz.*, (25) and (26).

Protection of the *two* free -OH moieties present in (23) and (24) by means of acetylation yielded the **stable diacetyl derivatives (27) and (28)** as given below:




- 27 : R = α -CH₃;
 28 : R = α -C₂H₅;

10.3.2.3 Cyanide and Methoxide Nucleophiles: 1, 6-Conjugate Addition

The probability of the interaction between **puupehenone (19)** with hydrogen cyanide (HCN) in the biological systems formed the basis of an extensive and intensive studies *in vitro* under different aqueous and methanolic experimental parameters. Investigations, revealed that 1, 6-conjugated nucleophilic addition of HCN was essentially accompanied by oxidation in an environment where there was no restriction with regard to the availability of air/oxygen to the reaction mixture.

Puupehenone (19) in a methanolic solution on being subjected to complete saturation with 100 molar excess of absolutely dried HCN gas produced a quantitative yield of 15 α -cyanopuupehenol (X). In order to render the compound (X) significantly stable, it was duly acetylated to give rise to **O-19, 20-diacetyl-oxy-15 α -cyanopuupehenol (29)**.




Interestingly, with a view to initiate and simulate the natural living environmental conditions of the sponge, **puupehenone (19)** was strategically combined to a *basic-sorbent*, **Florisil*** pH 8-8.5 in water, duly suspended in distilled-water and completely saturated with pure HCN gas. Thus, it resulted into the formation of **15 α -cyanopuupehenol (X)** and **15-cyanopuupehenone (30)** eventually.

It has been observed that **15 α -cyanopuupehenone (30)** was a secondary metabolite obtained along with **15 α -cyanopuupehenol (X)** in the **Verongid Sponge**. However, it was also synthesized successfully and quantitatively in a direct, spontaneous addition-oxidation of **15 α -cyanopuupehenol (X)** in an appropriate admixture of methanol and water at a slightly basic pH environment.

Quite contrarily, the acetylation of **15 α -cyanopuupehenone (30)** in the presence of dry pyridine gave rise to the formation of its corresponding **monoacetyl structural analogue (31)**.


The interaction of **puupehenone (19)** with **magnesium methoxide [(CH₃O)₂Mg]** resulted in the conjugate addition of methoxide nucleophile with the production of the corresponding monoacetylated product **15-methoxypuupehenol (32)**. Further acetylation of (32) with an appreciable large excess of acetic anhydride [(CH₃CO)₂O] gave rise to the corresponding diacetylated derivative (33), which was subsequently isolated as the exclusive major product. Interestingly, compound (33) may also be obtained by the direct addition of methanol to **puupehenone (19)** at an ambient temperature for a duration of 24 hours, and subsequently subjected to acetylation.



* **Florisil:** A basic-sorbent comprised of activated magnesium silicate-a hard porous granular substance available in variant pH ranges.

10.3.2.4 Nitroalkane Nucleophiles: Addition Reactions

Generally, the acidic α -hydrogen bearing compounds were logically employed as **Potential nucleophilic donors** for the addition reactions pertaining to the extension of 1, 6-conjugate system of puupehenone (19). The addition reactions were carried out by making use of **nitromethane** [$\text{CH}_3\text{—NO}_2$] and **nitroethane** [$\text{C}_2\text{H}_5\text{—NO}_2$] which were reacted with the stoichiometric proportions of **magnesium methoxide** [$(\text{CH}_3\text{O})_2\text{Mg}$] and the resulting generated nucleophiles were finally added to a solution of **puupehenone (19)** in pure dry benzene. The addition products thus obtained were subsequently subjected to acetylation with acetic anhydride [$(\text{CH}_3\text{CO})_2\text{O}$] and finally purified to obtain compounds **O-19, 20-diacetyl-Oxy-15 α -nitromethanepuupehenol (34)**; and **O-19, 20-diacetyl-Oxy-15 α -nitroethanepuupehenol (35)** as given below:



34 : R = H; O-19, 20-Diacetyl-Oxy-15 α -Nitromethanepuupehenol

35 : R = CH₃; O-19, 20-Diacetyl-Oxy-15 α -Nitroethanepuupehenol

10.4 SUMMARY

In short, the ‘**marine-world**’ evidently and explicitly enjoys the status of holding an enormous and tremendous potential towards the epoch making discovery of a plethora of altogether newer ‘**lead compounds**’ in the development of medicinally potent therapeutic agents that are active against a variety of parasites and infections ailments. The wide range of vat resources tapped from the marine fauna and flora that would certainly help in the evolution of previously known ‘**chemotypes**’ for stemming the influx of crucial drug-resistant microorganisms and insects. In order to explore, tap and above all exploit commercially the relatively virgin biological reserves exclusively depends on the use of rapid technological advancement towards their collection, preservation, identification, and characterization of trace quantum of essential secondary metabolites.

With the advent of recent development and notable advances legitimately accomplished in comparatively safer life-support systems; and amalgamated with most recent technologically advanced computer-aided **sophisticated a nalytical instrumentations***, such as: UPTLC; HPLC; LC-MS;

* **HPTLC** = High performance thin-layer chromatography.

HPLC = High performance liquid chromatography

LC-MS = Liquid chromatography-Mass spectroscopy

GC; GC-MS; NMR; 2 D-NMR; FTIR; UV; X-Ray Diffraction; and MS have turned this novel dream-like fiction into a stark reality.

FURTHER READING REFERENCES

1. Baslow, M.H.: '**Marine Pharmaceuticals**', Williams & Wilkins, Baltimore, 1969.
2. Baker, J., and V. Murphy: '**Handbook of Marine Sciences**', CRC Press, Cleveland, 1976.
3. Blunt JW and Munro MHG: '**Marinlit—A Database of the Literature on Marine Natural Products**', University of Canterbury, 1997.
4. Dewick, P.M.: '**Medicinal Natural Products**', John Wiley & Sons, Ltd., England, 2nd edn., 2002.
5. D. Chuck Dunbar: '**Discovery of Antimalarial Compounds from Marine Invertebrates**', Ph.D. Thesis, University of Mississippi, Oxford, 1996.
6. El Sayed KA *et al.*, *J. Agric. Food Chem.*, **45**: 2735, 1997.
7. Hoppe, H.A. and T. Levring: '**Marine Algae in Pharmaceutical Sciences**', Vol. 2, de Gruyter, Berlin, 1982.
8. Martin, D., and G. Padilla: '**Marine Pharmacognosy**', Academic Press, New York, 1973.
9. Nigrelli, R.F.: '**Biochemistry and Pharmacology of Compounds Derived from Marine Organisms**', Ann. NY Acad. Sc., New York, 1960.
10. Scheuer, P.J.: '**Chemistry of Marine Natural Products**', Academic Press, New York, 1973.
11. Tursch, B., J.C. : '**Chemistry of Marine Natural Products**', Vol.2., Academic Press, New York, 1978.
12. Webber, H.H. and G. D. Ruggieri: '**Food Drugs from the Sea Proceedings**', Marine Technol. Soc., Washington, DC, 1976.
13. Worthen, L.R.: '**Food Drugs from the Sea Proceedings**', Marine Technol. Soc., Washington, D.C., 1972.
14. Youngken, H.W.: '**Food Drugs from the Sea Proceedings**', Marine Technol. Soc., Washington, DC, 1969.

GC	= Gas chromatography
GC-MS	= Gas chromatography-Mass spectrometry
NMR	= Nuclear magnetic resonance
2D-NMR	= Two Dimensional-NMR (or COSY-NMR)
FTIR	= Fourier transform infrared spectrometry
UV	= Ultra-violet spectrophotometry
MS	= Mass spectrometry

11

Nutraceuticals

- Introduction
- Contemporary Nutraceuticals
- Phytochemicals as Nutraceuticals
- Further Reading References

11.1 INTRODUCTION

Nutraceuticals may be defined as—‘*any substance that may belong to a plant, food or an essential component of food providing definitive medicinal usefulness and health promotion as well as physiological benefits, and ultimately minimise the possible risk of the prevailing chronic disease significantly.*’

In the recent past, **nutraceuticals** have legitimately and overwhelmingly acclaimed enormous confidence, acceptance, and recognition amongst the mankind not only as a mere ‘**health promoting factor**’ but also as an excellent ‘**nutritional contributing factor**’. The ever-increasing comprehensive knowledge with respect to the ‘**natural inherent nutritive value**’ of a relatively large number of *food products*, their chemical constituents *vis-a-vis* biological values has virtually succeeded in combating the human ailments and sufferings to an appreciable extent across the globe.

The wonderful conceptualization of the phrase—‘**Mediterranean Cuisine**’ by the so called **Western Nutritional Wizards** have implanted ideas or habits by constant urging in their daily food-intakes of such items as: **srpouted beans and lentils, olive-oil-salad with raw vegetables, citrus juices, whole-wheat bread, red/white wines, and extensive usage of garlic, ginger, onion, mustard, and tomato pastes as food-additives.** On the contrary, the **Eastern Nutritional Experts** overwhelmingly advocate the profuse usage of such delicious food items and beverages as: **green-tea, unfermented palm-wine, brined and smoked fish, soyabean milk, soyabean yogurt (curd), fibrous raw fruits and vegetables.**

In a broader perspective one may consider the ‘**nutraceuticals**’ to exert their action synergistically to check deteriorating health conditions, protect cells, and finally to ward off human ailments significantly. In reality, the ‘**nutraceuticals**’ have strategically captured the noticeable professional curiosity of a host of disciplines, namely: health care, food and nutritional scientists, the processed food manufacturers, the pharmaceutical industries, and above all the dietary supplement conglomerates.

Relief from Major Diseases: Nutraceuticals are intimately associated with the **relief, prevention** and/or **treatment** of certain specific and well-known major diseases, such as: coronary diseases, diabetes, cancer, hypertension, nervous debility, impaired digestive problems, and the like. A few other related ailments are: acute arthritis, joint pains, osteoporosis, neural tube problems.

Emergence: Stephen Defelice* almost a decade ago vehemently opined that **nutraceutical** is a particular component, for instance: omega oil from salmon (a fish), chenopodium ambrosiodos, pudina oil, tulsi, neem, soy, and orange juice with calcium. It is an universal fact that both **'food'** and **'medicine'** are exclusively based upon *three* vital and important characteristic features, namely:

- (i) significance to health,
- (ii) efficacy, and
- (iii) safety.

Legacy: The history of **'nutraceutical'** dates back to more than 200 years, even though there have been significantly enormous wonderful discoveries accomplished in the field of medicines in the past six decades. The world recognizes the meaningful and excellent progress *vis-a-vis* tremendous achievements in the domain of **science** and **herbal medicines** specifically at a very early stage. The well-known Indian system of **'Ayurveda'** proved to be a main source and provided several techniques whereby many dreadful diseases could be eradicated/cured from a human being. An extensive survey of literature shall reveal that the **'nutraceuticals'** actually came into being around 1500 AD even when people were not really aware about the term **'nutraceutical'**. Several races in the world have generously and profusely contributed towards the phenomenal success, acceptance, and recognition of **'nutraceuticals'**, namely: Greeks, Africans, Chinese, Tibetans, Indians, Arabs, Japanese, Thais, Malaysians, Ceylonese, Burmese etc.

Functional Food Vs Nutraceutical

Functional Food may be defined as—**“the food usually prepared by the aid of ‘scientific intelligence’ using definite knowledge about its anticipated merit/usefulness.”** In this manner, the **functional food** essentially caters to the body with the necessary quantum of vital carbohydrates, fats, proteins and vitamins, etc, required for its normal as well as healthy survival.

Examples: Vegetables, rice, wheat, fruits, pulses, fish, eggs, poultry products, beef, meat etc.

Nutraceutical may be defined as—**“a food (or a portion of a food) that essentially provides distinct health and medical benefits, even including the prevention and/or treatment of a particular disease.”****

Examples: Fortified **'dairy products'** *eg.*, milk powder, baby food, malted milk food; **vitamic C enriched citrus fruits** *eg.*, orange juice, grape juice.

Dietary Supplement*:** Based on a number of criteria obtained from the **FDA/CF-SAN Resources**, one may define a **'dietary supplement'** as:

- intended solely to supplement the diet which essentially contains either one or more than one of such dietary requirements as: vitamin, mineral, herb or other botanical product, amino acid,

* Stephen Defelice MD., the Founder and Chairman of the private non-profit foundation for **'Innovation in Medicine'** Cranford, NJ, USA, first and foremost coined the term **'pharmaceutical.'**

** Brower V: **Nutraceuticals poised for a healthy slice of the healthcare market.**, *Nat. Biotechnol.*, **16:** 728-31, 1998.

*** FDA/CF SAN-Resources: Food and Drug Administration Web Site. Dietary Supplement Health and Education Act of 1994. [Available at: <http://vm.cfsan.fdagm/adms/dietsupp.html>.]

dietary ingredient for use by human being to supplement the diet by enhancing the total daily intake, or constituent extract, metabolite concentrate, or various combinations of these ingredients.

- represented for usage as a conventional food or as the exclusive component of a meal or diet.
- intended for ready ingestion in the form of a pill, tablet, capsule, or liquid form.
- labeled explicitly as a **‘dietary supplement’**.
- includes essentially such products as: approved **‘new drug’**, licensed biological product (*i.e.*, marketed solely as a dietary supplement etc.

11.2 PHYTOCHEMICALS AS NUTRACEUTICALS

In the recent past several newer terminologies have emerged gainfully, noticeably, and widely across the globe to represent strategically the innumerable **nutrients of the future**, namely: **‘Phytochemicals’**; **‘Nutraceuticals’**; **‘Phytonutrients’**; **‘Phytofoods’**, and **‘Functional Foods’**. Importantly, the advent of accumulated knowledge with respect to the **disease-preventing components** present specifically in the natural plant products used as foods are duly recognized as **‘nutraceuticals’**.

Nevertheless, **‘phytochemicals’** represent a rather more recent evolution of the terminology which categorically emphasizes the inherent plant-source of a relatively larger segment of these important **disease-preventing and protective** chemical entities. In true sense, the potential nutritional role for the **‘phytochemicals’** is gaining world-wide recognition and popularity based upon the aggressive on-going research activities specifically highlighting their wonderful and remarkable advantages. The term **‘phytonutrient’** expatiates the natural chemical compound’s status as **‘quasinutrient’**. May be in the near future one would soon encounter the so called **‘phytochemicals’** most justifiably as the **‘essential nutrients’**.

Researches and other authentic scientific evidences have overwhelmingly revealed that the longevity of our ancestors perhaps could be due to their **correct importance of diet to health** *i.e.* devoid of today’s so called ‘junk-food’, foods laced with chemical additives, preservatives, flavour enhancers, emulsifiers, stabilizers, and the like, which ultimately prevented them succumb to the **‘modern diseases’**. A lot of such evidences may be distinctly observed amongst the **‘centenarian tribes’** that live in remote **Andean villages** (*i.e.*, Andes Mountains), tribes living in Andaman-Nicobar Islands, tribes living in remote interior villages in African continent etc., who invariably embrace strict and rigid **‘traditional dietary practices’**. Obviously, these **‘people’** have been duly reported to live extraordinarily **much longer live-spans** which are certainly free of such dreadful human ailments like: heart diseases, cancer, arthritis, respiratory obstructions, impairment in eye-sight etc.

Based on the aforesaid statement of facts together with the stark reality that even as to date a certain small segment of people usually live as **‘naturally’** as do the said tribes in the remote Andean villages, researchers have adequately proclaimed **epidemiological*** evidence from **‘modern societies’** for definitive clues pertaining to the intricacies of the actual **diet-disease connection**. Importantly, such elaborated and intensive studies helped a long way to the **biochemical researchers** to identify

* **Epidemiology**: It is concerned with the traditional study of epidemic diseases caused by infections agents, and with health-related phenomena.

and establish certain '**phytochemicals**' that prominently aid the human body towards '**maintaining a perfect health**', and also '**combating diseases**'.

Health authorities* recommend an overall guideline for the consumption of such specific diets that are significantly rich in whole grains, fresh fruits, fresh vegetables, reduced fat intake, and animal-protein consumption.

The good number of **phytochemicals** that find their abundant and legitimate use as **Nutraceuticals** may be judiciously classified as enumerated under with typical examples and appropriate plausible explanations wherever necessary.

11.2.1 Terpenoids (or Isoprenoids)

Terpenoids (or Isoprenoid) may be defined as—'**any compound either biosynthesized from or containing isoprene** units, including terpenes, carotenoids, fat-soluble vitamins, ubiquinone, rubber and even some steroids.**'

The **Terpenoids** are classified into various categories as described below:

11.2.1.1 Carotenoid Terpenoids

Carotenoids refer to natural compounds that usually render corn yellow, carrots reddish-orange, and tomatoes red. Besides, they impart typical distinct natural characteristic colours to gold fish, flamingo, salmon, and also the autumn leaves (*i.e.*, when the nature's green **chlorophyll** get depleted significantly, the carotenoids and phenols remain). They are also found in **Bell-peppers** with different colours that invariably represent a **selection of carotenoids**, namely:

- (a) **Orange carotenoids** *e.g.*, α -, β -, and γ -carotene.
- (b) **Red carotenoids** *e.g.*, lycopene and astaxanthin.
- (c) **Yellow carotenoids** *e.g.*, lutein and zeaxanthin.

As to date approximately **600 carotenoids** have been duly isolated, purified, and characterized in the plant kingdom.

Important Points: The following **important points** need to be considered—

- Nearly half of approximately **fifty carotenoids** that are usually present in the '**normal human diet**' are adequately absorbed into the blood stream.
- Almost 30% of the **actual plasma carotenoids** are duly comprised each of **lycopene** and **β -carotene**.
- Conversion to **Vitamin A** is afforded exclusively by **α -carotene**, **β -carotene**, and a few **other carotenes** (but **not lycopene** or **lutein**).
- **Hypervitaminosis of Vitamin A** may not be caused due to the excessive intake of **α -carotene** or **β -carotene** by virtue of the fact that the ensuing conversion and absorption rates are exceedingly slow and sluggish.

Importantly, both **α -carotene** and **β -carotene** deemed to afford reasonable protection against two

* Drgsted LO *et al. Pharmacology and Toxicology*, 72 Suppl. 1: 116-35, 1993.


** **Isoprene:** An unsaturated branched chain, 5C hydrocarbon *i.e.*, the molecular unit of the isoprenoid compounds.

dreadful human diseases *viz.*, **liver cancer** and **lung cancer** as established by extensive and intensive animal and cell culture investigative studies.

- Free liberation of **carotenoids** *viz.*, **lycopene**, **β -carotene** from vegetables may be accomplished by heating, chopping and/or crushing.
- Significant absorption of **carotenoids** may be observed when they are associated with oils as these are practically insoluble in water.
- **Carotenoids** present in the blood-stream are found to be transported in the most lipid-rich (**LDL***) cholesterol particles; and, therefore, the tissues having relatively the most abundant **LDL receptors** invariably acquire the most **carotenoid**.
- **α -carotene** possesses essentially 50-54% of the **antioxidant activity** of **β -carotene**, whereas **epsilon carotene** bears 42-50% of the said activity.
- **Carotenes**, in general, also increase appreciably the **immune response** and afford adequate protection to **skin-cells against UV-radiation** (*i.e.*, incorporated in '**Sun-Creams**' profusely).

Following are some of the **typical examples** of **carotenoid terpenoids** together with their **characteristic features**, such as:

1. **Lycopene** [Synonyms: **$\psi\psi$ -Carotene**; (*all trans*-Lycopene):




Characteristic Features: These are as given under:

- (i) Red colouration of tomatoes, guava, papaya, watermelon, pink grape fruit
- (ii) **Lycopene** in the usual and common '**American Diet**' is often derived from **tomato-containing food products**.
- (iii) Naturally occurring ***trans*-Lycopene** gets poorly absorbed in the body.
- (iv) Conversion of the *trans*-isomer to the *cis*-isomer in the presence of **heat** and **light** actually enhances the latter's bioavailability.
- (v) Gets bound to the fibers intimately and tightly—as freed duly by high heat.
- (vi) **Lycopene** is found to be more soluble in oil than in an aqueous medium.
- (vii) **Tomato paste** affords four-fold '**bioavailability**' in comparison to the **fresh tomatoes**.
- (viii) A **powerful antioxidant** that categorically retards damage caused to DNA and proteins.
- (ix) Offers distinctly and appreciably much better **skin protection** against the **UV-light** than **β -carotene**.

* **LDL:** Low-density lipoprotein.

- (x) Almost represents **50%** of the total available **carotenoids** found in the **blood serum**.
- (xi) **Lycopene** specifically gets accumulated in the various segments of the human-body *viz.*, skin, adrenal glands, prostate glands, testes etc., thereby rendering adequate protection against **cancer**.
- (xii) Helps to reduce **LDL-cholesterol** levels considerably.
- (xiii) **Lycopene** noticeably arrests the **Insulin-like Growth Factor-1 (IGF-1)** stimulation of cancerous (tumour) growth.


2. **β -Carotene** [Synonym: **β , β -Carotene; Carotaben; Provatene; Solatene;]:**



Characteristic Features: The characteristic features are as given under:

- (i) It is a weak **antioxidant***, but prove to be strong against singlet oxygen.
- (ii) The supplements may enrich **LDL-cholesterol- β -carotene content** without affecting other **carotene variants** *e.g.*, **α -carotene; ϵ , ψ -carotene (or δ -carotene); γ -carotene**.
- (iii) It may significantly boost the activity of **Natural Killer (NK) immune cells**.
- (iv) It may appreciably cause **stimulation for the DNA-repair enzymes**.
- (v) It definitely provides distinctly better **cornea protection** against the **harmful UV-radiation** (from sun-rays or UV-tubes) in comparison to **lycopene**.

3. **α -Carotene:**



Characteristic Features: The characteristic features are:


- (i) It has proved to be almost **ten fold more anti-carcinogenic** in comparison to **β -carotene**.
- (ii) It distinctly increases the release of **immunogenic IL-1**** and **TNF- α *****

* **Antioxidant:** A naturally occurring or synthetic substance that helps protect cells from the damaging effects of oxygen free radicals, and highly reactive compounds formed during normal cell metabolism.

** **Immunogenic IL-1:** Immunogenic interleukin-1.

*** **TNF- α :** Tumour necrosis factor-alpha.


4. **Xanthophyll** [Synonyms: **Lutein**; **β , ϵ -Carotene-3, 3'-diol**; **Vegetable lutein**; **Vegetable luteol**; **Bo-Xan**;]:



Characteristic Features: The characteristic features of **xanthophyll (lutein)** are as stated under:

- (i) **Xanthophylls** are important by virtue of the fact that they seem to cause protection of **Vitamin A, Vitamin E** and **other carotenoids** from undergoing oxidation *in vivo*.
- (ii) Substantial evidence is emerging gradually that **xanthophylls** are highly tissue specific.*
Example: Cryptoxanthin, which appears to be highly protective of cervical, uterine, and vaginal tissues.
- (iii) **Lutein** and **zeaxanthin** almost comprise of nearly 50% of all **carotenoids** present strategically in the retina.
- (iv) Likewise, both **leutin** and **zeaxanthin** are the **carotenoids** exclusively present in the **macula**** of the eye.
- (v) It seems to protect the eye from macular degeneration and cataracts.
- (vi) It may also render protection against colon cancer.
- (vii) It imparts '**yellow colour**' to avagado, corn, and yolk of an egg.
- (viii) Abundantly found in **kale, spinach, watercress, and parsley**.

5. **Zeaxanthin** [Synonyms: **β , β -Carotene -3, 3'-diol**; ***all-trans*- β -Carotene -3, 3'-diol**; **Zeaxanthol**; **Anchovyxanthin**;]:




Characteristic Features: These are as given below:

- (i) **Zeaxanthin** and **lutein** are the only **carotenoids** that are located strategically in the macula of the eye.
- (ii) Both **zeaxanthin** and **leutin** are almost present in equal quantum in the macula.
- (iii) It essentially absorbs the damaging '**blue-lighty**'.
- (iv) It helps to protect the eyes from possible macular degeneration and cataracts.

* Parker RS, *J. Nutr.*: **119**: 101-4, Jan.-1989.

** A small spot of coloured area.

6. **Astaxanthin** [Synonyms: Ovoester; 3,3'-Dihydroxy-4, 4'-diketo- β -carotene]:



Characteristic Features: The various **characteristic features** of **astaxanthin** are as stated under:

- (i) It attributes a beautiful pinkish red colour to crab, salmon, and shrimp.
- (ii) **Astaxanthin** is found to be ten folds more powerful antioxidant in comparison to any other **carotenoid**.
- (iii) It not only augments the **T-cell production*** but also helps in the release of **cytokine**.
- (iv) It may also cross the **blood-brain barrier (BBB)** *i.e.*, it serves as a **brain-antioxidant**.
- (v) It possesses **water-soluble chemical entities** that eventually helps to release the trapped radicals to **Vitamin C**.


11.2.2 Non-Carotenoid Terpenoids

The **non-carotenoid terpenoids** usually do have rather simpler chemical structures. They may be categorized into the following sub-groups, such as:

11.2.2.1 Limonoids [or Terpene Limonoids]

In general, the **limonoids** are particularly found in the peels (*i.e.* outer skins) of several citrus fruits *e.g.*, oranges, mandarins, lemons, grape fruits etc., that specifically directed to the ultimate protection of the lung tissue. the above findings were duly established by means of an elaborated study employing a standardized extract of ***d*-limonene**, **pinene**, and **eucalyptol** which was found to be effective in removing grossly the '**congestive** mucus from the lungs of patients having a history of **chronic obstructive pulmonary disease**.

Example: *d*-Limonene [Synonyms: Cajeputene; Cinene; Kautschin]:




* **T-Cells:** T lymphocytes that are differentiated in the thymus and are virtually important in **cell-mediated immunity (CMI)**, besides in the regulation of **antibody-mediated immunity (AMI)**.

Characteristic Features: These are as stated below:

- (i) ***d*-Limonene** is found to be 45 times more anticarcinogenic in comparison to **hesperetin**.
- (ii) It specifically detoxifies the carcinogenic substances, and thereby promotes the ensuing **cancer cell apoptosis***.
- (iii) **Limonene** distinctly helps to promote the '**glutathione-S-transferase**' *i.e.*, refers to **detoxification by glutathione addition**.
- (iv) ***d*-Limonene** has a distinct **orange-like smell**; whereas ***l*-Limonene** possesses a **piney odour** (very much akin to **turpentine**).

11.2.2.2 α -Terpineol



Characteristic Features: These are as given below:

- (i) **α -Terpineol** gives rise to the peculiar carrot flavour to the fresh carrots.
- (ii) It causes critical arrest of the '**cell-cycle**' in the neoplastic cells (*i.e.*, cancer cells).

11.2.2.3 Saponins

Saponins represent a group of **amorphous colloidal glycosides** that usually form soapy aqueous solutions.

A number of '**Saponin Glycosides**' have been discussed in details in Chapter 4, and a few typical examples along with their cardinal usages have been given as under:

S.No. Name	Characteristic Features
(a) Shatavarin I-IV [Section: 4.2.8.1.3]	(i) Used as galactagogue to promote the flow of milk. (ii) Also employed as tonic and diuretic.
(b) Ginsenoside Rg [Section: 4.2.8.2.1]	(i) In Chinese System of Medicine as a general tonic, stimulant and carminative. (ii) It possesses ' antistress ' properties. (iii) It prolongs the life of elderly persons.
(c) Glycyrrhizin [Section: 4.2.8.2.2.]	(i) It is used as a flavouring agent in beverages and confectionary. (ii) Possesses remarkable expectorant properties.
(d) Amarogentin [Section: 4.2.10.2]	(i) It is used as a bitter tonic in anorexia. (ii) It also improves relatively the ' dull appetite '.

* **Apoptosis:** A pattern of cell death affecting single cells, marked by shrinkage of the cell, condensation of chromatin, and fragmentation of the cell into membrane-bound bodies that are eliminated by phagocytosis (*i.e.*, programmed cell death).

11.2.3 Polyphenolics [or Polyphenol Extracts]

Polyphenolics (or **Polyphenol Extracts**) essentially represent a host of **natural antioxidants**, used as **nutraceuticals**, and found in **apples, green-tea, and red-wine** for their enormous ability to combat cancer and are also thought to prevent heart ailments to an appreciable degree.

Polyphenolics may be judiciously classified as given below:


- (a) Flavonoid polyphenolics,
- (b) Phenolic acids,
- (c) Non-Flavonoid polyphenolics.

11.2.3.1 Flavonoid Polyphenolics


Flavonoids are actually **flavone-like** substances which are invariably **antioxidants** and sometimes as **anti-inflammatory agents**.

Mechanism of Action: Flavonoids exert their activity by carefully scavenging the **'free-radicals'** thereby giving rise to a fairly **'stable radical'** which in turn would undergo reaction with another **'flavonoid radical'** to yield **two non-radicals**.

Red wine contains *two* important constituents, namely: **Flavone**, and **Resveratrol** because it is prepared by carrying out the fermentation of the **'pulp'** along with the skin and seeds (though **'ultrafiltration'** is done to minimise both bitterness and astringency); whereas, the **white wine** lacks the said two components because it is usually made by pressing juice away from the solids.




Flavone



Resveratrol

Various **citrus fruits** invariably contain a host of **flavanoids**, such as: **rutin, hesperidin, and naringin**.



Rutin

Hesperidin

Naringin

A few typical examples of **flavonoid polyphenolics** are stated below together with their **characteristic features**.


11.2.3.2 Anthocyanins

The term **anthocyanin** was coined to assign the chemical substance actually responsible for attributing the typical colour of the cornflower (from the **Greek**: *anthos* = flower, and *kuanos* = blue) generally applicable to a group of water-soluble pigments contributing the **red, pink, mauve, purple**, blue, or violet colour of a plethora of fruits and flowers. All these wide-range of pigments invariably come into being as '**glycosides**' (*i.e.*, the **anthocyanins**), and their corresponding **aglycones** (*i.e.*, the **anthocyanidins**); and these are actually derived from the **2-phenylbenzopyrylium cation**, more frequently known as the **flavylium cation**. Interestingly, the nomenclature '**flavylium cation**' categorically emphasis the fact these molecules usually belong to the vast category of the '**flavonoids**' in the broader perspective of the said terminology.

Anthocyanins, whose distinct beautiful explicite colours invariably attract insects and birds; and, therefore, mostly play an important and vital role in the phenomenon of '**pollination** and '**seed-dispersal**'.

Anthocyanins, are water-soluble **glycosides** and **acyl-glycosides of anthocyanidins**. The structure of *six* major **anthocyanidins**, namely: **Cyanidin, Delphinidin, Malvidin, Pelargonidin, Peonidin, Petunidin**—are as given under:

1. **Cyanidin**: $R_1 = \text{OH}$; $R_2 = \text{H}$;
2. **Delphinidin**: $R_1 = R_2 = \text{OH}$;
3. **Malvidin**: $R_1 = R_2 = \text{CH}_3$;
4. **Pelargonidin**: $R_1 = R_2 = \text{H}$;
5. **Peonidin**: $R_1 = \text{OCH}_3$; $R_2 = \text{H}$;
6. **Petunidin**: $R_1 = \text{OCH}_3$; $R_2 = \text{OH}$;



Structures of Major Anthocyanidins


Proanthocyanidins (or **Pycnogenols**) are found to be the short-chained colourless polymers of **anthocyanidins** which usually release the **anthocyanins** either with the **application of heat** or on being subjected to **acidic hydrolysis**.

Characteristic Features: These are as stated under:

- (i) **Anthocyanins** usually occur in several varieties of **berries** *e.g.*, blackberries, blueberries, black raspberries. In **blueberries** their content actually increases as they get ripened.
- (ii) White grapes invariably are devoid of any colour due to the fact that they do not have **anthocyanins**.
- (iii) Abundantly present in **Green Tea** which is consumed profusely by the Chinese and the Japanese.
- (iv) **Anthocyanins** usually occur in conjunction with the **phenolic acids** in a good number of berries.
- (v) **Anthocyanins** remarkably protect the **endothelial cells*** from undergoing **oxidative damages** specifically.

* **Endothelial Cells:** The layer of epithelial cells that lines the cavities of the heart, the serous cavities, and the lumina of the blood and lymph vessels.

11.2.3.3 Catechin [Synonyms: Catechol; Catechinic Acid; Catechuic Acid; Dexcyanidanol; Cyanidol; Catergen;]




The family of **polyphenolic compounds** usually comprise of *two* types of molecules, namely:

- (a) **Monomeric molecules**—which includes: **catechin** and **epicatechin**, and
- (b) **Dimeric molecules**—that includes: **procyanidins B1 and B2**.

Characteristic Features: The various **characteristic features** of **catechin** are as enumerated under:


- (i) It is an **antioxidant** found in dark, chocolate.
- (ii) It is thermo-labile and hence gets lost while drying grapes to obtain raisins.
- (iii) **Catechin** particularly affords inhibition of the **catechol-O-methyl transferase norepinephrine degradation**.
- (iv) It categorically **enhances the metabolic rate** *i.e.*, helps to **'burn fat'** while substantially increasing the production of free-radical.
- (v) **Catechin** can prevent and stop the initiation/progression of neoplasms (cancer).
- (vi) It can protect against the DNA damage; and, therefore, may prove to be of great help for patients undergoing post-operative **radiation therapy** or **chemotherapy**.
- (vii) It is an **active constituent** present in **tea** [*Thea sinensis* (Linne) O. Kuntze] *e.g.*, **green tea solids** range between 15-20%, and **black tea solids** vary between 5-10%.
- (viii) It has been duly observed and established that (–) **epigallocatechin 3-O-gallate [EGCG]** is the **most abundant polyphenolic ingredient** found in **green-tea**.



EGCG


- **EGCG** may enhance the basal metabolic rate.
- **EGCG** inhibits the nitration of protein.
- **EGCG** is available in cranberries, and not in black tea.

- (ix) **Catechins** undergo polymerization to yield **tannins** present in **black tea**.
- (x) Tea drinking is intimately associated with lowered incidence of cancer in ovary, prostate gland, stomach, colon, and the buccal cavity.
- (xi) **Catechins** inhibit NF- κ B transcription of the ensuing **proinflammatory** and **antiapoptotic** (*i.e.*, **cancer-promoting**) **genes**.
- (xii) **Theaflavins*** and **Thearubigins**** are the **tannins** obtained from tea, and represent the orange-red/black polymers.



Theaflavine

11.2.3.4 Isoflavones [or 3-Phenylchromone]



Daidzein: $R_1 = R_2 = H$; $R_3 = OH$;

Genistein: $R_1 = R_3 = OH$; $R_2 = H$;

Glycitein: $R_1 = H$; $R_2 = OCH_3$; $R_3 = OH$;

Phytonutrients of this phenol subclass, **isoflavones**, usually are derived from **beans** and other **legumes**, which are more or less related to the **flavonoids**. Interestingly, the **isoflavones** invariably function very much similar to the **flavonoids** wherein they predominantly and effectively block the enzymes that essentially promote the tumour growth. The relatively important and well-known isoflavones are **Genistein** and **Daidzein** which are abundantly found in **soy products** and the herb *Pueraria lobata* (Kudzu). People who regularly consume traditional diets that are rich in soyfoods invariably do not suffer from **breast**, **uterine**, and **prostate cancers**.

Pueraria lobata, over the years, have gained immense recognition and popularity as an essential aid for those who consume 'alcohol' by virtue of the fact that it seems to change the prevailing activity of the specific **alcohol detoxification enzymes** *i.e.*, the speed and momentum at which the **alcohol dehydrogenase** converts **alcohol** into the corresponding aldehydes. The ultimate result is a reduced tolerance for alcohol, and lowering in the '**pleasure response**' to drinking it.***

* Obtained from black-tea extracts.


** Weakly acidic class of orange-brown phenolic pigments produced during the fermentation step of tea manufacture.

*** Xie CI *et al. Alcohol Clin Exp. Res.*, **18**: 1443-7, Dec.-1994.

Characteristic Features: These are as given under:

- (i) **Isoflavones** are found in most concentrated manner in **soy beans** viz., **daidzein, genistein,** and **glycitein** (upto 2-4 mg/g).
- (ii) They are commonly found in **legumes** and **pomegranate seeds; besides, parsley and grains.**
- (iii) The '**genistein**'—an **isoflavanone** inhibits specifically the **tyrosine kinases** involved in **tumorigenesis* (or oncogenesis).**
- (iv) They help in the elevation of **HDL cholesterol** i.e., **good cholesterol.**
- (v) They specifically lower the **LDL cholesterol** i.e., **bad cholesterol.**
- (vi) **Isoflavones** usually serve as **potent antioxidants** against both **hydrogen peroxide** and **superoxide dismutase** (it serves as an anti-inflammatory and radioprotective agent).
- (vii) **Isoflavone** possesses **estrogenic-like qualities** (i.e., regarded as **phytoestrogen).**
- (viii) In fact, the **lignans** and **isoflavones** are recognized as the two major groups of **phytoestrogens.**
- (ix) They also go a big way in reducing and management of **menopausal irregularities.**
- (x) **Isoflavones** cause prevention of **osteoporosis** (i.e., bone resorption) in **post-menipausal women.**
- (xi) **Isoflavones** derived from **Soy bean** shown to inhibit the **prostate cancer cells** by almost **30%.**
- (xii) **Genistein** may prevent breast cancer to a certain extent, but promote the existing breast cancer.

11.2.3.5 Hesperetin




Characteristic Features: These are as given under:

- (i) **Hesperetin** represents the major flavonoid in oranges and other citrus fruits.
- (ii) It serves as an '**antioxidant**' which helps to regenerate **Vitamin C** *in vivo*.
- (iii) It gradually slows down the proliferation of the neoplasm (cancer) cells.
- (iv) **Hesperetin** also specifically slows down the replication of various **viruses** e.g., **influenza, herpes,** and **polio.**

* The production or causation of tumours.


11.2.3.6 Naringin



Characteristic Features: The various **characteristic features** are as enumerated under:

- (i) **Naringin** provides the **grape fruit** its inherent characteristic bitter taste.
- (ii) It may also increase the ability to **‘taste’** by direct stimulation of the taste-buds.
- (iii) **Naringin** particularly reduces **LDL cholesterol** content in blood, but without affecting the **HDL cholesterol**.
- (iv) It may, however, interfere with the so called **‘intestinal enzymes’**, thereby increasing appreciably the **oral-drug absorption**.
- (v) **Naringin** is found to increase significantly the **lipid and alcohol metabolism** in the liver while enhancing the **liver antioxidant activity**.
- (vi) It helps in a big way to protect against the alcohol-induced stomach ulcers.
- (vii) **Naringin** protects against the radiation-induced damages in the body to a great extent.
- (viii) It also possesses **antiapoptotic properties***.

11.2.3.7 Rutin [Synonyms: Rutoside; Melin; Phytomelin; Eldrin, Ilixathin; Sophorin; Globularicitrin; Paliuroside; Osyritrin; Osyritin; Mytricolorin; Violaquercitrin; Birutan;]




Characteristic Features: These are as follows:

- (i) **Rutin** is usually found in buckwheat, asparagus, and a variety of citrus fruits and grapes.
- (ii) It has been observed that there is practically very little loss in drying grapes to raisins.
- (iii) **Rutin** is found to strengthen the capillary walls.

* Properties concerned with **programmed cell death**.


11.2.3.8 Quercetin [Synonyms: Meletin; Sophoretin; Cyanidenolon]



Characteristic Features: The various **characteristic features** of **quercetin** are as follows:

- (i) **Quercetin** is a '**flavonol**' and usually found in high concentrations in apple skins, red onions, red grapes, green tea, and buckwheat.
- (ii) It usually does not suffer any '**loss of content**' in drying grapes to raisins.
- (iii) It is established to serve as a structural backbone to the **citrus flavonoids**', such as: **rutin** and **hesperetin**.
- (iv) **Pycnogenols**, a colourless substance, are duly obtained *via* **oligomerization** of **quercetin**.
- (v) It serves as an efficient and **strong antioxidant**, and also causes reduction of **LDL oxidation** largely.
- (vi) **Quercetin** is found to act as a **blood thinner**, and as a vasodilator.
- (vii) It can kill certain **specific viruses** *e.g.*, **herpes**.
- (viii) **Quercetin** possesses '**antihistaminic activity**' and thereby may relieve **allergic symptoms** caused due to pollen grains, food allergens, dust, dust mite etc.
- (ix) It is found to specifically inhibit the **COMT* enzyme** thereby causing significant reduction of the ensuing **epinephrine breakdown.****
- (x) **Quercetin** evidently possesses **sirtunin-like** deacetylase activity.

11.2.3.9 Silymarin [Synonyms: Apihepar; Laragon; Legalon; Pluropon; Silarine; Silepar; Silirex; Silliver; Silmar;]



Characteristic Features: These are as stated under:

- (i) **Silymarin** is an antihepatotoxic principle isolated from the seeds of the **milk thistle** and **antichokes**.
- (ii) It serves as an excellent '**protective**' against skin-cancer.
- (iii) **Silymarin** serves as a strong **antioxidant**, **anti-carcinogenic**, and **anti-inflammatory**.

* **COMT:** Catechol-*O*-Methyl Transferase.

** An increased **epinephrine** enhances **fat oxidation** and **energy expenditure** *i.e.*, "**thermogenesis**".


- (iv) It also serves as an effective **anti-atherosclerotic*** agent.
- (v) It specifically aids in the mobilization and digestion of fat.

11.2.4 Phenolic Acids

Phenolic acids are present abundantly in **cranberry juice** that essentially help in the reduction of particular adherence of organisms to the cells lining the bladder, and the teeth, which ultimately lowers the incidence of **urinary-tract infections (UTI)** and the usual **dental caries**. It has been duly observed that sweetening slow down the adhesion characteristic properties of the **phenolic acids**. Besides, they categorically reduce the oxidation of the LDL-cholesterol. Importantly, the **phenolic acids** significantly minimize the formation of the specific **cancer-promoting nitrosamines** from the **dietary nitrites and nitrates**. However, the most vital and important **phenolic compounds** that are found in **grapes** (red wine, grape juice, raisins) invariably comprise of **proanthocyanidins, resveratrol, and ellagic acid**.

A few typical examples of ‘**Phenolic Acids**’ are briefly described below along with their characteristic features:


11.2.4.1 Ellagic Acid [Synonyms: Benzoic Acid; Lagistase;]



Characteristic Features: The **characteristic features** of **ellagic acid** are as given under:

- (i) **Ellagic acid** is found to rich in **strawberries**, but 50% more in **raspberries** (chiefly as **ellagitannins**).
- (ii) It particularly lowers the incidence of oesophagal and colon cancers.
- (iii) **Ellagic acid** specifically inhibits the formation of **DNA adducts**.
- (iv) It categorically causes the inhibition of **Phase-I Enzymes**, and potentiates **Phase-II Enzymes**.


11.2.4.2 Chlorogenic Acid



Characteristic Features: These are as enumerated below briefly:

* Inhibits expression of the adhesion molecules.


- (i) It is present in relatively higher concentration in **bell peppers, tomatoes, and blueberries**.
- (ii) **Chlorogenic acid** is usually found in the **pulp of grapes** together with **ellagic acid** as given below:



Ellagic Acid

- (iii) It is most frequently found as an ester of **caffeic acid**.
- (iv) In fact, **caffeic acid** (*i.e.*, hydroxycinnamic acid) helps to minimize the mutagenicity of polycyclic aromatic hydrocarbons.
- (v) It serves as a major contributor to the antioxidant activity of coffee.
- (vi) **Caffeic acid** can regenerate specifically the oxidized Vitamin E.
- (vii) Chlorogenic acid may prove to be a pro-oxidant in the propagation phase of **LDL-oxidation**.
- (viii) The **roasting of coffee beans** actually enhances the prevailing **antioxidant activity** to an appreciable extent.

11.2.4.3 *para*-Coumaric Acid [Synonyms: *p*-Hydroxy cinnamic acid; β -[4-Hydroxy phenyl] acrylic acid;]



Characteristic Features: These are as follows:

- (i) It is usually present in high concentration in both green and red bell peppers.
- (ii) ***para*-Coumaric acid** serves as an antioxidant for the color mucosa.
- (iii) It serves as a flavonoid precursor.
- (iv) It is found to get bound with nitric acid and its derivatives before they usually get combined with protein amines to result into the formation of '**nitrosamine**'.

11.2.4.4 Phytic Acid [Synonym: Alkaloverf]




Characteristic Features: These are as stated under, namely:

- (i) **Phytic acid** is invariably found in legumes and whole grains.
- (ii) It is usually found in high concentration in **flaxseed** and **wheat bran**.
- (iii) It especially **binds minerals** like: Ca^{2+} , and Fe^{2+} .
- (iv) Perhaps the **mineral chelation** would reduce the free radicals.
- (v) It helps to lower the **Ca^{2+} ions absorption** from the gut.
- (vi) It is observed that the digestion of starch *in vivo* gets reduced appreciably; and, therefore, lower the blood glucose-level considerably.

- (vii) The **iron-binding phenomenon** (*i.e.*, chelation with Fe^{2+}) helps to slow down the cancerous growth; and hence, minimises the cardiovascular disease significantly.

11.2.4.5 Cinnamic Acid [β -Phenylacrylic Acid]




Characteristic Features

- (i) It is found abundantly in balsam Peru or Tolu, coca leaves, and oil of cinnamon.
- (ii) It is responsible for attributing the typical cinnamon's characteristic odour and flavour.
- (iii) **Cinnamic acid** possesses antifungal, antibacterial, and antiparasitic properties.
- (iv) It has proved to be a '**building block**' for the **lignans**.
- (v) It is found in higher concentrations in the balsam tree resins, wood, and the inner bark.
- (vi) Importantly, when combined with **flavonoids** and **benzoic acid** derivatives to result into the formation of **tannins** and **pigments** that specifically impart to the '**vintage wines**' the *colour* and *bouquet*.

11.2.5 Non-Flavonoid Polyphenolics

The **non-flavonoid polyphenolics** do not contain the **benzopyran nucleus**. A few such compounds, such as: **curcumin**, **resveratrol**, and **lignans** shall now be discussed in the sections that follows:

11.2.5.1 Curcumin [Synonyms: Turmeric Yellow; CI Natural Yellow; CI 75300; Diferuloylmethane;]



Characteristic Features: The various **characteristic features** of **curcumin** are as stated under:

- (i) **Curcumin**, a phytochemical, is the major component of the spice '**turmeric**' occur naturally as rhizomes.
- (ii) **Curcumin** particularly helps in the inhibition of the '**gene**' that actually gives rise to the formation of the **inflammatory COX-2 enzymes**, and ultimately preventing their production overwhelmingly.*
- (iii) It is reported to be both strongly anti-inflammatory and strongly antioxidant.
- (iv) **Curcumin** causes appreciable inhibition for the release of the proinflammatory cytokine **TNF-alpha****
- (v) It is highly regarded to be the more effective anti-clotting agent in comparison to acetyl salicylic acid (*i.e.*, **aspirin**) without having any **ulcer-inducing stomach irritation** caused by the latter.


* The allopathic drug '**Celebrex**' only inhibits the **COX-2 enzymes**.

** **TNG-alpha**: Tumour Necrosis Factor-Alpha.

- (vi) It may cause prevention of the color cancer.
- (vii) It predominantly acts as a scavenger for the **peroxynitrite free-radical**.
- (viii) **Curcumin** blocks particularly the **amyloid*- β -aggregation** that may eventually help to prevent **Alzheimer's Disease, Hodgkin's disease, and carcinoma**.
- (ix) It distinctly inhibits **NF-kB transcription** of the **proinflammatory and antiapoptotic** (*i.e.*, neoplasm-promoting) genes.

11.2.5.2 Resveratrol

Resveratrol depicts the strongest **sirtuin-like deacetylase action** of any known chemical constituent derived from the plant kingdom. In fact, the **sirtuins** have been shown to overwhelmingly extend the ensuing lifespan of **yeast and fruit flies**. Contrary to excessive media propaganda and representations, there exist a plethora of other sources of **resveratrol** in addition to the alcoholic beverages (red wine) *e.g.*, **purple grape-juice**.



Resveratrol

Importantly, in diverse organisms, calorie restriction invariably slows the '**pace of ageing**' (*i.e.*, helps the rejuvenation process) and thereby enhances maximum lifespan significantly. It has been duly established that in the budding yeast *Saccharomyces cerevisiae*, the prevailing calorie restriction specifically extends the '**lifespan**' by increasing the activity of **Sir 2**, which being a bonafide member of the conserved **sirtuin family of NAD⁺ dependent protein deacetylases**. Also included in this family are **SIR-2, 1**, a *Caenorhabditis elegans* enzyme that eventually modulates '**lifespan**'; and **SIRT 1**, a **human deacetylase** which essentially promotes cell survival by negatively regulating the **p 53 tumour suppressor**.

However, it has been amply demonstrated that the **potent activator resveratrol**, a well-known polyphenol found in red wine, appreciably lowers the **Michaelis constant** of **SIRT 1** for the **acetylated substrate** as well as the **NAD⁺**, and increases cell survival by stimulating **SIRT 1-dependent deacetylation of p 53**. However, in yeast, resveratrol usually mimics the calorie restriction by stimulating **SIR 2**, thereby enhancing both **DNA-stability** and **extension in lifespan upto 70%**.

Characteristic Features: The **characteristic features** of **resveratrol** are given as under:

- (i) **Resveratrol** is usually found as the '**principle stilbene**' in grapes, in teas (*i.e.*, both green and black), peanuts, and berries.
- (ii) It is produced by plants within themselves as defense against fungi.
- (iii) **Resveratrol** serves as an anti-inflammatory agent, and also inhibits **COX-1 enzyme**.**

* **Amyloid:** A protein-polysaccharide complex having starchlike characteristics produced and deposited in tissues during certain pathological states.

** **COX-1:** It is a constitutive enzyme and plays a role in the production of essential **prostaglandins (PGE)**. Inhibition of this enzyme by all the older, **non-selective NSAIDs** is primarily responsible for a number of their side effects.

- (iv) It aids in the blockade of adhesion of blood cells to vessel walls.
- (v) It has been duly demonstrated that **resveratrol** distinctly reduces skin as well as breast cancer in mice.
- (vi) It also inhibits **NF-kB transcription** of proinflammatory and **antiapoptotic*** genes.

11.2.5.3 Lignans

Lignans refer to the plant products of low molecules weight produced primarily by the oxidative coupling of **para-hydroxyphenyl propene units**, wherein the two units may be linked together by an oxygen bridge. Nevertheless, the monomeric precursor units are, namely: **cinnamic acid, cinnamyl alcohol, propenyl benzene, and allylbenzene**. The terminology **Lignan** or **Haworth Lignan** is applied to such chemical entities that are derived from the coupling propenyl and/or allyl derivatives are termed as **Neolignans**.**

Characteristic Features: These are as enumerated under:

- (i) **Lignans** occur abundantly and have been obtained from the **roots, heartwood, foliage, fruit, and resinous exudates of plants**.
- (ii) They invariably represent the **dimer-stage intermediate** between **monomeric propylphenol units** and **lignin**.
- (iii) **Lignans** are found to be **optically active**.
- (iv) The **cinnamic acid dimers** do have the **2-unit composites**.
- (v) **Lignans** actually strengthens the **plant cell-walls** (*i.e.*, wood).
- (vi) They are mostly water-soluble, and not-oil soluble.
- (vii) As to date **flaxseed** is established to be the richest **dietary source for the lignans**.
- (viii) **Podophylotoxin lignan** *i.e.*, a recognized '**cytotoxic agent**', is mostly used to treat the venereal warts.
- (ix) It is regarded to be the '**phytoestrogens**'.
- (x) **Lignans**, in general, may reduce the '**risk of cancer**' in females.

11.2.6 Glucosinolates [or Thioglucosides]

Glucosinolates, formerly known as **thioglucosides**, are the '**anionic glycosides**' solely responsible for the potent and characteristic inherent flavours of numerous *Brassicaceae* (*viz.*, mustard, radish, rutabaga, cabbage); besides, a good number of species related to other botanically close families, such as: *Capparidaceae*, *Resedaceae*, and *Tropaeolaceae*. The exact content of **glucosinolate** markedly varies with respect to the species, the plant part(s), the cultivation, and the climate conditionalities. It invariably ranges, before cooking: from 0.5 to 1 g.kg⁻¹, and may reach upto 3.9 g.kg⁻¹ as could be seen in certain **Brussel Sprouts**.

Basic Structure: The basic structure of the '**glucosinolates**' essentially consists of: (a) **glucose residue**, (b) **sulphate function**, and (c) **variable aglycone**, and the resulting molecule occurring as the respective potassium salt. It has been duly observed that the structural diversity of the '**glucosinolates**' distinctly reflects that of their precursor amino acids. A few typical examples are

* **Antiapoptotic:** Neoplasm-promoting.

** Goltleib OR: *Fortschr.Chem.Org.Naturst.*, **35**: 1-72, 1978.

given below that expatiates the formation of **sinigrin**, **gluconasturtiin**, **glucobrassicin**, and **sinalbin** from the respective amino acids:


S.No.	Amino Acid	Intermediate	Glucosinolates	Source
1	Homomethionine	Allylglucosinolate	Sinigrin	Black Mustard
2	Homophenylalanine	Phenethylglucosinolate	Gluconasturtiin	Watercress
3	Tryptophan	3-Indolylmethyl-glucosinolate	Glucobrassicin	Cabbage
4	Tyrosine	<i>p</i> -Hydroxybenzyl-glucosinolate	Sinalbin	White Mustard

General Formula and Postulated Origin of Glucosinolates:

The **biosynthesis of the glucosinolates** are most probably and logically accomplished by the following sequential steps, namely:

- Decarboxylation of amino acids to the corresponding aldoximes,
- A '**sulphur atom**' (from cysteine) is duly incorporated into the aldoxime,
- Resulting product gets **glycosylated** in the presence of **UDP-glucose**, and
- The **glycosylated product** is finally **sulphated by phospho-adenosine-phosphosulphate**.

The various sequential steps as explained above in sections (a) through (d) may be summarized as given under:



Potentials of Glucosinolate: The various **glucosinolates** described earlier may prove to be of immense beneficiary help to human health. A survey of literatures* have amply revealed that the dietary intake of **glucosinolates** viz., from various vegetable sources as: broccoli, cabbage, cauliflower, and Brussel sprouts might exert a substantial protective support against the **colon cancer**. Importantly, the **isothiocyanates** and the **indole-3-carbinols** do interfere categorically in the **metabolism of carcinogens**.

Mechanism of Action: The **isothiocyanates** and the **indoles** cause inhibition of **procarcinogen activation**, and thereby induce the '**phase-II**' enzymes, namely: **NAD(P)H quinone reductase** or **glutathione S-transferase**, that specifically detoxify the **selected electrophilic metabolites** which are capable of changing the structure of **nucleic acids**.

Example: Regular consumption of **Brussel sprouts** by human subjects (upto $300 \text{ g}\cdot\text{day}^{-1}$) miraculously causes a very fast (say within a span of 3 weeks) an appreciable enhancement in the **glutathione-S-transferase**, and a subsequent noticeable reduction in the urinary concentration of a **specific purine metabolite** that serves as a **marker of DNA-degradation**.


Glucosinolates may be further classified into *two* major groups, namely: (a) **Isothiocyanates**, and (b) **Indoles**, which shall now be treated individually in the sections that follows:

11.2.6.1 Isothiocyanates

In general, the **isothiocyanates** are solely responsible for causing the hotness of **horseradish, radish, and mustard**. **Isothiocyanates** do have the $[-\text{N}=\text{C}=\text{S}]$ functional moiety present in the chemical constituents. **Mustard oil** essentially comprise of the **allyl thiocyanate**.

A few typical examples of the **isothiocyanates** shall be discussed briefly as under.

11.2.6.1.1 Phenethyl Isothiocyanate




Phenethyl Isothiocyanate

Characteristic Features: These are as given under:

- (i) **Phenethyl isothiocyanate** offers a '**bitter taste**' to watercress.
- (ii) It causes effective inhibition of **tumorogenesis** by the aid of polycyclic aromatic hydrocarbons.
- (iii) It also induces **apoptosis** by **caspase-8 activation**, and **not by p 53**.
- (iv) It is particularly good against the harmful **nitroamines** in **tobacco smoke**.
- (v) **Nitrosonicotine**, a major carcinogenic substance, is present in the **tobacco smoke**, and formed by the interaction of **nitric oxide** and **nicotine**.


11.2.6.1.2 Sulforaphane [Synonym: Raphanin;]



Sulforaphane

Characteristic Features: They are as stated under:


- (i) It is obtained from the seeds of radish *Ramphanus sativus* L., (*Cruciferae*)
- (ii) **Sulforaphane** is especially rich in **broccoli**.
- (iii) It has been proved to be an extremely potent **phase-2 enzyme inducer**.
- (iv) It predominantly cause specific **cell-cycle arrest** and also the apoptosis of the neoplasm (cancer) cells.
- (v) **Sulforaphane** categorically produces **δ -D-gluconolactone** which has been established to be a significant inhibitor of the **breast cancer**.

 δ -D-Gluconolactone

11.2.6.2 Indoles

The typical example of **indoles** is briefly discussed below:


11.2.6.2.1 Indole-3-Carbinol



Indole-3-carbinol

Characteristic Features The various **characteristic features** of **indole-3-carbinol** are as follows:

- (i) It is found to be the most vital and important indole present in **broccoli**.
- (ii) **Indole-3-carbinol** specifically inhibits the **Human Papilloma Virus (HPV)** that may cause **uterine cancer**.
- (iii) It blocks the **estrogen receptors** specifically present in the breast cancer cells.
- (iv) Interestingly **indole-3-carbinol** downregulates **CDK6**, and upregulates **p21** and **p27** in prostate cancer cells.
- (v) It affords **G₁ cell-cycle arrest** and **apoptosis of breast and prostate cancer cells** significantly.
- (vi) It enhances the **p 53 expression** in cells treated with **benzo [a]pyrene***.




Benzo [a] pyrene

- (vii) It also depresses **Akt, NF-kappaB, MAPK, and Bel-2** signalling pathways to a reasonably good extent.

* It is reasonably anticipated to be a human carcinogen: **Ninth Report on Carcinogens** (PB 2000-107509, 2000) p III-187.

11.2.7 Thiosulphinates [or Cysteine Sulphoxides]

The major flavour component of **garlic** (*Allium sativum*; Liliaceae/Alliaceae) is a **thiosulphinat**e called **allicin**. **Allicin** is duly formed when the garlic tissue is damaged due to the hydrolysis product of **S-allyl cysteine sulphoxide (alliin)** which is specifically produced by the **pyridoxal phosphate-dependent enzyme allinase**.



Allicin
[Diallyl Thiosulphinat)e]

The various garlic preparations used medicinally include steam-distilled oils, garlic macerated in vegetable oils (e.g., soybean oil), dried garlic powder, garlic oil in soft gelatine capsules (known as ‘**garlic pearls**’), and gel-suspensions of garlic powder. However, careful analyses amply indicate a wide variations in the nature as well as quantum of the ‘**active constituents**’ in the variety of available preparations. Therefore, the freshly crushed garlic gloves typically consists of **allicin** (upto 0.4%), and other **thiosulphinates** (upto 0.1%-chiefly **methyl thiosulphinat)e**).

Characteristic Features: The **characteristic features** of the **thiosulphinates** are as enumerated under:

- (i) **Thiosulphinates** are designated as ‘**organosulphur phytochemicals**’ that are abundantly present in **garlic*** and **onions**.
- (ii) It also comprises of **mercapto cysteins** together with **allylic** sulphides**.
- (iii) Actually the very presence of the ‘**allylic sulphides**’ contribute to the strong typical characteristic odour of garlic.
- (iv) **Allicin** protects garlic from the pests.
- (v) **Allicin** is found to be toxic to insects and the microorganisms.
- (vi) **Allicin** categorically affords protection against the ulcers by causing inhibition of *Helicobacter pylori*.
- (vii) **Allicin** is found to lose its stability as soon as it gets removed from **garlic**.
- (viii) **Allicin** particularly inhibits the proliferation of the cells present in the mammary glands, color, and endometrial.
- (ix) Garlic may reduce blood pressure in human beings.
- (x) Garlic can augment the induction of the **nitric-oxide synthetase activity**.
- (xi) **Garlic** distinctly causes the inhibition of the **platelet aggregation** by the **arachidonic acid, epinephrine, and other platelet agonists**.
- (xii) The incised onions usually releases a chemical component known as: **Propanethial-S-oxide** which gets converted to sulphuric acid in the eyes itself thereby causing a real ‘**burning sensation**’.

* **Garlic**—has more sulphur content in comparison to onions.

** **Allyl**—refers to a hydrocarbon bonded to a S-atom.

(xiii) Any type of heating (*i.e.*, cooking) totally destroys the **enzyme allinase** that certainly prevents the formation of many highly useful sulphur containing **chemical entities**.

11.2.8 Phytosterols

Phytosterols usually referred to as—‘**any sterol *i.e.*, steroidal alcohol, present in the vegetable oil or fat**’.

In a rather broader perspective **sterols** invariably occur in a large segment of plant species. It has been duly observed that both yellow and green vegetables do contain an appreciable quantum; and, of course, their respective seeds specifically concentrate the sterols. Importantly, an extensive and intensive research on these valuable **phytonutrients** has been adequately carried out upon the seeds of certain selected specimens, such as: **soybean, yams, rice, pumpkins, and specific herbs**.


Phytosterols invariably engage in competitive uptake of the **dietary cholesterol** in the **entire intestinal passage**. Besides, **phytosterols** have profusely demonstrated the capability to affect complete blockade in the uptake of cholesterol (to which they are intimately structurally related) and also facilitate its subsequent excretion from the body. In fact, **cholesterol** has been duly implicated ever since as a relatively dangerous risk factor associated with the cardiovascular disease.

However, it has been established beyond any reasonable doubt that the prevailing ratio between the **dietary phytosterols** and **cholesterols** was remarkably lower amongst the ‘**vegetarians**’ *vis-a-vis* the ‘**non-vegetarians**’. This particular observation overwhelmingly underlies the fact that **cholesterol**, *per se*, is not only the **marker of risk** for the aforesaid cardiovascular disease, but also its ensuing ratio with several other **modifying dietary components** may ultimately prove to be a definite better measure of risk factor*.

Interestingly, **phytosterols** invariably block the development of **tumours (neoplasms)** in **colon, breast, and prostate glands**. Although the precise and exact mechanisms whereby the said blockade actually takes place are not yet well understood, yet one may strongly affirm that **phytosterols** seem to change drastically the ensuing **cell-membrane transfer** in the phenomenon of neoplasm growth and thereby reduce the inflammation significantly.

β-Sitosterol seems to be the most befitting and typical example amongst the **phytosterols**, and this shall be dealt within the section that follows:

11.2.8.1 β-Sitosterol [Synonyms: α-Phytosterol; Cinchol; Cupreol; Rhamnol; Quebrachol; Sitosterin; Harzol; Prostasal; Sito Lande]



FOOD PHYTOSTEROLS	
FOOD	β-SITOSTEROL*
Peanut Butter	135
Cashew Nut	130
Almonds	122
Peas	106
Kidney Beans	91
Avocados	76

* milligrams per 100 gramms

* Nair P *et al.* *Am. Jr. of Clin. Nitri.*, **40**: (4 Suppl.): 927-30, Oct. 1984.


Characteristic Features: The various characteristic features of **β -Sitosterol** are as given below:

- (i) **β -Sitosterol** minimises the level of **cholesterol production** by the liver.
- (ii) It also blocks the **absorption of cholesterol** to a good extent.
- (iii) It appreciably **lowers the cancerous cell-growth** (as cholesterol is required by the cell membrane).
- (iv) **β -Sitosterol** causes specific inhibition of the **epithelial cell division** which may in turn drastically **reduce atherosclerosis**.
- (v) Its structure closely resembles to that of cholesterol.
- (vi) In fact, **β -stisterol** may be regarded as the **plant equivalent** of the **animal cholesterol**.


11.2.9 Anthraquinones

Anthraquinones are usually characterised by the presence of **phenolic** and **glycosidic** compounds, that are solely derived from **anthracene**. They do possess a variable degree of oxidation *viz.*, **anthrones**, **anthranols**: and designate the so called **anthraquinone glycosides**. In fact, these molecules *e.g.*, **chrysophanol**, **aloe-emodin**, **rhein**, and **emodin** have in common a double hydroxylation at positions C-1 and C-8.

The structures of some **anthraquinone** and **hydroxyanthraquinone** derivatives are as given below:



Structures of Anthraquinone and its Derivatives



Structures of Some Hydroxyanthraquinone Derivatives

However, the botanical distribution of the various species essentially containing the '**1, 8-dihydroxyanthraquinone glycosides**' is quite restricted, namely: *Liliaceae* (**aloe**); *Polygonaceae* (**rhubarbs**); *Rhamnaceae* (**cascara, buckthorn**); and *Caesalpinaceae* (**sennas**).


A few typical examples of **anthraquinones** *viz.*, **senna**, **barbaloin**, and **hypericin** shall now be treated individually in the sections that follows:

11.2.9.1 Senna

Senna the dried leaflets of *Cassia senna* L., essentially contains **Sennosides A and B, glucosides of rhein and chrysophanic acid**.*

* Fairbairne: *Planta Medica*, 12: 260, 1964.

Following is the structure of **Sennosides** *i.e.*, **anthraquinone glycosides** as observed in **senna** in almost in equal quantum, and also in the rhubarbs where the **Sennoside A predominates**.



Characteristic Features: The **characteristic features** of **sennosides** are as stated below:

- (i) **Sennosides** are **dianthrones**.
- (ii) They mostly act as purgative for the lower bowel.
- (iii) **Sennosides** specifically enhances the **peristaltic movement** in the colon (*i.e.*, large intestine).
- (iv) They possess an inherent nauseating taste.
- (v) **Sennosides** are distinctly contraindicated for the **hemorrhoids** or **inflammation**.

11.2.9.2 Barbaloin

The major constituents of the **Cascara Bark** are **cascarosides A and B**, that essentially contain both **O- and C-glucoside linkages**; and, therefore, represent a pair of optical isomers differing only in the stereochemistry of the C-glucoside bond. The acid hydrolysis does not cleave the C-glucose linkage, and instead gives rise to **barbaloin**, and a mixture of two diastereoisomeric forms, which have been named as **aloin A** and **aloin B**. Hydrolysis of the **O-glucose linkage** generates **chrysaloin**, sometimes also referred to as **deoxybarbaloin**. Both **barbaloin** and **chrysaloin** are also found in the bark, and are believed to be the breakdown products obtained by the enzymatic hydrolysis of the **cascarosides**.




Characteristic Features: These are as given below:

- (i) It acts as a laxative (lower bowel).
- (ii) It is largely obtained from the **Aloe Vera plant** which possesses a host of medicinal values, namely:
 - Controls diabetes
 - Protects Human Immune System
 - Prevents AIDS
 - Prevents Asthma
 - Protects Kidney Function
 - Protects Cough and Cold

- Improves Digestive System
- Cures Arthritis
- Manages Cholesterol
- Cleans up Toxins from Intestines
- Cures skin problems and Loss of Hair

11.2.9.3 Hypericin [Synonym: Hypericum Red]




Hypericin

Characteristic Features: The various **characteristic features** of **hypericin** are as enumerated under:

- It is obtained as a '**Red Pigment**' obtained from *Hypericum perforatum*, otherwise known as "**Saint John's Wort**".
- Hypericin** finds its usage as **analgesic** or to treat **neuralgic pain**.
- It is profusely used as a '**folk remedy**' against **anxiety, depression, and insomnia**.
- It is found to be free from **purgative properties**.
- It may sometimes be employed to treat **ulcers** and acute inflammation of the gut.

11.2.10 Glucosamine [Synonym: Chitosamine;]



Glucosamine

Glucosamine is found in **chitin, mucoproteins, and micropolysaccharides** *i.e.*, in the naturally occurring plant products. Besides, **glucosamine** is also found naturally in the human body, especially in **cartilage, tendons, and ligament tissues**. It forms an integral and essential part towards the production of **glycoaminoglycan (CAG)**, that actually constitutes a major segment of the cartilage tissue.* Importantly, it is not available in appreciable quantum in the diet; and, therefore, should be specifically synthesized in the body. As this ability declines progressively with the advancement in age, and hence predisposes the human body to **arthritis**. Several acute and serious types of disease conditions thus come into being, for instance: **osteoarthritis (OA), rheumatic arthritis (RA), and degenerative joint diseases** are becoming so common and widespread that they have more or less become a normal part of the **ageing process** and ultimately do affect the hips, knees, hands, and spine.

* Briffa J: *Int. J. Act. Comp. Med.*, **15**: 15-16, 1997.


Chondroprotection is the terminology invariably used when the GAGs, namely: **glucosamine**, are duly administered orally so as to protect these joints from undergoing further deterioration with the passage of time.*

Mechanism of Action: It has been duly established that **N-acetylglucosamine** causes inhibition of the enzyme elastase in a dose-dependent manner.** However, **elastase** plays a vital and important role in the breakdown of the **articular cartilage, ligaments, tendons, and bones in rheumatic arthritis (RA)**. Besides, **D-glucosamine**, the active principle of **glucosamine sulphate**, happen to be a comparatively small molecule which rapidly undergoes diffusion *via* all the **biological membranes**. It also exhibits a high degree of affinity for the specific cartilaginous tissue and is subsequently introduced right into the **proteoglycan molecules**. Thus, it serves as the most preferred building block for the meticulous synthesis of **GAGs**, and these in turn offer a reasonable and legitimate protection against the ensuing damaging effects of **NSAIDs** as well as steroids. Importantly, the overall manner whereby **glucosamine** acts against **joint degeneration and joint protection**.

Hexosamine Pathway gives rise to the formation of the **endogenous glucosamine**, that is intimately associated with the ensuing **insulin responses**. It has been duly established, based on experimental evidences, that **glucosamine** may go a long way to induce **insulin resistance via hexosamine pathway**. This net effect may be obtained even without the presence of high-glucose concentration, and fat-induced insulin resistance is similar affected by the presence of **glucosamine*****

From the survey of literature one may safely conclude that **glucosamine** is quite **effective, less toxic, safe**, and a **well-tolerated** alternative drug to **NSAIDs** in the particular treatment of joint degenerative ailments. The actual use of **glucosamine** is apparently widespread in Great Britain, United States, and Europe, and, hence, the pharmacists attached to hospitals and OTC in stores should certainly be informed with respect to its **nutritional supplement** for onward transmission to the '**actual consumers**' as its popularity is gaining momentum in geometrical proportion.

11.2.11 Octacosanol [Synonym: Octacosyl Alcohol]



Octacosanol is one of the constituents of the **vegetable waxes**. It is an exemplary of a '**nutraceutical**'. It is a 28 carbon-chain alcohol which is present in the superficial layers of fruit leaves, and skin of a plethora of plants as well as '**whole grains**'.**** In fact, a large segment of investigational studies based on **octacosanol** have widely made use of either **wheat germ oil** or **policosanols**. Interestingly, **policosanols** is a natural mixture of '**primary alcohols**' duly purified from the **sugar cane wax** having **octacosanol** as the **major component**.

Characteristic Features: These are as stated under:

- (i) **Athletes: Octacosanol** supplemented athletes distinctly showed a significant enhancement in the muscle girth measurements thereby ascertaining the formation of lean body mass.

* Gottlieb MS: *J Manipulative Physiolog. Ther.*: **20**: 400-14, 1997.

** Kamel M *et al. clin. Exp. Rheum.*, **9**: 17-21, 1991.

*** Hussain MA, *Eur. J. Endocrinol*, **139**: 472-5, 1998.

**** Kato S *et al. Br. J. Nut.*, **73**: 433-42, 1995.

Studies related to measured grip and chest strength (*i.e.*, an indication of body strength), and reaction time to both auditory and visual stimuli* revealed that exclusively the reaction time to a **visual stimulus** and **grip strength** after ingestion of 1,000 mcg of octacosanol for a duration of 60 days on regular basis.

(ii) **Parkinson's Disease (or Motor Neurone Disease):** Clinical studies with Parkinsonian disease subjects, at a dose level of 5 mg of **octacosanol**, for 90 days showed that they improved appreciably.**


(iii) **Lipid Metabolism:** Pons *et al.* (1993)*** showed that **hypercholesterolaemic patients** treated with **policosanol** 2 mg per day for 6 months showed extremely promising results, whereby the 'total cholesterol' was lowered significantly having an apparent safe-profile. Aneiros *et al.***** (1993) carried out the investigations to test the effect successive doses upon the **lipid profile** and **tolerability of treatment** on patient suffering from **primary hypercholesterolaemia** for a duration of 6 weeks initially with 2 tablets of **policosanol** (5g) **once daily**, and followed by 2 tablets **twice daily** for another phase of 6 weeks. They observed that the patients' **blood-lipid profile** were as given below:

Lower-dose level: Significant reduction in **total cholesterol level** and **LDL-cholesterol level**;
Higher-dose level: Further reduction of these *two* levels in patients.

Torres *et al.****** (1995) studied the effect of **policosanol** in patients having a history of **non-insulin-dependent diabetes mellitus (NIDDM)**, and enhanced **LDL levels** which may render a major **coronary-artery-disease risk factor**. Patients having **stable glycaemic control** were administered with **policosanol** (5 mg) twice daily for 3 months; and they showed a significantly lowered levels of total **cholesterol** and **LDL-C**. Hence, the above investigative studies evidently shows a prominent, plausible, and possible place of **policosanol** as a **definitive cholesterol lowering nutraceutical** in such patients having **controlled NIDDM**.

(iv) **Thromboxane Pathways:** Arruzazabala *et al.****** (1993) investigated the possibility that **policosanol** may substantiate the action of certain **lipid-lowering drugs** causing an effect on the **platelet aggregation**. Furthermore, **policosanol** at a dose level of 200 mg. kg⁻¹ also appreciably lowered the mortality with respect to the cerebral infarction in **gerbils**.***** Conclusively both of these studies would indicate the close involvement of **policosanol** in the **prostaglandin** and the **thromboxane pathways**.

11.2.12 Carnitine [Synonym: γ -Trimethyl- β -hydroxybutyrobetaine;]



Carnitine

* Saint-John M and Mc Naughton L: *Int. Clin. Nutr. Rev.*, **6**: 81-7, 1986.

** Snider SR: **Octacosanol in Parkinsonism**: *Ann. Neurol.*, **16**: 723, 1984.

*** Pons P *et al.* *Curr. Ther. Res.*, **53**: 265-9, 1993.

**** Aneiros E *et al.* *Curr. Ther. Res.*, **54**: 304-12, 1993.

***** Torres O *et al.* *Diabetes Care*, **18**: 393-7, 1995.

***** Arruzazabala ML *et al.* *Leukot. Essent. Fatty Acids*, **49**: 695-7, 1993.

***** Arruzazabala ML *et al.* *Ibid.*, **69**: 321-7, 1993.

Carnitine refers to an essential cellular component, which is sometimes referred to as **Vitamin BT**, although it is not generally considered as a Vitamin. In reality, if an amino acid derivative duly synthesized from the amino acids **lysine** and **methionine** present in the liver and kidney, from where it gets adequately released into the **systemic circulation**. Brain also partially synthesizes this chemical entity. Out of the total **carnitine** present in a human body, almost 98% is found in the specific cardiac and skeletal muscle. Importantly, **carnitine** may also be derived from the 'diet' viz., chiefly food of animal origin, and the subsequent quantum ingested therefrom virtually determines the **rate of absorption**.*

Stereochemistry: **Carnitine** is available in *two isomeric forms* viz., *d-* and *l-*forms, and the *l-isomer* only occurs in the nature. However, the racemic mixture *i.e.*, a mixture of *d-* and *l-*forms are invariably marketed as a **nutritional supplement**, which is claimed to be less safer in comparison to the *l-form*.**

Mechanism of Action: The **mechanism of action** of **carnitine** is solely governed by the fatty acid metabolism. The **biochemical reactions** of this **nutraceutical** are predominantly based on the reaction taking place between **carnitine** and the **acyl moieties** as given under:



Perhaps the above cited reaction would strongly support the analogy that **carnitine** is intimately associated with a plethora of **Coenzyme A-dependent pathways**.***

The first and foremost recognized action of **carnitine** was its direct and articulated involvement in the long-chain fatty acid oxidation at the very **mitochondrial level** thereby providing energy. Importantly, **carnitine** serves as a carrier of the **acyl and acetyl** moieties just across the mitochondrial membrane before the β -oxidation can effectively materialize so as to provide 'energy'.**** It is, however, pertinent to state at this point in time that the **emergence of this energy** represents the main pivotal source of energy in the **cardiac** as well as the **skeletal muscle**, thereby legitimately ascertaining the most important role of **carnitine**.***** Besides, **carnitine** is strategically responsible for possessing *two* other **cardinal metabolic functions**, namely:

- (a) Branched chain α -ketoacid oxidation, and
- (b) Specific detoxification of potentially toxic **Acyl-coA metabolites** from other pathways.

Carnitine is observed to be excreted as 'free carnitine' or as 'acyl carnitine' by the kidneys, whereby more than 90% being reabsorbed by the **proximal renal tubules** genuinely.*****

Advantage of Carnitine Supplementation in Haemodialysis Patients:

It has been duly established that **carnitine deficiency** in haemodialysis patients may also lead to serious cardiac problems, that can be even severe. Sakurabayashi *et al.****** (1999) meticulously carried out a clinical study thereby comparing the oral **carnitine** administration in **haemodialysis**

* Kletzmayer J *et al. Kidney Int.*, **55** (Suppl. 69): S 93-S 106, 1999.

** Li Wan Po A: *Pharm. J.*, 245: 388-89, 1990.

*** Brass EP and Hiatt WR, *J. Am. Coll. Nutr.*, **3**: 207-15, 1998.

**** Grandi M *et al. Int J Clin Pharm Res.*, **17**: 1437, 1997.

***** Kelly GS, *Alt Med Rev.*, **3**: 345-60, 1998.

***** De Vivo D *et al. Epilepsia*, **39**: 1216-15, 1998.

***** Sakurabayashi T *et al. Am J Nephrol.*, **19**: 480-4, 1999.


(HD) patients and the **controls** having neither renal nor cardiac diseases. As anticipated, before the actual study was commenced the **observed carnitine levels** in the patients undergoing HD were significantly lower in comparison to the controls. Two months after the regular **carnitine administration** in the said two groups of patients—there was a significant increase in the plasma levels in the patients, and exceeded those of the controls.

Besides, the faulty **myocardial fatty acid metabolism** as observed before the treatment started, that may eventually result in a heart failure in chronic renal patients, was rectified magnificently, thereby putting forward another plausible suggestion with respect to the advantage for the carnitine supplementation in the HD patients particularly.

Other Important Therapeutic Applications of Carnitine: Carnitine finds its abundant vital therapeutically variant applications, and a few important ones shall be enumerated as under:

- **Heart disease:** beneficial usage of **carnitine** for several heart conditions.
- **Angina:** supplementation with **carnitine** may be of great help in patients with **ischaemic heart disease***.
- **Congestive heart failure:** prolonged usage of **carnitine**, at a dose level of 2 g twice daily upto 12 months, showed distinct positive results in terms of improved cardiac events together with greater life-expectancy.
- **Cardiogenic shock:** **carnitine** acts as a protective means in the **cardiogenic shock****, thereby showing a marked enhancement in the overall survival time.
- **Future applications:** **carnitine** in the '**glucose metabolism**' and '**insulin deficiency**'; in **Rett Syndrome*****; HIV; muscle weakness in chronic fatigue symptom; beneficial to transfusion-dependent β -thalassemia major subjects, anorexia; diptheria; and male infertility.****

11.2.13 Capsaicin [Synonyms: Axsain; Mioton; Zacin; Zostrix;]



Capsaicin

Capsaicin represents the pungent principle in the fruits of various species of *Capsicum* (*Solanaceae*).

Characteristic Features: These are as follows:

- (i) **Capsaicin** is obtained from **paparika** and **cayenne**.
- (ii) It makes chilli peppers '**hot**'.
- (iii) It finds its usage as a '**pepper spray**' for riot control and self defense.

* When blood flow to the heart is reduced, **carnitine levels** in the myocardial muscle may get reduced even upto 40%.


** 'A state of severe tissue hypoperfusion resulting from underlying pump function.

*** 'A neurological disorder, affecting girls, which involves the progressive loss of intellectual and motor skills, resulting in severe mental retardation.

**** Kelly GS, *Alt Med Rev*, **3**: 345-60, 1998.

- (iv) It causes a **'burning sensation'** for the mammals, but not birds.
- (v) **Capsaicin** helps to stimulate the **neurons** for burning and abrasion sensation.
- (vi) It is insoluble in water, but soluble in oil and fat.
- (vii) It can impart a **'cool sensation in mouth'** when mixed with cold milk, alcohol, or ice-cream.
- (viii) The strength/potency of **capsaicin** never gets lowered either by cooking or freezing.
- (ix) **Capsaicin** specifically promotes apoptosis in the **pancreatic neoplasm cells**.
- (x) It exerts practically little effect on the normal pancreatic cells.
- (xi) **Capsaicin** may critically **relieve chemotherapy-induced neuropathy**.
- (xii) **Capsaicin** predominantly inhibits **NF-kB transcription** of the proinflammatory and antiapoptotic (*i.e.*, neoplasm-promoting) genes.

11.2.14 Piperine




Piperine

Piperine is obtained from **black pepper** (*Piper nigrum L*), and also in *P. longum L*, *P. retrofractum* Vahl. (*P. officinarum* C.D.C.);

Characteristic Features: The **characteristic features** of **piperine** are as stated below:

- (i) **Piperine** is abundantly found in **hot jalapeno peppers, peppercorns (black pepper)**.
- (ii) It distinctly enhances the intestinal absorption of foods *i.e.*, better digestion.
- (iii) It is also employed traditionally to mask the taste of the spoiling (putrifying) meat.

11.2.15 Chlorophyll



Chlorophyll a


Chlorophyll b

Chlorophyll is the nature's green pigment specifically found in the plant kingdom. However, the **higher plants** and **green algae** contain **Chlorophyll a** and **Chlorophyll b** in the approximate ratio

of 3 : 1. In fact, **Chlorophyll C** is found together with **Chlorophyll a** in a variety of **marine algae**.

Characteristic Features: These are as follows:

- (i) It is regarded to be the most abundant naturally occurring green pigment in plants.
- (ii) **Chlorophyll** represents the principal light-absorbing pigment used profusely in the **photosynthesis**.
- (iii) **Chlorophyll** seeks its name from the **Greek: chloros = ‘yellowish-green’**.
- (iv) **Chlorophyll** essentially contains the porphyrine ring which is very similar to the **‘heme’** (*i.e.*, **haemoglobin**) but contains **Mg** as the central element instead of **Fe**.
- (v) It forms certain definitive **compact molecular complexes** with some **carcinogenic substances**, such as: **aflatoxin-B1**, **polyaromatic hydrocarbons** (tobacco smoke), and the **heterocyclic amines** (smoked meat).
- (vi) It absorbs the red and the violet light more strongly.
- (vii) **Chlorophyll** present in the leaves undergoes the decaying process in autumn thereby leaving behind the **carotenoid** colours (*e.g.*, yellowish red, yellow).
- (viii) Chemically, **Chlorophyll a** bears a methyl ($-\text{CH}_3$) side chain, whereas **Chlorophyll b** has an aldehydic ($-\text{CHO}$) side chain.
- (ix) Plants generally comprise of both **Chlorophyll a** and **Chlorophyll b**.
- (x) The **cyanobacteria*** particularly is devoid of **Chlorophyll b**.
- (xi) In the UV-region the **chlorophyll a** absorbs **red light** more strongly, whereas **chlorophyll b** absorbs the **violet light** more predominantly as given below:



Chlorophyll Light Absorption Spectrum

* **Cyanobacteria:** These are the toxin-producing pond-seum organisms termed as **‘blue-green algae’**.

11.2.16 Pectin

Pectin refers to a **polysaccharide** substance usually located in the cell walls of all plant tissues that essentially functions as an **intercellular cementing material**. One of the richest sources of **pectin** is **lemon rind** or **orange rind** that comprises of nearly 30% of this particular polysaccharide.

Characteristic Features: These are as stated under:

- (i) It is available as the soluble fiber in **apples** which perhaps gives the feeling of fullness when eaten.
- (ii) **Pectin** invariably serves as an antidiarrheal agent.
- (iii) It possesses an unique property of getting bound to the sugars, and subsequently releasing them gradually as and when required thereby maintaining the blood sugar levels steady almost.
- (iv) It is found to lower cholesterol in the body.

11.2.17 Dominant Phytochemical Pigments


There are quite a few **dominant phytochemical pigments** or **phytochemical class** that critically provides the most exclusive source of the colouring matter attributed solely to the host of natural **fruits** and **vegetables** which directly or indirectly contribute a lot more as **nutraceuticals**. A good number of such well-known and **dominant phytochemical pigments** are as listed below:

Dominant Phytochemical Pigments


S.No.	Colour	Pigment	Fruit or Vegetable
1	Black	Thearubigens Anthocyanins	Black Tea Black berries
2	Blue/Purple	Anthocyanins	Blueberries, Concord grapes, Eggplant, and Plums
3	Green	Chlorophyll	Asparagus, Broccoli, Cabbage, Green Tea, Kale, and Spinach
4	Orange	β -Carotene β -Cryptoxanthin	Apricots, Carrots, Cantelope, Mangoes, Sweet Potatoes and Pumpkin Oranges, Tangerines
5	Red	Anthocyanins	Apples, Cherries, Crauberies, Pomegranates, Raspberries, Red grapes, and Strawberries
		Lycopene	Pink Grape fruit, Tomatoes, and Watermelon
		β -Cyanins	Beets
6	Yellow	Lutein, Zeaxanthin, Curcumin	Avocado, Corn, Turmeric (Rhizome)

11.2.18 Tocotrienols and Tocopherols


Tocotrienols usually occur in nature *viz.*, in **grains**, **palm oil** together with their related cousins—the **tocopherols**.




α -Tocotrienol
[From Wheat Bran]



β -Tocotrienol
[From Bran and Wheat Germ Oil]



α -Tocopherol
[From Green Vegetables,
Grains, Palm oil]



β -Tocopherol
[Naturally occurring form of
Vitamin E]


Characteristic Features: These are as enumerated under:

- (i) **Tocotrienols** seem to cause inhibition of the growth of breast cancerous cells, whereas the **tocopherols** fail to do so.
- (ii) The biological functionalities of **tocotrienols** and **tocopherols** are quite different.
- (iii) Biological activity of **tocopherols** resembles to that of **Vitamin E**.
- (iv) **Tocotrienols** have explicitly shown their distinct **cholesterol-lowering effect**.

11.2.19 α -Lipoic Acid and Ubiquinones

A. **α -Lipoic acid (Thioctic Acid) and Ubiquinones (Coenzymes Q)** have gained ample cognizance as most vital and important **antioxidants** which in turn exhibit their wonderful overall effect to extend the ensuing effects of other **antioxidants** as well.

α -Lipoic Acid [Synonyms: **Thioctic Acid; Protogen A; Acetate Replacing Factor; Pyruvate Oxidation Factor; Biletan; Thioctacid; Thioctan; Tioctan;**]:



α -Lipoic Acid (*d*-Form)


Characteristic Features: The characteristic features of **α -Lipoic acid** are as follows:

- (i) In terms of research, **α -lipoic acid**, is regarded to be the '**new kid on the block**' *i.e.*, **nutraceuticals**.
- (ii) It efficiently serves as a '**hydroxy moiety quencher**' (because of its ability to form '**esters**' rapidly).
- (iii) The sulphur-sulphur bond in the pentagonal ring system caters as the most reactive segment of the molecule.

- (iv) It is found to be active on both lipids and tissue fluids.
- (v) Besides, the available **hydroxyl moieties**, it also strategically scavenges **peroxyl, ascorbyl, and chromanoxyl moieties**.
- (vi) **Thioctic acid** serves as a '**protective agent**' of both **Vitamin E** and **Vitamin C** by virtue of the fact that it critically functions in both **lipid** and **water phases**.
- (vii) **α -Lipoic acid** articulately affords protection to **catalase, superoxide dismutase (SOD)**, and **glutathione**, which play a vital and important role in the **liver-detoxification activities**.*
- (viii) It essentially plays an important role in energy production.

B. Ubiquinones [Synonyms: coenzymes Q; Q-275; Sa;]:

Ubiquinones refer to a group of lipid-soluble benzoquinones directly involved in electron-transport in mitochondria *i.e.*, in the **oxidation of succinate** or **reduced nicotine adenine dinucleotide (NADH)** *via* the **specific cytochrome system**. It predominantly occurs in the majority of aerobic microorganisms, from bacteria to higher plants and animals.



Ubiquinone

Characteristic Features: These are as given under:

- (i) Coenzyme Q is recognised as a recently discovered antioxidant.
- (ii) It serves as an important means of energy production.
- (iii) The clinical aspects of Coenzyme Q reveals that it may be an useful cardiogenic.

11.3 CONTEMPORARY NUTRACEUTICALS

The modern state-of-the-art **nutraceuticals** *i.e.*, **contemporary nutraceuticals** are penetrating the world market based entirely on their legitimate merit(s), safer usages, high degree of efficiency, and above all the superb and excellent means of their therapeutic efficacy across the globe. One may observe the goodness of '**black raspberries**' which are found to be extremely rich in '**antioxidants**'; and therefore, may serve as a powerful tool in the fight against the most dreadful human ailment '**cancer**'. In the same vein one may also consider the so called '**green leafy vegetables**' that are used extensively in the **modern research** essentially comprising of fibre with an exceedingly high content of **galactose**—a carbohydrate that is earnestly believed to help categorically prevent the proteins known as '**lectins**' from getting bound to the inner lining of the colon (large intestine) and thereby causing serious and permanent damage.

In this specific context, a few important '**contemporary nutraceuticals**' shall be discussed briefly as stated under:

* Sumathi R *et al.* *Pharmacol Res*, 27: 309-318, May-June-1993.

11.3.1 Spiruline


Spiruline (*Spirulina platensis*)—a cyanobacterium having a beneficiary action upon the alimentary biochemical processes for the undernourished and convalescent subjects—is profusely employed in the form of its dry powder capsulated in hard-gelatin capsules as a nutritional supplement. The reasonably significant effect of the **spiruline supplement** is on account of its **relatively higher iron content**.*

The mechanism of action of **spiruline** may be attributed by virtue of the high amount in the lipid fraction of ω derivative *e.g.*, **γ -linolenic acid**.** However, the singular exclusive presence of ω -6 vividly records a distinct metabolic gain, because **disaturase enzyme** may not be available in sufficient quantum in the undernourished subjects.***

Spirulina, a commercialized seaweed, should pass the stringent toxicological criteria *viz*, Maximum levels of iodine [$< 5 \text{ g. kg}^{-1}$], toxic minerals [As: $< 3 \text{ mg. kg}^{-1}$; cd: $< 0.5 \text{ mg. kg}^{-1}$; Sn and Pb: $< 5 \text{ mg. kg}^{-1}$; Hg : $< 0.1 \text{ mg. kg}^{-1}$], and the dried seaweeds should pass the following microbiological criteria (per g): faecal *E. coli* < 10 , anaerobic microorganisms < 100 , aerobic microorganisms $< 10^4$, and *Clostridium* < 1 .****

11.3.2 Broccoli

Broccoli [*Brassica oleraceae italica*; **Cruciferae/Brassicaceae**] essentially contains **glucoraphanin**, the glucosinolate precursor of **sulphoraphane**, as given below, has been amply demonstrated to possess very useful medicinal characteristic features.



It has been observed that **glucoraphanin** helps to induct the specific **carcinogen-detoxifying enzyme systems**, and thereby accelerates the removal of the **xenobiotics**.***** Nevertheless, the young sprouted seedlings do comprise of approximately 10-100 folds as much **glucoraphanin** as the fully grown plant; and, therefore, **broccoli** may be regarded as the most valuable dietary vegetable supplement.

Glucoraphanin could prevent **breast cancer** and may also sabotage the uncontrolled cell division of **colon cancer cells**. However, the purely **synthetic compound** exerts its action very much akin to its **natural counterpart** usually known as **oxomate, sulphoraphane**, duly identified and recognized as a cancer preventive agent in **broccoli**.

* Kapoor R and Mehta U: *Indian J Exptl. Biol.*, **30**: 904-7, 1992.

** Decsi T and Koletzko B: *Nutrition.*, **16**: 447-53, 2000.

*** Koletzko B *et al.* *Eur J Pediatr.*, **145**: 109-15, 1986.

**** **Bulletin du Ministere des Affaires Sociales**: Text No: 1705, Nov. 28-1990.

***** **Xenobiotic**: A drug or other substance not normally found in the body.

Mechanism of Action: The body's production of **Phase II enzymes*** is substantially increased by the two aforesaid compounds whereby the enzyme has a tendency to cause detoxification of cancer-causing chemicals, and thus minimise the risk of **cancer**. Oxomate could be administered together with other '**antineoplastic agents**' like **tamoxifen** *i.e.*, a combination of **phytonutrients** and **drugs**, in an obvious attempt to **muster maximise protection**. Interestingly, **tamoxifen** is at present the only US-FDA approved drug recommended for **breast cancer prevention** particularly in the high-risk women. It actually exerts its therapeutic action by an altogether different mechanism than that of **oxomate**. It has been observed that **tamoxifen** aids a female subject essentially having an **oestrogen-dependent tumours**. However, it may further be expatiated that a drug exclusively based upon the **oxomate** would certainly help prevent cancer formation irrespective of the fact whether the particular tumour is either an **oestrogen-dependent** one or a **non-oestrogen dependent** one.

Stomach Cancer and Ulcers: Sulphoraphane, an active principle found in **broccoli** and **broccoli sprouts** has been observed to cause a bactericidal effect on the specific bacterium responsible for the plethora of **stomach cancers**. In fact, the causative organism, *Helicobacter pylori*, for specifically debilitating **stomach ulcers** as well as **stomach cancers** could be cured with antibiotics. Jed Fahey** has rightly advocated that—

“If future extended clinical studies reveal that a nutraceutical can either relieve or at least prevent diseases intimately associated with *H. pylori* in people, it could have significant public health implications not confined to the United States only but also around the world.”

Sulphoraphane showed its proven ability to protect cells against cancer by augmenting their production of **phase 2 enzymes**, *i.e.*, a family of proteins that are responsible for carrying out the detoxification of cancer-causing substances and also damaging the ensuing free-radicals. Nevertheless, its **antibiotic profiles** are not quite explicitly understood as to date, and are likely to take place *via* certain other mechanism.

Brassica Protection Products (BPP) a joint venture by **Fahey and Johns Hopkins University** developed and marketed specialized **chemoprotective food products** and also the **broccoli sprouts**.

11.3.3 Aloe Vera Gel and Aloe Juice

There are *two* botanical sources of '**Aloe**', namely:

- (a) **Cape Aloe** [*Aloeferox*. Miller].
- (b) **Curacao Aloe** [*A. vera* (L.) Burm. f: **Asphodelaceae**].

In the usual traditional manner, the juice which is made to flow almost spontaneously right from the incised leaves gets duly collected and carefully concentrated by simple boiling. Thus, the concentrated and thickened juice comprises of the **dark brown masses (Curacao Aloe)** and with distinct **greenish reflections (Cape Aloe)**. However, one may obtain the **Aloe Gel** after duly eliminating the outermost tissues of the leaf meticulously.


* A family of proteins which detoxify certain cancer-causing agents and damaging free radicals.

** **Jed Fahey:** A plant physiologist in the Department of Pharmacology and Molecular Sciences at the Johns Hopkins School of Medicine, USA.


Composition: The **Aloe Vera Gel** composition consists of a rich content of water, besides certain not-so-specific compounds as: **amino acids, enzymes, lipids, sterols**, and most of all, **polysaccharides** (*e.g.*, **pectins, hemicelluloses**).

Characteristic Features: The **characteristic features** of **Aloe Vera Gel** are as stated under:

- (i) It* contains upto 20 + 1% **hydroxy anthraquinone derivatives** *viz.*, Aloin A; α -L-Rha Aloinoside A; Aloin B; and α -L-Rha-Aloinoside B, as given under:



Aloin A: R = H;
Aloinoside A: R = α -L-RHa;



Aloin B: R = H;
Aloinoside B: R = α -L-RHa;

- (ii) **Aloe Vera Gel** appears to contain various antibacterial and antifungal chemical entities which may potentially delay or inhibit the growth of such microorganisms which are solely responsible for **food borne illness in humans** and **food spoilage** generally.
- (iii) It helps to preserve several fresh highly perishable fruits and vegetables *viz.*, table grapes, bananas, strawberries, apricot, peaches, plums, oranges and the like.
- (iv) The **Aloe Vera Gel** adequately affords **potential environmental benefits**. Alternatively, it could provide a greener perspective to sulphur dioxide (SO₂) and several other synthetic food preservatives which are invariably used on the agricultural produce thereby increasingly causing serious health hazards.
- (v) Manufacturers of this **naturaceutical** as health product make a wide spectrum of therapeutic advantages in humans ranging from: diabetes, human immune system, AIDS, digestive system, arthritis, cholesterol management/control, asthma, kidney protection, cough and cold, cleansing of toxins from intestines, skin manifestations, and hair loss.
- (vi) It is employed extensively in **cosmetic products** to serve as an extremely hydrating ingredient in liquid or creams, sun-lotions, shaving creams, lip-guards, face packs, healing ointments, and various protective creams.
- (vii) It may also be employed skillfully in the composition of **phytomedicines** traditionally utilized as an adjunct in the **emolient** and **antipruriginous** treatment of various skin disorders, as a trophic protective agent for cracks, abrasions, chaps, frost bite, insect-bites, superficial and limited burns, sunburn, and also for diaper rash (in babies).

Mechanisms: Several mechanisms have been duly invoked so as to explain the aforesaid activities, such as:

- stimulation of the complement linked to polysaccharides
- high water content imparting various sequential phenomena *viz.*, **hydrating, insulating, and protective properties to the gel.**

11.3.4 Soyfoods

Soyfoods are processed or semi-processed products prepared from **soya bean** or **soja bean** or **Lincoln bean** or **Manchurian bean** or **Chinese pea**. Soya bean is the seed of *Glycine max* (L.) Merrill, and several other species *viz.*, *G. Soja* Sieb & Zucc., *G. hispida* (Moench) Maxim., and *Soja hispida* Moench, belonging to the natural order **Leguminosae**.

A Chinese investigative study has adequately established the fact that regular consumption of '**Soyfoods**' may drastically reduce the risk of fracture in **postmenopausal women** specifically amongst those who are in the early years following the state of menopause.

It has also been proved beyond any reasonable doubt that there exists a definite linkage between the '**soyfood intake**' and **bone-mineral density**.

In the recent past, **soyboods** are distinctly enjoying an overall strong appeal as an integral component by virtue of the fact they are relatively **free** of the **artery-clogging trans fats**, that are duly formed when these fats are suitably hydrogenated to render them more solid and also extend their shelf-life significantly. Besides, the oil also continues to benefit from an ever-increasing awareness of the innumerable health properties of the antioxidant-rich oil.

11.3.5 Omega-3 Fatty Acids

The **omega-3 fatty acids** represent a class of fatty acids invariably found in '**fish-oils**'. It is especially and abundantly obtained from **Salmon** and some other **cold-water fish**. It has been amply proved and established that the **omega-3 fatty acids** duly help to lower the levels of **cholesterol** and **LDL** (*i.e.*, low-density lipoproteins) in the blood. **LDL** usually represent the so called '**bad-cholesterol**' present in the blood. It serves as a nutritional supplement as well as a natural source of the marine product. The **Omega-3 fish oil** is regarded to be a **nutraceutical**, which is a food providing health supplement. They provide an antiinflammatory action very much within the body that may prove to be extremely beneficial for the particular relief of **inflammatory disorders**, namely: **rheumatoid arthritis**. Generally, eating fish has been reported to cause protection against the age-related **macular degeneration** *i.e.*, a common eye ailment.

Commercial Product

Blackmores Anti-inflammatory Fish Oil-1000

[Manufactured By: Blackmores LTD., 23-Rosebery Street, Balgowlah, Auckland, New Zealand]

Active Ingredients Per Capsule:

Fish Oil (Natural)	1 g (1000 mg)
Containing Omega-3 marine triglycerides	300 mg as:
Eicosapentaenoic Acid [EPA]	180 mg
Docosahexaenoic Acid [DHA]	120 mg

11.3.6 Pomegranate Juice


The **pomegranate juice** is remarkably quite rich in **antioxidants**, for instance: **anthocyanins**, tannins, and soluble **polyphenols**, that articulately scavenge the ‘**free radicals**’ and thereby help to prevent specifically DNA damage which may ultimately lead to a number of serious health conditions.

It also possesses **antiatherosclerotic activities** *i.e.*, preventing the thickening of the arteries, and thereby slowed down the **oxidation of cholesterol** to almost 50%*.

Pomegranate Juice helps to prevent the ischemic **Coronary heart disease (CHD)****, thereby showing improved flow of blood to the heart by almost 17%. However, no apparent negative effects on lipids, blood-glucose level, haemoglobin A 1c, blood pressure, and above all the body weight of the subject.

11.3.7 Walnuts

In the recent past, the health-conferring benefits of the **walnuts** have shown that are a prominent source of the natural **antioxidant hormone melatonin**; and, therefore, its consumption would certainly help to boost up the blood levels of **melatonin** significantly.



Melatonin

Russel Reiter’s findings adequately demonstrated that the **Walnuts** substantially contain **melatonin**, which is duly absorbed when it is eaten. It also improves in patients the ability to **resist oxidative stress** caused on account of certain toxic molecules usually termed as ‘**free radicals**’.

Melatonin content in **walnuts** ranges between 2.5 to 4.5 ng per g.

The **melatonin supplements** are widely employed by the consumers (*i.e.*, patients) whose sleep patterns are found to be **irregular, engaged in shift work** or **suffering from jet lag**.

11.3.8 Certified Organic Mushroom Nutrace

In a broader sense, **Mushrooms** are quite valuable health food products, which are low in calories, high in vegetable proteins, iron, zinc, chitin, and profusely used in the **Traditional Chinese Medicine (TCM)**. In fact, their **legendary** effects on the promotion of good and sound health rightly suggest that the **Mushrooms** are **probiotic** in nature *i.e.*, they help in restoring **bodies health, balance, and natural resistance to disease**. The active principles present in them exert to boost the **immune system** *i.e.*, they do possess immune system enhancement properties.

A few typical examples of the **certified organic mushroom nutrace** are as stated under:

* *Clin. Nutr.*: 23(3): 423-33, June-2004.

** Researches carried out at: (a) California Pacific Medical Centre, and (b) University of California’s Non-profit Preventive Medicine Research Institute. **CHD** is the biggest cause of death in several Western countries.

11.3.8.1 Trimyo-Gen™ [Composed of: 33% *Cordyceps sinensis* (Winter Worm: Summer Grass), 33% *Ganoderma lucidum* (Reishi), and 33% *Schizophyllum commune*]

Importantly, the strain of *Cordyceps* was found high in the **Himalayan Mountains** for a short duration in summer, and growing on its natural host—a caterpillar. However, it is now scientifically cultured on organic whole grain substrates, and produced under stringent sterile quality control environment.

11.3.8.2 MycoPlex-7™

It is essentially a **Seven Mushroom Formulation** with a base of the most established **nutraceutical mushrooms**. These mushrooms are of immense use in both **Oriental and Chinese Traditional Medicine (TCM)**. It has been shown they exhibit **legendary** effects both as an **adaptogen** and **vitality enhancer**.

FURTHER READING REFERENCES

1. Arruzazabala ML *et al.*: *Leukot. Essent. Fatty Acids*, **49**: 695-7, 1993.
2. Brower V: **Nutraceuticals Poised for a Healthy Slice of the Healthcare Market**, *Nat. Biotechnol.*, **16**: 728-31, 1998.
3. **Bulletin du Ministere des Affaires Sociales**: Text No: 1705, p. 103, Nov. 28-1990.
4. *Clin. Nutr.*: **23**(3): 423-33, June-2004.
5. Hussain MA: *Eur. J. Endocrinol.* **139**: 472-5, 1998.
6. Kato S *et al.*: *Br. J. Nutr.*, **73**: 433-42, 1995.
7. Kelly GS: *Alternative Med. Rev.*, **3**: 345-60, 1998.
8. Parker RS: *J. Nutr.*, **119**: 101-4, Jan-1989.
9. Snider SR: **Octacosanol in Parkinsonism**, *Ann. Neurol.*, **16**: 723, 1984.
10. Sumathi R *et al.*: *Pharmacol. Research*, **27**: 309-318, May/June-1993.
11. Torres O *et al.*: *Diabetes Care*, **18**: 393-7, 1995.
12. Xie CI *et al.*: *Alcohol. Clin. Exp. Res.*, **18**: 1443-7, 1994.

12

Enzyme and Protein Drug Substances

- | | |
|---|---|
| <ul style="list-style-type: none">• Introduction• Enzyme Variants• Enzymes of Pharmaceutical Relevance and Utility• Brief Description of Enzymes Used as Drugs | <ul style="list-style-type: none">• Protein as Drug Substances• Introduction• Protein Variants• Brief Description of Proteins Used as Drugs• Further Reading References |
|---|---|

A. ENZYME AS DRUG SUBSTANCES

12.1 INTRODUCTION

An **enzyme** is usually defined as—'an **organic catalyst produced by the living cells but capable of acting outside cells or even *in-vitro***. In a broader sense, the **enzymes** may be regarded as **proteins** which categorically alter the rate of **chemical reactions** without requiring the aid of an **external energy source** or being **charged themselves**. Importantly, an enzyme may be able to catalyze a particular reaction several times effectively.

Specific Characteristic Features: The various **specific characteristic features** of **enzyme** are as enumerated under:

- (1) Generally, **enzymes** are reaction specific in that they act exclusively on certain substances (known as '**substrates**').
- (2) **Enzyme** and its corresponding substrate or substrates invariably give rise to an **enzyme-substrate complex**, which involves not only **physical shape** but also **chemical bonding**.
- (3) **Enzyme** helps in promoting the '**creation of bonds**' either between altogether separate substrates, or induces the cleavage of bonds in a single substrate to result into the formation of the product or products of reaction.
- (4) **Metabolism:** Numerous **enzymes** present in human body, whereby each catalyzing one of the several reactions which essentially occur as part of **metabolism**.
- (5) **Functionality:** It has been duly observed that each **enzyme** acts at an **optimum temperature** and a **pH**, at which it does function most efficaciously. For most **human enzymes**, these shall be particularly confined to such factors as: **pH of cells, body temperature, tissue fluid, and blood**.

- (6) **Impaired Activity: Impaired activity of enzymes** may be caused due to extremes of **pH, temperature, dehydration, UV-radiation**, and the presence of **heavy metals** *viz.*, Pb or Hg.
- (7) **Specific Requirements:** Certain enzymes specifically require the dire presence of **coenzymes** (*i.e.*, non protein molecules *e.g.*, **Vitamins**) to enable them function properly; whereas, still others require some **critical minerals**, such as: Fe, Cu, Zn).
- (8) **Proenzyme:** It has been observed that certain **enzymes** are obtained as **proenzyme** *i.e.*, in an inactive form; and, therefore, must be duly activated by appropriate means *viz.*, **inactive pepsinogen** is suitably converted to **active pepsin** by the help of hydrochloric acid (HCl) present in the **gastric juice**.
- (9) **Activity: Enzymes** do possess a variety of vital and important **activities**, a few of them shall now be discussed briefly as under:
- (a) **Digestive Enzymes:** These are usually most common and familiar. They are basically the '**hydrolytic enzymes**' which specifically catalyze the addition of water molecules to relatively bigger food-molecules to help them split into rather simpler chemical entities. Quite often the very name of the enzyme explicitly indicates the '**substrate**' with addition of the suffix-**ase**.
- Examples:** (i) **Lipase**—It splits **fat** (triglycerides) into the corresponding fatty acids and glycerol respectively.
- (ii) **Peptidase**—It splits **peptides** to the corresponding amino acids.
- Exceptions:** Certain **enzymes** *e.g.*, **pepsin** and **trypsin** do not usually end in **-ase**, because they were duly baptized much before this method of **nomenclature** was actually instituted.
- (b) **Enzymes for Synthesis Reactions:** The **enzymes for synthesis reactions** help to synthesize a host of **biological products**, such as: **glycogen, hormones, nucleic acids (DNA and RNA), phospholipids for cell membranes, and proteins**, most of them need one if not several enzymes.
- Examples: DNA Polymerase**—is essentially required for DNA-replication, that actually precedes **mitosis**.
- (c) **Energy Production:** It also specifically requires a plethora of enzymes.
- Examples:** Each and every step related to cell respiration needs essentially a particular **enzyme**, for instance: **cytochrome transport system, glycolysis, and Krebs cycle**.
- (d) **Deamination Reactions:** The **deamination reactions** are usually carried out by **deaminases** which critically remove the **amino moieties** from the available pool of excessive **amino acids** so that they may exclusively utilized for **energy**.
10. **Miscellaneous Activities of Enzymes:** These categorically include certain highly specific **enzymes** to perform a definite purpose *in vivo*, A few such typical examples are as given under, namely:
- (a) **Cessation of Long-chain Fatty Acids:** Specific **enzymes** aid in the splitting of long-chain fatty acids into relatively smaller compounds which in turn used up in the **cell respiration** mostly.
- (b) **Maintenance of Blood Pressure:** Specific enzymes are usually required for '**blood clotting**', and also for the formation of **angiotensin II** solely required to **maintain and raise the blood pressure**.

Another school of thought describes the ‘**enzyme**’ as—‘**a biocatalyst that essentially accelerates specific biological reactions.**’ Nevertheless, the recognized concept of biocatalysts is rather broad-spectrum, and generously embraces a variety of **pure substance(s), cell product(s), and cell extract(s)**, namely:

- pure enzymes,
- plant cells,
- animal cells,
- crude cell extract,
- microbial cells,
- intact non-viable microbial cells.

The sources of **enzymes** that are viable commercially range from **animals, higher-plants, and microorganisms.**

Following are certain prevalent and important **enzymes** exploited commercially that belong to the aforesaid categories as stated under:

- (a) **Animal Enzymes** *e.g.*, **lipases, rennets, trypsin** etc.
- (b) **Higher-plant Enzymes** *e.g.*, **amylases, papain, proteases, and soybean lipoxygenase.**
- (c) **Microorganisms** *e.g.*, *Acetobacter lacti, Clostridium aceticum.*

Sasson* (1984) critically advocated that certain **enzymes** are employed overwhelmingly in the **Food and Beverage Industries**, such as:

- Papain** : obtained from **papaya fruit** used mostly as **meat tenderizer**, and in making ‘**wort**’ from malt (to solubilize residual protein) in breweries to prepare **Beer.**
- Protease** : used profusely in the manufacture of **detergents**, and in **softening of leather** in **tanning industry.**

In the recent past, the **microbial enzymes** have legitimately acclaimed a wide popularity and overwhelming recognition. In fact, the spectacular production of both **primary and secondary metabolites** by the microorganisms could only be feasible by virtue of the involvement of a host of specific **enzymes.** Thus, one may classify the **enzymes** based upon their **site of action** as given below:

- (a) **Endoenzymes [or Intracellular Enzymes]:** The **enzymes** that are solely secreted very much within the cell are termed as **endoenzymes.** They essentially are involved in the apt synthesis of different **cellular components, food reserves,** and also serve as **bioenergetic materials.**** Importantly, as these various processes do occur in the intracellular zones, the **enzymes** involved are also strategically located in the **intracellular region.**
Examples: Isomerases, Phosphorylases, Synthetases etc.
- (b) **Exoenzymes [or Extracellular Enzymes]:** The **enzymes** that are exclusively secreted outside the cell are invariably known as **exoenzymes** or **extracellular enzymes.** They usually exert a digestive feature in their overall activity and function. Interestingly, they help in the hydrolysis of relatively complex molecules into much simpler compounds.

* Sasson A: **Biotechnology: Challenges and Promises**, UNESCO, Paris, 1984.

** **Bioenergetic Materials:** Such substances that liberate energy from food stuffs.

Examples: **Amylases**—hydrolyse **starch** components;
Lypases—hydrolyse **lipids** (*i.e.*, triglycerides); and
Proteoses—hydrolyse **proteins** into **amino acids**.

There is a plethora of **endoenzymes** that are specifically generated by **pathogenic** as well as **saprophytic* microorganisms**.

Examples: **Cellulase**—converts **cellulose** to **cellobiose**;
Polygalacturonase—
Pectinmethylesterase—
Polymethylgalacturonase—

Riviere (1977)** observed that the **endoenzymes**, namely: **asparaginase**, **invertase**, and **uric oxidase** are noticeably of much higher economic value, and also are quite difficult to undergo extraction because they are critically produced very much inside the cell.

12.2 ENZYME VARIANTS

Over the years a good number of **enzyme variants** have been duly identified and recognized for their specific usages as summarized in the following Table 12.1.

Table 12.1 Enzyme Variants and their Applications

S.No.	Type of Enzyme (Abbreviation)	Applications
1	Activating Enzyme	Catalyzes the attachment of an amino acid to the suitable transfer ribonucleic acid [t RNA].
2	Allosteric Enzyme	Alteration in activity caused due to certain types of effectors , known as allosteirc effectors, bind to a nonactive site on the enzyme.
3	Amylolytic Enzyme	Catalyzes the conversion of starch to sugar.
4	Angiotensin-converting Enzyme [ACE]	Converts angiotensin I (a segment of the renin-angiotensin-aldosterone mechanism of the kidney) to angiotensin II <i>i.e.</i> , the ultimate and final step in the renin-angiotensin mechanism . The latter helps to stimulate aldosterone secretion , and hence Na retention.
5	Autolytic Enzyme	Produces autolysis or cell digestion.
6	Branching Enzyme	Transfers a carbohydrate unit from one molecule to another <i>e.g.</i> , glycosyl transferase .
7	Brush Border Enzyme	Serves as the lining of the small intestine and produced by the cells of the villi and microvilli (brush border) .
8	Coagulating Enzyme	Catalyses the conversion of soluble proteins into insoluble ones (<i>Synonym: Coagulase</i>).
9	Debranching Enzyme	Removes a carbohydrate unit from molecules which essentially contain short carbohydrate units usually attached as side-chains <i>e.g.</i> , dextrin-1, 6-glucosidase .
10	Deamidizing Enzyme	Splits amine off amino acid chemical compounds.
11	Decarboxylating Enzyme	Separates specifically carbon dioxide [CO ₂] from organic acids <i>e.g.</i> , carboxylase .
12	Digestive Enzyme	Controls digestive processes in the alimentary canal.

(Contd.)

* **Saprophytic:** Any organism living on decaying or dead organic matter *e.g.*, **higher fungi**.

** Riviere J: **Industrial Application of Microbiology**, Survey Univ. Press, London, 1977.

13	Fermenting Enzyme	Brings about fermentation especially of the carbohydrates and produced by organisms or yeasts .
14	Glycolytic Enzyme	Catalyses the oxidation of glucose .
15	Hydrolytic Enzyme	Catalyses the phenomenon of hydrolysis.
16	Inhibitory Enzyme	Blocks or inhibits a chemical reaction.
17	Inverting Enzyme	Catalyzes the hydrolysis of sucrose.
18	Lipolytic Enzyme	Catalyses the hydrolysis of fats (triglycerides) <i>e.g.</i> , lipase .
19	Mucolytic Enzyme	Depolymerizes mucous by splitting mucoproteins , <i>e.g.</i> , lysozyme , mucinase (or hyaluronidase).
20	Oxidising Enzyme	Catalyses oxidative reactions [<i>Synonym: Oxidase</i>].
21	Proteolytic Enzyme	Catalyses the conversion of proteins into peptides.
22	Redox Enzyme	Catalyzes oxidation-reduction reactions.
23	Reducing Enzyme	Removes oxygen [O ₂] [<i>Synonym: Reductase</i>].
24	Respiratory Enzyme	Acts within tissue cells to catalyze oxidative reactions by releasing energy.
25	Splitting Enzyme	Facilitates removal of a portion of a molecule.
26	Transferring Enzyme	Facilitates the moving of one molecule to another chemical entity. [<i>Synonym: Transferase</i>].
27	Uricolytic Enzyme	Catalyses the conversion of uric acid to urea.
28	Yellow Enzyme	Involves particularly in the cellular oxidations <i>viz.</i> , a group of flavoproteins .

12.3 ENZYMES OF PHARMACEUTICAL RELEVANCE AND UTILITY

A survey of literature would reveal that there exist quite many enzymes that specifically possess well recognized pharmaceutical relevance and utility. Interestingly, most of them find their highly critical and pivotal role in the therapeutic armamentarium to serve as useful means to provide enormous help in curing human diseases. The following Table 12.2 includes a number of such enzymes that are invariably employed as ‘**drugs**’ along with their particular type, source(s), and applications.

Table 12.2 Drugs (Enzymes) with Pharmaceutical Relevance

S.No.	Drugs (Enzymes)	Enzyme Type(s)	Source(s)	Applications
1	Bromelain	Proteolytic	Stem of pineapple plant, <i>Ananas comosus</i> , [<i>Family: Bromeliaceae</i>].	Soft tissue anti-inflammatory agent.
2	Chymotrypsin	—do—	Pancreas of Ox, <i>Bos taurus</i> (<i>Family: Bovidae</i>).	In ophthalmology; and also as anti-inflammatory agent.
3	Collagenase	—do—	Fermentation of <i>Clostridium histolyticum</i> .	Debridement of skin burns and derma ulcers.
4	Deoxyribonuclease	Nucleolytic	Pancreas of Ox, Cow, Buffalo	Lowering the viscosity of bronchopulmonary secretions.
5	Fibrinolysin	Proteolytic	Human plasminogen;	Cure of thrombotic disorders.
6	Hyaluronidase	Amylolytic	Mammalian (Bovine) testes.	Enhancing of IM injections.
7	Muramidase	Mucolytic	Serum, tears, lungs of animals.	Antibacterial and antiviral agent.
8	Papain	Proteolytic	Latex of unripe fruits of tropical melon tree (Papaya) [<i>Carica papaya</i> (Caricaceae)].	Clarification of fruit juices, beer, and as meat tenderizer.
9	Pancreatin	Proteolytic, Lipolytic, and Carbolytic	Pancreas of Hog [<i>Sus serofa</i> (<i>Suidae</i>)].	Digestive agent for fat, protein and polysaccharides.
10	Pancrealipase	—do—	—do—	Chronic pancreatitis, and cystic fibrosis.

(Contd.)

11	Pepsin	Proteolytic	Glandular layer of fresh stomach of hog [<i>sus scrofa</i> (Suidae)].	Conversion of protein into peptone as well as proteose
12	Rennin (Chymosin)	—do—	Glandular layer of fresh stomach of calf [<i>Bos taurus</i> (Bovidae)].	Commercial manufacture of different types of processed cheese.
13	Seratiopeptidase	Proteolytic	Organism belonging to genus <i>Serratia</i> .	To increase the effects antibiotic due to its inherent anti-inflammatory activity. In thromboembolic disorders.
14	Streptokinase	Plasminogen activator	Culture filtrates obtained from β -hemolytic <i>Streptococci-Group C</i> .	
15	Urokinase	Fibronolysis	Kidney-tissue or human urine cultures.	Lysis of blood clots or fibrin in pulmonary embolism.
16	L-Asparaginase	Hydrolytic	Obtained from <i>Escherichia coli</i> , plant, animal tissues, fungi and yeast.	Interferes with growth of malignant cells.

12.4 BRIEF DESCRIPTION OF ENZYMES USED AS DRUGS

In the following sections a brief description of certain **enzymes** that are specifically used as **drugs** shall be dealt with relevant therapeutic applications;

12.4.1 Bromelain

Bromelain is a proteolytic enzyme particularly present in the pineapple plant. It is usually available as a buff coloured, odourless amorphous powder having poor solubility in aqueous and organic solvents. It is mostly indicated in the cure and treatment of oedema caused due to injury or surgery, and in acute soft-tissue inflammation.

12.4.2 Chymotrypsin

Chymotrypsin refers to the digestive enzyme produced duly by the pancreas and functioning in the small intestine that, with **trypsin**, hydrolyzes proteins to peptones or further. [**Trade names: Avazyme^(R); amd Enzeon^(R)**]. Each mg of **chymotrypsin** contains not less than 1000 USP units. **Chymotrypsin** is largely employed in **ophthalmology** for the critical dissection of zonule of eye-lens for the specific **intracapsular cataract extraction**. It is also indicated for the topical usage to reduce soft-tissue inflammation from abscesses, ulcers, fistulas, as well as necrotic injuries.

12.4.3 Collagenase

Collagenase represents an enzyme that induces certain specific changes in **collagen** to cause its respective degradation. It is invariably obtained from *Clostridium histolyticum* by its fermentation; and also bears the capability for the necessary digestion of both **denatured and native collagen**.

The actual potency of **collagenase** is usually given by its capability to digest **normal bovine collagen** *in vitro*, and displays its optimum activity between pH 7 and 8.

Collagenase is mostly used in the form of its '**ointment**' for the specific dedebriment of burns, dermal ulcers, and necrotic ulcers.

12.4.4 Deoxyribonuclease [DNase]

Deoxyribonuclease refers to an enzyme that hydrolyzes and thus depolymerizes **deoxyribonucleic acid (DNA)**.

DNase usually loses its activity in an aqueous medium. The optimum activity is exhibited between pH 6 to 7. It has been duly observed that Mg^{2+} ions are an absolute must for its activation.

DNase finds its abundant applications in:

- cure of haematomas and localized abscess formation,
- minimise the specific viscosity of the pulmonary secretions as its '**aerosol preparations**' in **Inhalers**, and
- increase the flow of the expectoration of sputum in typical bronchopulmonary infections.

12.4.5 Fibrinolysin

Fibrinolysin designates a **proteolytic enzyme** duly obtained from the activation of **human plasminogen** by the presence of **streptokinase**. It essentially aids a complicated system of biochemical reactions for the lysis of clots in the vascular system. The principal physiological activator of the **fibrinolytic system is tissue plasminogen activator**. It particularly converts plasminogen in a fibrin-containing clot to plasmin. The fibrin polymer is duly degraded by plasmin into fragments which are subsequently scavenged by **monocytes** and **macrophages**.

Fibrinolysin finds its use in the critical treatment of **thrombotic disorders** essentially caused due to its fibrinolytic inherent nature.

12.4.6 Hyaluronidase

Hyaluronidase is observed duly in the testes and semen. It specifically depolymerizes **hyaluronic acid**, thereby enhancing the ensuing permeability of the connective tissues by dissolving the various substances that hold body cells together. It also acts to disperse the cells of the **corona radiata** about the newly ovulated ovum, thus largely facilitating entry of the sperm.

A few other typical **applications** are as follows:

- reduces the viscosity of tissue cement thereby rendering the tissues easily permeable to tissue fluids; and this characteristic feature has been duly exploited to increase the rate of absorption of both IM and subcutaneous injectables in humans.
- employed in hypodermolysis as an essential aid to the specific subcutaneous administration of relatively large and excessive volume of parentals in patients.

12.4.7 Muramidase

Muramidase is normally found in blood cells of the **graunulocytic** and **monocytic** series. Its serum and urine level is enhanced in patients having acute or chronic leukemia. It is also mostly present in tears, sweat, and saliva.

Muramidase is observed to hydrolyze **mucopolysaccharides**, and is invariably found to be active in the transformation of **insoluble polysaccharides** of the cell wall to the corresponding **soluble mucopeptides**, particularly in the **Gram-positive organisms**.

By virtue of this unique inherent activity **muramidase** is usually administered IV for the treatment of bacterial or viral infections in human beings.

12.4.8 Papain

Papain refers to a **proteolytic enzyme** derived from the fruit of the **papaya**, *Carica papaya* (**Family: Caricaceae**). It is soluble in water and glycerine and possesses optimum activity ranging between pH 5 to 6. It is usually available as an admixture with **chymopapain**.

Papain finds its use in **medicine** as enumerated under:

- anti-inflammatory agent
- relieve symptoms of **episiotomy***

Papain finds its enormous usage in **industry**, such as:

- clarification of beverages *e.g.*, beer, fruit juices etc.,
- meat tenderizer,
- cheese processing as a substitute of **renin**,
- degumming of silk fibres in textile industry, and
- dehairing of animal skins and hides in leather industry.

1 NF unit of Papain \equiv **Activity released by 1 mcg of Tyrosine derived from a Standard Casein Substrate.**

12.4.9 Pancreatin

Pancreatin represents one of the active ferments of the pancreas essentially containing a mixture of **enzymes**, namely: **amylase**, **lipase**, and **protease**. It exhibits its maximum activity in an **alkaline medium**.

Pancreatin finds its abundant uses as stated under:

- used chiefly as a digestant,
- as it becomes **inactive** in an acidic medium, it should be administered orally in combination with a mild alkaline substance *e.g.*, NaHCO_3 (sodium bicarbonate),
- for the preparation of predigested or peptonized food, products, and
- conversions of **starch** into **dextrin**; **proteins** into **amino acids**; and **fats** into **fatty acids** and **glycerols**.

Potency of Pancreatin Its has been established that:

1 g of **Pancreatin** \equiv 12,000 Units of **Amylase Activity**;
 \equiv 15,000 Units of **Lipase Activity**; and
 \equiv 10,000 Units of **Protease Activity**.

* **Episiotomy**: Incision of the perineum at the end of the second stage of labour to avoid spontaneous laceration of the perineum and to facilitate delivery.

12.4.10 Pancrealipase

Pancrealipase represents a rather more concentrated version of **pancreatin** whereby the specific **lipase activity** is enhanced significantly. It is derived from the pancreas of the hog, *Sus scrofa* var. **domesticus** belonging to the natural order **Suidae**.

Potency of Pancrealipase: Pancrealipase has the following potencies:

1 g of **Pancrealipase** \equiv 100 Units of **Amylase Activity**,
 \equiv 24 Units of **Lipase Activity**, and
 \equiv 100 Units of **Protease Activity**.

Pancealipase finds its various applications in therapy, such as:

- cystic fibrosis,
- chronic pancreatitis, and
- pancreatectomy (*i.e.*, surgical removal of pancreas).

12.4.11 Pepsin [*Greek*: Pepsis = digestion]

Pepsin refers to the chief enzyme of gastric juice, that essentially converts proteins into **proteoses** as well as **peptones**. It is usually formed by the major cells of gastric glands, and produces its **optimum activity** at a pH of 1.5 to 2. It is mostly obtainable in the granular form. It has been observed that in the presence of HCl, it helps to digest proteins *in vitro*. **Pepsin** on being heated with either **pancreatic enzymes** or **mild alkali** aptly **loses its biological activity**. It exerts an optimum activity at pH 1.8.

Potency of Pepsin: The potency of **pepsin** is as given under:

Pepsin: capacity to digest **2500 times** its weight of coagulated egg protein;

Modified Pepsin: capacity to digest **10,000 folds** its weight of coagulated-egg protein (**albumin**).

12.4.12 Rennin [or Chymosin]

Rennin (Chymosin) designates the **enzyme** that specifically curdles milk, and usually present in the gastric juice of young ruminants. In fact, **chymosin** is the most preferred terminology used for **rennin** due to the possible confusion with **renin**.

Renin is obtained either from the glandular layer of the digesting stomach of the **calf**, *Bos taurus* (**Family: Bovidae**) or by microbiologically monitored fermentation of *Bacillus cereus*, *Endothia parasitica*, and *Mucor pusillus*.

Renin is employed for making cheese and junket*. It is also recommended for patients under convalescence and weak in physical status in order to digest milk rather easily.

12.4.13 Seratiopeptidase

Seratiopeptidase is obtained from the microorganism belonging to the **genus Serratia (E₁₅-Species)** which is exclusively present in the gut of silk-worm, but now-a-days solely generated by biotechnology

* **Junket:** A dish of sweetened curds of milk.

based fermentation. It is regarded to be a better, more effective, and lesser toxic **microbial enzyme** that certainly has an edge over other enzymes *viz.*, **trypsin** and **chymotrypsin**.

Serratopeptidase has the following applications, namely:

- possesses **bradykinin** and **histamine** proteolytic and hydrolyzing characteristic activity,
- accelerates and supports the '**wound-healing phenomenon**' which specifically lowers the capillary permeability and cleavage of exudates and proteins,
- retards inflammation,
- liquefaction of thick sputum (*i.e.*, lowering viscosity) on account of proteolytic effect, and
- enhancement of '**antibiotic effects**' by virtue of the effective removal of inflammatory barrier thereby substantially increasing the **actual antibiotic transfer** to the strategically located infected areas.

12.4.14 Streptokinase

Streptokinase refers to an **enzyme** produced by certain specific strains of **Streptococci*** which is capable of converting **plasminogen** to **plasmin** exclusively.

It is found to be water-soluble and exhibits optimum activity at pH 7.

The various **applications** of **streptokinase** are as stated under:

- used as a **fibrinolytic agent** to help in the removal of **bifrin thrombi** from the arteries (*i.e.*, **reperfusion**).
- employed profusely in the treatment of **thromboembolic disorders** pertaining to the lysis of **arterial thrombus, acute coronary artery thrombosis, deep vein thrombosis, and pulmonary emboli**.

Mechanism of Action: **Streptokinase** exerts its activity on account of the particular activation of **plasminogen** into a **proteolytic enzyme** *e.g.*, **plasmin**, that critically carries out the degradation of the **fibrin clots, fibrinogen, and certain specific plasma proteins**.

12.4.15 Urokinase

Urokinase is an enzyme obtained solely from human urine, which is used experimentally for dissolving **venous thrombi** and **pulmonary emboli**. It is usually administered IV.

Urokinase** also exerts its action as an activator of the specific **endogenous fibrinolytic system** that eventually help in the conversion of **plasminogen to plasmin**; and also causes degradation of **fibrin clots, fibrinogen, and plasma proteins**. It is invariably employed to carry out the dissolution of **fibrin** or **blood clots** present strategically in the **anterior chamber of an eye**, and in **acute massive pulmonary emboli**.

* Culture filtrates of β -hemolytic **Streptococci Group C**.

** **Urokinase**: It is found to be less **antigenic** than **enzymes** due to the fact that this is derived solely from the human source.

12.4.16 L-Asparaginase

L-Asparaginase refers to an **enzyme** that serves as an **antineoplastic agent** derived from the organism *Escherichia coli*. It helps to catalyze the hydrolysis of **L-asparagine** to the corresponding **L-aspartic acid** and NH_3 .

Potency Each 1 mg of L-Asparaginase \equiv 250 Units

The *two* chief uses of **L-asparaginase** are as given below:

- interferes directly with the growth of **cancerous (malignant) cells** that are incapable of synthesizing **L-asparagine** for their necessary metabolism; and, therefore, it is mostly employed in the usual **chemotherapy of very serious lymphocytic leukemia** in preferred sequential combination with other **antineoplastic drugs**,
- beneficial for induction of required remission in children having relapse of **acute lymphocytic lymphoma**, and
- exhibits **immunosuppressive** therapeutic profile.

B. PROTEIN AS DRUG SUBSTANCES

12.5 INTRODUCTION

Protein [*Greek: protos* = first] refers to a class of rather complex nitrogenous compounds that are synthesized by all living organisms, and yield respective array of amino acids when hydrolyzed. Importantly, **proteins** essentially provide the amino acids required for the growth and subsequent repair of impaired animal tissue.

Composition of Proteins: **Proteins**, are composed of a host of vital elements, such as: C, H, O, N, P, S, and Fe, which ultimately make up the greater segment of the animal and plant tissue. In fact, the **amino acids** do represent the basic structure of **proteins**. Generally, foods invariably contain protein with varying numbers and types of amino acids. However, one may recognize a '**complete protein**' as one that predominantly contains **all the essential amino acids viz., arginine, histidine, isoleucine, lysine, leucine, methionine, phenylalanine, threonine, tryptophan, and valine**. In human beings, they are indeed an absolute must for the maintenance of body weight and required growth.

Sources: Interestingly, the various known and important sources of proteins are, namely: cheese, milk, eggs, meat, fish, and certain vegetables *viz.*, soybeans are recognized as the best sources. Nevertheless, the proteins are invariably found in both animal and vegetable sources of food. It has been duly observed that there are many '**incomplete proteins**' which are found in vegetables; and they do contain some of the so-called **essential amino acids**. Thus, a '**vegetarian diet**' may judiciously make up for this by combining various vegetable groups which complement each other in their basic amino acid groups. This ultimately provides the body with '**complete protein**'.

The **major animal proteins** are as enumerated under in Table-12.3:

Table 12.3 Major Animal Proteins and Their Respective Sources

S.No.	Name of Protein(s)	Source(s)
1	Lactalbumin; Lactoglobulin;	Milk; Milk products;
2	Ovalbumin; Ovaglobulin;	Eggs; Egg powder;
3	Serum albumin;	Blood serum;
4	Myosin; Actin;	Striated muscle fibers/tissues;
5	Fibrinogen;	Blood;
6	Serum globulin;	Serum;
7	Thyroglobulin;	Thyroid gland;
8	Globin;	Blood;
9	Thymus histones;	Thymus gland;
10	Collagen; Gelatin;	Connective tissues;
11	Keratin;	Epidermis; Hairs; Nails; Horny Tissue;
12	Chondoprotein;	Tendons; Cartilage;
13	Mucin; Mucoids;	Secreting glands and animal mucilaginous substances;
14	Caseinogen;	Milk; Milk products;
15	Vitallin;	Egg-yolk;
16	Hemoglobin;	Red blood cells [RBCs];
17	Lecithoprotein;	Blood; Brain; Bile secretions;

Functionality of Proteins The ingested **proteins** serve as an important source of **amino acids** essentially required to synthesize the **body's own proteins**, that are quite essential not only for the growth of **new tissue** but also the **repair of damaged tissue**. Importantly, **proteins** constitute an important segment of all the cell membranes. It is proved beyond any reasonable doubt that the excess of amino acids derived from the diet may be conveniently converted to rather **simpler carbohydrates**, and ultimately get oxidized to produce **adenosine triphosphate (ATP)** as well as heat. Thus, 1 g of protein gives rise to 4 kcal of heat.

12.6 PROTEIN VARIANTS

Interestingly, a wide spectrum of **protein variants** have been duly recognized and summarized in Table-12.4 together with their applications:

Table 12.4 Protein Variants and Their Applications

S.No.	Type of Protein	Applications
1	Acute Phase Protein	Specific role in fighting pathogens is not so clear, but they are believed to influence the erythrocyte sedimentation rate (ESR) significantly. Examples: Complement factor C₃ and C-reactive protein.
2	Blood Protein	Integral component of blood, including hemoglobin in RBCs and serum proteins.
3	Carrier Protein	It elicits an immune-response when coupled with a hapten.
4	Complete Protein	They contain all the essential amino acids.
5	Conjugated Protein	They usually contain the protein molecule along with certain other molecules. Examples: Chromoproteins [e.g., hemoglobin]; glycoproteins [e.g., mucin]; lecithoproteins, nucleoproteins, and phosphoproteins [e.g., casein]
6	C-Reactive Protein	It designates an ' abnormal protein ' detectable specifically and exclusively in blood during the active phase of some human diseases viz., rheumatic fever.
7	Denatured Protein	Deformation of the amino acid composition and stereochemical structure of amino acid(s) caused due to chemical or physical means.

(Contd.)

8	Derived Protein	Protein derivative duly achieved either by the action of chemical alteration or a purely physical process <i>e.g.</i> , heat.
9	G-Protein	Protein which determines the activation of a specific physiologic event. It invariably acts at the cell surface to couple receptors for the neurotransmitters , namely: hormones, epinephrine, odorants, and light photons .
10	Immune Protein	An immunoglobulin or an antibody produced by the plasma cells that essentially label foreign antigens and initiate the process for their ultimate destruction (death).
11	Incomplete Protein	Protein that essentially lacks one or more of the essential amino acids.
12	Native Protein	Protein which occurs in its natural state <i>i.e.</i> , it relates to such a protein that has not yet been denatured.
13	Plasma Protein	Protein which is essentially present in the blood plasma <i>viz.</i> , albumin, or globulin .
14	Serum Protein	Protein that forms an integral component present specifically in the serum portion of the blood.
15	Simple Protein	Refers to such proteins which gives rise to the genesis of only α -amino acids upon hydrolysis <i>e.g.</i> , albumins, albuminoids, globulines, gluteline, histones, prolamines, and protamines .
16	Protein C	Represents plasma protein which specifically inhibits coagulation Factor V and Factor XIII, thereby preventing excessive clotting. Its deficiency may cause thrombosis.
17	Protein Kinase	Enzyme that constitute an integral part of the immune reaction; and usually after activation by cytokines strategically mediates cellular processes <i>viz.</i> , motility and secretion.

12.7 BRIEF DESCRIPTION OF PROTEINS USED AS DRUGS

There are quite a few unique and remarkable proteins that are essentially present in the humans which attribute highly efficacious and critical functions *in vivo*. Some proteins that belong to this category, used normally as therapeutic agents (drugs), have been described briefly in the sections that follows:

12.7.1 Complement Protein (Complement Factor C-3) [Latin; *Complere* = to Complete]

Complement protein refers to a **group of proteins** in the blood which play a vital role in the **body's immune defense mechanisms** *via* a **cascade of interactions**. However, the components of complements are labeled C1 through C9. Nevertheless, C3 and C5 are the most important of these. Complement invariably acts by killing directly the organisms; by opsonizing an antigen, thereby stimulating phagocytosis; and by stimulating inflammation and the B-cell mediated immune response. Importantly, all **complement proteins** usually lie **inactive** in the blood unless and until activated by either the classic or the recognized alternative pathways.

An observed abnormality or deficiency in complement protein strategically refers to an **autosomal recessive trait**. The lack of factor C3 enhances susceptibility to common microbial infections, whereas deficits in C5 through C9 are invariably associated with enhanced incidence of **autoimmune ailments**, such as:

- glomerulonephritis, and
- systemic lupus erythematosus.

It has been duly observed that the lack of any of the more than 25 proteins intimately involved in the complement system may affect the body's defense mechanism adversely.

12.7.2 Gelatin [*Latin: Gelatina = Gelatin*]

Gelatin refers to a derived protein duly obtained by the hydrolysis of **collagen** strategically located in the connective tissues of the skin, bones, and joints of animals. It is used profusely in food products *e.g.*, fruit jellies; in the preparations of pharmaceutical dosage forms *e.g.*, **soft-gelatin capsules** for Vitamin E, garlic pearls etc., and **hard-gelatin capsules** for chloramphenicol, tetracycline, acetaminophen (paracetamol) **Tylenol (R)** in US; and also as a medium for the culture of certain microorganisms.

Gelatin is also employed as a vehicle for some highly specific pharmaceutical injections *e.g.*, **Pitkin's menstrum**—which comprises of **heparin, gelatin, dextrose, acetic acid** and **water**.

Gelatin is also used for the treatment of '**brittle finger nails**', and '**non-mycotic defects**' of the nails in humans.

Types of Gelatin: **Gelatin** is normally available in *two* distinct forms, namely:

- (a) **Absorbable Gelatin Sponge:** It is a sterile, white, tough, and finely porous spongy, water-insoluble, and absorbable substance. Even though it is water-insoluble but it is adequately absorbed in body fluids. Nevertheless, it usually takes upto not less than 30 folds its equivalent weight of water.

It has been observed that 9 g of **absorbable gelatin sponge** takes upto 405 g (*i.e.*, 45 times) of **well-agitated oxalated whole blood**.

The various uses of **absorbable gelatin sponge** are as follows:

- as an effective haemostatic,
- as a **localized anticoagulant**, and
- when placed upon a surgical incision after being duly moistened with sterile NaCl solution, it gets slowly absorbed within a span of 4-6 weeks.

- (b) **Absorbable Gelatin Film:** **Absorbable gelatin film** refers to a light amber coloured, sterile, non-antigenic thin film invariably produced from a especially prepared **gelatin-formaldehyde solution** by careful drying followed by subsequent sterilization.

Absorbable gelatin film is largely employed in the form of saline-soaked rubber-like thin sheets chiefly in surgical repair of such observed defects in membranes, such as: **dura and pleura matter**, where it grossly serves as a **mechanical means of protection, replacement matrix**, and **temporary supportive structural wall**.

12.7.3 Collagen [*Synonym: Ossien*]: (*Greek: kolla = glue, + gennan = to produce*)

Collagen refers to a strong, fibrous insoluble protein found in the connective tissue, including the dermis, tendons, ligaments, deep fascia, bone, and cartilage. **Collagen** is the protein typical of dental tissues (except the enamel of teeth), thereby forming the matrix of **dentin, cementum**, and alveolar bone proper. **Collagen fibers** also form the **periodontal ligament**, that eventually attaches the teeth to their respective bony sockets of the lower and upper jaws.

It has been duly observed that there are two important amino acids, namely: **glycine** and **proline** that are strategically located in the central core of the triple helical molecule of the **collagen**. However, **collagen** may be easily differentiated from other accompanying fibrous proteins *e.g.*, **elastin**,* **reticulin** etc. Interestingly, **collagen** is duly characterized by the presence of a host of **vital amino acids**, such as: **glycine**, **hydroxyproline**, **hydroxylysine**, **proline**, and **tyrosine**; whereas, **elastin** essentially comprises of absolutely **non-polar amino acids**, for instance: **isoleucine**, **leucine**, and **valine**. Nevertheless, one may come across a plethora of **collagen variants** which solely depends upon the presence of the '**amino-acid sequence**'. When **collagen** is carefully boiled with water it gets duly converted into gelatin.

Collagen finds its typical applications in the preparation of **photographic emulsions**, **sutures**, and also as a 'gel' in **food casings**.

12.7.4 Casein [Latin: *caseus* = cheese]

Casein designates the principal protein in milk. It is present in **milk curds**. It essentially provides all the amino acids that are necessary for the growth and development in humans. When milk is subjected to coagulation by **rennin** or **acid**, **casein** becomes one of the principle ingredients of **cheese**.

Casein actually represents the **phosphoprotein** with a composition of **0.85% P** and **0.75% S**.

Characteristic Features—are as follows:

- (i) % N = 15 to 16
- (ii) Sulphated Ash (%) = NMT** 1.5
- (iii) Loss on Drying (%) = NMT 6
- (iv) Isoelectric Point = 4.7
- (v) Specific Gravity = 1.25 to 1.31
- (vi) Molecular Weight = 75 K*** to 370 K

Casein Variants: **Casein** has several known variants, such as:

- (a) **Lactalbumin:** **Lactalbumin** refers to the albumin of milk and cheese; and it is a **soluble simple protein**. It is present in relatively higher concentration in **human milk** in comparison to the **cow's milk**. When milk is heated, the **latalbumin** aptly coagulates and appears as a film on the surface of the milk.
- (b) **Lactoglobulin:** **Lactoglobulin** refers to a protein found most abundantly in milk. Both **casein** and **lactoglobulin** are the most common proteins invariably seen in the cow's milk.
- (c) **Acid Casein:** The warm skimmed milk when acidified with a diluted mineral acid, the whey usually gets separated. The solid curd is duly separated by any suitable means, residual solid mass is now washed thoroughly, dried and pulverized to obtain **acid casein powder**.
- (d) **Rennet Casein:** The skimmed milk is adequately treated with an enzyme, **rennet extract**, whereby the product is first separated carefully, and subsequently purified to obtain the **Rennet Casein**.

* **Elastin:** It represents a highly cross-linked protein with a distinct hydrophobic character.

** **NMT** = Not More Than;

*** **K** = Thousand;

- **Casein** (or **soluble casein**) is usually recommended as a dietary supplement for protein in both **pre** and **post-operative** care of the patients.
- **Casein** is also employed as a 'base' in the **proper standardization of the proteolytic enzymes**.
- **Casein** is also exploited as an **emulsifying agent**.
- **Casein** is used in **sizing** of *paper* and *textile*.
- **Casein** is employed as an **unique adhesive agent** for the preparation of large-scale casein paints and casein plastics.

12.7.5 Lectins [Synonyms: Agglutinins; Affinitins; Phasins; Protectins;]

Lectin refers to one of the several plant proteins that specifically stimulate the **lymphocytes** to undergo **proleferation**.

Examples: Phytohemagglutinin; Concanavalin.

Alternatively, **lectins** are **proteins** or **glucoproteins** without having an **immune origin**. **Lectins** may be isolated from various natural sources, such as: bark, fungi, fresh-eggs, roots, microorganisms, body fluids of lower-vertebrates, invertebrates, sea-weed and sponges, and the mammalian cell membranes.

Importantly, the **lectins** are not used directly as a medicine, but they do have the following usages elsewhere, namely:

- For determining **blood-groups**; and for carrying out **erythrocytic polyagglutination investigative studies**.
- For performing **histochemical studies** related to either **normal and pathological status**.
- For establishing **structural elucidation studies of the carbohydrate bearing molecules**.
- For carrying out the **mitogenic stimulation of lymphocytes**.

As tools for studying **cell-surface properties in cancer research**.

Natural Sources of Lectins: a few typical **natural sources of lectins** are as given under:

- **abrin:** *Abrus precatorius*;
- **concanavalin A:** *Conoivalia ensioformis*;
- **green marine algae:** *Codium fragile*;
- **red kidney bean:** *Phaseolus vulgaris*; and
- **horse gram:** *Dolichos biflorus*.

12.7.6 Yeast

Yeast invariably refers to any of several **unicellular fungi** of the genus **Saccharomyces**, that particularly reproduce by budding. They are capable of **fermenting carbohydrates**. **Yeasts**, especially *candida albicans*, may cause **systemic infections** as well as *vaginitis*. It has been generally observed that 'yeast infections' are frequently present in patients with **malignant lymphomas, AIDS, severe diabetes mellitus**, and several other conditions causing **immunocompromise**.

Types of Yeast: There are in fact *two* types of **yeast**, namely:

- (a) **Brewer's Yeast:** **Brewer's Yeast** refers to the specific yeast obtained duly during the **brewing of beer**'. However, it may also be used in its '**dried form**' as a good source of **Vitamin B**.
- (b) **Dried Yeast:** **Dried Yeast** designates the particular '**dried yeast cells**' obtained from the strains of *Saccharomyces cerevisiae*. It is mostly used as a viable source of **proteins** and **Vitamins** (especially **Vitamin B Complex**).

Interestingly, as to date approximately **350 yeast variants** have been duly isolated, purified, characterized for their specific uses. In a rather broader sense, the **yeasts** have been duly classified according to their actual usages to which they are based upon their **typical morphological characteristic features**, such as:

- **For making Wines** : Winer's Yeast;
- **For making Bakery Products** : Baker's Yeast;
- **For making Lager Beers** : Brewer's Yeast;
- **For making Alcohol from Malt Wort, Molasses** : Distiller's Yeast;
- **For making Drugs** : Brewer's Yeast; Baker's Yeast;

Sources of Yeast: The natural habitats of this specific type of microorganism (*i.e.*, **yeast**) are namely: **fruit juice, bread, fermented media, and fermenting media**. On a commercial scale the **yeast** is usually produced by making use of **citrus fruits, molasses, grain wort, malt wort, molasses wort** etc.

Chemical Composition of Yeast: **Yeast** usually contains a wide range of vital and important chemical constituents, namely: nitrogenous ingredients (*i.e.*, **proteins**); vitamins (*e.g.*, **thiamine, riboflavine, folic acid, pantothenic acid, biotine** etc.); enzymes (*e.g.*, **diastase, maltase, zymase** etc.); and **glycogens** and **minerals (ash)**.

Applications of Yeast: Following are some of the vital applications of **yeast** in various **food, beverage, and pharmaceutical** industries:

- (1) **Beverage Industries** *viz.*, Bears, Wines, Alcoholic Beverages such as: Gins, Whiskies, Rums, Vodkas etc.,
- (2) **Food Industries** *viz.*, Bakery Products, Biscuit Industries;
- (3) **Pharma. Industries** *viz.*, antibiotis, papain etc.

12.7.7 Thaumatin [Synonym: Talin;]

Thaumatin is the sweet-tasting basic protein duly extracted from the fruits of the tropical plant, *Thaumatococcus danielli* Benth., *Marantaceae*, found extensively in West Africa from Sierre Leone to Zaire, in Sudan and Uganda. It is mostly composed of *five* distinct forms *viz.*, thaumatins I, II, III, b, and c. However, **thaumatins I and II** predominate invariably. Importantly, all of them are almost 100,000 times sweeter than sucrose, and do have molecular weights around 22,000.

Characteristic Features: The **various characteristic features** of **thaumatin** are as enumerated under:

- (i) It has increasingly sweet taste with a licorice-like distinct after taste.
- (ii) It is strongly cationic having isoelectric point greater than or equivalent to 11.7.

- (iii) It exhibits UV_{max}: 278 nm at pH 5.6; and 283,290 nm at pH 13.0.
- (iv) It is about 750-1600 times sweeter than sucrose on a weight basis; and 30,000 to 100,000 times on a mole basis.
- (v) Its threshold values are very close to 10⁻⁴%.
- (vi) The **proteins** usually lose sweetness upon heating, whereby the disulphide-bridges undergo strategical cleavages in their basic structures.
- (vii) **Thaumatococcus** also loses its sweetness at pHs <2.5, which evidently points to the importance of the very presence of the **tertiary structure** for its inherent sweetness properties.

Thaumatococcus finds its enormous usage as a potential low-calorie sweetener. It is also used largely in such products as: **chewing gums, breath fresheners. Talin** is its branded product being marketed largely in Japan, United States, and the European countries as well.

FURTHER READING REFERENCES

1. **British Herbal Pharmacopoeia**: British Herbal Medicine Association, London (UK), 1983.
2. Cervoni P *et al.*: **Cardiovascular Agents—Thrombolytic Agents**, In: *Kirk Othmer Encyclopedia of Chemical Technology*, 4th edn., Vol. 5, Wiley, New York, pp. 289-293, 1993.
3. **Database on Medicinal Plants Used in Ayurveda**: Vol. 1 (2000), Vol. 2 (2001), Vol. 3 (2001), Central Council for Research in Ayurveda and Siddha, Department of ISM and H, Ministry of Health and Family Welfare (Govt. of India), New Delhi.
4. **Japanese Pharmacopoeia—XIII**, Ministry of Health and Welfare, Tokyo (Japan), 1996.
5. Kaufman PB *et al.*: **Natural Products from Plants**, CRC Press, New York, 1999.
6. **Report on the Task Force on Conservation and Sustainable Use of Medicinal Plants**: Planning Commission, Govt. of India, 2000.
7. **The Ayurvedic Pharmacopoeia of India**, Vol. 1. Pt. 1. Rep. 2001, Vol. 2 (1999), Vol. 3 (2001): Govt. of India, Ministry of Health and Family Welfare, Department of Indian Systems of Medicine and Homeopathy, New Delhi.
8. Wiseman A: **Better by Design: Biocatalysts for the Future**, *Chem. Brit.*, 571-573, 1994.
9. Wong S: **Immunotherapeutic Agents**, In: *Kirk-Othmer Encyclopedia of Chemical Technology*, 4th edn, Vol 14, Wiley, New York, pp: 662-676, 1995.
10. Yokotsuka T: **The Quality of Food and Beverages**, Academic Press, New York, Vol. 2, pp. 171-196, 1981.

13

Biomedicinals From Plant-Tissue Cultures

- | | |
|--|---|
| <ul style="list-style-type: none">• Introduction• Profile of Plant-Tissue Cultures• Biomedicinals in Plant-Tissue Cultures | <ul style="list-style-type: none">• Bioproduction of Commendable Secondary Metabolites• Further Reading References |
|--|---|

13.1 INTRODUCTION

In the recent past, the enormous utility and qualified success of the **plant-tissue cultures** in the skilful and meticulous **bioproduction** of a host of **natural biomedicinals** under stringent aseptic laboratory/pilot-scale conditions has virtually opened up several altogether new approaches in the ever-expanding new era of **biotechnology**. Bearing in mind the aforesaid factual statements one may legitimately exploit the application of the interesting **tissue-culture technique** exclusively for carrying out the biosynthesis of '**secondary metabolites**' prevailing particularly in the plants of pharmaceutical importance and significance, musters an excellent promise for the generation of need-based **plant constituents**.

Importance of Biomedicinals from Plant-Tissue Cultures: The importance of biomedicinals from plant-tissue cultures are enumerated under:

- (1) To obtain a constant and ample supply of good quality '**biomedicinals**' in comparison to the *conventional older methods*.
- (2) To avoid unpredictable factors, such as: absolute neglect of the conservation of environment, careless exploitation of plant species, ever-enhancing labour costs, and unavoidable technical as well as economical constraints associated with the cultivation of the desired medicinal plant species.
- (3) To circumvent such critical problems [as in (2) above] the application of plant-tissue cultures may prove to be a great asset and boon.
- (4) To obtain '**biomedicinals**' from the **higher plants** are still regarded to be a vital source, inspite of the excellent developments in the field of **pure synthetic chemistry**.

At this material time it has become an absolute necessity to understand the *two* terminologies and their functionalities to a certain extent, namely: (a) plant-tissue culture; and (b) biomedicinals.

Plant-Tissue Culture

Plant-tissue culture refers to the specific growth of plant tissue *in vitro* on artificial culture media for the experimental research exclusively.

The quantum leap in 'biotechnology' has virtually opened up the flood-gate related to the latest available techniques with respect to the various cardinal utilities of **plant-tissue cultures** in the specific bioproduction of several natural plant constituents under stringent aseptic parameters using such techniques that are found to be very much akin to the ones employed for the systematic culture of microbes.

Plant tissue cultures (cell suspension cultures) are theoretically regarded to be the most ideal medium for carrying out an exhaustive studies with regard to **metabolism** and also the underlying **biosynthetic pathways** yielding specifically the **secondary metabolites**. In actual practice, such **cultures** are invariably grown from the **callus tissue** in a duly sterile liquid medium containing the necessary nutrients. At the initial stage the growth is **exponential**, but eventually attains a **stationary phase**; and at this point the batches of cells may be transferred conveniently into a new medium. Interestingly, this particular process helps the number of cells increased continually, which may prove to be of great and distinct advantage, provided the **plant tissue cultures** are employed on a **commercial scale** for various **biomedicinals** judiciously.

As to date commercial application of a plethora of **plant tissue culture technologies** have been duly demonstrated and recognized; however, several teething problems and constraints with regard to high labour costs, and other relevant expenses pertaining to composition of nutrient medium, hardening, and precise delivery of tissue cultured plants still remains to be overcome.

Nevertheless, the **plant-tissue culture technique (PTCT)** has a good number of chief advantages *vis-a-vis* the large-scale **cultivation technique (CT)** which may be summarized briefly as under:

- (i) **PTCT** affords the advantage of absolutely uniform biomass achievable at every occasion under strictly modulated as well as reproducible parameters, that may be obtained with remote possibility while dealing with the living potential materials.
- (ii) Importantly, one may conveniently make use of the **plant-cell culture technique (PCCT)** to accomplish the much desired synthesis of such **therapeutically viable agents** (*i.e.*, **medicinal compounds**) that are found to be rather difficult or quite cumbersome to synthesize chemically.
- (iii) Possible to obtain required useful compounds invariably under rigid and controlled environmental parameters that are certainly quite independent of the unavoidable variants, such as: alterations in climatic profile, and soil conditions.
- (iv) **Biotransformation reactions** are feasible using **plant-cell cultures**. One may carry out particular desired alterations/modifications in the '**chemical structures**' of certain plant constituents by this specialized technique in comparison to either pure **chemical synthesis** or **microbiologically** (as in **antibiotics**).
- (v) Plant cells derived from both temperate or tropical environments may be attenuated (multiplied) suitably in order to cause the production of particular metabolites generated by them duly.
- (vi) **PTCT** may be exclusively used to undertake an elaborative study of the **biosynthesis of secondary metabolites**. However, it may be absolutely feasible to induct labelled precursors strategically to the cell cultures, and subsequently deduce logical and plausible interpretations (or explanations) with respect to the various metabolic pathways of the pre-determined chemical entity.
- (vii) Possible to preserve and maintain the '**cultured-cells**' quite free from any attack by insects or contamination by various microorganisms.

(viii) **PTCT** specifically helps to the ensuing **immobilization of cells*** that may be utilized effectively for carrying out various **biochemical reactions** *i.e.*, biotransformations.

Historic Genesis of Plant-Tissue Culture Technique (PTCT): The genesis of actual **plant-tissue culture technique (PTCT)** dates back to almost 100 years *i.e.*, the very beginning of the twentieth century, whereas its enormous benevolent applications were noticed and recognized across the globe in the past three decades; and its **pharmaceutical potential** duly hailed and appreciated for the benefit of mankind.

One may get a glimpse of the **historic genesis of PTCT** commencing from early 1900 in a sequential manner as enumerated under:

- Haberlandt (1902)** : It has been hypothesized that the **isolated plant cells** must be able to undergo propagation (*i.e.*, cultivation) on an **artificial culture medium**. Such investigative studies revealed that one may have an actual assessment of their ensuing capacity as well as typical specific properties as pure **elementary organisms** that are found to be absolutely free from **multicellular system of plant**.
- Robbins (1922) and Kotte (1922)** : These *two* researchers have independently and **successfully cultured the excised plant roots** for the first time, and hence supported the earlier hypothesis of **Haberlandt** put forward two decades ago.
- White (1934)** : Further substantiated the researches carried out by both **Robbins** (USA) and **Kotte** (Germany) by successfully performing the culture of **excised tomato root tips**.
- White (1939)** : Meticulously and successfully reported the **uninterrupted (*i.e.*, continuous) culture of the excised-tomato-root-tips**.
- Gautheret (1939)** : Demonstrated the classical research carried out upon the **cambial tissue cultures**.
Observed the critical behaviour of the **carrot explants** on a medium essentially comprised of **dextrose** (with cystein hydrochloride + aneurine hydrochloride), inorganic salt mixture, and 3-indole-acetic acid** (in low concentration).
- Muir, Hilderbrandt, and Ricker (1954)** : Made an epoch making discovery in the growth of **liquid cultures** distinctly comprising of single cells together with small clusters of cells of two plant species, namely: *Nicotiana tabacum* and *Tagetes erecta*.
- Reinert (1956)** : Demonstrated the occurrence of distinct and highly specific cell divisions in an almost identical preparations having either suspension of single cells or group of cells duly obtained from *Picea glauca*.

* **Immobilization of Cells:** It refers to a specific strain of cells duly obtained from suspension cultures which in turn mobilized by suspending in a solution of sodium alginate thereby causing precipitation of the **alginate** together with the entrapped cells by the aid of CaCl₂ solution, followed by **pelleting**, and allowing the product to get hardened in due course.

** A plant-growth hormone.

- Jones, Hilderbrandt, Ricker, and Wu (1960)** : Critically observed the spectacular growth of plant tissues exclusively in a hanging-drop culture of separated cells duly accomplished from the **callus*** of certain hybrid species of tobacco *viz.*, *Nicotiana tabaccum*, and *Nicotina glutinosa*;
- Rao et al. (1999)** : Ascertained that the '**synthetic seed technology**' would find its application in both **plant propagation**, and **delivery of tissue cultured plants.****

Biomedicinals:

Biomedicine usually refers to the clinical medicine based upon the principles of the so-called **natural sciences** *viz.*, biology, biochemistry, biotechnology etc.

Nevertheless, the present scenaria across the globe entails not only the stability, but also the ensuing viability and potential of newer range of pharmacobiotechnologically developed products. This **new millenium** may eventually see quite a few of them duly approved and okayed by the US-FDA. A few typical examples of '**biotechnology-based products**', otherwise referred to as **biomedicinals**, have already gained world-wide cognizance, namely:

- **Leupine^(R)** : For autologus bone marrow transplantation.
- **Proleukin^(R)** : For renal cell carcinoma.

13.2 PROFILE OF PLANT-TISSUE CULTURES

The profile of **plant-tissue culture** may be adequately expatiated and elaborated in terms of the following different aspects, for instance:

- (a) Types of Cultures,
- (b) Composition of Culture Medium,
- (c) Surface Sterilization of Explants, and
- (d) Preparation of Tissue Cultures.

Each of the aforementioned aspects shall now be treated individually in the sections that follows:

13.2.1 Type of Cultures

In a broader perspective, one may rightly observe that the **plant tissue cultures** which initially took off as an intellectual curiosity has recently attained quite a respectable status by virtue of its significant contributions to basic knowledge and biotechnology. Its judicious usage has ultimately led to several practical gainful advantages to pharmaceutical industry and agriculture.

As to date there are *three* major cardinal categories of highly specific and articulated development areas associated with the **plant-tissue culture**, such as:

- (a) Culture of specific **isolated plant organs,*****

* **Callus**: A mass of unorganized cells.

** Rao PS *et al.* '*Synthetic Seed Technology as a Method for Plant Propogation and Delivery of Tissue Cultured Plants*, **In: Trends in Plant Tissue Culture and Biotechnology**, Agro Botanical Publishers (India), 47, 1999.

*** **Isolated Plant Organs**: Roots, leaf primordia, immature embryos, flower structures, and stem-tips.

- (b) Growth of **callus masses** upon **solidified media**, and
 (c) Growth of **mixed suspensions of separated cells/small cell clusters** in **liquified media**.

Technique Variants: The **plant-tissue culture** may be duly accomplished using different techniques, namely:

- (1) Tissue may be conveniently cultured upon either **liquid nutrient medium** or **solid agar medium**.
- (a) **Liquid suspension cultures:** These usually comprise of an admixture of **cell aggregates, cell clusters (groups), and even single cells**. It has been duly observed that the actual growth rate of the '**liquid-suspension cultures**' are mostly **much higher** in comparison to the **solid-agar medium**. Besides, the said technique essentially gives rise to much elegantly **superior control over the growth of biomass** due to the fact the cells are surrounded by the nutrient medium completely.
- (b) **Solid-agar Medium:** In this particular instance, the plant-tissue when grown on **solid-agar medium** always gives rise to a **callus**. It is, however, pertinent to state here that the wonderful technique of **callus culture** seems to be much more easier as well as convenient for not only initially and maintaining of cell-lines, but also for carrying out the investigational studies related to **organogenesis** and **meristem culture**.

Importantly, both these aforesaid techniques *viz.*, **liquid suspension culture**, and **callus culture** may be adequately obtained from tissues of several species; however, the ease of initiating the said culture exclusively depends and varies along with the specific type of plant, and the tissue organ employed. Interestingly, one may make use of **any portion of a plant** for the **initial induction** to produce either a **suspension culture** or a **callus**. The required necessary tissue may be duly obtained from various segments of a plant *viz.*, leaf, root, stem, pollen, and seedling, which eventually is made to grow distinctly as a mass of **undifferentiated cells** upon previously enriched solidified in an aseptic environment.

13.2.2 Composition of Culture Medium

Since 1939 various researchers have put forward specific well-defined **culture medium** for the growth of cell cultures that essentially comprise of a host of ingredients, such as:

- Carbon source** : From carbohydrates, starch, steep liquors.
Vitamins : To augment the growth of **callus**, and **liquid-suspension cultures**; besides, **meristem culture**.
Plant growth hormones : To serve as growth regulators.
Organic supplements : To aid the overall growth of certain specific cultures.

Murashige and Skoog* (1962) found an altogether revised medium for rapid growth and bioassay with tobacco tissue cultures; Grambow *et al.*** (1972) studied the cell-division and plant development from protoplasts of carrot cell suspension cultures; Chang and Hsing*** (1980) carried out studies

* Murashige T and Skoog F: *Plant Physiol.*, **15**: 437-497, 1962.

** Grambow HJ *et al. Planta*, **103**: 348-355, 1972.

*** Chang WC and Hsing WI, *Nature*; **284**: 341-342, 1980.

of *in vitro* flowering of embryoids derived from mature root callus of **ginseng** (*Panax ginseng*; Shillito *et al.** (1983) investigated agarose plating and a **bead-type culture technique** and stimulate development of protoplast-derived colonies in a number of plant species; Wang (1990)** reported callus induction and plant regeneration of **American ginseng** Hort.; and Arya *et al.**** (1993) described the rapid multiplication of adventitious somatic embryos of *Panax ginseng* (CA Meyer).

Following are the *five* cardinal categories of important ingredients that are absolutely essential as an **ideal nutrient medium** required for **propagation of plant-tissue cultures**, namely:

- (1) **Source of Carbon:** Two commonly and abundantly employed source of carbon are **glucose** and **sucrose** invariably used at a concentration ranging between 2-4% (w/v).
- (2) **Inorganic Salts:** Importantly, the concentration of K^+ ion (cation) and NO_3^- ion (anion) must be strategically adjusted within a range of 20-25 mM (*i.e.*, **millimole**) for each of the said ions; and the concentrations of certain other ions *viz.*, Mg^{2+} , PO_4^{3-} , SO_4^{2-} should vary between 1-3 mM. However, there are some vital and **important micronutrients** that are usually recommended for the **culture medium**, such as: boric acid (H_3BO_3), Γ , besides certain bivalent salts of Zn, Mn, Cu, Co, and Fe which are duly inducted in their chelated form [*e.g.*, sequestered chelates obtained with **ethylene-diamine tetra-acetic acid (EDTA)**].
- (3) **Vitamins:** The most essential vitamin required for the **culture medium** is **thiamine hydrochloride** (or **Vitamin B₁**); whereas, **pyridoxine hydrochloride** (or **Vitamin B₆**), **inositol** (*i.e.*, a part of the **Vitamin B complex**, and **niacin** (or **nicotinic acid**) are invariably employed in **culture medium** to improve significantly the desired **cell growth**.
- (4) **Organic Additives:** It has been duly established that the particular incorporation of such **organic additives** as: fresh/pasteurized **coconut milk** (containing **liquid endosperm**), **malt extract**, **protein hydrolyzates** (*e.g.*, **soy-protein hydrolyzates**), and **yeast extract** are used predominantly to cause an appreciable increase in the actual growth rate of the cells present in the **biomass**.****
- (5) **Growth Regulators (or Modulators):** **Growth regulators** are essentially required to cause the induction of the desired **cell division** in an affective manner. The various chemical compounds which are mostly used as **growth regulators** together with their optimum operative concentrations are given in the following Table 13.1.

Table 13.1 Chemical Compounds Used as Growth Regulators

S.No.	Chemical Compound (Abbreviation)	Concentration Used (mM)	Remarks
1	2,4-Dichlorophenoxyacetic acid [2, 4-D]	$3 \times 10^{-5} - 5 \times 10^{-5}$	Degraded gradually in plant cells; and are quite stable to autoclaving.
2	Naphthalene acetic acid [NAA]	$3 \times 10^{-5} - 5 \times 10^{-5}$	— do —
3	Cytokines***** <i>viz.</i> Kinetin, Benzyladenine	$10^{-7} - 10^{-5}$	Gives good callus production when used in conjunction with either 2, 4-D or NAA .

* Shillito RD, Paszkowski J, and Potrykus I, *Plant Cell Rep.*, **7**: 418-420, 1983.

** Wang AS: *Science*, **25** (5): 571-572, 1990.

*** Arya S, Arya ID, and Eriksson T: *Plant Cell Tissue and Organ Culture*, **34**: 157-162, 1993.

**** **Biomass:** All of the **living organisms** located in a specified area.

***** **Cytokines:** One of more than 100 distinct proteins produced primarily by WBCs. They provide signals to regulate immunological aspects of **cell growth** and function during both inflammation and specific immune response.

13.2.3 Surface Sterilization of Explants*

Sterilization essentially refers to the process of completely removing or destroying all microorganisms on a substance by exposure to chemical or physical agents, exposure to ionizing radiation or by filtering gas or liquids *via* porous materials which essentially remove microorganisms.

For carrying out extensive and intensive research on **tissue culture propagation** it is almost important and necessary to afford **surface sterilization** of the organ from which the **specific tissue** is required to be excised aseptically or the particular seed material the germination of which shall yield ultimately the **tissue explant**. In actual practice there are quite a few tried and tested frequently employed compounds that find their abundant application as surface-sterilizing agents, namely:

S.No.	Name of Compound	Effective Strength (%)
1	Bromine water	1 to 2
2	Hydrogen peroxide	10 to 12
3	Mercuric chloride	0.1 to 1
4	Silver nitrate	1
5	Sodium hypochlorite	1 to 2

Procedure: The various steps adopted for the **surface sterilization of explants** are as enumerated under:

- (1) Seeds are adequately treated with ethanol (70% v/v) for a short span of 2 minutes; and subsequently washed thoroughly with distilled water.
- (2) The resulting seeds are duly subjected to '**sterilization**' by **surface sterilizing agents** for a known prescribed duration.
- (3) Seeds thus treated are rinsed once again with enough sterile distilled water, and kept for germination under rigid aseptic environment for a stipulated period.
- (4) **Germination** of seeds may be accomplished in several tried and tested methods, namely:
 - by placing them on double layers of pre-sterilized filter paper in petri-dishes, duly moistened with sterile DW (distilled water),
 - in petri-dishes adequately soaked with sterile cotton plugs, and
 - in pre-sterilized culture tubes.
- (5) Seeds are duly germinated in complete darkness preferentially at a temperature ranging between 26 to 28 °C.
- (6) Finally a small segment of the resulting seedlings are duly utilized for the actual starting of the **callus culture**. Importantly, this step needs to be carried out under perfect sterile conditions so as to avoid any possible risk of contamination inadvertently.
- (7) **Aerial Segments of Plant Materials**** are invariably sterilized by subjecting them to sterilization by submerging the specific segments in ethanol (70%, v/v) for a duration of 2-3 minutes in sterile distilled water.

* **Explant:** To remove a piece of living tissue from the body and transfer it to an **artificial culture medium** for growth, as in **tissue-culture**.

** **Aerial Segments of Plant Materials:** These are buds, leaves, and stem portions.

13.2.4 Preparation of Tissue Cultures

The actual preparation of **tissue cultures** is accomplished by adopting well-defined methodologies in a sequential manner. The said **tissue cultures** may be prepared either from **suspension cultures** or from **solid cultures**, which shall now be treated separately in the sections that follows:

13.2.4.1 From Suspension Cultures

The various steps adopted generally for the preparation of '**Suspension Cultures**' are as given under:

- (1) **Suspension cultures** are first and foremost initiated by careful transference of a previously established **callus tissue** into a sterile agitated liquid nutrient medium* held in an Erlenmeyer Flask (size: 500 mL, or 250 mL) usually termed as '**culture vessels**'.
- (2) Composition of the medium for the **suspension cultures** must be exactly the same as meant for the corresponding **callus cultures**, except the addition of agar (for solidification).
- (3) **Soft callus** thus generated gives rise to a desired **suspension culture** quite readily without any hazles at all.
- (4) The subsequent release of **tissue cells** as well as **tissue fragments** from not so easily crumbling **callus masses**. Thus, one may afford the critical maintenance of a significant extent of **cell separation** *via* sustaining a relatively high level of **auxin**** concentration in the prevailing **liquid medium**.

It is, however, pertinent to mention here that it is absolutely necessary to maintain a suitable and feasible balance between **auxin and yeast extract** or **auxin and kinetin**.

- (5) **Incubation** of the **suspension cultures** are carried out invariably at 25 ± 1 °C either in **total darkness** or in **dim and diffused fluorescent light**.
- (6) Constant and continuous agitation of the flask cultures is generally accomplished by the help of a **motorized-speed-controllable horizontal shaker** that rotates between 100 to 200 RPM.
- (7) The labeled and sterile culture flasks are adequately sealed either with **solid-paraffin wax** or with **thick aluminium foil** so as to check the evaporation to a bare minimum level. It takes almost 4 to 6 weeks to obtain a reasonably appropriate and good **cell suspension**.
- (8) Cells grown in **suspension cultures** are found to be **meristematic** in nature. These are usually undifferentiable; besides there exists no vivid and apparent evidence that the ensuing cells belonging to either root or shoot tend to be altogether different metabolically.
- (9) Finally, the resulting **suspension cultures** are duly subjected to further subculturing by the actual transfer at regular gap of time of both the duly **fractionated known amounts of the suspension culture** and the **untreated suspension culture** into the **fresh sterilized medium**.

13.2.4.2 From Solid Cultures

The various steps involved in the preparation of **tissue cultures** from the **solid cultures** are as enumerated under:

* Usually 30 to 50 mL for each 250 mL Erlenmayer flask.

** A substance that promotes growth in plant cells and tissues.

- (1) Surface sterilized plant material is adequately transferred onto the solidified nutrient medium taken in Erlenmayer flasks, culture tubes, and glass jars aseptically.
- (2) These inoculated glass containers (*viz.*, flasks, jars, tubes) are duly incubated between 26 to 28 °C for a period ranging between 3-4 weeks in complete darkness aseptically.
- (3) At the end of the incubation period (*i.e.*, 3-4 weeks) the size and shape of the **callus** must be at least 5 to 6 times bigger than the corresponding size of the **explant**.
- (4) Importantly, certain **tissue explants** specifically exhibit **some extent of polarity** which evidently gives rise to the formation of **callus** most rapidly at one surface particularly.
- (5) **Stem portions** do exhibit the formation of **callus** wherein the specific surface *in vitro* is pointed towards the **root**.
- (6) Invariably, the **callus** gets developed rather swiftly originated either from the tissue not immersed in the solidified culture medium or from the tissue that are found to be not in contact with the said medium.
- (7) Regular **maintenance of growth in the callus tissue** may be duly accomplished by the aid of ‘**subculturing**’ which essentially requires the effective transfer on each occasion of a **specific piece of healthy viable tissue** after each 30 days into the flask having freshly prepared, sterilized, and solidified nutrient medium.
- (8) One may evidently observe that even when the usual standard incubation temperature is brought down between 5 to 10°C from 26°C, several cultures would remain healthy and extend their growth usually at a much lower pace but for longer durations without the necessity of ‘**subculturing**’.
- (9) Finally, one would distinctly notice that the normal growth of a plethora of tissue cultures, and specifically of those wherein **chlorophyll gets formed** is overwhelmingly stimulated by a certain **low-intensity illumination**.*

13.3 BIOMEDICINALS IN PLANT-TISSUE CULTURES

The intensive and extensive investigative studies related to the inherent biosynthetic capabilities of **plant-tissue cultures** meticulously derived and articulately accomplished from a variety of **higher plants** has been carried out in a systematic and methodical fashion stretched over the past three decades across the globe. It has been duly observed that in several instances the much desired specific chemical entities could not be generated for one reason or the other; whereas, in others extremely low quantum of such compounds of interest could be achieved precisely.

The observed significantly **poor yields** of the **secondary metabolites**** seem to be caused on account of the prevailing cardinal factors, such as:

- (1) Due to apparent differences between duly dispersed, young, and quickly dividing **culture cells vis-a-vis** the relatively mature and slow-growing cells pertaining to the variants in product generation based upon a plethora of such vital factors as: regression (or dormancy) in

* **Light source** having a 12 hr cycle or continuous exposure is invariably provided in most of the **Incubation Chambers** by the usual fluorescent tubes.

** A molecule derived from a primary metabolite, many serve protective functions.

biosynthetic pathways, non-excretion of viable products from the **culture cells**, and the significant function of time in culture.

(2) These above mentioned drawbacks (in. 1.) may be circumvented by adopting one of the following effective procedures, namely:

- development of immobilized cell systems,
- development of highly specific hairy-root cultures, and
- evolution of such methodologies which may evidently encourage the process of excretion of the desired viable product right into the **culture medium**.


It is indeed worthwhile to mention at this point in time that there are several proven and glaring examples which obviously show that the **callus** as well as the **suspension cultures** are magnificently capable of carrying out the synthesis of several desired **secondary metabolites** having **overwhelming and satisfactory yields** that are found to be fairly comparable to the **intact plant** itself.

A survey of literature would reveal that during the past two decades there have been an excellent progress with regard to the promising discoveries in the field of **biomedicinals** (*i.e.*, medicinally potent substances), a few of them may have the potential to be scaled up to the **commercially viable extent**.*

13.3.1 Secondary Metabolites

In the context of the present day knowledge there exists no clear cut, distinct, and sharp line of division between **primary** and **secondary metabolites**. The **secondary metabolites** are now regarded to be an absolute must to the **plant life**; and, therefore, several of them do provide an effective **defence mechanism** against **bacterial, viral, and fungal** attack very much analogous to the **immune system of animals**. Importantly, the actual presence of a chemical entity derived from the **secondary metabolism** exclusively rests upon the sensitivity of the analytical procedure detecting its presence precisely.

Example: Azetidine-2-carboxylic acid (I), an amino acid, present in the sugar beet [*Beta vulgaris*; **Family: Chenopodiaceae**], was first and foremost detected and analysed and found to be a minor constituent of the plant. Interestingly, there exists a copious evidence indicating that **(I)**, which was once regarded restricted to the natural order **Liliaceae**, is perhaps a ubiquitous (*i.e.*, found everywhere) plant constituent.







Function of Secondary Metabolites: Nevertheless, it would be quite unfair to consider the **plant secondary metabolites** as the '**waste products**', as was once even concluded, but now it has been proved beyond any reasonable doubt that they do possess **several most useful functions**. A few such vital functions shall be enumerated as under:

* **Commercially viable extent** refers to the productivity for economic considerations, and may be defined in terms of **gram product/culture volume (L)/day**.

(a) **Plant Hormones:** In fact, there are *two* types of **plant hormones**, namely:

- (i) **Plant growth hormone**—responsible for the growth of plant(s) *e.g.*, **Auxin** [indolylacetic acid] (II); **cytokinins** [zeatin] (III); **ethylene** [abscisic acid] (IV); and **gibberellins** [gibberellic acid] (V).





- (ii) **Wound hormone**—responsible for the healing of damaged tissue *e.g.*, **traumatic acid** (VI):



Traumatic Acid (VI)


(b) **Flavonoids and Carotenoids:** A good number of coloured **flavonoids** and **carotenoids** invariably serve to attract specifically **insect and bird pollination** of even the **seed dispersal agents**. However, a few other **secondary metabolites** *e.g.*, the **volatile terpenoids**, repel predominant invaders. It is worthwhile to state here that a relatively large segment of the **secondary metabolites** do exert their marked and pronounced action as **vital fungicides** or **potential antibiotics**, which in turn cause protection of various plants from **fungal or bacterial attacks**.

Examples: Some typical examples belonging to this particular class are, namely: **pterocarpan** [*e.g.*, **pisatin** (VII)], and **sesquiterpenoid** [*e.g.*, **ipomeamarone** (VIII)] which are nothing but **phytoalexins** usually generated in response to **specific fungal attack**.




13.3.2 Usefulness of Secondary Metabolites


The **usefulness of the secondary metabolites** play an important role either as such to the diet of human beings, such as: **Vitamin C** [*e.g.*, **ascorbic acid (IX)**, **Vitamin E**] *e.g.*, α -tocopherol (X), Vitamin K [*e.g.*, **phylloquinone (XI)**] that are remarkably biosynthesized by plants; or β -carotene [*e.g.*, **precursor of Vitamin A (XII)**], **ergosterol** [*e.g.*, precursor of Vitamin D (XIII)] which are also designated as **potential secondary plant metabolites**.





Vitamin C (IX)



Vitamin E (X)




Vitamin K (XI)


 β -Carotene (XII)

Vitamin A

**1 mole of β -carotene
yields 2 moles of
Vitamin A**




Ergosterol (XIII)



Vitamin D


In addition to the above interesting applications of the **secondary plant metabolites (SPMs)** there are quite a few other noticeable usefulness as enumerated below:

- (a) **Precursors for Drug Synthesis**—SPMs are also vital and important as precursors for the synthesis of certain drugs *e.g.*, **steroids** and **hormones**.
- (b) **Plant Poisons (Insecticidal agents)**—Several plants do contain poisonous materials used as insecticidal agents; however, only a few have been duly exploited on a commercial scale, such as: **Nicotine** [*Nicotiana* species: Family: **Solanaceae**]; **rotenone** [*Derris* and *Lonchocarpus* species: Family: **Leguminosae**]; and **Pyrethrins** [*Chrysanthemum cinerariaefolium*: Family: **Compositae**].



- (c) **Natural Dyes**—As to date quite a few **natural dyes** do find their usefulness in foods and drugs, such as:

- **Carthamin** : *Carthamus tinctorius* (**Compositae**);
- **Bixin** : **Annatto** [*Bixa orellana* (**Bixaceae**)];
- **Croctin** : **Saffron** [*Corcus sativa* (**Iridaceae**)];
- **Curcumin** : **Turmeric** [*Curcuma longa* (**Zingiberaceae**)].



13.3.3 Secondary Metabolites in Chemosystematics

It has been amply demonstrated that the **secondary plant metabolites (SPMs)** may contribute profusely and overwhelmingly to **plant taxonomy, systematics, and the study of evolution**. One

may observe that the evolution of **chemical** and **morphological characteristic features** are interrelated to an enormous extent; and, therefore, a large segment of these studies related to the **secondary metabolites** invariably serve to confirm the '**morphological classifications**'. Nevertheless, in such specific instances wherein the available morphological relationships seem to be unclear, one may heavily depend upon the so called '**chemotaxonomic markers**' which usually give rise to valuable and critical information(s).

Interestingly, the **biosynthetic pathway** plays an important role while taking into consideration the **chemosystematics** and **plant revolution**, due to the fact that the same kind of **secondary metabolite** may be the **product of two altogether different pathways**. However, the specific enzymes catalyzing the metabolic reactions are of immense importance *vis-a-vis* the products of the reactions, by virtue of the fact that **evolution** solely depends upon the alterations in the **enzyme profiles**.

Plants that essentially possess identical morphological and chemical characteristic profile should have the same ancestor may not prove to be always true perhaps on account of the ensuing **convergent (parallel) revolution**.

Example: Pyrrolizidine alkaloids—are produced by *two* altogether different plant species, namely: *Crotolaria* (Family: **Leguminosae**); and *Senecio* (Family: **Compositae**) by the help of similar pathways; however, these *two genera* are absolutely unrelated morphologically. In the light of the above statement of facts it is significantly important to take into consideration all relevant **secondary metabolites** (and not just one particular class) while **classifying the plants** based upon the **biochemical data**.

13.3.4 Newer Products Developed

It has been ascertained that the **secondary metabolites** duly produced from the **plant-tissue cultures** give a yield which is either equal to or greater than that of the **parent plants**.* In fact, almost **thirty natural products** have so far been produced *via* the **plant-cell culture**; and a few such well-known drugs are as follows:

- **Diosgenin** : steroid hormone precursor.
- **Opium Alkaloids** : morphine and its natural congeners
- **Digitalis Glycosides** : digitoxin, digoxin,
- **Essential Oils** : peppermint oil, lemon oil, and
- **Catherenthus Alkaloids** : vincristine, vinblastine.

Shikonin, an Asiatic Drug, obtained from *Lithospermum erythrorhizon* has been adequately generated *via* **plant-cell culture** significantly in larger amounts than the usual extraction procedures.** Robins (1987) have duly reported the presence of certain major secondary products from the **hairy-root cultures**, for instance:***

- **Atropine** : from *Atropa belladonna* L,

* Staba JE: *J. Nat. Prod.*, **48** (2): 203, 1985.

** Evans WC: **Pharmacognosy**, WB Saunders, London, 15th edn., 2004.

*** Robbins R: *Planta Medica*: **53**: 474, 1987.

- **Hyoscyamine** : from *Datura stramonium* L.,
- **Nicotine** : from *Nicotiana tabacum*,
- **Cantharathene** : from *C. roses*, and
- **Quinoline Alkaloids** : from *Cinchona ledgeriana*.

Based on the scientific evidences one may observe that each **undifferentiated cell** essentially comprises of the necessary genetic information responsible for the production of **secondary metabolites**. It is quite possible to produce commercially **specific plant cells** by integrating their genetic information; and also to increase the production of **secondary metabolites appreciably**. It has been duly observed that the **cell cultures** of several **vital medicinal plants** may effectively produce their **secondary metabolites in cultures**; however, the actual yield is comparatively lower to the conventional plant. Last three decades have witnessed the successful achievement of increased yield of desired **secondary metabolites in culture**. Thus, extremely promising results have been achieved for the production of **alkaloids, anthraquinones, and cardioactive glycosides**.*

13.4 BIOPRODUCTION OF COMMENDABLE SECONDARY METABOLITES

Plant tissue culture has proved to be a viable, feasible and productive alternative method to carry out the production of a plethora of commendable **secondary metabolites** used as **drugs**. In the past *two* decades a copious volume of research has been done in this direction, of which a few typical and important **drugs** shall now be treated in the sections that follows:

- (a) **Taxol and Related Compounds:** Christen *et al.* (1991)** patented the production of **taxol** from cell cultures. The very next year Fetto-Netto *et al.* (1992)*** reported the cell cultures of *Taxus* species *Taxus cuspidata* and *Taxus canadensis*. It has been observed that the **callus cultures** of *T. cuspidata* could generate **taxol** in quantities almost equivalent to the **stem of the intact plant** (0.02% DW) after 60 days of culture. However, the **immobilized cell cultures** of *T. cuspidata* produced **taxol** very much equivalent to those found in the **bark of the tree** (0.012% DW) after 180 days of culture. All these findings obviously indicated the possibility of making use of **tissue cultures** towards the genuine production of **taxanes**.
- (b) **Camptothecin:** Wall *et al.* (1966)**** duly reported the presence of **camptothecin**, a *quinoline alkaloid*, from the chinese tree *Camptotheca acuminata*; and subsequently Govindachari *et al.* (1972)***** discovered the same in the indigenous tree *Nothapodytes foetida*. It is an **antitumour alkaloid**.
- (c) **Medicinal Plants with Anticancer Compounds:** Misawa and Nakanishi (1988)***** reported the presence of certain anticancer compounds in a few well known medicinal plants, such as:

* Verpoorte R: **Metabolic Engineering of Plant Secondary Metabolism**, Academic Press, Netherlands, 1999.

** Christen *et al.* United States Patent No.: 5: 019, 504 (1991).

*** Fetto-Neto AG *et al. Bio. Technol.*, **10**: 1572, 1992.

**** Wall *et al.*: *J. Am. Chem. Soc.*, **88**: 3889, 1966.

***** Govindachari TR and Vishwanathan N., *Phytochemistry*, **11**: 3529, 1972.

***** Misawa M and Nakanishi TM: *Antitumour Compounds Production by Plant Tissue Cultures*. IN: **Biotechnology in Agriculture, Forestry, Medicinal and Aromatic Plants II**, Springer Verlag, Berlin, pp. 192-207, 1988.

- *Cephalotaxus harringtonia*,
- *Podophyllum peltatum*,
- *Trypterigium wilfordii*, and
- *Putterlickia*.

The **tissue cultures** of these medicinal plants have been duly established, and the **cultures** essentially comprised of the desired compounds of the parent plant. Generally, apparent differences were adequately noticed with respect to the yield of the compounds. Kupechan (1972)* observed that in certain specific instances *e.g.*, *T. wilfordii* the **cultures** were able to synthesize appreciably much higher amounts of **tripliolides** when compared with the intact plant. Kadkade (1982)** carefully observed the **podophyllotoxins** from *P. peltatum* were significantly higher [7.1×10^{-1} % DW] in comparison to the actual plant obtained from the field (*i.e.*, **natural environment**) which stands at [6.4×10^{-1} % DW]. Likewise, the **cephalotoxine** content present in the **cultures** of *C. harringtonia* (in **differentiated tissues**) were found to be much higher in concentrations than the **suspension cultures**.

Conclusively, one may observe that the **potential antineoplastic compounds** may be produced substantially by **tissue cultures** provided the large-scale production could be made feasible if the process can be scaled up in relatively **large bioreactors** followed by critical optimization of experimental parameters required essentially for the **active production**.

- (d) **Antimalarials:** Fulzele *et al.* (1991)*** reported 0.012% DW of **artemisinin** present in the **shoot cultures of *Artemisia annua***, whereas He *et al.* (1983)**** reported only 0.008% DW of the same.

Woerdenberg *et al.* (1983)***** duly reported an yield of 0.16% DW of **Artemisinin** in the shoot cultures of *A. annua* growing on MS 1/2 + BA (0.2) + NAA (0.05) medium containing 1% sucrose. However, the respective **Artemisinin** content enhanced with **0.5% sucrose**. Meanwhile, supplementing the **culture medium** with GA (10 mg. L^{-1}), CH (0.5 mg. L^{-1}) and 10 or 20 mg. L^{-1} of **naftifine** remarkably increased the production of **Artemisinin** in the shoot cultures of *A. annua*. Importantly, these wonderful observations do suggest that the duly prepared **shoot cultures** may ultimately prove to be a logical, feasible and viable alternative method for the production of **Artemisinin**.

- (e) **Adenylate Cyclase Activity Stimulator: Forskolin** is a potent **diterpenoid** isolated from the roots of *Coleus Forskohlii* (an Indian Medicinal Plant) which distinctly possesses **adenylate cyclase activity stimulating profile**. Sen *et al.* (1993)***** duly reported the presence of **forskolin** in the untransformed root cultures of *C. forskohlii*. The roots were carefully grown in various **basal media** and altogether variable **culture conditions**. In actual practice, the **culture media** employed duly for the growth of **root cultures** essentially consisted of MS 25%, sucrose 1%, and either IAA, IBA, or 2-chlorinated derivatives of IAA *viz.*, 4-Cl-IAA

* Kupechan SM *et al.* *J. Am. Chem. Soc.*, **94**, 7194, 1972.

** Kadkade PG: *Plant Sci. Lett.*, **25**: 107, 1982.

*** Fulzele DP *et al.* *Phytotherapy Research*, **5**: 149, 1991.

**** He XC *et al.* *Acta. Bot. Sin.*, **25**: 87, 1983.

***** Woerdenbag HJ *et al.*: *Plant Cell Tissue and Organ Culture*, **32**: 247, 1993.

***** Sen J *et al.*: *Phytochemistry*, **34**: 1309, 1993.

and 5, 6-C 12-IAA. The **tissue cultures** grew very satisfactorily in all the dosages of the chlorinated IAAs. Besides, it also generated more quantum of **forskolin** in comparison to the **cultures** grown in IAA or IBA singly. It has been observed that in a medium comprising 50 mcg. L⁻¹ of Cl-IAA the **forskolin** content stood at 0.09% DW, that was evidently comparable to **forskolin** content present in the **wild plant** (0.1% DW). However, at a concentration level of 500 mcg. L⁻¹ of IAA or IBA exclusively small amounts of **forskolin** was obtained.

To sum up one may add that the **plant-tissue cultures** continue to remain as a distinct major source for several novel **biomedicinals (phytochemicals)**, and the subsequent strategic development in the **Medicinal Plant Biotechnology** together with **Plant Tissue Culture and Technology** have virtually opened up newer avenues both for the propagation of **elite medicinal plants** and **mass cultivation of cells in sophisticated PC-based automated Bioreactors** for superb and novel products.

FURTHER READING REFERENCES

1. Brar DS and Khush GS: **Cell and Tissue Culture for Plant Improvement**. In: Basra AS (ed.): *Mechanisms of Plant Growth and Improved Productivity: Modern Approaches*, MerceL Dekker Inc., New York, pp. 229-278, 1994.
2. Brown DCW and Thorpe TA: **Plant Regeneration by Organogenesis**. In: Vasil IK (ed): *Cell Culture and Somatic Cell Genetics*, Academic Press, New York, Vol. 3, 1986.
3. Congress, Office of Technology Assessment: **New Developments in Biotechnology: US Investments in Biotechnology—Special Report**. OTA-BA-360. Supdt. Documents, Washington DC, 20402-9325, 1988.
4. Debergh P and Zimmerman R (eds.): **Micropropagation**, Kluwer Academic Publication, New York, 1991.
5. Eliet U: **Cell Culture and Somatic Cell Genetic Implants**, Academic Press, San Diego, 1987.
6. Imrie BC and Hacker JB (eds): **Focused Plant Improvement: Towards Responsible and Sustainable Agriculture**, Pergamon Press, London, 1993.
7. Ingram DS and Helgeson JP (eds.): **Tissue Culture Methods for Plant Pathologists**, Blackwell, Oxford, UK, 1980.
8. Marmorosch K and Sato GH (eds.): **Advances in Cell Culture**, Academic Press, California, Vol. 7. 1989.
9. Mohan Ram HY and Kapoor A: **In vitro Growth and Development of Hornwort**. In: *3rd International Conference: Plant Tissue and Cell Culture*, Leicester, UK, 1974.
10. Rao AN (ed.): **Tissue Culture of Economically Important Plants**, Asian Network for Biological Sciences, Singapore, 1981.
11. Reinert J and Bajaj YPS (eds.) **Applied and Fundamental Aspects of Plant Cell, Tissue, and Organ Culture**, Springer, Berlin, 1977.
12. Staba EJ: **Plant Tissue Culture as a Source of Biochemicals**, CRC-Press, Boca Raton, 1987.
13. Verporate R: **Metabolic Engineering of Plant Secondary Metabolism**, Academic Publishers, Netherlands, 1999.
14. White PR: **A Handbook of Plant Tissue Culture**, Jaques, Catell, England, 1943.

14

Hi-Tech Products from Plant Sources

- Introduction
- High Throughput Screening (HST)
- Success of HTS of Plant Source Materials for New Lead Chemical Entities
- Hi-Tech Products
- Further Reading References

14.1 INTRODUCTION

Since more than three centuries the World has witnessed the enormous and highly specific therapeutic applications of both **traditional and orthodox medicaments**. Invariably, the use of ‘**crude natural plant extracts**’ comprising of multi-component products duly exerted either **extremely potent/active entities**, such as: **Digitalis leaf** or **extremely weak/active** components, for instance: **Cinnamon bark**. Interestingly, the **orthodox medicaments** exclusively depend upon mainly single (*i.e.*, a very small quantum of) well defined, duly characterized active-chemical entities predominantly showing highly specific activities at, in a plethora of instances, well-established and known ‘**biological targets**’.

It is, however, pertinent to state here that these **hi-tech products** (or **medicines**) are invariably found to be extremely potent; therefore, most of them usually **display pretty narrow gap** prevailing between an **effective dose (ED₅₀)** and a **lethal (toxic) dose (LD₅₀)**. Importantly, one may duly observe that the so called **orthodox medicaments** are meticulously formulated and designed into a good number of dosage forms which may be judiciously ‘**standardized**’ for their ultimate bioavailability *in vivo*.

Following are some of the most **invaluable orthodox drugs**, that are exclusively obtained from chemical compounds frequently occurring in higher plants, are basically ‘**analgesic**’ in character, such as: **morphine** and **codeine**; **antineoplastic agents** *viz.*, **taxol** and **vincristine**; **antimalarials**, for instance: **quinine** and **artimisinine**; and **anti-asthmatics** *e.g.*, **cromoglycate**.

As to date, certain naturally occurring materials from the plant sources (or plant kingdom) still continue to enjoy the most viable and vital commercial source of the ‘**active principle**’. **Examples**: The *two* prominent and exclusive examples are as stated below:

- (a) **Poppy capsule**: Glaxo Wellcome still harvests up to 10,000 MT dry weight of ‘**poppy capsule**’ every year (under license) to provide an authentic genuine source of the ‘**opiate alkaloids**’, and

- (b) ***Digitalis lanata* leaf:** Contractors hired by Glaxo Wellcome also produce 170 MT per annum dry weight of *Digitalis lanata* leaf to serve as the prime source of ‘**digoxin**’—a potent **cardiotonic**.

14.2 HIGH THROUGHPUT SCREENING (HTS)

In an attempt to explore and discover ‘**new medicines**’ that essentially give rise to an altogether distinct and remarkably significant advantages upon the **prevailing practised therapies**, the primary crucial and key starting point is to identify, recognize, and exploit the ‘**novel biological targets**’, which do possess a critical role in the control and management of a plethora of **disease processes**. Interestingly, the immediate text protocol would be to evolve and involve discovery intimately associated with the ‘**biological assays**’, that are quite capable of detecting such substances which may eventually cause articulated modification of the ensuing activity of the ‘**target**’. It is, however, pertinent to mention at this point in time that the evolved ‘**lead chemicals**’ are predominantly such compounds that are absolutely amenable to a broad-spectrum of ‘**chemical manipulations**’ to achieve the following *two* major objectives, namely:

- (a) To substantially optimize their existing **bioactivity** and **bioavailability** profile, and
(b) To look for and develop such rare compounds that may serve as ‘**potential candidates in their own right**’.

High Throughput Screening (HTS) process is presently the most important and highly procreative power for the useful and ultimate discovery of a variety of **newer lead chemical entities** in the **pharmaceutical industry**.

Miniaturized Assay: HTS indeed makes use of extremely specific **miniaturized assay formats** having the following salient features:

- Invariably uses **microtitre plates** capable of handling **384 sample variants** that may be assayed most conveniently at < 50 μ L total assay volume per run effectively,
- Based upon 100% fully automated sophisticated device one may carry out the assay of hundreds of thousands of samples **against each biological target of interest**, and
- Final numbers of samples actually being assayed exclusively depend upon the overall cost involved per assay, that may usually range between INR1 per well to INR 100 per well approximately.

HTS being a complex phenomenon that exclusively demands the following cardinal aspects, such as:

- (a) In-depth knowledge with respect to the critical role of **specific biological targets** in the domain of disease progression,
(b) The systematic development of **well-defined bioassays** that are solely capable of discovering desired regulators of the target,
(c) Soft targets with automation-friendly techniques with regard to the respective design, miniaturization, and ultimate ‘**automation of bioassays**’,
(d) Adequate thorough understanding of the **micro-** as well as **macro-structure** of the ensuing

'**biological target**' in such a manner that the prevailing '**sample-selection strategy**' is optimized appropriately,

- (e) Engineering excellence and master-piece of meticulously designed **custom-built robots** that may exploited judiciously for desired **storage, retrieval, and bioassay** of exceptionally large excess of samples within a span of 1 year, and
- (f) Aggressive development of '**numerous software systems**' that may go a long way in helping the scientists intimately engaged and associated with the handling of the copious volume of '**data**' that emerges eventually.

The **five** vital and important aspects of **HTS** enumerated as under:

- (a) **HTS** and Bioassays,
- (b) Access to Plants *vis-a-vis* Natural Source Materials,
- (c) **HTS** and Selection for Plant Materials,
- (d) Identification Process of Plants for Targeted Sets, and
- (e) Dereplication and Isolation of Active Compounds.

The aforesaid aspects of **HTS** shall now be discussed briefly in the sections that follows:

14.2.1 **HTS and Bioassays**

It has been duly observed that the '**ideal bioassays**' for **HTS** are those which critically enable **identification of chemical compounds** in *six* different manners, namely:

- (a) exerting action upon **specific biological targets**,
- (b) engage a minimum number of '**reagent addition steps**',
- (c) act **predictably and reliably**,
- (d) conveniently **amenable** to both **automation and miniaturization**
- (e) essentially involve **low-cost ingredients**, and
- (f) make use of **effective detection technology**.

Importantly, the '**biochemical targets of interest**' in the context of the most enterprising **pharmaceutical lead discovery** which invariably range from **enzymes** to **receptors*** to **ion channels**; however, in the specific instance of infectious disease to the entire microbial cells.

A few typical examples of **HTS and bioassays** are described as under:

Example: (a) Bradykinin (BK): A tissue hormone belonging to a group of hypotensive peptides invariably termed as **plasma kinins**. It was first ever obtained by incubation with the venom of *Bothrops jararaca* or with crystalline trypsin. It is used as a vasodilator.

In fact, **bradykinin (BK)**, an endogenous neuropeptide gets duly involved in the control and management of different kinds of pain associated with mammalian central nervous system (CNS). Therefore, systematic approach in inducing adequate antagonism of **BK** to its corresponding receptors is indeed a potential target towards the unique development of altogether new breeds of **analgesic agents**.

* **Receptor:** Both nuclear and transmembrane.

(b) *Candida albicans* cell-free translation system based upon **Polyurethane** as a **synthetic template** has been adequately proved and established to look for such chemical entities (*i.e.*, compounds) that essentially causes the inhibition of **critical fungal protein synthesis**.*

14.2.2 Access to Plants *vis-a-vis* Natural Source Materials

Ten Kate and Laird (1999)** reported duly that the **United Nations Convention on Biodiversity (CBD)** has been ratified in 1992 by almost 175 countries of the world. The most vital and important objectives of **CBD** are as stated under:

- (1) To ensure the conservation of biological diversity.
- (2) To implement sustainably its various cardinal components, and
- (3) To affect equitable and fair sharing of usefulness emerging from its application and usage.

In order to fully respect the '**letter and spirit**' of the said **Convention (CBD)** within the framework are several essential issues, such as:

- (a) Sovereignty of countries over their genetic resources, and
- (b) Country's reciprocal obligation to facilitate an easy access.

In fact, the *two* mutual contracting parties must establish appropriate means and ways for legitimately honouring the '**benefit sharing**' in the instance of **commercial utilization** of the genetic resources. The concerted efforts of the so-called **three agencies**, namely: (a) **collector**, (b) **source country**, and (c) **industrial collaborator** must prevail in all respects.

Glaxo Wellcome (UK) has come up with a **documented legal policy** to acquire '**natural source materials**' evidently lays down the various cardinal issues, essentially come across by a Pharmaceutical Company, to address such vital aspects as enumerated below briefly:

- that a Pharmaceutical Company shall solely collaborate with such firms or organizations who may amply demonstrate the qualified expertise together with genuine capability to provide **natural source materials** according to the schedule agreed upon in the '**mutual agreement**',
- that only comparatively small quantum of **plant materials** are duly collected from the **sustainable** natural sources,
- that '**endangered species**' *i.e.*, such species which are on the verge of extinction must not be accessed intentionally
- that a well defined 'materials transfer agreement is suitably drawn up which clearly ascertains that **collectors** are duly reimbursed towards the actual cost of collection together with their expertise,
- that the terms and conditions of the agreement should also specify expatiately **all financial benefits** in the event of **commercialization**,
- that the '**agreement**' must clearly state that a significant component of the ensuing '**royalty payment**' should be duly paid back to the source country so as to continue and sustain the **education and training** programme at the basic community level, and
- that major portion of **payments** must be affected duly during different successful stages of the on-going '**drug-development process**' (by Glaxo Wellcome).

* Kinsman OS *et al.* *J. of Antibiotics*, **51**(1): 41-49, 1998.

** Ten Kaite K and Laird SA: **The Commercial Use of Biodiversity**, Earth Scan Publications, London (UK), 1999.

14.2.3 HTS and Selection for Plant Materials

The actual impact of **HTS** and the suitable selection for plant materials always prevails predominantly in the event when a **'target'** belongs exclusively to a **'specific class'** that is rather difficult to find **small molecule hits**, such as:

- (a) protein-protein interactions,
- (b) occurrence of a strong precedent, and
- (c) rationale for natural product-derived activities.

Importantly, the complete compliance of the above three aspects legitimately command the desired **natural product input**.

Examples: The following typical examples would be further useful in the adequate clarifications of the above statement of facts:

- (a) **Antimicrobial Activity** *i.e.*, the track-record of various drug discoveries from a wide spectrum of **microbial sources**, and
- (b) **Analgesic Medicaments** *i.e.*, the same rationale shall hold good for the adequate track record of plant species in the production of analgesic medicaments.

Salient Features: The various **salient features** with regard to **HTS and selection for plant materials** are as stated under:

- (1) An easy and convenient access to diversified and huge collections of **natural plant materials**.

Examples:

- (a) Samples that are collected skilfully in order to add varying diversity to the important collection,
 - (b) Collections may include such samples that are specifically selected based upon various logical reasons *viz.*, microbial producer of a certain chemical entity, or a plant prominently employed **ethnomedically** for a certain prescribed parameter.
- (2) A diversity-based point of view certainly requires adequate gaining possession of pre-selected **taxonomic* groups**. Thus, one may make use of a variety of time-tested techniques to critically analyze the **natural taxonomic spread of a plant collection** which may subsequently be extended to minimise the existing gaps so that the ultimate collection distinctly reflects the **'available diversity'** more exhaustively.
 - (3) **'Chemical Targeting' and 'Biological Targeting'**: Recently, a much more critically focused approach exclusively based upon the **'prior available knowledge pertaining to some selected samples'** amply suggests that they invariably comprise of a good number of:
 - (a) highly specific **chemical classes of interest**, and
 - (b) essentially possess **desirable biological characteristic features**.

Interestingly, the aforesaid **'approach'** may be justifiably considered under *two* categories, namely:

* **Taxonomic:** Concerning the laws and principles of classification of living organisms.

- (i) **Chemical Targeting:** It accomplishes its cardinal objectives in *two* different manners, namely:
- makes use of **natural plant materials** as the prime sources of particular chemical compounds of great interest to a specific disease regimen, and
 - provides genuine and authentic sources of chemical class of compounds predicted to possess appropriate '**pharmacophore moieties**'.

In this manner, it would be quite feasible to identify the ensuing **chemical species** that are actively under due consideration in a **prevailing sample collection**.

- (ii) **Biological Targeting:** It may be regarded as to pursue a disease driven process. In actual practice, one may even select **plant samples** that may be utilized for the '**biological evaluation**' thereby providing some sort of relevant information associated with them which in turn could throw ample light with respect to their precise relevance for evaluation *vis-a-vis* a **given therapeutic target**.

Examples:

- (a) The **ethnobotanical** reports of **traditional medicinal applications** of plant materials, and
 (b) Commercially available **orthodox medicinals** duly discovered by **definite leads** given by indigenous knowledge.*

14.2.4 Identification Process of Plants for Targeted Sets

As to date a good number approaches have been duly pursued, worked out, and adopted in the process of adequate assimilation of the relevant information required necessarily so as to specifically select **plant materials** of particular relevance for a **given disease target**.

Table: 14.1. summarizes the various strategies adopted for the actual identification process of plants for the targeted sets:

Table 14.1 Strategies Adopted for Identification Process of Plants for Targeted Sets

S.No.	Research Group	Adopted Methodology	Comments
1	Ethnobotanical Network	Worked intimately with indigenous colleagues and traditional doctors in different countries.	Low output of actual plant samples for evaluation in laboratory. High output of valuable information(s) of their usage.
2	Pharmaceutical Companies	Make use of information(s) reported in Books, Journals etc.	Chinese Traditional Medicines; Indian Traditional Medicines.
3	Natural Products Alert [NAPRALERT] Database**	The system is maintained at the University of Illinois at Chicago (USA).	Contains huge number of references related to: <ul style="list-style-type: none"> • Ethnobotanical reports, • Reports of biological activity in scientific literature, and • Phytochemical data.
4	Chemical Information Databases	Dictionary of Natural Products [Chapman and Hall, New York]	Database contains informations on more than 100,000 natural plant products, including the plant species from where the chemical compound actually originates.
5	Literature Survey	To generate semi-purified plant extracts or chemical group of specific interest, or extracts that are enriched in the chemical entities.	Plants having an ethnomedical application the extracts may be prepared using recommended traditional medicine .

* Cox P.: **Ethnobotany and the Search for New Drugs**, Wiley, Chichester, pp. 25-41, 1994.

** Loub WD *et al.*: *J. Chem. Inform. and Comput. Scs.*, **25**: 99-103, 1985.

In short it may be added that these typical ‘**targeted**’ approaches invariably give rise to the actual involvement of rather smaller quantum of **natural plant samples** *vis-a-vis* a **high throughput random screening programme** [HTRSP].

14.2.5 Dereplication and Isolation of Active Compounds

From actual hands-on experience one may observe critically that before starting a complete **bioassay monitored fractionation of the active samples** it is absolutely important and necessary to carry out the review with respect to the **tolerance of a given bioassay** of either a **semi-purified** or a **crude** extract of **natural plant materials**. However, the cardinal objective of evaluating such samples in a **planned bioassay** is to identify **chemical entities** that eventually interact with a **specific biological target**, such as: a **receptor** or an **enzyme**.

However, from actual practice it has been duly observed that there are several unwanted, problematic, and trouble-shooter components that directly interfere with the **bioassay** *viz.*,

- **Polyphenolics**—forming complexes with a wide range of proteins, antioxidants, and UV quenchers,
- **Detergent-like Compounds**—causing disruption of cell membranes.

Van Middlesworth and Cannell (1998)* strongly advocated that it is very important and necessary to detect and subsequently get rid of all such **active chemical compounds** as early and swiftly as possible.

Importantly, the physico-chemical characteristic features of a pure isolated compound may provide extremely vital and useful clues with regard to its identity. In fact, the *two* most abundantly used properties, namely: (a) **HPLC** -retention time**, and (b) **UV-spectral data**, are invariably accomplished *via* well-known **standard analytical procedures**. However, one may easily compare these data *vis-a-vis* the known **reference compounds**; and, therefore, possibly characterize the various components present in a **mixture** without even going through the cumbersome process of **complete isolation** of each individual component.

Preparative HPLC, based on solvent partitioning, perhaps is the most common and widely used method for the isolation of pure chemical entities from various plant source materials.

Meticulous exhaustive studies towards the elucidation of the chemical structure of the pure isolated compound from various plant source materials shall render enormous valuable information(s) together with certain definitive indication of the superb and classical ‘**drug-like**’ qualities present in the molecule.

14.3 SUCCESS OF HTS OF PLANT SOURCE MATERIALS FOR NEW LEAD CHEMICAL ENTITIES

In the recent past, the discovery of a plethora of extremely potent and elegantly novel **euphane triterpenes** amply proves and demonstrates the actual enormous ability and potential of several plant

* Van Middlesworth F and Cannell RJP: *Dereplication and Partial Identification of Natural Products*: In: Cannell RJP (ed.): **Natural Products Identification**, Humana Press, Totowa, NJ, pp: 343-363, 1998.

** **HPLC**: High performance liquid chromatography [Kar, A: **Pharmaceutical Drug Analysis**, New Age International, New Delhi, **2nd** edn., 2005].

extracts to produce highly beneficial ‘**chemical leads**’ in a defined **HTS-programme**.

The above factual observations may be duly substantiated with the help of the following important investigative experimental results, namely:

- (a) **Inhibitors of Human Thrombin:** In the usual course of a ‘**random**’ screening exercise to look for certain ‘**novel inhibitors of human thrombin**’ that essentially help in the critical blockade of the actual formation of **blood clots**; and, therefore, may be duly exploited in the treatment, control, and prevention of the **deep-vein thrombosis**. For this meticulous and herculian task, a sizable (approx. 1,50,000) samples adequately derived from both natural sources *viz.*, **plant extracts, microbial extracts, fungal extracts**, and purely **synthetic chemical compounds** were subjected to **vigorous investigative evaluations**. The interesting outcome of this big-job revealed that the **methanolic extracts of *Lantana camara*** leaves, belonging to the natural order **Verbenaceae**, showed **remarkable potent activity**.

Sharma and Sharma (1989)* attributed the inherent toxic effects of the leaves of *Lantana camara* due to the presence of **lantadenes** *i.e.*, a series of **pentacyclic triterpenes**. Later on, Uppal *et al.* (1982)** reported the specific haematological manifestations occurring in sheep with ‘**lantana poisoning**’ associated with the following vital biological effects, such as:

- Enhancement in **blood-coagulation time** and **prothrombin time**,
- Reduction in **blood-sedimentation rate**,
- Total **plasma-protein content**, and
- Total **fibrinogen content**.

The aforesaid observation ascertains the corresponding **thrombin-inhibitory translactone** having the **euphane triterpenes**.

- (b) **Biological Activity** of the **euphane triterpenes** has been studied both extensively and intensively by Weir *et al.* (1998)***. This investigative study amply revealed their actual ‘**mechanism of action** as specific inhibitors for **blood clotting** through the strategic acylation of the available **active site(s)** *i.e.*, **Ser 195** residual segment of **thrombin**. It is pertinent to state here that the observed ‘**acylating activity**’ as the very root cause for being genetic against several other particular **serine protease enzymes**. Interestingly, this critical and vital observation forms the basis for both exhaustive exploration and extended exploitation in the ever-expanding domain of ‘**drug discovery**.’

14.3.1 Use of MS**** for Identification of Potent Biologically Active and Important Drug Molecules

After having isolated the pure chemical constituents from the **natural plant source** it becomes absolutely necessary to analyze such **compound(s)** or **protein(s)** of general interest, to elucidate and identity unambiguously their **exact positions** or **precise retention times (RTS)** by subjecting

* Sharma OMP and Sharma PD, *J. Scientific Industrial Research*, **48**: 471-478, 1989.

** Uppal RP and Paul Bs, *Ind. Veterinary J.*, pp 18-24, 1982.

*** Weir MP *et al. Biochemistry*, **37**: 6645-6657, 1998.

**** Mass Spectrometry.

them *via* the **selected adsorption phase** invariably employed in any known sophisticated type of **chromatography**.

Example: HPLC-Chromatograms: Each individual '**sample run**' shall give rise to a cluster of several sharp peaks, designating separate chemical compounds present, which may be systematically identified by the help of '**known standards**' under identical experimental parameters. Ultimately, it will certainly determine the **Rt*** of the pure **chemical entity** or the **protein of interest**. However, the presence of '**unknown compounds**', whose **RTs** are fairly comparable to those of the **standards (reference compounds)** may be identified by the aid of '**multiple forms of chromatography**' quite tentatively.

It is pertinent to mention here that either **Mass Spectrometry (MS)** or **Nuclear Magnetic Resonance (NMR) spectroscopy** analysis of the '**unknown peak**' should be carried out very carefully and painstakingly so as to identify the '**actual compound of interest**' unambiguously.


Identification of Taxol: **Taxol**, a natural product, obtained from *Taxol brevifolia* Nutt (*Fam. Taxaceae*), abundantly used for breast cancer (see Chapter 1) enjoyed the cogent application for its adequate and explicit characterization meticulously by **MS** by McClure *et al.* (1992)** and by **NMR** by Falzone *et al.* (1992)***.

A survey of literature would certainly reveal a good number of chemical compounds, derived from the natural plant sources and invariably used as '**drugs**', that have been duly characterized by the said spectral analyses *viz.*, MS, NMR.

14.4 HI-TECH PRODUCTS

Since 1990s a plethora of **hi-tech products** duly obtained from plant source materials have gained qualified success and well-deserved recognition across the globe for the therapeutic usages in humans. It will be worthwhile to discuss a few such **hi-tech products** at this point in time hereunder.

14.4.1 Genistein [*Syn. Genisteol; Prunetol;*]



4',5,7-Trihydroxy isoflavone;

Genistein is the phytoestrogen normally found in **soy products**. It represents the **aglucon of genistin** and of **sophoricoside******. It serves as a **specific protein kinase inhibitor**.

* **Rt:** It is the time (minutes) recorded duly on the **HPLC-chromatogram** where a given peak of interest occurs.

** McClure TD *et al. J. Am. Soc. Mass Spect.*, **3**: 672-679, 1992.


*** Falzone CJ *et al.: Tetrahed. Lett.*, **33**: 1169-1172, 1992.

**** **Sophoricoside:** It is genistein-4'-glucoside.

Characteristic Features: The various characteristic features of **genistein** are as enumerated under:

- (1) It lowers the incidence of breast, prostate, and several other carcinomas.
- (2) It minimises the incidence and number of cancerous tumours; and also to enhance the latency in the animal models of cancer.
- (3) In cell-culture models it particularly inhibits proliferation of certain types of cancer cells.
- (4) Various sophisticated well-defined techniques have been adopted *viz.*, **in vitro techniques**, **cell-culture techniques**, so as to decipher (unfold) the mechanism whereby **genistein** might change the ensuing **cancer-cell kinetics**.
- (5) **Genistein** causes the inhibition of **angiogenesis**,* steroid hormone receptors, inhibition of tyrosine kinase, inhibition of radical O₂-species formation, and above all interaction with **topoisomerase**.**

14.4.2 Camptothecin




Camptothecin is an **antitumour alkaloid**, which essentially serves a **prototype DNA topoisomerase I*** inhibitor**.

In fact, quite a few products are now the focus of attention, and have virtually undergone extensive preclinical and intensive clinical investigations, or even both of them together, namely:

- 9-amino-20S-camptothecine;
- 9-dimethylaminomethyl-10-hydroxy-20(S)-camptothecine [Topotecan[®]];
- 7-ethyl-10-[4-(1-piperidino) carbonyloxy-camptothecine [Irinotecan[®]]; and
- 9-nitro-20(S)-camptothecine.


14.4.3 Rhein [Syn: Monorhein; Rheic Acid; Cassic Acid; Parietic Acid; Rhubarb Yellow]



* Fotis T *et al.* *J. Nutr.*, **125**: 790S-79S, 1995.

** Barnes S. and Peterson TG: *Proc. Exp. Biol. Med.*, **208**: 109-115, 1995.

*** **Topoisomerase I**: An enzyme intimately involved in the uncoding of DNA, a prerequisite for **replication** and **transcription**.



Homoharringtonine

It is extensively employed in the treatment and control of malignancy.* Zhou *et al.* (1995)** reported promising results of **HHT** for the control treatment, and management of **acute nonlymphoblastic leukemias** and **chronic myelogenous leukemia**.

Mechanism of Action: Dwyer *et al.* (1986)*** duly observed that the **HHT** exerts its specific **cytotoxic effects** in the G_1 and G_2 phases of the prevailing cell cycle, which eventually synchronizes with the actual times of **intense protein synthesis**.

The protein synthesis really takes place in *two* distinct major stages, namely:

Stage-I: Initiation: In the course of '**initiation**' *mRNA* which essentially possess the code for the '**new protein**', gets hooked on to the **ribosome**. In doing so the *first tRNA* subsequently gets attached to the *mRNA* thereby **commencing the initial amino acid building block process for the protein**.

Stage-II: Elongation: Elongation refers to the particular phenomenon whereby the ensuing *tRNA* securely get attached to the *mRNA*. Thus, the resulting bonds are duly formed amongst the amino acids to give rise to the production of the desired **polypeptide protein**. Here, the **HHT** plays a vital role in causing inhibition of the '**elongation step**'. This unique phenomenon perhaps comes into being by actually competitively inhibiting the particular enzyme, **peptidyl transferase**, that rightly catalyzes the formation of the polypeptide bond, and rather remotely from inhibiting the bonding of *tRNA* to *mRNA* [Zhou *et al.* (1995)]

The aforesaid hypothesis expatiating the probable mechanism of action amply ascertains that **HHT** may cause induction of both **apoptosis** and adequate differentiation of the neoplasm cells, thereby rendering it to be a **potent newer antineoplastic drug**.

FURTHER READING REFERENCES

1. Bajaj YPS (Ed.): **Biotechnology in Medicinal and Aromatic Plants**, Springer Verlag, New York, 1988.
2. Bensky D and Foster S: **Chinese Herbal Medicine, Materia Medica**, Eastland Press, Seattle WA, 1986.

* Ohnuma T and Holland JF: *J. Clin. Oncol.*, **73**: 604-606, 1985.

** Zhou DC *et al. Bull. Canc.*, **82**: 987-995, 1995.

*** Dwyer PJ *et al. J. Clin. On col.*, **4**: 1563-1568, 1986.

3. Bisset NG (Ed.): **Herbal Drugs and Phytopharmaceuticals**, Medpharm, Stuttgart (Germany), and CRC-Press, Boca Raton, FL., 1994.
4. Combs SP: *Res. Staff. Phys.*, **43**: 54-57, 1997.
5. Markovits J *et. al. Biochem. Pharm.*, **50**: 177-186, 1995.
6. Olin BR (Ed.): **Review of Natural Products: Facts and Comparisons**, Wolters Kluwer Co., St. Louis MO, 1995.
7. Peterson G.; *J. Nutr.*, **125**: 784 S-789 S, 1995.

15

Indian Traditional Herbal Drugs

- Introduction
- Indian Traditional Herbal Drugs
- Further Reading References

15.1 INTRODUCTION

A large number of natural plant species, specifically those used extensively in various **Indian Traditional Herbal Drugs**, have been, and are still being investigated for ascertaining their specific inherent vital pharmacological and microbiological activities. However, quite a few notable overviews intimately associated with certain specific areas of intensive and extensive research are adequately dealt with practically in most of the chapters in the present compilation.

In the recent past, stretched over to almost two decades the spectacular thrust generated enough interest, inquisitiveness, and incredible latest scientific approach to the search for '**new drugs**' of tremendous potential value and worth in comparison to the modern allopathic system of medicine.

Based upon the high quality, proper standardization procedures, ultra-modern packaging concepts and ideas, exhaustively informative drug-usage literatures, and above all the broad-spectrum methodical promotions both in India and abroad, the **Indian Traditional Herbal Drugs** have undoubtedly made their presence felt amongst the valued consumers. An overwhelmingly plausible and sound confidence amongst the consumers to make use of such available drugs as : OTC products, prescribed medications, long-term usage in chronic ailments, have really turned them into a widely accepted alternative saga of safer and effective medications not only in India but also across the entire globe.

The importance of '**medicinal plants**' right from the very dawn of civilization up to the last couple of decades have witnessed a tremendous cumulative, informative, and educative volume of researches carried out in the ever-expanding field of pharmaceutically significant naturally occurring plant products. Interestingly, the better understanding of the plants as a whole *vis-a-vis* their important chemical constituents have undoubtedly broadend and strengthened one's acceptability and overall confidence in their usages amongst the consumers. Hence, the prevailing biodynamism of the '**active principles**' strategically located in the plant kingdom would certainly provide the mankind with an eternal store-house of clinically beneficial **herbal drugs**.

15.2 INDIAN TRADITIONAL HERBAL DRUGS

The **Indian Traditional Herbal Drugs** have grown in length and breadth with the advent of most sophisticated means of separation *viz.*, **preparative HPLC, preparative HPTLC, GLC**, and the like, ultimately gave birth to a host of '**phytopharmaceuticals**'.


A few very important and highly promising **phytopharmaceuticals** have been duly described in the following sections, wherein each individual class of drugs essentially deals with their name, chemical structure, natural source(s), and pharmacological activity.

15.2.1 Cardiovascular Drugs

The following '**herbal drugs**', *viz.*, **escin, curcumin, digoxin, and forskolin**, shall now be treated individually in the sections that follows:

15.2.1.1 Escin [Syn: Aescin; Aescusan; Reparil;]

Escin usually obtained as a mixture of **saponins** occurring in the seeds of the **horse chestnut tree**, *Aesculus hippocastanum* L. (Fam. *Hippocastanaceae*)




Escin

Escin is commonly used in the treatment vascular disorders, anti-inflammatory agent.

15.2.1.2 Curcumin [Syn: Turmeric Yellow;]

Curcumin is the natural dye stuff obtained from the root of *Curcuma longa* L. [Fam: **Zingiberaceae**]




Curcumin


It exerts anti-inflammatory activity.

15.2.1.3 Digoxin [Syn: Digacin; Dilanctin; Endigox; Lanacordin; Lanicor; Lanoxin; Lanoxicaps; Lenoxin; Neo-Dioxanin; Rougxin;]

Digoxin is the secondary glycoside obtained from *Digitalis landa* Ehrh., or *D. orientalis* Lam., (Fam. **Scrophulariaceae**). **Digoxin** serves as a potent cardiotonic.



Digoxin

15.2.1.4 Forskolin [Syn: Colforsin; Boforsin;]

Forskolin [or Colforsin]

Colforsin is a diterpene isolated from *Coleus forskohlii*, Briq. (Fam: Labiatae).


Forskolin possesses both vasodilating and cardiostimulatory properties.

15.2.2 Immunomodulators and Adaptogens

Indian traditional herbal drugs which specifically serve as **immunomodulators and adaptogens** are discussed as under:

15.2.2.1 Sitoindoside VII and VIII

The two **acylstearyl glucosides**, namely (a) **Sitoindoside VII**, and (b) **Sitoindoside VIII** have been duly isolated from the roots of *Withania somnifera* L., Dual., [Syn: *Physalis flexuosa*] belonging to the natural order **Solanaceae**.



Sitoindoside VII:

R = Glu (6'-O-acyl stearyl)


Sitoindoside VIII:

R = Glu (4'-O-acyl stearyl)

These two naturally occurring **glucosides** are invariably used to relieve the 'stress' in humans *i.e.*, they act as an **antistress drug**.

15.2.2.2 Syringin [Syn: Syringoside; Ligustrin; Lilacin; Methoxyconiferine;]

Syringin is obtained from the bark of *Syringa vulgaris* L., (**Lilac Bark**), and also from the **cambial sap** of spruce.



Syringin


Syringin exerts its action as an immunomodulator.

15.2.3 Antidiabetic Drugs

There are quite a few largely used **antidiabetic drugs** that are used in the **Indian Traditional Herbal Medicines**; however, only a couple of these drugs shall be discussed briefly as under, such as: **charantin**.

15.2.3.1 Charantin

Charantin is obtained from the fresh green fruits of the plant *Momordica charantia* L., (Fam. Cucurbitaceae).



Charantin

It is used for lowering the level of **'blood sugar'** significantly; and, therefore, used profusely for the control and treatment of diabetes mellitus.

Importantly, there exists an appreciable scope for the interaction with various reported **oral hypoglycemic drugs**.^{*} A survey of literature reveals that there are quite many allied herbal drugs related to *M. Charantia* (**Karela**) that are employed profusely both by the **Vaids**^{**} and **Hakims**^{***} for the effective, control, and management of **diabetes melitus** very much within the **Asian community** in particular, and quite recently in the **European community** in general.

15.2.4 Antineoplastic Drugs

The **Indian Traditional Herbal Drugs** that are exclusively employed as **'antineoplastic drugs'** are:

15.2.4.1 Bryostatins


In fact, the **Bryostatins** belong to a family of seventeen **biologically active macrolides** viz., Bryostatins 1 through 15, A and B. These **macrolides** are duly isolated from the **marine bryozoans**, *Bugula neritina* L., and other related organisms.

Bryostatins are usually employed as antineoplastic drugs, which specifically exert a partial agonist effect upon the protein **kinase C**.

* Aslam *et al. Lancet.* 1: 607, 1979.

** **Vaids**: The professional practitioners of Ayurvedic System of Medicine.

*** **Hakims**: The professional practitioners of Unani System of Medicine.




Bryostatin 1

Bryostatins are invariably used as immunomodulators.

15.2.4.2 Nicandrin B

Nicandrin B is obtained from the roots and seeds obtained from *Nicandra physaloides* L. Gaertn. (Fam: **Solanaceae**)*.



Nicandrin B

It is primarily responsible for showing *in vitro* cytotoxic activity.**

15.2.5 Antiviral Drugs

It is pertinent to mention here that the phenomenal success accomplished in the management of bacterial infection by the cautious usage of natural antibiotics (see Chapter 9) solely derived from microorganisms was not adequately matched to the same extent in the serious search for **antiviral**

* Rastogi *et al.* **Indian Medicinal Plants**, Vol. 2., PID, New Delhi, 1991.

** *Planta Medica*: 43: 389, 1981.


drugs'. Evidently, the various viral ailments still remain an important segment of medicine for which critical particular therapeutic treatments are lacking.

In the recent past, there are quite a few vital '**antiviral drugs**' have been identified, tested, and tried safely on humans which essentially belonged to the **Indian Traditional Herbal Medicine**.

Following are some of these '**drugs**' discussed briefly, such as:

15.2.5.1 (+)-Calanolide A

One may observe a very significant discovery in this field to date being the development of a series of **coumarins**—the **Calanolide A** and **Calanolide B** took place in a rather relatively smaller yield from the leaves and twigs of the forest tree *Colophyllum langigerum* (Fam: **Guttiferae**).




(+)-Calanolide A

(+)-Calanolide A possesses anti-HIV activity.

15.2.5.2 6-*epi*-Castanospermine

Bell and co-workers conducting research, at the London University, London, on certain specific and probable pesticidal non-protein amino acids, observed in the seeds of *Castanospermum australe* (Fam: **Leguminosae**) an altogether new **alkaloid** known as **6-*epi*-castanospermine**, which essentially has the **tetrahydroxyindolizidine moiety**.




6-*epi*-Castanospermine

It exerts its action as a potent inhibitor of **HIV**. The probable mechanism of action of **6-*epi*-castanospermine** against **HIV** is due to its ability to inhibit **glucosidase I and II**, that eventually monitor the formation of **glycoproteins** in the **viral coat**, and thereby without the presence of the essential enveloped structure of the ensuing virus to be able to infect the healthy **WBCs**.

15.2.5.3 Salaspermic Acid

Salaspermic acid is a pentacyclic terpenoid obtained from the roots of *Tripterigium wilfordii* (Fam: Celastraceae).



Salaspermic Acid

It shows inhibition of **HIV reverse transcriptase** and **HIV replication** in the HG lymphocyte cells.

FURTHER READING REFERENCES

1. Evans WC: **Trease and Evans Pharmacognosy**, 15th edn., Saunders, New York, 2004.
2. Mathee G *et al. Planta Medica*, **65**(6): 493-506, 1999.
3. McKee TC *et al. J. Nat. Prod.*, **60**(5): 431-438, 1997.
4. Rastogi RP [Ed.]: **Compendium of Indian Medicinal Plants**, Vol. 1 to 3, CDRI-Lucknow & PID, New Delhi, 1993.
5. Vlientinck AJ *et al. Planta Medica*, **64**(2): 97-109, 1998.

Index

Symbols

- (-)-Epicatechin 34, 289
(-) Arabinose 101
(-) Epigallocatechin 3-O-gallate [EGCG] 746
(-)-Rhamnose 101
(-)-Catechin 289
(-)-Cathinone 409
(-)-Ephedrine 409
(-)-Ergotamine 379
(-)-Form of santonin 551
(-)-Hyoscyamine 473
(-)-Lupinine 449, 452
(-)-Norephedrine 409
(-)-Norpseudoephedrine 660
(-)-Phellandrene 327
(-)-Quinine 379
(-)-Sparteine 452
(-)-Zingiberene 327
(+) – Galactose 101
(+) Glucuronic acid 101
(+)–Coniine 405
(+)–d-Tubocurarine chloride 379
(+)–Calanolide A 833
(+)–Citronellal 327
(+)–Cytisine 452
(+)–Ergotamine 379
(+)–Form of santonin 551
(+)–Lupanine 452
(+)–Norpseudoephedrine 409
(+)–Pinene 318
(+)–Pseudoephedrine 409
(+)–Quinine 379
(-)– α -Bisabolol (Bisabolane) 232
(-)–Melacacidin 168
(\pm)–ar–Turmerone 332
(\pm)–Ephedrine 407, 477
(\pm)–Form of santonin 551
(\pm)–Lupanine 450
(\pm)–Octopamine hydrochloride 703
(8 α , 9S)-Cinchonan-9-ol) 530
(9R)-Cinchonan-9-ol 526
(R)-Reticuline 495, 496
(S)-Form reticuline perchlorate 493
1, 5-Anhydrosorbitol 193
1, 8-Dihydroxyanthraquinone glycosides 761
1, Dihydro-ergotamine methane sulphonate 539
1-Dioxide 639
1-Isoborneol 220
1-Methyl β -carboline 503
1-Methyl-4-acetyl-1-cyclohexene 332
1-Perillaldehyde- α -anti-oxime 277
1-Perillaldehyde- α -syn-oxime 277
1-Rhodosamine 594
10-Deacetylbaecatin 824
10-Undecenoic acid 684
12, 34-Oxamanzamine F [18] 729
1,2-Dihydropyridine 460, 461
13-Cis Retinoic acid 240
1,3-Dihydroxy-N-methylacridone 436
1,4-Terpin 260
15-Cyanopuupehenone 732
15-Methoxypuupehenol 732
15 α -Cyanopuupehenol 731
15 α -Cyanopuupehenol (X) 732
15 β -Ethylpuupehenol 731
15 β -Methyl puupehenol 731
16-Acetate derivative of germine 424
19-Hydroxyoleanolic 338
19-NOR steroids 185
19:R=H;puupehenone 730
2-Chloro-4-amino-5-fluoropyrimidine 686
2-Cyano-4, 5-dibromopyrrole 717
2-Deoxstreptamine 590
2-Desoxyribose 201
2-Fluorovidarabine 699
2-Methyl- β -carboline 501
2-Oxosparteine 450
2-Phenylethanol 360
2-Thiazolidinone moiety 721
(20S)-Protopanaxatriol 188
22-33-Dihyrostigmasterol 190
2,4-Dichloro-5-fluoropyrimidine 686
26-Hydroxycholesterol 419
28-Deoxyzoanthenamamine 721, 724
2 β -Carbomethoxy 466
2D-NMR 734
2D-NMR spectroscopic studies 727
2F-Ara A 699
3, 4, 7, 15, 16-Pentaacetate derivative of germine 424
3, 4-Dihydroxycinnamic acid 341
3—Phenyl chroman 167
3-Acetate derivative of germine 424
3-Carene 258
3-Deoxykanosamine [Nabrosamine] 590
3-Dihydro-benzopyran tubic acid lactone 560
3-Formylrifamycin SV 658
3-Hydroxy-p-cymene 225
3-Methyl imidazole 437
3-Methyl-4-carboline 501
3-p-Cymenol 225
3-Phenylchromone 747
3-Phosphoglyceraldehyde 460
3,4-Dimethoxy cinnamic acid 192
3,5-Dinitrobenzoate 225
3 α -Tropanyl S-(–)-Tropate 469
3 β -Benzoxytropane 466
3D Picture of the ‘Lock’ and ‘Keys’ 78

- 4-Acetyxy-3-methoxy benzaldehyde 282
- 4-Benzenediol 700
- 4-Carene 258
- 4-Dimethy amino-intermediate 657
- 4-Keto intermediate 657
- 4-Methyl quinoline 429
- 5-Dehydro-quinic acid 208
- 5-Dehydro-shikimic acid 208
- 5-FC; 686
- 5-Fluorocytosine 681, 686
- 5-Fluorouracil 686, 687
- 5-HT 500
- 5-Hydroxy-L-Trp 500
- 5-Hydroxytryptamine 500
- 5-Methoxy-psoralen 171, 172
- 5-Nitroimidazole 668
- 5-Phospho-shikimic acid 208
- 5'-Deoxy-5-iodoberbericidin 708
- 6, 7-Dihydroxy-coumarin 346
- 6, 7-Dimethoxy-2 560
- 6- (β -D-xyloside) 141
- 6-Aminopenicillanic acid [6-APA] 631
- 6-APA 634, 640
- 6-Conjugate addition 731
- 6-Demethyl tetracycline 655
- 6-epi-Castanospermine 833
- 6-Hydroxyoleanoic acids 338
- 6-Methoxy-7-hydroxy coumarin 346
- 6-Methoxyumbelliferone 348
- 6-Methylene-5-hydroxytetracycline 653
- 6-Methylpreteramide 657
- 6-n-Tridecyl salicylic acid 720
- 7-Aminocephalosporanic 601
- 7-Chloro-4-keto intermediate 657
- 7-Chlorotetracycline 651
- 7-Deoxy-7(S)-chloro-lincomycin 617
- 7-Glucoapigenin 160
- 7-Hydroxy 6-methoxycoumarin 348
- 7-Hydroxy coumarin 346
- 7 β , 8 α -Dihydroxydepoxy-sarcophine 727
- 8-Acetate 203
- 8-Methoxy-psoralen 171, 172, 351
- 8-MOP 351
- 8-MP 351
- 9'-Rubanol 430
- A 3-Carene 262
- α and β glycosides 122
- α , γ -diaminodutyric acid (DABs) 643
- α - and β -glucogallin 133
- α - and β -heerabomyrrholic acids 333
- α -, β -, and γ -carotene 738
- α -, β -, and γ -commiphoric acids 333
- α -6-Deoxy-5-hydroxytetracycline 654
- α -Aminoadipic acid δ -semialdehyde 455
- α -Amyrin 186
- α - and β -guaiaconic acids 316
- α - and β -peltatins 319
- α -Borneol 220
- α -Carotene 738, 740
- α -Chymotrypsin 635
- α -Configuration 123
- α -Curcumene 327
- α -Cyclodextrin 96
- α -Ergocryptine 542
- α -Estradiol 12
- α -Farnesene 229
- α -Glucogallin 133
- α -Interferon 62
- α -Ionone 274
- α -Kainic Acid 722
- α -Lignan 362
- α -Lipoic acid 771, 772
- α -Lobeline 443
- α -Methylene γ -lactone system 228
- α -Methyl 437
- α -Peltatin 319
- α -Phellandrene 255
- α -Phytosterol 760
- α -Pinene 26, 216, 218, 219, 250, 260, 261, 328
- α -Pyrone 553
- α -Quinidine 529
- α -Santalene 229
- α -Santonin 31, 550, 551
- α -Terpene 255
- α -Terpinene 260, 270, 299
- α -Terpineol 268, 270, 271, 743
- α -Thujane 259, 260
- α -Tocopherol 771
- α -Tocotrienol 770
- α/β -Unsaturated carbonyls 228
- β , β -Carotene -3, 3'-diol 741
- β , ϵ -Carotene-3, 3'-diol 741
- β -Amygdalin 123, 124
- β -Amyrin 186, 210
- β -Barbaloin 129
- β -Bisabolene 327
- β -Bitter acid 549
- β -Cadinene 228, 229
- β -Carboline 500
- β -Carboline carboxylic 503
- β -Carboline ring system 500
- β -Carotene 13, 32, 81, 738, 739, 740, 808
- β -Carotene oxygenase 240
- β -Carotenes 239
- β -Carotenoids 240
- β -Caryophyllene 228
- β -Cytosine arabinoside 698
- β -Cocaine 466
- β -Configuration 123
- β -Cyclodextrin 96
- β -D-glucose 155
- β -Ergocryptine 542
- β -Ergoptine 542
- β -Ergosine 542
- β -Farnesene 229
- β -Galactosidase 49
- β -Galactosidase activity 49
- β -Glucogallin 133
- β -Glucopyranoside 36, 180
- β -Hydroxyethylbenzene 360
- β -Ionone 239, 274
- β -Lactam 611
- β -Lactam antibiotics 611, 621, 639, 647, 648
- β -Lactam bond 638
- β -Lactam variants 611
- β -Lactamase 600, 636, 638, 639
- β -Lactamase combination 630
- β -Lactamase inhibitors 638, 579, 640
- β -Lactamase-producing strains 605
- β -Lactams 579, 611, 614
- β -Lupulic acid 549
- β -Metacapaivic acid 324
- β -Methyl adipic acid 275, 276
- β -Methyl- β -carboxy propanol 437
- β -Methylesculetin 348
- β -Murcene 334
- β -Myrcene 25, 252, 334
- β -Myrecene 217

- β -Peltatin 319
 β -Phellandrene 255
 β -Phenylacrylic acid 753
 β -Phenylethyl alcohol 360
 β -Pilocarpine 439
 β -Pinene 252, 260, 328
 β -Quinine 527
 β -Salicin 123, 124
 β -Sesquiphellandrene 327
 β -Sitosterol 760, 761
 β -Terpene 255
 β -Terpinene 260
 β -Tocopherol 771
 β -Tocotrienol 771
- A**
- Aaptamine 708
 Abbocin 652
 Abietaceae 321
 Abietic acid 216, 310, 311
 Abricycline 650
 Abridged phenylpropanoids 340, 354
 Abrifam 658
 Abrin 794
 Abscisic acid 228, 807
 Absorbable gelatin sponge 792
 Absorbent cotton 98
 Abyssinian coffee 367
 Acacia 101, 244, 315
 Acanthifolicin 718, 719
 Acef 602
 Acerola 37
 Acetate 210, 267
 Acetate pathways 205
 Acetate replacing factor 771
 Acetate-derived alkaloids 401
 Acetate-malonate pathway 364
 Acetate-mevalonate pathway 364
 Acetic Acid 206
 Acetone 273, 276, 284
 Acetyl salicylic acid (Asprin) 11
 Acetylation of puupehenone 730
 Acetylcholine 704
 Achro 650
 Achromycin 650
 Achuma 479
 Acid casein 793
 Acid phthalate 225
 Acid quinine hydrochloride 524
 Acid resins 313
 Acid saponins 184
 Acid sulphate of quinine 431
 Acidic bitter principles 548
 Acipen-V 631
 Aclacinomycin A 590, 594
 Aclarubicin 594
 Aconine 400, 412, 413
 Aconine hydrobromide sesquihydrate 413
 Aconine hydrochloride dihydrate 413
 Aconite 341, 412, 450, 477
 Aconitine 376, 389, 396, 397, 400, 412, 413, 414
 Aconitine hydrobromide hemipentahydrate 414
 Aconitine nitrate 414
 Aconitine-skeleton 415
 Acontine-type 414
 Acquired Immunodeficiency Syndrome (AIDS) 65
 Acridine alkaloids 434, 435
 Acridine-ring-system 434
 Acridines 396
 Acronize 651
 Acronycine 435, 436
 Acsulin 169
 Actimmune(R) 42, 79
 Actin 790
 Actina 680
 Actinia equina 705
 Actinidine 412
 Actinitine 412
 Actinomycetaceae 201
 Actinomycetales 570
 Actinomycete antibiotics 649
 Actinomycetes 570
 Actinomycetin fungi 17
 Actinomycin 692
 Actinomycin A_{IV} 692
 Actinomycin C₁ 692
 Actinomycin D 692
 Actinomycin I₁ 692
 Actinomycin X₁ 692
 Activase (R) 79
 Activating enzyme 782
 Active chemical compounds 820
 Active constituents 21, 24
 Active metabolite 659
 Acute phase protein 790
 Acyclic 228
 Acyclic (Aliphatic) alcohols 266
 Acyl carnitine 766
 Acyl moieties 574
 Acyl-coA metabolites 766
 Acyl-glycosides of anthocyanidins 745
 Acylating activity 821
 Acylfilicinic acid 325
 Adaptogens 830
 Adenimine 44
 Adenine 48, 201, 202
 Adenosine 125, 126, 706, 707
 Adenosine deaminase (ADA) 78, 707
 Adenosine deaminase deficiency (ADD) 78
 Adenosine kinase 708
 Adenosine triphosphate (ATP) 790
 Adenylate cyclase activity stimulator 812
 Adotodai 427
 Adrenaline 481, 482, 704
 Adriacin 591
 Adriamicina 653
 Adriamycin 6, 591, 593
 Adriamycinone 591, 592
 Adriblastina 591
 Adultrant 266
 Advantages of PCR-Technique 58
 Aerial segments of plant materials 803
 Aeroplysinin 718
 Aeroplysinin-1(-) 718
 Aeroplysinin-1(+) 718
 Aerosporin 643
 Aerugipen 637
 Aescin 828
 Aescusan 828
 Aesculetin 346
 Affinitins 794
 Aflatoxin-B1 769
 Aflorix 689
 African rue 501
 Agar 114, 115
 Agents for mycobacterium tuberculosis infections 665, 678
 Agents for systemic UTIs 665, 676
 Agglutinins 794

- Aggregates of fungal cells 573
Agit 539
Aglucon 33
Aglucon of genistin 822
Aglycone 122, 124, 146, 183, 184
Aglycone moiety 127
Agristrep 587
Ague wood 367
AIDS 69
AIDS test 67
AIDS vaccines 66
Ajmalicine 504, 505, 506, 509, 510
Ajmalicine hydrobromide 506
Ajmalicine hydrochloride 506
Ajmaline 381, 501
Ajowan oil 225
Aklaviketone 592, 593
Aklavinone 593, 594
Akuaminicine hydrochloride dihydrate 507
Akuaminicine perchlorate monohydrate 507
Akuammicine 504, 505, 506, 507
Akuammicine hydroiodide monohydrate 507
Akuammicine methiodide 507
Akuammicine nitrate 507
Ala tet 650
Alamycin 652
Albamycin 663
Albasapogenin 187
Albaspidin 325, 326
Albipen 633
Albistat 689
Albumin 21, 368, 787
Alcohol 1, 4, 26
Alcohol dehydrogenase 747
Alcohol detoxification enzymes 747
Alcohol volatile oils 251, 264, 266
Alcomycin 582
Aldehyde glycosides 127, 195, 210, 214
Aldehyde volatile oils 251, 271
Aldehydes 27
Aldobionic acid 106
Aleuritic acid 320, 321
Alexandria senna 137
Alexandrium tamarense 711
Alfalfa 34, 463
Alfamox 633
Alfifetin 645
Algin 35, 113
Alginic acid 113
Alicyclic hydrocarbon 26, 254
Aliphatic ketones 284
Aliphatic terpene aldehydes 272
Alizarin 203
Alkaloid biosynthesis 531, 542
Alkaloid-precipitating reagents 384
Alkaloidal 200
Alkaloids 4, 21, 199, 340, 372, 373, 374, 375, 377, 378, 380, 381, 382, 383, 384, 385, 387, 388, 390, 391, 393, 394, 395, 396, 397, 400, 401, 405, 427, 462, 542, 543, 724, 811, 824
Alkaloids derived from amination reactions 400, 401
Alkaloids derived from anthranilic acid 400, 427
Alkaloids derived from histidine 400, 436
Alkaloids derived from lysine 400, 441
Alkaloids derived from nicotinic acid 401, 454
Alkaloids derived from ornithine 401, 461
Alkaloids derived from tryptophan 401, 495
Alkaloids derived from tyrosine 401, 475
Alkaloids in tissue cultures 542
Alkalover 752
Alkamine 423
Alkamine aglycones 420
Alkyl group 382
All-trans retinoic acid 240
All-trans- β -Carotene -3, 3'-diol 741
Allelochemicals 217
Allicin 759
Allied drugs 156
Allied drugs of digitalis 148
Alliin 759
Allose 35
Allosteric enzyme 782
Allspice 297
Allyguaicol 28
Allyl iso-thiocyanate 181, 246, 757
Allylamines 685
Allylic 759
Allylic sulphides 759
Almodan 633
Alocresinotannol 312
Aloe emodin 129
Aloe emodin-8-glucoside 138
Aloe juice 774
Aloe vera gel 774, 775
Aloe vera plant 762
Aloe-emodin 128, 132, 138, 761
Aloe-emodin anthrone diglycoside 138
Aloes 127
Aloesin 129, 130
Aloin 122, 126, 129, 775
Aloin-like C-Glycosides 134
Aloinose 775
Alpen-N 634
Alpha (TNF-a) 77
Alpha-glycoside bondage 90
Alteration in cellular membrane function 664
Alternate drug 664
Altracin 645
Aluminium isopropoxide 660
Amarogenin 198
Amarogentin 198, 743
Amaroswerin 199
Amaryllidaceae alkaloids 475
Ambisome 683
Ambracyn 650
Ambramicina 650
Ambramycin 650
Amcill-S 634
American boxwood 367
American hellebore 420, 425, 426
American oil of turpentine 249
American podophyllum 319
American storax 334, 335
American unplanted cotton 498
Ametycine 693
Amfipen 633
Amicardine 553
Amidazine 680
Amides 396
Amikacin 18, 576, 580, 581
Amikavet 581
Amikin 581
Amiklin 581
Amines 396

- Amino acids 36, 401
Amino alkaloids 396
Amino glycosides 18, 579, 580, 589, 606
Aminodeoxy 583
Aminoform 677
Aminoglycoside antibiotics 576, 580
Aminoglycosidic-production-paths 17
Aminopenicillins 627, 629, 633, 635
Ammicardine 553
Ammipuran 553
Ammivin 553
Ammivisnagen 553
Ammoform 677
Ammoidin 351
Ammonia 438
Ammoniacum 310, 329, 330, 331
Ammoniated glycyrrhiza 191
Ammonium vanadate 517
Ammoresinol 330
Ammoresinotannol 312
Amocilline 633
Amoksiklav 639
Amolin 633
Amopenixin 633
Amoram 633
Amoxicillin 579, 629, 633, 639
Amoxicillin trihydrate 639
Amoxidin 633
Amoxipen 633
Amoxycillin 633
Amoxyphen 633
Ampho-moronal 683
Amphoteric alkaloids 383
Amphoteric glycopeptide antibiotic 646
Amphotericin 689
Amphotericin B 681, 683
Amphozone 683
Ampicillin 600, 602, 629, 633, 635
Ampicillin sodium 634
Ampicilin 634
Ampipenin 633
Amudane 681
Amygdalin 4, 35, 36, 122, 175, 176, 177, 246
Amylases 781, 782, 786
Amyloid 754
Amyloid- β -aggregation 754
Amyolytic enzyme 782
Amylopectin 90
Amylose 90, 91
Amymicin 623
Anabactyl 636
Anabasine 454, 457, 458, 459, 461
Anabasine hydrochloride 458
Anaerobic 669
Analetica pharmacognostica 1
Analgesic 124
Analgesic agents 816
Analgesic medicaments 818
Analogue 595
Analogue of nereistoxin 725
Analytical chemistry 369
Anaprel 513
Anaptivan 604
Ancef 602
Anchovyxanthin 741
Ancobon 686
Ancotil 686
Andergin 689
Anemolin 633
Anemonia Toxin-II (ATX-II) 702
Anethole 251, 291, 292, 293, 344
Anethole-rich oils 292
Aneurine hydrochloride 37
Angelica 352, 353, 556, 557
Angelicin 556, 557, 558
Angiogenesis 823
Angionorm 539
Angiotensin II 780
Angiotensin-converting Enzyme [ACE] 782
Anglicin 171, 172
Angular furanocoumarin 207
Anhalamine 481, 484, 485
Anhalamine hydrochloride dihydrate 483
Anhaline 478
Anhalonidine 481, 483, 484, 485
Anhalonine 481, 483, 484, 485
Anhalonine hydrochloride 483
Animal enzymes 781
Animal proteins 789
Anionic glycosides 755
Anisaldehyde 292, 293
Anisaldehyde test 563
Anise 347
Anise camphor 291
Anise fruit 292
Anise oil 248
Aniseed 292
Anisole 292
Anisole propionaldehyde 292
Anisole-p-(1-chloropropane) 292
Anjir 350
Ansamycin antibiotics 658
Ansamycins 658
Anthelmintic 552, 723
Anthelminticum 299
Anthocyanidin glycosides 159, 167, 212
Anthocyanidins 157, 158, 745
Anthocyanins 167, 777
Anthopleurin-A (AP-A) 702
Anthopleurins A, B and C 701
Anthracene aglycones 203
Anthracene glycosides 127, 211
Anthracycline 579, 590, 592, 595
Anthracycline antibiotics 590
Anthracycline drugs 68
Anthracyclinones 590, 595
Anthranilic acid 427
Anthraniloyl-CoA 428, 436
Anthranol glycosides 135
Anthranols 129, 761
Anthraquinone 203, 590, 761, 811
Anthraquinone glycosides 29, 762
Anthrocyanidine 167
Anthrones 129, 761
Anti-asthmatics 814
Anti-LPS Mab 80
Anti-rheumatic 124
Antiapoptotic 755
Antibacterials 568
Antibiotic 631
Antibiotic 16, 21, 568, 569, 575, 579, 638, 657, 696
Antibiotic activity 571
Antibiotic complex 582, 584, 589
Antibiotic component 570, 571
Antibiotic development 569
Antibiotic glycosides 201, 214
Antibiotic level 578
Antibiotic MA 144 A1 594
Antibiotic masking 577
Antibiotic masking of microbial contaminants 577
Antibiotic of second choice 642

- Antibiotic production phase 572
 Antibiotic substances 696, 576, 718
 Antibiotics based on mechanism of action 657
 Antibiotique EF 185, 585
 Antibodies 75
 Antibody-mediated immunity (AMI) 742
 Anticancer antibiotics 665, 690
 Antichokes 750
 Antidiabetic drugs 830
 Antifeedants 217
 Antifungal agents 665, 680
 Antifungals 568
 Antigens 75
 Antiinflammatory 696, 720
 Antiinflammatory and antispasmodic agents 696, 720
 Antilirium 533
 Antimalarials 812
 Antimetabolites 667
 Antimicrobial 568
 Antimicrobial activity 818
 Antimicrobial drugs 696, 716
 Antimony trichloride test 147
 Antimycobacterial agents 679
 Antimycobacterial drugs 678
 Antineoplastic 16, 696
 Antineoplastic agent 362, 699, 744, 789, 814
 Antineoplastic drugs 831
 Antioxidant 239, 740, 771, 777
 Antirheumatic 824
 Antirobe 617
 Antispasmodic 720
 Antitumor necrosis factor 80
 Antitumour alkaloid 811, 823
 Antiviral 696
 Antiviral drugs 832, 833
 Antril 80
 AP-A 701, 702
 AP-A, B, C 701
 AP-C 701
 Aparika 36
 Aphrodine 511
 Aphrodyne 511
 Apigenin 159, 160
 Apigenin (C) 206
 Apihepar 750
 Apiin 159, 160
 Apioaldehyde 297
 Apiole 29
 Apiole 29, 291, 296, 297, 300
 Apiole tribromide 296
 Apioic acid 297
 Apiose 159, 160
 Aplysinopsin 721, 724
 Aplysistatin 699
 Apoptropine 388
 Apoptropine hydrochloride 380
 Apocarotenoids 239
 Apocynaceae 515
 Apomorphine 388, 488
 Aporphine 399
 Aporphine alkaloids 396
 Apoterin S 513
 Applications of yeast 795
 Apsin VK 631
 Ara-A 723
 Ara-C; Alexan 698
 Arabian coffee 367
 Arabian senna 139
 Arabica coffee 367
 Arabinose 107, 333
 Arabinosyl nucleoside 707
 Arabitin 698
 Aracati jaborandi 437
 Arachidonic acid 759
 Aracytidine 698
 Aracytine 698
 Arariba rubra mart 501
 Arbutin 139, 140, 141, 204
 Arcacin 631
 Arecoline 389, 398
 Arginine 712, 789
 Aribine 501
 Arilin 669
 Arithromycin 619
 Arnyl 656
 Aromatic hydrocarbon 26
 Aromatic ketones 284
 Aromatic terpene aldehyde 280
 Arrow poison 8
 Artemisinin 19, 6, 230, 231, 812, 814
 Artery-clogging trans fats 776
 Artificial culture medium 799
 Asafetida 359
 Asafetida 329
 Asafoetida resin 329
 Asafoetida 310, 329, 343, 347
 Asant 329
 Asaresinotannol 329, 347
 Ascabin 302
 Ascabiol 302
 Ascaridole 299, 300
 Ascaridole 291, 299, 300
 Ascaridole glycol 300
 Ascarisin 299
 Ascorbic acid 37
 Ashwagandha 349, 374, 397, 462
 Asiatic acid 193
 Asistic cotton flower 34
 Asparaginase 782
 Asparagine 190
 Aspenil 633
 Aspergillus niger 358
 Asperidol 700
 Aspidinol 325
 Aspidosperma 505
 Astaxanthin 738, 742
 Asthma weed 39, 443, 444
 Astrosaponins 705
 ATCC 11137 729
 ATCC 11925 729
 ATCC 7601 729
 ATCC-11145 727
 ATCC-24795 727
 Atiine-type 414
 Atirin 602
 Atisine 400, 414
 Atisine-skeleton 415
 Atropine 10, 12, 374, 380, 381, 388, 389, 396, 398, 464, 465, 470, 810
 Atropine hydrochloride 465
 Atropine methyl bromide 465
 Atropine methylnitrate 465
 Atropine sulphate 380, 387
 Atropine sulphate monohydrate 465
 Atropisol 465
 Attaway 710
 Attractants for polliniferous 217
 Attrition mill 88
 ATX-II 701, 702
 Augmentin 639
 Aureine 474
 Aureococina 651
 Aureomycin 651
 Aurichloride 422
 Aurones 157, 158
 Autolytic enzyme 782

- Automation and miniaturization 816
Automation of bioassays 815
Autonomium chloride 702, 704
Auxin 804, 807
Auxotrophic mutants 369
Ava-ava 322
Aveicularin 161
Averon-1 601
AW-105-843 685
Axsain 323, 410, 768
Ayfivin 645
Azactam 613
Azadirachtin 238
Azasteroid 425
Azetidine-2-carboxylic acid (I) 806
Azithromycin 622, 623
Azoles 687
Azonom 613
Azthronam 613
Aztreon 613
Aztreonam 611, 612, 613
Azudoxat 654
- B**
- B-cells 46, 75
B-lymphocytes 78
Bacampicillin 629, 635
Bacampicine 635
Baccatin 824
Baccidal 673
Bacitracin 573, 644, 645, 646
Bacitracin A 645, 546
Bacitracin F 646
Bacitrasin 641
Bacogenin A 193
Bacogenin B 193
Bacopa 193
Bacoside A 193
Bacoside B 193
Bacterial mammalian tissue culture 58
Bacterial vaccine 72
Bactericidal 647, 668
Bacteriophage lambda 53
Bacteriostasis 668
Bacterophage 48
Bactopen 632
Bad cholesterol 748, 776
Baldo 303
Baljet reagent 147
Baljet test 147, 155
Balsam of Peru 303, 309, 335, 346, 359, 753
Balsam of Tolu 307, 310, 336
Balsam storax 334
Balsam tree 318
Balsamic acids 336
Balsamic resin 337
Balsams 218, 314
Banan 609
Banisterine 502
Bantron 444
Barayata-saponin complex 184
Barazan 673
Barbados aloe 341
Barbaloin 122, 126, 129, 134, 135, 761, 762
Barbaloin (Aloin) 134
Bareon 673
Barettin 721, 725
Bark of poplar (populus) 359
Barqui simento jaborandi 437
Base-sugar unit 201
Basergin 538
Basic cephalosporin ring system 610
Basic structures of cinchona alkaloids 429
Bassorin 104
Bavachromanol 172
Baxan 602
Bay 493
Bay oil 287
Baycip 672
Bayciper 637
Bayer 5312 680
Baypen 637
Bdellium 310
Bead-type culture technique 802
Bearberry 357
Bedome 37
Beef-steak plant 290
Begoniaceae 236
Bekanamycin 583
Belladonna 348, 397, 462, 465
Belladonna alkaloids 386
Belladonna leaves 469
Belladonna 374
Benzaldehyde 176, 246, 280, 406
Benzaldehyde test 335
Benzathine 626
Benzene 553
Benzeneethanol 360
Benzo [α]pyrene 758
Benzo- α -pyrone 553
Benzo- γ -pyrone 553
Benzoic acid 751
Benzofuran 553
Benzoic acid 178, 337, 356, 357, 753
Benzoin 303, 307, 310, 334, 337, 338, 356
Benzopyran 345
Benzoresinol 312
Benzoyl acetic acid 446
Benzoyl-CoA 473
Benzoylacetate 407
Benzoylacetate-CoA 447
Benzoylmethylecgonine 466
Benzyl benzoate 302
Benzyl carbinol 360
Benzyl cinnamate 334
Benzylhydroxymethylketone 477
Benzylisoquinolines 396
Benzylpenicillin 596
Benzyltetrahydroisoquinoline 485
Berberidaceae resins 313
Berberine 380, 381, 382, 398
Bergacef 603
Bergamot 302, 350
Bergamot oil 302
Bergamot peel 243
Bergapten 171, 172, 350, 352, 353, 556, 558
Bergapten (e) 352
Bergaptol 558
Berkfurin 677
Beromycin 631
Bertrand's reagent 385
Beta (TNF-b) 77
Beta interferon 80
Beta lactamase combinations 627
Beta seron (R) 79
Beta-lactamase combinations 638
Betabactyl 639
Betacef 610
Betaine 372
Betamaze 640
Betamox 633
Betapen VK 631
Betaseron (R) 42

- Bethacil 640
Betony 463
Betula oil 30
Bewon 37
Bialatan 653
Biazolina 602
Bicyclic 228
Bicyclic monoterpene hydrocarbons 258
Bicyclic monoterpenes 259, 287
Bicyclic pyrrolizidine nucleus 471
Bicyclic terpene aldehydes 279
Bicyclic terpene ketones 287
Bidocef 602
BIIIB 725
Biklin 581
Biletan 771
Biliter apple 342
Biliter principle 324
Bio-tetra 650
Bioactivity 815
Bioavailability 739, 815
Biochemical reactions 799
Biochemical targets of interest 816
Biociclin 604
Bioconversion studies 728
Bioenergetic materials 781
Biofurex 604
Biogenesis 24, 119
Biogenesis of carotenoids 238
Biogenetic isoprene rule 216
Biogenetic relationship of flavonoids 364
Biological activity 821
Biological assays 815
Biological evaluation 819
Biological source 653
Biological target 816, 814, 818, 819
Biological-amines 372
Biologically active macrolides 831
Biologically-active-prototypes 4, 10
Biomass 801, 802
Biomedicinals 797, 806
Biomitsin 651
Bioproduction 797
Biosol 585
Biosynthesis 264, 418, 588
Biosynthesis of aconitine-type alkaloids aconite 414
Biosynthesis of ajmalicine, vindoline and catharan 509
Biosynthesis of aldehyde glycosides 210
Biosynthesis of alizarin 204
Biosynthesis of aminology 18
Biosynthesis of anhalamine 484
Biosynthesis of anhalamine, anhalonine and anhalon 485
Biosynthesis of anthracene glycosides 203
Biosynthesis of brevetoxin [Pb Tx-1] 714
Biosynthesis of capsaicin 411, 412
Biosynthesis of castanospermine and swainsonine 454, 455
Biosynthesis of chloramphenicol 662, 663
Biosynthesis of chlortetracycline 656, 657
Biosynthesis of cinchonine, quinidine and cinchoni 530, 531
Biosynthesis of coumarin and furanocoumarin glycos 205
Biosynthesis of cyanogenetic glycosides 207
Biosynthesis of doxorubicin and epirubicin 592
Biosynthesis of elaeagnine 503
Biosynthesis of ephedrine alkaloids 409
Biosynthesis of ergotamine 540, 541
Biosynthesis of flavonoid glycosides 204
Biosynthesis of γ -coniceine and coniine 405
Biosynthesis of griseofulvin 682
Biosynthesis of hordenine and mescaline 481, 482
Biosynthesis of hygrine 471, 473
Biosynthesis of imidazole alkaloids 441
Biosynthesis of lobeline and lobelanine 447
Biosynthesis of lupinine, lupanine and sparteine 451, 452
Biosynthesis of morphine 494, 497
Biosynthesis of nicotine 459
Biosynthesis of phenol glycosides arbutin 204
Biosynthesis of phenylpropanoids 369, 370
Biosynthesis of physostigmine 534
Biosynthesis of piperine 448
Biosynthesis of psoralen 558
Biosynthesis of quinine 433, 434, 524, 530
Biosynthesis of retronecine and senecionine 475, 476
Biosynthesis of rotenone 561
Biosynthesis of rutacridone 436
Biosynthesis of saponin glycosides 209
Biosynthesis of secondary metabolites 798
Biosynthesis of serotonin 499
Biosynthesis of solasodine 419
Biosynthesis of the glucosinolates 756
Biosynthesis of various monoterpenoids 265
Biosynthesis of vasicine 428
Biosynthesis of yohimbine 518
Biosynthetic classification 396
Biosynthetic pathway 520, 810
Biosynthetic pathway of components of streptomycin 589
Biosynthetic pathway of doxorubicin and epirubicin 593
Biosynthetic penicillins 626
Biotechnology 41, 60, 61, 79, 82, 571, 797, 798
Biotechnology and modern drug discovery 79
Biotechnology based pharmaceuticals 72
Biotechnology drugs 71
Biotechnology medicines 79
Biotechnology-based products 42, 800
Bioterciclin 652
Biotexin 663
Biotransformation reactions 798
Biotransformations 799
Bioxima 604
Biozolene 690
Biphenylpropanoid 363
Biphenylpropanoid derivatives 340, 361

- Biquinate 524
Birutan 749
Bistriazole antifungal agent 690
Bisulphite adduct 282
Bitter almond 174, 175
Bitter apple 341
Bitter compounds 547
Bitter glycosides 127, 197, 198, 199, 214, 277
Bitter lettuce 469, 564
Bitter nightshade 397, 415
Bitter orange 163
Bitter principles 21, 199, 547, 561
Bitter principles of cucurbits 236
Bitter sweet 200, 397, 415
Bitterwood 357
Bivatin 37
Bixin 809
Black balsam 335
Black cohosh 357, 367
Black locust 305
Black mustard 180, 181
Black mustard seeds 36, 343
Black nightshade 416
Black pepper 294, 295, 373, 768
Black raspberries 772
Black snakeroot 357, 367
Black tea 747
Black tea solids 746
Blepharitis 661
Blood 61
Blood clotting 64
Blood protein 790
Blood-brain barrier (BBB) 742
Blue rocket 341, 412, 450, 477
Bo-Xan 31, 741
Boat and chair form of isomers 256
Boiling range 250
Boldine 396, 397, 399
Boldo 346
Bonapicillin 629, 633
Bone-mineral density 776
Boneset 367
Bor-cefazol 602
Borax test 102, 130
Bornane 264, 265
Borneol 219, 220, 232
Borneol camphor 219
Bornyl alcohol 261
Bornyl chloride 219, 261
Borrachero 397
Boxberry 304, 342, 343
Bradykinin 788
Bradykinin (BK) 816
Brahmic acid 193
Brain-antioxidant 742
Branching enzyme 782
Brassia alba 182
Brassica campestris 125
Brassica juncea 209
Brassica protection products (BPP) 774
Brazilian arrowroot starch 91
Brentan 689
Brevetoxin 712, 713, 714
Brevetoxin (Pb Tx-1) 714
Brewer's yeast 795
Bridchilli 323
Bristaciclina 650
Bristagen 582
Britacil 633
Broad spectrum semi-synthetic antibiotic 637
Broad-spectrum antibiotics 569, 637
Broccoli 773, 757, 774
Broccoli sprouts 774
Bromelain 783, 784
Bromination 286
Bromine reaction 288
Bromine test 130
Brouadrel-boutmy test 488
Brown algae 716
Brucine 4, 389, 517
Brush border enzyme 782
Brussel sprouts 757
Bryostatins 831, 832
Bubble boy 78
Buckthorn 367, 761
Buckwheat leaves and flowers 161
Buckwheat plant 161
Buffers 578
Burseraceae 307
Butanedione 266
Butolic acid 321
Butyrate 267, 268
Butyric acid 325
Bicyclic terpenoid ketone 220
Bykomycin 585
- C**
- C-9 monoterpene iridoid glycosides 197
C-glycosides 124, 126, 130, 134, 135, 159
C-reactive protein 790
Cabermox 633
Cacao beans 2
Cacao seeds 2
Cadaverine 451, 461
Cade oil 33
Cadinene 216
Caechin 34
Caffeic acid 29, 341, 342, 751, 752
Caffeic acid ester 751
Caffeine 2, 3, 383, 384, 389, 396, 399
Caffeine-base 380
Caftazidime 599
Cajeput 297
Cajeputene 742
Cajeputol 297
Cajuput oil 221
Calabar bean 532
Calamus 294
Calamus oil 241
Calanolide A 833
Calciopen 631
Calcitonin 62
Calcitonin gene-related peptide (CGRP) 62
Calcium oxalate 133, 151
California bay 297
California laurel 297
California laurel 294
California poppy 486, 490
California sassafras 297
Callus 801
Callus culture 801, 803, 804
Callus masses 804
Callus tissue 804
Calonolide B 833
Calumba 547
Cambial tissue cultures 799
Camomile 297, 346
Camphane 259, 262, 263
Camphene 263, 220, 328
Camphor 1, 27, 219, 220, 221, 225, 261, 287, 294, 341

- Camphora 219
Camphorsulphonic acid 660
Camptothecin 811, 823
Cananga 294, 302, 304, 305, 356
Canary dextrin 100
Cancer-cell kinetics 823
Candicidin 664
Candicidin D 665
Candivon 664
Canex 559
Cannabichromene 317
Cannabidiol 23, 316, 317
Cannabigerol 317
Cannabinaceous alkaloids 397
Cannabinoid 316
Cannabinol 216, 317
Cannabis 316
Cannabol 317
Cannapis 316
Cannin 317
Cantaxin 37
Cantharathene 811
Cantharanthus 9
Cantharides 122, 170, 172, 173, 561, 566, 567
Cantharides beetles 173
Cantharides camphor 566
Canthinones 396
Canton rhubarb 343
Cantrex 583
Capaiba 323, 324
Capaivic acid 324
Cape aloe 127, 128, 131, 774
Cape periwinkle 505, 506, 507, 508, 512
Caper spurge 341, 343, 349
Caperomycin 679
Caphamycin C 598
Caprifoliaceae 168
Caprylic acid 332
Capsaicin 36, 323, 324, 376, 398, 405, 410, 411, 412, 768
Capsanthin 324
Capsicum 323, 411
Capsicum fruits 411
Carane 259, 261, 262, 264, 265
Caraway oil 241
Carbamoyl ecgonine 473
Carbapen 636
Carbecin 636
Carbenicillin 629, 635, 637
Carbenicillin disodium 636
Carbenoxolone 190, 191
Carbohydrate biogenesis 119
Carbohydrates 84, 85
Carbomycin 625
Carbopenem 612
Carboxy benzyl penicillin 636
Carcinogen 591
Cardelmycin 663
Cardiac glycosides 144, 152, 157, 204, 701
Cardio-khellin 553
Cardioactive glycosides 811
Cardioquin 528
Cardiotonic 124, 705, 815
Cardiotonic activities 702
Cardiotonic compounds 701
Cardiotonic effect 704
Cardiotonics 701
Cardiotoxin-II 701
Cardiovascular 696
Cardiovascular active drugs 696, 701
Cardiovascular activities 701
Cardiovascular drugs 828
Caribbean sponge 698, 706, 707
Carline thistle 297
Carnitine 765, 766, 767
Carotaben 740
Carotenes 13, 240, 739
Carotenoid terpenoids 738
Carotenoids 31, 218, 238, 239, 738, 739, 740, 807
Carpidine 440
Carpiline 440
Carrageenan 116
Carrier displacement method 645
Carrier protein 790
Carrot 28
Carrot explants 799
Cartap 725
Carthamin 165, 166, 809
Cartric 513
Carum capticum 225
Carvacrol 227, 287, 288
Carveol 257
Carvestrene 258
Carvone 27, 257, 284, 285, 286
Caryophyllene 325, 334
Casanthranol 135
Cascara bark 127, 761, 762
Cascara buckthorn 367
Cascara sagrada 134, 367, 133, 135
Cascarilla 359
Cascaroside 126, 762
Cascarosides (A, B, C and D) 126, 135, 762
Casein 793, 794
Casein powder 793
Casein variants 793
Caseinogen 790
Caspase-8 activation 757
Cassava starch 91
Cassia oil 344
Cassie 304, 346, 356
Castanospermine 453, 455
Catechin 289, 290, 746, 747
Catechinic acid 746
Catechol 34, 289, 290, 354, 355, 356, 746
Catechol methyl ether 282
Catecholamines 481
Catechu 34
Catechuic acid 746
Catergen 746
Catharanthine N-oxide 520
Catharanthine 504, 505, 508, 509, 510, 519, 520
Cathartics 127
Catheranthus alkaloids 810
Cathine 409, 410
Cathinone 409
Cathocin 663
Cathomycin 663
Cat's hair 39
Caulosapogenin 187
Cayenne 36, 768
Ceara jaborandi 437
Cedar wood Oil 26
Cedol 603
Cefa-drops 602
Cefadroxil 598, 602, 603
Cefalexin [cephalexin] 598
Cefalotin 600, 601
Cefam 603
Cefamandole 603, 605
Cefamandole nafate 603
Cefamedin 602
Cefamezin 602
Cefamox 602

- Cefatoxime 606
Cefazil 602
Cefazina 602
Cefazolin 600, 607
Ceffoprim 604
Cefiran 603
Cefixime 599
Cefmonoxime 600
Cefodox 609
Cefonicid 605
Cefonicid disodium 605
Ceforal 602
Ceforanide 600
Cefotaxime 600, 605, 606, 608
Cefoxitin 599, 600, 610
Cefpodoxime 599, 609
Cefpodoxime proxetil 598, 600, 609
Cefracycline 650
Cefradine [cephradine] 598
Ceftazidime 600, 607, 608
Ceftazidone 614
Ceftizoxime 600
Ceftriaxone 599, 600, 606
Ceftriaxone disodium hemiheptahydrate 607
Ceftriaxone sodium 607
Cefumax 604
Cefurex 604
Cefurin 604
Cefuroxime 600, 604
Cefuroxime axetil 600
Cefuroxime sodium 604
Cefuroxime-axetil 598, 599, 609
Celery 159, 347, 350
Cell growth 802
Cell suspension 804
Cell suspension cultures 798
Cell-culture techniques 823
Cell-cycle arrest 757
Cell-mediated immunity (CMI) 742
Cell-surface-receptors 46
Cellulase 782
Cellulose 21, 86, 97, 98
Cellulose to cellobiose 782
Cemado 603
Cemandil 603
Cemastin 601
Cembranes 696, 697
Cembranoids 699
Cementum 792
Cenomycin 610
CentoRX 80
Centraureo 651
Cephaeline 391
Cephalosporin 596, 597, 605
Cephalosporin antibiotics 598
Cephalosporin C 596, 597, 600
Cephalosporin P 596
Cephalosporinase 600
Cephalosporins 16, 579, 576, 600, 603, 609, 611, 626, 638
Cephalothin 600
Cephalothin sodium 601
Cephamycin C 600, 610
Cephamycins 598, 599, 600, 610
Cephapirin 600
Cephatin 601
Cephazolin 601
Cephazolin sodium 602
Cephem 611
Cephos 602
Ceporacin 601
Cepovenin 601
Cerelac 91
Cerezyme(R) 79
Certified dye 643
Certified organic mushroom nutraceutical 777
Certomycin 586
Cerubidin 690
Cerubidine 6, 691
Cetacidal 602
Cevadiline 421
Cevadine 32, 420, 421, 422, 423, 424
Cevalin 37
Ceveratrum alkaloids 420, 425
Cevine 421
Ceylon cinnamon 297, 302, 344
CEZ 601
Chalcone 164, 205, 206, 366
Chalcone glycosides 159, 164
Chalone glycosides 212
Chalones 158, 164, 165
Chamomile 232
Chamomile oil 241
Chancroid 588
Chandra 506, 511, 512, 513
Charantin 830, 831
Chelidonine 377
Chelidonine acid 377
Chellah 553
Chemical additives 574
Chemical characteristics of volatile oils 250
Chemical composition of resins 309
Chemical composition of yeast 795
Chemical properties of pesins 308
Chemical-engineered-flow process 576
Chemiofuran 677
Chemistry of resins 309
Chemotherapy 47, 746
Chemotypes 733
Chenopodiaceae 299
Chenopodium ambrosioides 299, 736
Chenopodium oil 299
Chibroxin(e) 673
Chibroxol 673
Chicory plant 348
Chili 323, 348, 415, 418
Chimono 673
China oil 335
Chinaberry tree 238
Chinese anise 290, 292, 294, 297, 356
Chinese pea 776
Chinese rhubarb 343
Chinese sacred lilly 356
Chinidin 528
Chiorella 120
Chiral centres 124
Chirata 196, 199
Chiratin 199
Chitin 22, 117, 118, 763
Chitin sulphate 118
Chitosan 118
Chlorambon 662
Chlorambucil 574
Chloramphenicol 38, 659, 660, 661, 663
Chloramphenicol monosuccinate sodium salt 662
Chloramphenicol palmitate 662
Chloramphenicol succinate/palmitate 578
Chlorination 650
Chlorogenic acid 29, 342, 751, 752
Chlorohydrin 669
Chlorolifarina 662

- Chloromycetin 16, 38
 Chloropal 662
 Chlorophyll 392, 738, 768
 Chlorophyll a 768, 769
 Chlorophyll b 768, 769
 Chlorophyll c 769
 Chloroquine 12, 24
 Chlortetracycline 577, 649, 651, 657, 659
 Cholesterol 419, 760, 761
 Choline 372
 Chondoprotein 790
 Choriocarcinoma 692
 Chorionic vili sampling and amniocentesis 69
 Chroman 157
 Chromone 553
 Chromone bitter principles 547, 553
 Chromosantonin 551
 Chromosomal genes 17
 Chromosomes 44
 Chrysaloin 134, 762
 Chrysotropic acid 348
 Chrysoeriodictyol 315
 Chrysoeriol (G) 206
 Chrysomykine 651
 Chrysophanic acid 136, 761
 Chrysophanol 132, 761
 CHX-3311 698
 Chymopapain 786
 Chymosin 787
 Chymotrypsin 783, 784, 788
 CI 75300 753
 CI natural yellow 753
 Ciblor 639
 Cichorin 169
 Cichorlin 348
 Ciclobiotic 653
 Ciclolysol 656
 Cicutine 401, 442
 Cidomycin 582
 Ciflox 672
 Ciguatera toxins 709, 714
 Ciguatoxin 714, 715
 Cilastatin sodium 614
 Cilleral 634
 Ciloxan 672
 Cimetidine 675
 Cin-Quin 527
 Cinchol 760
 Cinchona 397, 521, 547
 Cinchona alkaloids 387, 429, 431, 530
 Cinchona bark 389, 391, 429, 521
 Cinchonamine 531
 Cinchonidine 429, 430, 431, 432, 433, 521, 522, 524, 529, 530, 531
 Cinchonidine dihydrochloride 529
 Cinchonidine hydrochloride dihydrate 529
 Cinchonidine sulphate trihydrate 529
 Cinchonidine tartrate 523
 Cinchonidinone 434
 Cinchonine 4, 429, 430, 431, 433, 521, 522, 524, 526, 530, 531
 Cinchonine dihydrochloride 433, 526
 Cinchonine hydrochloride dihydrate 434, 526
 Cinchonine sulphate dihydrate 434, 526
 Cinchoninone 525, 531
 Cinchonione 530
 Cinchovatine 529
 Cinene 26, 742
 Cineol 15, 332
 Cineole 221, 291, 297, 298, 300
 Cinnaloid 513
 Cinnamaldehyde 251, 280, 344, 345
 Cinnamein 336, 337
 Cinnamic acid 204, 206, 210, 369, 447, 753
 Cinnamic ester 334
 Cinnamon 814
 Cinnamon bark 195, 814
 Cinnamon oil 33, 241, 243,
 Cinnamonum camphora 219
 Cinnamoyl cocaine 467, 464
 Cinnamoyl ecgonine (methylecgonine) and hyoscyamin 471
 Cinnamoyl-methylecgonine 467
 Cinnamoylcocaine 467
 Cinnamoylcocaine [(E)-form] 467
 Cinnamyl cinnamate 302, 303, 338
 Cinoxacin 670
 Cinocin 671
 Ciprinol 672
 Cipro 672
 Ciprobay 672
 Ciprofloxacin 671, 672, 673, 674, 676
 Ciprofloxacin monohydrochloride monohydrate 672
 Ciprofloxacin 673
 Ciproxan 672
 Ciproxin 672
 Circolene 505
 Cis-1, 4-terpin 300
 Cis-anethole 292
 Cis-b-ocimene 253
 Cis-isomer 256
 Cis-o-coumarinic acid lactone 346
 Cis-terpin hydrate 268
 Citral 27, 270, 291, 272, 273
 Citral-a 272
 Citral-b 272
 Citral-enol-acetate 273
 Citron 163
 Citronella oil 241, 274
 Citronellal 271, 272, 274, 275, 276
 Citronellol 276
 Citronin 163
 Citrous juices 735
 Citrus flavonoids 750
 Citrus fruits 37
 Cladinose 619
 Clamoxyl 633
 Clarithromycin 619, 621
 Classification and spectrum 626
 Classification of glycosides 127
 Classification of resins 313
 Classification of volatile oils 251
 Claviceps 534, 535
 Clavine alkaloids 535
 Clavos 344
 Clavulanate amoxicillin 630
 Clavulanate amoxicillin trihydrate 639
 Clavulanate potassium 639
 Clavulanate ticarcillin disodium 639
 Clavulanate-ticarcillin 630
 Clavulanic acid 611, 638
 Cleocin 617
 Cliaacil 631
 Clindamycin 614, 616, 617, 618
 Clindamycin hydrochloride monohydrate 618
 Clindamycin phosphate 618
 Clinimycin 652
 Clinimycin (rescinded) 617
 Cloned CGRP 62

- Cloned DNA 52
Cloned DNA segments 52
Cloning 56
Cloning cDNAs 53, 54
Cloning eukaryotic DNA 53
Cloning eukaryotic DNAs phase genomes 53
Cloning of genes 79
Cloning of mice 81
Cloning plasmid in bacterial cells 51
Cloning process 50
Cloning vector 48, 53
Clont 669
Clortetrin 652
Closina 641
Clotrimazole 688, 689
Clove 287, 350
Clove oil 33, 241, 281, 283, 287
Clove pepper 297
Clovers 344
Cloxacillin 628, 631, 632
Cloxacillin sodium monohydrate 632
Cloxapen 632
Cloxyphen 632
CMT 603
Co-amoxiclav 639
Coagulating enzyme 782
Coal tar 322
Coaxin 601
Coca 304, 397
Coca kola 2
Cocaine 12, 374, 381, 389, 391, 398, 464, 466, 468, 471, 473
Cocaine hydrochloride 467
Cocaine nitrate dihydrate 467
Cocaine permanganate 467
Cocaine sulphate 467
Cocalne 473
Cocculin 561
Coclanoline 493
Codeine 23, 381, 388, 389, 486, 490, 494, 497, 814
Codeine acetate 491
Codeine hydrobromide 491
Codeine hydrochloride 491
Codeine methyl bromide 491
Codeine phosphate 491
Codeine salicylate 491
Codeine sulphate 491
Codeinone 488, 495, 497
Codicept 490
Coenzyme A-dependent pathways 766
Coenzymes 780
Coenzymes Q 772
Coffearin 37
Coffee bean 2, 29
Coffee seed 2
Coflizoxime 608
Cogmine 532
Cohoba 355
Cohobation 284
Cohobation of distillation waters 247
Colchicine 373, 375, 380, 381, 382, 389, 400, 405
Colchicum 375
Cold Water sponge 725
Colforsin 233, 829
Colimycin 644
Colisticina 644
Colistimethate sodium 645
Colistin 641, 644
Colistin A 642, 645
Colistin B 642
Colistin formaldehyde-sodium eisolphite 645
Colistin sodium methanesulphonate 645
Collagen 784, 790, 792, 793
Collagen variants 793
Collagenase 783, 784
Colloidal glycosides 743
Colocynth 341, 342
Colophony 39, 283, 310, 314, 315, 324
Colophony resin 314
Colour test 408, 523
Colouring matters 21
Coltericin 583
Coltsfoot 358, 367, 474
Coly-mycin 644
Commercial penicillins 626
Commiphora 332
Commiphoric acid 310
Common foxglove 341, 342
Common lavender 302
Common licorice 367
Common milkweed 456
Common precursor 574
Common thyme 302
Comomile 349
Comphor oil 294
Complement factor C-3 791
Complement protein 791
Complete protein 789, 790
Complex chemical constituents 5
Complex polysaccharides 90
Complex tricyclic aglycone 646
Compocillin VK 631
Components 588
Components of streptomycin 588
Compound 83405 603
Compound tincture of benzoin 131
Compylobacter jejuni 672
COMT enzyme 750
Concanavalin 794
Concanavalin A 794
Condylactis gigantea 705
Condyline 30
Conessine 389
Conhydrine 401, 403, 404
Conicine 401, 442
Coniferae 260
Coniferin 144
Coniferous resins 313
Coniferyl alcohol 365
Coniferyl benzoate 338
Coniine 401, 402, 404, 442, 443
Coniine hydrobromide derivative 443
Coniine hydrochloride derivative 443
Conjugated protein 790
Connesine 400
Connine 398
Conoderm 689
Conofite 689
Conotoxins 721, 725
Conquinine 527
Constituents of resin 314
Contact poison 457
Contemporary nutraceuticals 772
Convention (CBD) 817
Convolvulaceae 201
Copaiba 310, 324
Copaivic acid 310
Copies of genes 575
Coriander 301
Corilagin 368
Corkwood tree 397, 456, 457, 469

- Cornanincol 531
 Cornocentin 537
 Coronary heart disease (CHD) 777
 Coronin 553
 Corticosteroids 183
 Corticosterone 14
 Corvone 287
 Coryanthe 505
 Corynantheal 525, 530
 Corydaline 398
 Corynantheal 434
 Corynine 511
 Cosmegem 692
 Cosmopen 627
 Cotton 86
 Cotton seed 161
 Coughwort 358, 367, 474
 Coumaric acid 345, 346
 Coumarin 168, 169, 205, 345, 346, 553
 Coumarin and furanocoumarin glycosides 127, 168, 212
 Coumarin eitter principles 547, 556
 Coumarin derivatives 190
 Coumarin glycosides 168, 169, 212
 Coumarinic acid 345, 346
 Coumarinic anhydride 346
 Coumarins 168, 340, 345, 833
 Coumarone 553, 559
 Coumarone bitter principles 547, 559
 Coumestrol 34
 Coupling of a flavonoid 365
 Couramin 168
 Cowgrass 341, 342
 Cox-1 enzyme 754
 Cox-2 enzymes 753
 CP-45899 639
 CP-45899-2 640
 Cranberry juice 751
 Cranebill 357
 Crassin acetate 697
 Cresol with soap 179
 Critical fungal protein synthesis 817
 Crocetin 809
 Crocin 277
 Cromoglycate 814
 Crotalaria 471
 Cruciferae 180, 236
 Crude alkaloidal extract 393
 Crude antibiotic 576
 Crude natural plant extracts 814
 Crustacean shells 117
 Cryophyllene 324
 Cryptotethia crytpa 706
 Cryptoxanthin 741
 Crystal gum 105
 Crystalomicina 583
 Crystamin 677
 Crystapen 627
 CTX 715
 Cube roots 559
 Cucurbitacin-B 236, 237
 Cultivated carrot 295
 Cultivation technique (CT) 798
 Culture cells 806, 798
 Culture vessels 804
 Cumarin 346
 Cumaru 347
 Cuminaldehyde 280, 333
 Cupraloin test 131
 Cupreine 433, 524
 Cupreol 760
 Cuprocyanate test 176
 Curacao aloe 127, 130, 774
 Curaeao aloes 128
 Curare 8
 Curcuma oil 332
 Curcumin 332, 753, 754, 809, 829
 Curcuminoids 332
 Curling factor 681
 Curocef 604
 Curoxim 604
 Cuscohygrine 462, 463, 471, 473
 Cuspareine 398
 Cutin 21
 Cuxacillin 633
 Cyanide and methoxide nucleophiles: 1 731
 Cyanidenolon 750
 Cyanidin 167, 745
 Cyanidin choride 35
 Cyanidol 34, 746
 Cyanoacetic acid 273
 Cyanogenesis 207
 Cyanogenetic glycoside 173, 174, 176, 177, 178, 179, 207, 209, 213, 340
 Cyanogenetic precursors 207
 Cyanohydrins 207
 Cyantin 677
 Cyclamycin 624
 Cyclic terpene aldehyde 276
 Cyclic terpene ketones 284
 Cyclodextrins 86, 95, 96
 Cycloheximide 570
 Cyclomycin 650
 Cyclopar 650
 Cyclopentane 217
 Cyclophosphamide 574
 Cycloserine 641, 678
 Cyclosporine 675, 690
 Cymarin 154, 155
 Cymarose 84, 155
 Cynoglossum 471
 Cyonogenetic glycosides 127
 Cysteine sulphoxides 759
 Cystit 677
 Cystogen 677
 Cystospaz 469
 Cytarabine 698
 Cytarabine (Ara-C) 697
 Cytarbel 698
 Cytochalsins 722
 Cytochrome P-450 pathways 690
 Cytochrome P450-mediated pathways 675
 Cytochrome system 772
 Cytochrome transport system 780
 Cytokines 63, 742, 802
 Cytokinins 807
 Cytosar 698
 Cytosine 44, 48, 201, 686
 Cytotoxic/antineoplastic agents 696
- D**
- D(-)-Ephedrine 378, 407
 D(-)-Ψ-Ephedrine 407
 D(+) ephedrine 378
 D-α-phellandene 332
 D-Camphor sulphonate 225
 D-Camphor 28
 D-Carene 262
 D-Carotene 740
 D-Carvone 27, 286
 D-Catechin 133
 D-Cefactor 600
 D-Cefadroxil 600

- D-Cefalexin 600
 D-Cefamandole 600
 D-Cefonicid 600
 D-Cefoperazone 600
 D-Cefradine 600
 D-Cephalexin 600
 D-Cephradine 600
 D-D-Gluconolactone 757, 758
 D-Fenchone 264, 287
 D-Form 261, 262, 277
 D-Glucosamine 764
 D-Glucose 140, 141, 142
 D-Limonene 257, 742, 743
 D-Lupanine 450
 D-Mycinose 618
 D-Prunasin 178
 D-Pseudoephedrine 408, 409
 D-Pseudo-ephedrine hydrochloride 408
 D-Sabinene 332
 D-Tryptophan 525
 D-Tubocurarine 379
 D-Xylose 141
 D.H.E. 45 539
 D1-Piperidine 451
 D3-Carene 258
 D4-Carene 258
 D6 and W2 clones 728
 D8-tetrahydrocannabinol 317
 Dab-residues 642
 Dactinomycin 16, 690, 692
 Daidzein 561, 747, 748
 Dairy products 736
 Daktarin 689
 Dalacin 618
 Dalacin C 617
 Dammar 310
 Daphentin 349
 Daphnin 169, 349
 Datisceae 236
 Datura 377
 Daturine 469
 Daughter cells 51
 Dauno-rubicinol 691
 Daunoblastina 691
 Daunomycin 593, 690
 Daunomycinone 690, 691
 Daunorubicin 16, 6, 593, 594, 690, 691
 Daunorubicin hydrochloride 691
 Daunorubicinol 691
 Daunosamine 591, 592, 690, 691
 Deacetylated chitin 118
 Deadly nightshade 348, 397, 462
 Deamidizing enzyme 782
 Deamination reactions 780
 Debranching enzyme 782
 Decarboxylating enzyme 782
 Declid 684
 Declomycin 652
 Deepoxysarcophine 727
 Deertongue 346
 Definite leads 819
 Deflamon 669
 Defleurage 245
 Deganol 652
 Dehydroabietic acid 311
 Dehydrogeissoschizine 509, 510
 Dehydrogenation 217
 Dehydrogessoschizine 518
 Dehydrogeseofulvin 682
 Dehydropodophyllotoxin 319
 Dehydrorotenone 560
 Delacillin 633
 Delinting 98
 Delivery of tissue cultured plants 800
 Delphinidin 167, 745
 Demeclocycline 649, 650, 652, 654
 Demeclocycline hydrochloride 652
 Demeclocycline sesquihydrate 653
 Demethoxy vindoline 508
 Demethylchlortetracycline 652
 Demethylmunduserone 561
 Demethylsuberosin 558
 Demetraciclina 652
 Demulcent 124
 Denatured protein 790
 Dendalone-3-hydroxybutyrate 720
 Dentin 792
 Dentojel 524
 Deoxybarbaloin (chrysaloin) 134
 Deoxyribonuclease 783, 785
 Depasan 451
 Dephentin 346
 Depipeptides 697
 Deralbine 689
 Dergotamine 539
 Derivatives of ergonovine 537
 Derivatives/salts of quinine 524
 Derived protein 791
 Dermonistat 689
 Derris roots 559
 Des-thia-carbapenam nucleus antibiotics 612
 Desacetyl vinblastine 68
 Deschlorobiomycin 650
 Deserpidine 512, 519
 Desired secondary constituents 16
 Desmethoxy yangonin 322
 Desosamine 619
 DET MS 539
 Detravis 652
 Devil's dung 329
 Devil's shoe string 559
 Devil's tobacco 443
 Dexcyanidanol 746
 Dextran 40, 70, 75, 86, 92, 94, 95
 Dextran-sucrases 95
 Dextrin 86, 99
 Dextro methorphan 381
 Dextrose 792
 Dhurrin 207, 208, 209
 Diacetyl 284
 Diacetyl derivative 730
 Diacetyljervine 426
 Diagnostic agent 94
 Diagnostic products 61
 Dianthrones 762
 Diastase 795
 Diazocin alkaloids 396
 Dicloxacillin 628
 Dicot plants 183
 Dicotyledons 375
 Dicoumarin 33
 Dicoumarol 33
 Diergo 539
 Dietary supplement 736, 737
 Diferuloylmethane 753
 Different modes of distillation 243
 Differentiated tissues 812
 Diflucan 690
 Digacin 829
 Digenic acid 722
 Digest 47
 Digestive enzymes 780, 782
 Digilanid A 148
 Digilanid B 148
 Digilanid C 148
 Diginin 146

- Digitalin 146
 Digitalis 145, 146, 147, 148, 342
 Digitalis glycosides 705, 810
 Digitalis lanata 148
 Digitalis leaf 814
 Digitalis lutea 149
 Digitalis thapsi 149
 Digitalose 85
 Digitonin 146
 Digitosaponin 146
 Digitoxin 32, 122, 204
 Digitoxose 84
 Digoxin 19, 829, 675, 815
 Dihalides 258
 Dihydergot 539
 Dihydric phenols 288, 289
 Dihydro-p-cyrenes 255
 Dihydrochalone 164
 Dihydrodisulphonic derivative 275
 Dihydroergotamine mesylate 539
 Dihydrokawain 322
 Dihydromethysticin 322
 Dihydromyrcene 252, 253
 Dihyronicotinic acid 459, 460
 Dihydropyridine 726
 Dihydroveratramine derivative 425
 Dihydrochalcones 164
 Dil 295
 Dill oil 28, 29, 295, 296
 Dill seed 295
 Dimeric alkaloids 543
 Dimethylpodoophyllotoxin 319
 Dimethylpyruvic acid 535
 Dimorphic fungi 680
 Dimyrcene 334
 Dinoflagellate species 711
 Dinoflagellates 711, 715
 Diocimex 654
 Diocyclin 650
 Dioscin 184
 Dioscorea 183, 184
 Diosgenin 13, 184, 210, 810
 Diosmin 159, 160
 Dipentene 257, 270
 Dipentene dihydrochloride 257
 Dipterocarpaceae 307
 Direct ELISA 70
 Direct steam distillation 242, 243
 Direktan 459
 Dirgotarl 539
 Disaccharide (+) – maltose 90
 Disaccharides 167
 Disaturase enzyme 773
 Dish-cloth gourd 236
 Disomin 161
 Disruption of DNA-metabolism 658
 Distakaps V-K 631
 Distaquaine V 631
 Distillation water 271
 Distribution of resins in plants 306
 Diterpene 251, 268
 Diterpene alkaloids 396
 Diterpenoid alkaloids 414
 Diterpenoid structural analogues 824
 Diterpenoids 215, 218, 233
 Diuretic steroid 183
 dl-Limonene 257
 dl-Camphor 28
 dl-Carvone 27, 286
 dl-Catechin 290
 dl-Ecgonine trihydrate 469
 dl-Ephedrine 478
 dl-Ephedrine hydrochloride 478
 dl-Ephedrine sulphate 478
 dl-Form lupinine 450
 dl-HCl 408
 dl-Hydrochloride 410
 dl-Hyoscyamine 465
 dl-Menthone 285
 dl-Sulphate 408
 dl-Tropyl tropate 465
 DNA adducts 751
 DNA fragment 51
 DNA Polymerase 780
 DNA probe 68
 DNA probes and RFLP analysis 68
 DNA-cloning 50
 DNA-ligase 48
 Dog senna 139
 Dogwood 367
 Doktacillin 629, 633
 Dolcontin 486
 Domicillin 629, 633
 DOMOI 723
 Domoic acid 721, 723
 Domoic acid dihydrate 723
 Dopamine 482
 Dorema ammoniacum 330
 Doridosine 707
 Dowpen VK 631
 Doxatet 654
 Doxorubicin 6, 590, 591, 592, 593, 596, 690
 Doxorubicin hydrochloride 591
 Doxychel hyclate 654
 Doxycycline 649, 650, 654
 Doxycycline hyclate 654
 Doxycycline hydrochloride hemie-
 thanolate hemihydra 654
 Dracoese 312
 Dragendorff's reagent 716
 Dragon's blood 356
 Dried yeast 795
 Dronabinol 23
 Drug biosynthesis 24
 Drug discovery 821
 Drug molecules 695
 Drug targetting 68
 Drug targetting and tumour imaging
 68
 Drug-development process 817
 Drug-plant-cultivation 16
 Drying oil 179
 Duboisine 469
 Dufalone 33
 Dumocyclin 650
 Dumopen 633
 Duphacycline 652
 Duracef 602
 Duradoxal 654
 Duragentum 582
 Duricef 602
 Duriles 528
 Duromorph 486
 Duxima 604
 Dyer's broom 450

 e, γ -carotene 740
 e-rhodomyacinone 593
 Ecgonine 468
 Ecgonine cinnamate methyl ester
 467
 Ecgonine hydrochloride 469
 Ecgonine methyl ester 467
 Ecgonine methyl ester benzoate
 466
 Ecgonine monohydrate 468
 Econazole 688, 689

- Ecuelle a piquer 243, 244
 Edible burdock 341
 Effect of enzyme induction 574
 Effective dose (ED50) 814
 Efpenix 633
 Egotamine 535
 Egyptian privet 357
 Ehrlich reagent 386
 Either 306
 Ekkert's colour test 439
 Ekomine 465
 Elaeagnine 501, 503
 Elastin 793
 Elazor 690
 Eldoisin 705
 Eldrin 749
 Elementary organisms 799
 ELISA 69
 ELISA methodologies 70
 Ellagic acid 751, 752
 Elongation 825
 Elongation step 825
 Elyzol 669
 Elzogram 602
 Emetine 374, 382, 389, 398
 Emodin 29, 129, 132, 135, 203, 761
 Emodin anthranol 129
 Emodin anthrone 129
 Emodin dianthrone 132
 Emodin oxanthrone 129
 Emodine anthrone 132
 Emperical antibiotic therapy 608
 Empiric observations 573
 Empyreumatic products 308
 Emulsin 124, 140, 142, 175, 176
 Endocarditis 577, 588
 Endoenzymes 781, 782
 Endogenous fibrinolytic system 788
 Endogenous glucosamine 764
 Endomixin 585
 Endophleban 539
 Endothelial cells 745
 Endotoxins 709
 Enflourage method 245
 Engemycin 652
 Engerix-B(R) 79
 Engineered T-cells 78
 English camomile 297, 346, 349
 Enriched citrus fruits 736
 Ent-8-hydroxymanzamine 728
 Ent-8-hydroxymanzamine A[16] 729
 Ent-kaurene 414
 Enterfram 585
 Enterokanacin 583
 Enzymatic hydrolysis 242, 246
 Enzyme 70
 Enzyme as drug substances 779
 Enzyme immunoassays (EAI) 67
 Enzyme inhibitors 696
 Enzyme linked immunosorbant assay (ELISA) 69
 Enzyme variants 782
 Enzyme-substrate complex 779
 Enzymes for synthesis reactions 780
 Enzymes-cascade technique 67
 EPEG 362
 Ephedra 397
 Ephedral 478
 Ephedrine 372, 373, 381, 389, 396, 397, 398, 405, 406, 407, 408, 409, 410, 477, 478
 Ephedrine (Adrenomimetic) 11
 Ephedrine hydrochloride 387
 Ephedrine 406
 Ephedrine oxalate 477
 Ephetonin 478
 Epicatechin 133, 746
 Epicinchonidine 530
 Epicinchonine 526
 Epidemiology 737
 Epidermal growth factor 80
 Epinephrine 481, 759
 Epirenor 703
 Epirubicin 590, 592
 Epirubicin hydrochloride 592
 Epogen(R) 42, 79
 Epoxidations 714
 Erdmann's reagent 386, 444
 Eremfat 658
 Eremursine 478
 Ergate 539
 Ergine 536, 540
 Ergine methane sulphonate 540
 Ergobasine 536
 Ergobine 542
 Ergobutine 542
 Ergobutyryne 542
 Ergocornine 535, 542
 Ergocorninine 535
 Ergocristine 535, 542
 Ergocristinine 535
 Ergocryptine 535
 Ergocryptinine 535
 Ergoklinine 536
 Ergolines 535
 Ergomar 539
 Ergometrine 396, 399, 536
 Ergometrine tartrate hydrate 537, 538
 Ergomimet 539
 Ergonine 542
 Ergonovine 397, 536, 537, 538, 539
 Ergonovine maleate 537, 538
 Ergonovine tartrate hydrate 538
 Ergont 539
 Ergoptine 542
 Ergosine 535, 542
 Ergosinine 535
 Ergostat 539
 Ergosterol 33, 686, 680, 808
 Ergostetrine 536
 Ergostine 542
 Ergot 374, 534, 535, 541
 Ergot alkaloids 23, 386, 495, 534, 535, 537, 538, 541
 Ergot oil 538
 Ergot peptide alkaloids 535
 Ergotamine 10, 19, 23, 216, 374, 381, 389, 399, 535, 536, 538, 539, 540, 541, 542
 Ergotamine group 542
 Ergotamine hydrochloride 539
 Ergotamine tartrate 539
 Ergotaminine 535
 Ergotartrat 539
 Ergotocine 536
 Ergotonin 539
 Ergotrate 536
 Ergotrate maleate 537
 Ergots 535
 Ergovaline 542
 Ergoxine group 542
 Ericaceae 127
 Ericolin 315
 Eriodictin 315
 Eriodictyon 315
 Eriodictyonic acid 315
 Eriodonol 315
 Ermetrine 537
 Erotoxine group 542

- Erpalfa 698
 Erytaurin 565
 Erythro 478
 Erythrocentaurin 565
 Erythromycin 16, 619, 620, 621, 622, 625
 Erythromycin A 621, 622
 Erythromycins B and C 619
 Erythroquinine test 433
 Erythrose-4-P., 208
 Escin 828
 Esculetin 346, 348
 Esculin 348
 Eserine 532
 Eseroline 533
 Eskacillin 627
 Eskel 553
 Essential (or volatile) oils 25, 39, 240, 810
 Essential amino acids 789
 Ester alkaloids of veratrum 420
 Ester resins 313
 Ester volatile oils 251, 301
 Esterification provides better leaving group 496
 Esters 30
 Esters containing nitrogen 301
 Esters of aliphatic acids 301
 Esters of aromatic acids 301, 302
 Estimation of ascaridole 300
 ESTM 80
 Estradiol 12
 Ethambutol 678
 Ethereal oils 240
 Ethimide 680
 Ethionamide 679, 680
 Ethnobotanical 819
 Ethyl cinnamate 334
 Ethylene-diamine tetra-acetic acid (EDTA) 802
 Ethylnarceine hydrochloride 481
 Ethylphenyl acetate 361
 Ethylvanillin 335
 Etoposide 362, 363
 Eucalyptol 23, 219, 221, 222, 297, 742
 Eucalyptus oil 225, 264, 297
 Eucodin 491
 Eudesmanolide 230
 Eugenol 15, 22, 27, 28, 233, 246, 251, 281, 282, 288, 289, 344
 Eukaryote DNA 75
 Eukaryotic DNA segment 53
 Eukaryotic fragment 53
 Eumydrin 465
 Eunicin 718
 Euphane triterpenes 821
 Euphorbia 39
 Euphorbiaceae 127, 236
 European angelica 352, 353, 556
 European goldenrod 342
 European mistletoe 342, 343
 European squill 150, 153
 Evaporation residue 250
 Eviodictyol 315
 Evramycin 624
 Exal 514
 Examples of convergence 544
 Examples of divergence 544
 Excin 674
 Exmigra 539
 Exocyclic lactone 697
 Exoderil 685
 Exoenzymes 781
 Exogonium purga 201
 Exons 53
 Explant 803
 Exposure to air and light 241
 Expression 242, 243
 Expression cloning 56
 Extended-spectrum 629
 Extended-spectrum (antipseudomonas) penicillins 627, 635, 638
 Extentabs 527
 External metabolites (ectocrines) 709
 Extracellular enzymes 781
 Extraction 242, 244
 Extraction and isolation of alkaloids 389
 Extraction of alkaloids 390
 Extraction of volatile oil 244, 245
 Extraction with acid solution 393
 Exudate gums 101
 Eyebright 342
- F**
- Fabianol 581
 Factor VIII 62
 Faculty genes 76
 Fado 603
 False acacia 305
 Farex 91
 Farmiserina 641
 Farmorubicin 592
 Farmoxin 610
 Fasigin 670
 Fasigyn 670
 Fat 133
 Fat-soluble vitamins 738
 Fehlings test 354
 Felonwood 397, 416
 Fenchane 259, 264, 265
 Fennel fruits finocchio 292
 Fennel oil 264
 Fenospin 631
 Fenoxypen 631
 Fermenting enzyme 783
 Fermergin 539
 Ferric chloride test 489
 Ferriferrocyanide test 176
 Ferrocyanide test for quinidine 528
 Ferulic acid 30, 329, 341, 342, 343, 347, 411, 412
 Festamoxin 608
 Fibrin clots 788
 Fibrinogen 788, 790
 Fibrinolysin 783, 785
 Fibrinolytic system 785
 Field horsetail 367
 Figs 350
 Fijian sponge 721
 Filamentous dermatophytes 680
 Filamentous fungi 681
 Filicic acid 325, 326
 Filicin 326
 Filicinic acid 325
 Firmacef 602
 First aminopenicillin antibiotic 634
 First generation 597
 First generation cephalosporins 600, 601
 First-choice-chemotherapy 691
 First-line agents 642
 First-line drugs 678
 Fish-oils 776
 Fixed oil 241
 Flacanone glycosides 212
 Flagroot 294
 Flagyl 669
 Flaran 290

- Flash point 250
 Flavan 34
 Flavan based constituents 34
 Flavanoids 744
 Flavanone glycosides 159, 163
 Flavanones 157, 158, 164, 165, 205, 206, 366
 Flavaspidic acid 325, 326
 Flavone 157, 158, 163, 205, 744,
 Flavone glycosides 159, 212, 234
 Flavonoid glycosides 157, 159, 190, 205, 212, 364
 Flavonoid biosynthesis 205, 364
 Flavonoid polyphenolics 744
 Flavonoid radical 744
 Flavonoid ring system 364
 Flavonoids 151, 157, 319, 364, 365, 366, 747, 753, 807
 Flavonol glycosides 159, 161, 212
 Flavonolignans 365
 Flavonols 157, 158, 205
 Flemoxin 633
 Flexibilide 720
 Flobacin 674
 Flociprin 672
 Floral concretes 245
 Florentine flasks 242
 Florid 689
 Floripondio 397
 Florisil 732
 Flower volatile oil 245
 Floxacin 673
 Floxil 674
 Floxin 674
 Fluavil 312
 Fluconazole 681, 688, 690
 Flucytosine 686, 687
 Fludarabine 699
 Fludarabine and aplystatin 697
 Fluorescence immunoassays (FIA) 67
 Fluorescence test 432
 Fluorinated quinolone antibacterial 674, 676
 Fluorinated quinolone antibiotic 671
 Fluorocoumarin 171
 Fluorocytosine 686
 Fluoroquinolones 671, 675, 676
 Fluoroquinones 671
 Flustramine A and B 720
 Foetal DNA 76
 Folic acid 667
 Folk medicines 5
 Food of the gods 329
 Foreign DNA 51
 Foreign genes 46
 Formaldehyde test 567
 Formate 267
 Formation of phthaleins 288
 Formin 677
 Formononetin 561
 Foromacidin A 625
 Foromacidin C 625
 Forosamine 619
 Forpen 627
 Forskohlii 233
 Forskolin 234, 812, 813, 828, 829
 Forsythia 161
 Fortaz 607
 Fortracin 645
 Fossypol 669
 Found 709
 Frachde's reagent 143
 Fractional distillation 298
 Frademicina 617
 Fragrance 241
 Framycetin 584, 585
 Framygen 585
 Frangula 127
 Frangula bark 136, 137
 Frangulin 29
 Frangulin A 136
 Frangulin B 136
 Fraquinol 585
 Fraxetin 346, 349, 350
 Fraxin 169, 350
 Free triterpenoids 236
 Free-cinnamic acid 334
 Free-radicals 744
 Freezing and thawing 114, 115
 French lavender oil 271
 French parsley 28
 Fresh tea-leaves 37
 Friar's balsam 131
 Froehde's test 156
 Froehd's reagent 386, 444
 Fructan 86
 FTIR 734
 Fuamed 677
 Fulgram 673
 Function of alkaloids in plants 377
 Function of secondary metabolites 806
 Functional food 82, 736, 737
 Functional food revolution 82
 Fungi 127
 Fungicidal 682
 Fungiderm 689
 Fungilin 683
 Fungizone 683
 Funtumine 400
 Furachel 677
 Furadantin 677
 Furadantine MC 677
 Furadoin 677
 Furalan 677
 Furan 553
 Furan ring 169
 Furan-based constituents 33
 Furanochembranoid diterpene 727
 Furanocoumarin 168, 169, 170, 205, 207, 345, 350
 Furanocoumarin glycosides 169
 Furanocourmarin 212
 Furantoina 677
 Furfural 33, 266, 283
 Furobactina 677
 Furophen T-caps 677
 Furoquinolines 396
 Fused b-lactam containing oxygen 638
- G**
- g-Coniceine 130, 401, 402, 403, 405
 g-Coniceine hydrochloride 403
 g-Cyclodextrin 96
 g-Diaminobutyric acid (L-Dab) 642
 g-Linolenic acid 773
 g-Protein 791
 g-Pyrone 553
 g-Strophanthidin 156
 g-Terpinene 260
 g-Trimethyl- β -hydroxybutyro-
 betaine 765
 GABA 716
 Galactoin 528
 Galactose 333, 772
 Galacturonic acid 107

- Galangin 161
Galbanum 310
Galbaresinotannol 312
Galcodine 491
Gallic acid 133, 140, 356, 357, 358, 369
Gallotannin 141
Galls 367
Garden angelica 352, 353, 556
Garden basil 292, 294
Garden dill 295
Garden rue 350, 353, 435
Gardenia 244
Garlic 735, 759
Gaultherin 139, 141, 247, 304
GC 734
GC – analysis 250
GC-MS 734
Gein 246
Geissoschizine 434
Gelatin 115, 368, 792
Gelseminic acid 348
Gelstaph 632
Gene cloning 73
Gene cloning techniques 48
Gene coding 17
Gene splicing 79
Gene splicing and DNA recombinant procedures 73
Gene therapy 75, 77, 78
Gene-cloning 72
Gene-splicing 73, 74
General properties of terpene phenols 288
Generation 626
Genetic code 44
Genetic combination 73
Genetic composition 22
Genetic effects 22
Genetic engineering 48, 49, 61, 73
Genetically engineered potato 82
Genetically engineered vaccines 72, 79
Genetically nonmutant ADA gene 78
Genetically–engineered–drugs 17
Genin 122
Genistein 747, 748, 823
Genome 50, 53
Genomic fragment 53
Genoptin 582
Gent-ophtal 582
Gentacin 582
Gentak 582
Gentalline 582
Gentalyn 582
Gentamicin 581, 585, 586
Gentamicin C1 581, 582
Gentamicin C1a 581, 582
Gentamicin C2 581, 582
Gentamicin complex sulphate 582
Gentamicin congeners 582
Gentamicin hydrochloride 582
Gentamycin 16, 581
Gentamycin 16, 576, 580, 581, 634, 637
Gentian 196, 197, 547
Gentianic acid 199, 566
Gentianin 199, 566
Gentibioptal 582
Genticin 582
Gentiin 199, 566
Gentinin 199
Gentiocrucine 199
Gentiogenin 198
Gentiopicrin 198, 199
Gentiopicroside 198
Gentioside 199
Gentisic acid 199
Gentisin 199, 566
Gentisin diacetate 566
Gentocin 582
Gentogram 582
Gentrasul 582
Genus 397
Geographutoxins 725
Geometrical isomerism 267
Geometrical isomerism 256
Geomycin 652
Geopen 636
Geranial 272, 273
Geranial-cyanoacetic anhydride 273
Geranic acid 273
Geraniol 26, 27, 217, 266, 267, 268, 269, 270, 271, 504, 505
Geranoil 273
Geranyl acetate 301
Geranyl pyrophosphate 233
Geranyl-1 700
Geranylhydroquinone 700
Germ cell 77
Germacranolide 230
German chamomile 347
German chamomile 232
German rue 350, 353, 435
Germiciclin 653
Germination 803
Germine 420, 423
Gernebcin 589
Geroquinol 700
Gerrard reaction 470
Gibberellic acid 807
Gibberellins 31, 807
Gibicef 604
GIIIA 725
Gildemeister Hoffmann's method 275
Ginger 323, 326, 327, 735
Ginger oil 241
Gingergrass oil 285
Gingerols 327
Ginkgo 6
Ginkgolide 19
Ginkgolide–A 6
Ginkgolide–B 234
Ginkgolides A,B, C and M 235
Ginning 98
Ginseng 7, 187, 188, 802
Ginsenoside 19, 7
Ginsenoside Rg 743
Ginsenoside Rg1 188
Ginsenosides 188
Githagenin 187
Gitin 146
Gitoxin 204
Glabra 189
Glandulifera 189
Gastrointestinal cancer 67
Globe fish 712
Globin 790
Globularicitrin 749
Glory lilly 356
Glucan 86
Glucarophanin 773
Gluco-alkaloids 377
Glucobrassicin 756
Glucocorticoid 644
Glucofrangulins A and B 136
Glucoheptonate 578
Gluconapin 180
Gluconasturtiin 756

- Gluconic acid quinidine salt 528
 Glucoproteins 794
 Glucosins 306
 Glucorhein 132
 Glucosamine 579, 763, 764
 Glucosamine sulphate 764
 Glucose 35, 159, 176, 190, 246, 348, 588, 589
 Glucose coniferin 144
 Glucose metabolism 767
 Glucosidase 35
 Glucoside (salicin) 142
 Glucosides of rhein 761
 Glucosinolate compounds 179
 Glucosinolate precursor 773
 Glucosinolates 209, 340, 755, 756, 757
 Glucotropaeolin benzyl isothiocyanate 375
 Glucovanillin 139, 144, 195
 Glucovanilline alcohol 196
 Glucoxylose 304
 Glucuronic acid 333
 Glutathione 743, 772
 Glutathione-S-transferase 743, 756, 757
 Glycan 86
 Glycerine of starch 91
 Glycerophosphoric acid 36
 Glycitein 747, 748
 Glycoaminoglycan (CAG) 763
 Glycogen 35, 795
 Glycolysis 780
 Glycolytic enzyme 783
 Glycopeptide antibiotic 648
 Glycoprotein (Purified) 63, 240, 833
 Glycopyrrolate 12
 Glycoresins 314
 Glycosidal resins 201, 308
 Glycosides 21, 135, 146, 122, 236, 238, 246, 364, 400, 745
 Glycosides in natural plant sources 211
 Glycosidic linkages 124
 Glycosidic veratrum alkaloids 420
 Glycosidically 646
 Glycosylation 59, 205
 Glycyrrhetic acid 190, 191
 Glycyrrhetic 190
 Glycyrrhetic acid 191
 Glycyrrhiza 189, 190, 191
 Glycyrrhizic acid 190
 Glycyrrhizin 190, 191, 743,
 Glyoxaline 437
 Glyoxylic acid 485
 Glyoxylic acid reagent test 537
 Gnoscopine 480
 Golden groundsel 474
 Gonorcin 673
 Good cholesterol 748
 Gorgonian corals 716
 Gossypin 161
 Gram-negative bacillii 605
 Gram-negative organisms 659
 Gram-positive bacteria 656
 Gram-positive organism 659
 Gramaxin 602
 Granular 105
 Granular honey 87
 Grape juice 751
 Greanyl acetate 301
 Great burdock 341
 Great scarlet poppy 486, 490, 491, 494
 Great scouring rush 342, 456
 Grecian laurel 493
 Green bay 297, 493
 Green coffee 342
 Green marine algae 794
 Green tea 735, 745
 Green tea solids 746
 Grepafloxacin 675
 Grifulvin 681
 Grignard reaction test 176
 Grignard reagent 176, 730
 Grignardization of puupehenone 730
 Grinsil 633
 Gripe water 297
 Gris-PEG 681
 Grisactin 681
 Griseofulvin 681
 Griseofulvin 681, 682
 Griseophenone C 682
 Grisovin 681
 Growth factor-1 740
 Growth phase 573
 Growth phase of the organism 572
 Growth regulators 802
 Grysio 681
 GTXs 713
 Guadeloupe 437
 Guaiac 315, 316
 Guaiac resin 315
 Guaiacic acid 316
 Guaiacol 27, 282, 355
 Guaiaconic acid 310
 Guaiacum 39
 Guaiacum resin 307, 315
 Guaianolides 230
 Guaiarresinol 312
 Guaiazulene 229
 Guaic 39
 Guanine 44, 47, 48, 201
 Guar gum 111
 Guaran 2, 111, 112
 Gum asafoetida 329
 Gum benzoin 356
 Gum guaiac 315
 Gum opium 7
 Gum rosin 314
 Gum tragacanth 105
 Gum-resins 309
 Gum-turpentine 328
 Gumlich aldehyde 509
 Gums 100, 281
 Gurjuresinol 312
 Gutta-percha 313
 Gymnodinium catenatum 711
 Gynergen 539
 Gynesine 37
 Gyno khellan 553
 Gypsogenin 187
- H**
- Habitrol 456
 Hager's reagent 385, 438
 Halichondrin 720
 Halogen acids (HX) 726
 Hard-gelatine 792
 Harmaline 501, 502, 503, 504
 Harmaline hydrochloride dihydrate 502
 Harman 501, 503, 504
 Harman hydrochloride 501
 Harmel 427, 501
 Harmine 501, 502, 503, 504
 Harmine hydrochloride dihydrate 503
 Harvatrate 465

- Harzol 760
Haworth lignan 361, 755
Hawthorn 348
HDL cholesterol 748
Hederagenin 187
Helch's reaction 439
Helch's violet-colour test 439
Helianthic acid 643
Heliotrope 474
Heliotropin 295
Heliotropium 471
Helper t-cells 46
Helvamoxy 633
Hematopoietic functions 77
Hemiacetal 621
Hemiketal 621
Hemiterpene 251
Hemoglobin protein sequences 75
Hemoglobinulin 790
Hemp 397
Henbane 346, 397, 462, 469
Henblain 346, 397, 462
Henblain jusquaime 469
Heparin 792
Hepatic 128
Hepatic glucuronidation 675
Hepatitis 69
Hepatitis B 72
Hepatitis B surface antigen (HBSAG) 72
Heptane macrolide antifungal antibiotic complex 665
Herbal drugs 827, 828
Herbal medicines 736
Heroin 389
Herpathite 433
Herpathite test 433, 523
Hesperetin 743, 748, 750
Hesperidin 163, 164
Hesperidin chalone 164
Hesperidin methyl chalone 164
Hetastarch 86, 92
Heterocyclic terpene aldehyde 283
Heteroglycan 85, 86, 100
Heterotrimeric glycoproteins 702
Hexahydrocymene 254
Hexamethylamine 677
Hexamine 677
Hexosamine pathway 764
Hexosans 86
Hexosulose 589
Hi-tech products 814, 822
Hibiscitrin 161
High molecular weight phenylpropanoids 340, 366
High performance liquid chromatography (HPLC) 389, 395
High performance thin-layer chromatography (HPTLC) 389, 395
High throughput random screening programme [HTRSP] 820
High throughput screening (HTS) 815
Higher-plant enzymes 781
Hindu datura 456, 462, 469
Hip berries 37
Histamine 788
Histidine 789
Historic genesis of PTCT 799
HIV 70
HIV antigens 70
HIV replication 834
HIV reverse transcriptase 834
HIV-antibodies 70
HNO₃ test 352
Holothurins 705
Holothurins and astrosaponins 705
Holothuroids 705
Homatropine 12
Homoallytic isomerization 518
Homocarotenoids 238
Homoeriodictyol 315
Homoglycan 85, 86
Homoharringtonine (HHT) 824, 825
Homopilopic acids 438
Homospermidine 451, 476
Homotropine 389
Hon-Sho 294, 341
Honduras balsam 335
Honey 86
Hordenine 398, 475, 477, 478, 479, 482
Hordeum vulgare 398
Hormones 45, 61, 63, 809
Horse chestnut 350
Horse chestnut tree 828
Horse gram 794
Horse raddish root 36
Horse-hoof 358, 367, 474
Horsemint oil 225
Horseradish 757
Horsetail kelp 35
Host 47
Hostacyclin 650
Hot jalapeno peppers 768
Hrodenine 479
HTS and bioassays 816
HTS and selection for plant materials 818
HTS-programme 821
Huang 477
Huisache 304, 346, 356
Human clone 80
Human deacetylase 754
Human DNA 51
Human fertilization and embryology authority (HFEA) 81
Human Immunodeficiency Virus (HIV) 65
Human insulin 61
Human proteins as drugs 61
Humatrope (R) 79
Humulin (R) 61, 79
Humulon 548, 549
Hungarian camomile 347
Hyaluronic acid 785
Hyaluronidase 783, 785
Hyasorb 627
Hybrid-antibiotics 575
Hybridization probes 49
Hybridoma technology 66
Hybrimycins 18
Hydramycin 654
Hydrangea 161
Hydrangin 33, 347
Hydrastine 374
Hydrate rule 84
Hydrocarbon volatile oils 251
Hydrochloride 410
Hydrocortisone 14
Hydrocoumarone rotenone 560
Hydrocyanic acid 176
Hydrogen gas 477
Hydrogen sulphide adduct 286
Hydrogenation 223
Hydrolytic enzyme 783, 780
Hydrolyzable tannins 38
Hydrolyzing effect 391
Hydromorphone 11

- Hydroquinone 140, 204
Hydrorhombinine 450
Hydrosarpan 505
Hydrosulphoric derivative 275
Hydroxy anthraquinone derivatives 775
Hydroxycinnamic acids 340, 341
Hydroxycoumarins 345, 346
Hydroxygentamycin 18
Hydroxylated steroidal glycosides 705
Hydroxynitrilase 175, 176
Hydroxynitriles 207
Hygric acid methylbetaine 463
Hygrine 374, 396, 398, 462, 471
Hylenta 627
Hymenin 708, 709
Hymenotera 86
Hyoper 636
Hyoscyne 389
Hyoscyamine 375, 376, 381, 388, 389, 398, 464, 469, 470, 471, 473, 811
Hyoscyamine hydrobromide 470
Hyoscyamine hydrochloride 470
Hyoscyamine methyl bromide 470
Hyoscyamine oxalate 470
Hyoscyamine sulphate dihydrate 471
Hypericin 761, 763
Hypericum red 763
Hypnea valentiae 708
Hypotensive compounds 701, 706
Hypotensive peptides and other compounds 708
Hypothetical intermediate 510
- I**
- I-carrageenan 116
I-indolizidinone 455
Ibiamox 633
Ibuprofen 11
Ichthyotoxic actions 705
Idamycin 595
Idarubicin 590, 594, 595
Idarubicin hydrochloride 595
Ideal bioassays 816
Ideal nutrient medium 802
Identification of taxol 822
Idiophase of the organism 572
Ikaran 539
Ilimaquinone 731
Ilixathin 749
Illurinic acid 324
Imidazole 396, 399, 687, 688
Imidazole alkaloids 396, 441
Imidazoles 396
Imipem 614
Imipenem 611, 612, 613, 614
Immobilization of cells 799
Immobilized cell cultures 811
Immune protein 791
Immune surveillance 75
Immune system 46
Immuneron 80
Immunizing agents 72
Immunoactive drugs 64
Immunoassays 67
Immunocompromised hosts 680
Immunogenic IL-1 740
Immunomodulators 830
Imperatorin 350, 353, 556
Important penicillin drugs 627
Improved strains 574
In vitro fertilization (IVF) 80
Inactive 780
Inamycin 663
Incomplete protein 789, 791
Indian apple 318
Indian balsam 335
Indian bdellium 329, 333
Indian ginseng 187
Indian podophyllum 319
Indian squill 152
Indian tobacco 443, 444
Indian traditional herbal drugs 827, 828, 831
Indian traditional herbal medicine 830, 833
Indian tragacanth 104
Indirect ELISA-test 70, 71
Indocybin 499
Indole 399, 531
Indole alkaloidal 544
Indole alkaloids 396, 427, 521
Indole nucleus 535
Indole-3-carbinol 758, 756
Indole-related alkaloid 725
Indole-ring system 495
Indoles 757
Indolizidine 453, 455
Indolizidine alkaloids 453
Indolizidine-alkaloid biosynthesis 454
Indolquinazolines 396
Inert constituents 21
Inflatine 443
Influenza vaccine 73
Inhibition of antibiotic production 574
Inhibition of cell-wall formation 663
Inhibition of protein synthesis 659
Inhibitors of human thrombin 821
Inhibitory enzyme 783
Initiation 825
Insect propellents 217
Insect tissue culture 58
Insecticidal agents 809
Insertional mutagenesis 17
Insulin 45, 61, 740, 764
Intact plant 543
Intense protein synthesis 825
Interleukin 62
Interleukin-2 46
Intermediate acting cephalosporin 610
Intermediate-acting second generation cephalosporin 605
Intermediate-spectrum fluoroquinolone 675
Intoxicating mint 463
Intracellular enzymes 781
Intron-A(R) 79
Intybin 565
Inulin 86, 92, 93, 151
Invaluable orthodox drugs 814
Invertase 782
Inverting enzyme 783
Iodic acid test 488
Iodine ointment 179
Iodine value 179
Iodo-quinine 433
Iodoquinol 669
Iodosulphate of quinine 433
Ion channels 816
Ion-exchange 394
Ionone 274
Ipaccef 604

- Ipomea 536
 Ipomeamarone 807
 IR-Spectroscopy 250
 Ircinala [17] 729
 Iretin 698
 Iridocin 680
 Iridoid 412
 Iridoid materials 412
 Iron-binding phenomenon 753
 Isflavones 747
 Isinglass 115
 Isipen 638
 Isoapiole 297
 Isoarteril 505
 Isobarbaloin 129
 Isobergapten 171, 172
 Isoborneol 220
 Isocarhamin 166
 Isocaryophyllene 324, 325, 334
 Isocinnamic acid 335
 Isoeugenol 282
 Isoflavone 167, 158, 561, 747, 748
 Isoflavone compound 559
 Isoflavonoid glycoside 159, 166, 212
 Isoflavonoid ring system 364
 Isoflavonoids 166, 561
 Isolated plant cells 799
 Isolated plant organs 800
 Isoleucine 789
 Isomelacidin 168
 Isomenthone 285
 Isomerases 781
 Isomeric menthanes 254
 Isomerism 378
 Isomerization 217
 Isomerization of carvone to carvacrol 287
 Isomyristicin 295
 Isoniazid 642, 679, 678
 Isonitrosochloride 217
 Isopentenyl diphosphate (IPP) 712
 Isopilocarpine 437, 438, 439
 Isopilocarpine hydrochloride 439
 Isopilocarpine hydrochloride hemihydrate 440
 Isopilocarpine nitrate 440
 Isopilosine 440
 Isopimpinellin 558, 557
 Isoprene 215, 738
 Isoprene (2-Methyl buta 1,3 Diene) 216
 Isoprene units 215, 250, 251, 712
 Isoprenoid 215, 738
 Isopropanol 226
 Isopropylethiogalactoside (IPTG) 49
 Isopsoralen 171
 Isopulegol 275, 276
 Isoquinoline 398, 399
 Isoquinoline alkaloids 396
 Isorhamnetin 138
 Isorhamnetin (H) 206
 Isosafrole 294, 295
 Isothiocyanate glycosides 179
 Isothiocyanates 757
 Isotopically labelled organic compounds 119
 Itraconazole 688
 Ituran 677
- J**
- Jaborandi 367, 437
 Jaclacin 594
 Jalap 201, 309
 Jamaica dogwood 559
 Jamaica pepper 297
 Jamaican quassia 563
 Japanese ginseng 187
 Jasmine 244, 305
 Java lemon oil 274
 Jellyfish 37
 Jenacylin 654
 Jequerity 357
 Jervine 420, 424, 425, 426
 Jervine hydrochloride 426
 Jeveratrum alkaloids 424, 425
 Jimson weed 465, 469
 Jusquaime 346, 397, 462
 Justicia adhatoda 428
- K**
- K-carrageenan 116
 K-strophanthidin β 154, 155
 K-strophanthoside 154, 155
 Kaempeerol(D) 206
 Kaempferin 161
 Kaempterol 138
 Kafur 219
 Kainic acid 721, 722, 723
 Kaminax 581
 Kamycin 583
 Kamynex 583
 Kanabristol 583
 Kanacedin 583
 Kanamycin 18, 576, 580, 582, 583, 584
 Kanamycin A 582, 583
 Kanamycin A sulphate 583
 Kanamycin B 583, 590
 Kanamycin B sulphate 583
 Kanamycin C 583
 Kanamytrex 583
 Kanaqua 583
 Kanasamine residue 590
 Kanasig 583
 Kanatrol 583
 Kanendomycin 583
 Kanendos 583
 Kanesein 583
 Kanicin 583
 Kannasyn 583
 Kano 583
 Kantres 583
 Kantrexil 583
 Kantrox 583
 Kaolin 109
 Karaya gum 101, 104, 105
 Katine 410
 Katum 397, 506
 Kautschin 742
 Kava 322
 Kava pyrones 323
 Kava-kava 322, 356
 Kawain 322, 323
 Kedde test 147
 Kedde's reagent 147
 Kefadol 603
 Kefandol 603
 Keflin 601
 Kefroxil 602
 Kefurox 604
 Kefzol 602
 Kelamin 553
 Kelgin 35
 Kelicor 553
 Kelicorin 553

- Kellar-kiliani test 147, 155
 Keller's test 537
 Kellin 553
 Keloid 553
 Keratin 22, 790
 Keratin precursor cells 682
 Kerrolic acid 321
 Kesint 604
 Keto-alcohol 406
 Ketoconazole 688
 Ketone volatile oils 251, 284
 Ketones 27
 Khat plant 410
 Khelfren 553
 Khella 553
 Khellin 170, 171, 553, 554, 555, 556
 Khellinin 555
 Khellol 170
 Khellol glucoside 170, 535
 Khellol glucoside tetracetate 555
 Kibbling machine 110
 Kiditard 528
 Killer T-cells 46
 Kinase C 831
 Kinetin 804
 Kinichron 528
 Kinidin durules 528
 Kippenberger's process 393
 Klavocin 639
 Klebcil 583
 Klimicin 617
 Klinomycin minomycin 655
 Klion 669
 Klorita 38
 Klunge's isobarbaloin test 131
 KMnO_4 oxidation 438
 KoGENate (R) 42
 Kohlenhydrates 84
 Kolanuts 2
 Korean ginseng 187
 Krantum 506
 Kratum 397
 Kraut's reagent 385
 Krebs cycle 780
 Kusum tree 320
 Kutkoside 197
 Kutum 397, 506
 Kyocrystine 515
- L**
- L(+)-Ephedrine 407
 L-2-Deoxyfucose 594
 L-alanine 540
 L-arginine 418, 713
 L-asp 428
 L-asparaginase 784, 789
 L-aspargine 789
 L-aspartic acid 789
 L-camphor 28
 L-carrageenan 116
 L-carvone 27, 286
 L-catechin 290
 L-cladose 618
 L-cocaine 466
 L-dopa 482
 L-ecgonine 468
 L-ephedrine hydrochloride 478
 L-ephedrine 409, 478
 L-ephedrine sulphate 478
 L-form lupinine hydrochloride derivative 449
 L-HCl 408
 L-istidine 436, 441
 L-hyoscyamine 469
 L-lupanine 450
 L-lupinine 449
 L-lys 455
 L-lysine 441
 L-menthone 284, 285
 L-mycarose 618
 L-oleandrose 618
 L-ornithine 441, 461
 L-phenylalanine 405, 409, 412, 540
 L-pipecolic acid 454, 455
 L-proline 540
 L-santonin 550
 L-sparteine 450
 L-streptose 587
 L-sulphate 408
 L-tropic acid ester with tropine 469
 L-tropine tropate 469
 L-tryptophan and secologanin 525
 L-tyrosine 405
 Labetalol 234
 Lac 320
 Lac Z gene (B) 49
 Lacca 320
 Laccic acid 321
 Lactalbumin 790, 793
 Lactobionate 578
 Lactoglobulin 790, 793
 Lactone bitter principles 547, 550
 Lactone coumarin 345
 Lactone moiety 698
 Lactones 30
 Lactucin 564
 Lactucopicrin 565
 Lagistase 751
 Lambda phage (I) 50
 Laminin 702
 Lamisil; SF-86-327 685
 Lamoryl 681
 Lamoxactam 608
 Lampocef 602
 Lampomandol 603
 Lamsporin 604
 Lamuran 505
 Lanceleaf periwinkle 508, 511
 Landomycin 623
 Lanosterol 216
 Lantadenes 821
 Lantana poisoning 821
 Lappa 341
 Laragon 750
 Larger segments of DNA 49
 Largest biotech startups 61
 Lastet 362
 LAT-A 722
 LAT-B 722
 Latamoxef 608
 Latex 39
 Latrunculin A (LAT-A) 721, 722
 Latrunculin B (LAT-B) 721, 722
 Latrunculins 721
 Lauraceae 219
 Laurel leaf oil 221
 Lavender 301, 302
 Lavender oil 241, 302
 Lavender-cotton 342
 Lavondula 264
 LCR 515
 LDL cholesterol 740, 748, 751
 LDL oxidation 750, 752
 Lead acetate test 102
 Lead compounds 733
 Lecithins 36
 Lecithoprotein 790

- Lectins 772, 794
Lectucin 565
Lectucin para-hydroxyphenylacetate hydrate 565
Ledercillin VK 631
Ledermycin 652
Lee's method 727
Legal test 147, 155
Legalon 750
Legumes 748
Leguminosae 168, 169, 307
Leguminosae 127, 137
Leishmaniasis 69
Lemon 37, 243
Lemon grass oil 272, 301
Lemon, lime, orange and ginger root 272
Lemon oil 274
Lemon verbena 297
Lemonol 26
Lentisk 318
Leopard's-bane 341, 348
Leprosy 69
Lesch-nyhan syndrome 77
Lethal (toxic) dose (LD50) 814
Leucine 789
Leucoanthocyanidins 168
Leucocyanidin 34
Leucocyanidol 34
Leucoharminine 502
Leukaemomycin C 690
Leukine (R) 79
Leupine (R) 42
Leurocristine 515
Levant storax 334
Levant wormseed 551
Levomycetin 38
Levorin 664
Levorin A2 665
Levsin 469
Lexinor 673
Libitatae 264
Licheniformin 573
Lichenin 86, 94
Lichens 127
Licorice root 367
Liferoot 474
Ligases 47
Lignan 361, 755
Lignan glycosides 319
Lignans 316, 319, 361, 362, 748, 753, 755
Lignin 21, 27, 97, 281, 366, 367
Lignin waste 281
Ligustrin 830
Likuden 681
Liliaceae 150, 375
Lily of the valley 341, 342
Lime 301, 350
Lime tree 302, 359, 360
Limonene 26, 216, 217, 250, 255, 256, 257, 258
Limonoids 742
Linaloe oil 269
Linalool 269, 270, 271, 273, 302
Linalyl acetate 271, 301, 302,
Linamarin 174, 179
Linase 179
Lincocin 617
Lincolcina 616
Lincoln bean 776
Lincolnensin 616
Lincomycin 616, 617
Lincomycin hydrochloride hemi-hydrate 617
Lincosamides 579, 616
Linden tree 301, 302, 359, 360
Linear furanocoumarin 207
Lingraine 539
Lingran 539
Linseed 178
Linustatin 174
Lipases 781, 786
Lipid metabolism 765
Lipolytic enzyme 783
Liquid cultures 799
Liquid nutrient medium 801
Liquid suspension cultures 801
Liquid-liquid extraction 576
Liquiritigenin 165
Liquiritin 163
Liquorice 189, 191
Liquorice compound powder 191
Live vaccines 47
Livery 128
Liviqlina 602
Lobelanidine 445, 446
Lobelanidine hydrobromide 446
Lobelanidine hydrochloride 446
Lobelanine 442, 444, 445, 446, 447
Lobelanine hydrobromide 445
Lobelanine hydrochloride 445
Lobelanine nitrate 445
Lobelia herb 443
Lobeline 389, 396, 397, 398, 442, 443, 444, 445, 446, 447
Lobeline hydrochloride 444
Lobeline sulphate 444
Lobeton 444
Lobidan 444
Lobron 444
Locumin 174
Locust bean gum 110
Loganin 504, 505
Lomebact 673
Lomefloxacin 671, 672, 673
Lomefloxacin monohydrochloride 673
Londomycin 653
Loricin 640
Lospoven 601
Lotaustralin 174
Loturine 501
Loveapple 397, 462, 469
Lugacin 582
Lukadin 580
Luminescence immunoassays (LIA) 67
Lupanine 396, 449, 450, 451
Lupeol 186
Lupin beans 450
Lupinidine 450
Lupinine 399, 449, 451
Lupulon 548, 549, 550
Lutein 741, 738
Luteolin (E) 206
Lycocetone 400
Lycopene 239, 738, 739, 740
Lycopodine 543
Lymecycline 649, 650, 656
Lymecycline sodium 656
Lymphogranuloma 661
Lymphokines 61, 62
Lypases 782
Lyphocin 646, 647
Lyphomed 646
Lysergamde 536
Lysergic acid 539
Lysergic acid amides 535
Lysergic acids 535

Lysine 449, 574, 766, 789
Lythreaceae 127

M

m-conotoxins 725, 726
m-cresol 28, 226, 333
m-thymol 225
Ma huang 410
MABs in diagnostics 66
MABs in imaging and therapy 67
Mace 28, 295
Maceration method 246
Macrobid 678
Macroclant 677
Macrolactone 578
Macrolide antibiotics 618, 622
Macrolides 579, 618, 624, 696
Macrophages 77, 785
Madagascar 505, 506, 507, 508, 512
Mael 345
Magic-bullet 45
Mahogany wood 34
Maitotoxin 715
Maitotoxin (MTX) 714
Maizena 91
Major cinchona alkaloids 530
Malabar nut 427
Male fern 323, 325
Maleric anhydride 293
Malimyxin 645
Malonate 405
Malonyl-CoA 436
Maltase 795
Malvidin 167, 745
Mammalian liver enzyme systems 727
Manaca 348, 397
Manacan 348, 397
Manchurian bean 776
Manchurian liquorice 191
Mandalin's reagent 386
Mandalin's test 157
Mandarin peel oil 26, 257, 285
Mandelin's reagent 517
Mandelonitrile 175
Mandelonitrile glucose 176, 177
Mandokef 603
Mandol 603

Mandolsan 603
Mandrake 397, 462, 469
Mandrakes root 318
Mannich reaction 447
Mannich-like mechanism 484
Mannich-like reaction 503
Mannich-like reaction using keto acid 504
Mannich/pictet-spengler type reaction 500
Mannitol 190
Manzamine A 728
Manzamine A [15] 729
Manzamine A and ent-8-hydroxymazamine A 728
Manzanilla 347
Maranhm jaborandi 437
Marijuana 397
Marine algae 769
Marine biomedicines 706
Marine biotoxins 709
Marine black sponges 719
Marine bryozoans 831
Marine glycosides 705
Marine gums 112
Marine natural products 726
Marine neurotoxin 712
Marine nucleosides 706
Marine organisms 233, 695, 696, 701, 706, 708, 709, 717, 720, 724
Marine peptides 701, 702
Marine toxins 696, 709
Marine-based chemical substances 721
Marine-chemists 718
Marine-derived antibiotic substance 718
Marine-world 733
Marme's reagent 385
Marmesin 558
Marqui's reagent 354, 386
Martec 30
Mask root 347
Mass spectrometry 250, 822
Mastic 310, 318
Mastiche 318
Mastichic acid (a-resin) 318
Masticin (b-resin) 318
Masticoresene 312
Mastimyxin 643

Mastisol 318
Mastix 318
Mate 2
Matricarin 232
Maw seed 7
Maxaquin 673
May apple 318
May pop 501
Mayer's reagent 385, 438
Maypop 501
MDL-507 648
Meadosweet 304, 359
Meceration method 245
Mechanical process 244
Mechothramine hydrochloride 574
Meciclin 652
Mecke's reagent 386
Meconic acid 4, 34, 376
Medicinal chemistry 4
Medicinal compounds 798
Medicinal herbs 354
Medicinal plant 20
Medicinal plant biotechnology 813
Medicinal plants 811, 813, 827
Medium-acting sulphonamide 668
Medoxim 604
Mefoxin 610
Mefoxitin 610
Melacine 80
Meladinine 351
Melatonin 777
Meletin 750
Meliaceae 238
Melicopicine 435, 436
Melilotus alba 205
Melin 749
Melitoxin 33
Meloxine 351
Memory cells 46
Menthane 254, 264, 265
Menthol 22, 27, 219, 223, 224, 249
Menthone 224, 226, 275, 276, 285
Menthoxo acetate 225
Mephacyclin 650
Mephalan 574
Meractinomycin 692
Mercapto cysteins 759
Mercaptothiazole 574
Meristem culture 801

- Meronom 614
Meropenem 612, 614, 615
Meropenin 631
Meroterpenoids 216
Merrem 614
Merxin 610
Mescal buttons 479, 483
Mescaline 381, 398, 477, 479, 480, 482
Mescaline acid sulphate 480
Mescaline hydrochloride 480
Mescaline sulphate dihydrate 480
Meta-menthadiene 258
Metabolic engineering 46
Metabolin 37
Metabolism studies 726
Metabolites in culture 811
Metacycline 653
Metadomns 653
Metanite 465
Metel 456, 462, 469
Methacolimycin 645
Methacycline 649, 650, 653, 654
Methacycline hydrochloride 653
Methadone 11
Methenamine 677
Methicillin 628
Methicillin-resistant 570
Methionine 574, 766
Methocillin-S 632
Methoxsalen 350, 351, 352, 556, 558
Methoxsalen (Xanthotoxin) 354
Methyl amide 438
Methyl amine hydrogen 406
Methyl anthranilate 305, 306
Methyl arbutin 140
Methyl chalone 163
Methyl D-glucosides 122
Methyl ergometrine 428
Methyl hydrate betaine 463
Methyl salicylate 30, 141, 247, 301, 302, 304, 305, 356, 358
Methyl thiosulphinate 759
Methyl- α -D-Glucoside 123
Methyl- β -D-Glucoside 123
Methyl-ecgonine 473
Methyl-isopropyl cyclohexane 254
Methylamine 446, 477
Methylarbutin 140
Methylated cyclohexanes 217
Methylation 650
Methylation of the hesperidin chalone 163
Methylatropine nitrate 465
Methylergonovine mealeate 538, 539
Methylionone 274
Methylmorphine 490
Methysticin 322, 323
Metilenbiotic 653
Metrogel 669
Metrolag 669
Metrolyl 669
Metronidazole 614, 669, 670
Metropine 465
Mevalonate 210
Mevalonic acid 203, 204
Mexocine 652
Mezereon 347, 349
Mezlin 637
Mezlocillin 629, 635, 637
Mezlocillin sodium 638
Mezlocillin sodium monohydrate 637
MIC 664
Micatin 689
Miconazole 681, 688, 689
Miconazole nitrate 689
Micoserina 641
Micotef 689
Microaerophilic 669
Microbe 568
Microbial cultures 727
Microbial gums 116
Microbial metabolites 16
Microbial transformations 726
Microbiology 571
Microchemical test of cineole 298
Microcillin 636
Micronutrients 802
Microorganism 47, 726
Micropolysaccharides 763
Microtin 601
Mignonette 357
Migranal 539
Mikavir 581
Milk thistle 750
Mimosa 244
Mineral chelation 752
Mineralocorticoid activity 190
Miniaturized assay: HTS 815
Minimum inhibitory concentration (MIC) 571
Minimum threshold concentration 578
Minocycline 649, 650, 655
Minocycline hydrochloride 655
Minocyn 655
Mioton 36, 323, 410, 768
Miscellaneous antibiotics 579, 657
Miscellaneous bitter principles 547, 561
Miscellaneous compounds 697
Miscellaneous glycosides 127, 200, 214
Miscellaneous pharmacologically active substances 696
Mithramycin 16
Mitocin-C 693
Mitomycin C 16, 690, 693
Mitosis 51
Mitoxantrone 590, 595, 596
Mitoxantrone dihydrochloride 596
MMC 693
Moderil 513
Modified benzyltetrahydroisoquinoline alkaloids 485
Modified borntrager's test 130, 135, 139
Modified dragendroff's reagent 386
Modified pepsin 787
Modified tetrahydro iso-quinoline alkaloids 475
Modifying dietary components 760
Mole plant 341, 343, 349
Molecular biology 575
Molecular probing 75
Molecular refraction 250
Molecular scissors 47
Molecular vehicle 46
Monalide 720
Monapen 637
Monasirup 291
Mondamin 91
Monistat 689
Monkshood 341, 412, 450, 477
Monoacetyl puupehenone 730
Monoacetyl structural analogue 730
Monobactam 612
Monobenzoyl glucose 143

- Monocid 605
 Monocidur 605
 Monoclonal antibodies (MABs) 66
 Monocot plants 183
 Monocotyledons 375
 Monocyclic 228
 Monocyclic phenol 28
 Monocyclic terpene 27
 Monocyclic terpene aldehydes 276
 Monocyclic terpene ketones 284
 Monocyclic terpenes 254, 256
 Monocytes 785
 Monohydric phenols 288
 Monopen 627
 Monosaccharide D-(+) – glucose 90
 Monosaccharides 167
 Monoterpene 141, 251
 Monoterpenoids 215, 218, 219, 287
 Moraceae 171
 Moray eel 715
 Morena 539
 Moreton bay chestnut 453
 Mormon tea 406, 477
 Morphia 486
 Morphina 486
 Morphinan 497
 Morphinan alkaloids 486
 Morphine 4, 10, 11, 19, 23, 25, 7, 374, 376, 377, 380, 381, 382, 383, 388, 389, 391, 396, 397, 480, 486, 487, 488, 489, 490, 494, 513, 814
 Morphine 3-methyl ether 490
 Morphine acetate trihydrate 489
 Morphine hydrochloride 380, 387
 Morphine monohydrate 489
 Morphine monomethyl ether 490
 Morphine tartrate trihydrate 489
 Morphinone 486, 494, 495, 497
 Morphinum 486
 Morphological classifications 810
 Mostly containing nitrogen 36
 Motherwort 463
 Mountain tobacco 341, 348
 Moxalactam 600, 608, 609
 Moxalactam disodium 608
 Moxaline 633
 Moxam 608
 MS 734
 MTX 715, 716
 Mucin 790
 Mucoids 790
 Mucolytic enzyme 783
 Mucomycin 656
 Mucoproteins 763
 Mugwort 297
 Multicellular system of plant 799
 Multimycine 645
 Multiple cloning site 49
 Muramidase 783, 785, 786
 Murein 642
 Muscatel sage 297, 302
 Mustard 757
 Mustard oil 757
 Mustard seeds 125
 Mutagenesis 56
 Mutamicins 18
 Mutamycin 693
 Mutant 53
 Mutant strains 650
 Mutation and strain selection: mutation 574
 Myacine 585
 Mycaminose 619
 Mycarose 619
 Mycinose 619
 Mycivin 617
 MycoPlex-7™ 778
 Mycosis fungoides 351
 Myoinositol 589
 Myoscint 80
 Myosin 790
 Myrcane 264
 Myrcene 216, 218, 250, 253
 Myrcia oil 287
 Myristicin 28, 291, 295, 300, 344
 Myristicin aldehyde 296
 Myristinic acid 296
 Myrosin 182
 Myrrh 310, 329, 332, 333
 Myrtenal 279
 Mytricolorin 749
- N**
- N-formylmethionine 59
 N-acetylthranilic acid 428
 N-acetyl derivative 585
 N-acetylglucosamine 764
 N-acetylharmaline 502
 N-acetylmescaline 480
 N-atom 382
 N-benzoylmescaline 480
 N-glycosides 124, 125
 N-glycosylation 59
 N-lysinomethyl tetracycline 656
 N-methyl anthraniloyl-CoA 436
 N-methyl conine 401, 404
 N-methylmescaline 480
 N-methylpyrroliniumcation 460
 N-oxide derivative 459
 NAD(P)H quinone reductase 756
 NADPH 525, 530
 Nafcil 632
 Nafcillin 628, 632
 Nafcillin sodium 632
 Naftifine 681, 685, 812
 Naftifine hydrochloride 685
 Naftifungin 685
 Naftin 685
 Naftopen 632
 Nalidixic acid 670, 671
 Narceine 380, 383, 397, 398, 477, 480, 487
 Narcissus 244
 Narcotine 374, 383, 384, 388, 480
 Narcyl 481
 Nareceine 480
 Naringenin 366
 Naringin 163, 165, 744, 749
 Naringin chalcone 165
 Naticardina 528
 National cancer institute (NCI) 696
 National sea grant office (NSGO) 696
 Native collagen 784
 Native dextran 95
 Native protein 791
 Natural antioxidants 744
 Natural biomedicinals 797
 Natural camphor 221
 Natural dyes 809
 Natural gums 100
 Natural killer (NK) cells 46
 Natural penicillins 627, 628
 Natural plant samples 820
 Natural product input 818
 Natural products 681
 Natural resins 309
 Natural sources of lectins 794
 Naturally occurring alkaloids 388

- Naturally occurring resins 310
Neamine 585
Nebactam 613
Nebramycin Factor 6 589
Necic acids 471
Necine 474
Neck squamous 693
Neem 736
Neem tree 238
Nematodes 552, 723
Neo-duplamox 639
Neo-fulcin 681
Neobreltin 585
Neocefal 603
Neofermergen 538
Neogermitrine 420
Neolignans 361, 755
Neomix 585
Neomycin 576, 584, 644, 646
Neomycin A 585
Neomycin A hydrochloride 585
Neomycin B 584, 585
Neomycin B hydrochloride 585
Neomycin B sulphate 585
Neomycin C 584, 585
Neomycin complex 584
Neonicotine 457
Neopinone 497
Neopogen (R) 79
Neoprotoveratrine 420, 421
Neosamine 579
Neosulf 585
Nepenthe 486
Nephrotoxicity 577
Neral 27, 272
Nereistotoxin 721, 725
Nerol 267, 268, 269, 270, 271
Neroli oil 305
Nerolidol 336
Netilmicin 576, 580, 585, 586
Netilmicin sulphate 586
Netilyn 586
Netromicine 586
Netromycin 586
Nettacin 586
Nettilin 586
Neurophysiological 696
Neurotransmitters 45
Neutral hydroiodide of quinidine 528
Neutral or slightly acidic alkaloids 384
Neutral salt of quinine 431
Neutral saponin 184
Nevada jointfir 406, 477
New antibiotic 571
New immunodiagnostic agents 66
New medicines 815
Newer antibiotics 18, 570
Newer fluoroquinolone 676
Newly fertilized embryo 81
NF 6 589
NF-kB transcription 754, 755
Niacin 454, 458, 459
Niacin sodium salt sesquihydrate 459
Nicabate 456
Nicandrin B 832
Nichogencin 582
Nicodeerm 456
Nicolan 456
Nycopatch 456
Nicotell TTS 456
Nicotiana 377
Nicotiana tabacum 543
Nicotine 376, 444, 454, 456, 458, 459, 460, 757, 809, 811
Nicotine bitartrate 457
Nicotine dihydrochloride 457
Nicotine hydrochloride 456
Nicotine salicylate 457
Nicotine sulphate 457
Nicotine zinc chloride double salt monohydrate 457
Nicotinell 456
Nicotinic acid 454, 459, 460, 461
Night blooming cereus 479
Ninhydrin reagent 716
Niopo 355
Nisotin 680
Nitric acid test 131, 354, 489
Nitric oxide 757
Nitric-oxide synthetase activity 759
Nitroalkane nucleophiles 733
Nitroamines 757
Nitroethane 733
Nitrofurantoin 677, 678
Nitrofurantoin monohydrate 678
Nitrogen sources 572
Nitroimidazoles 665, 668
Nitromethane 733
Nitromycin C 693
Nitrosamine 752
Nitrosochloride 217
Nitrosochloride adduct 217
Nitrous acid test 130
Nivemycin 585
NMR 734
NMR – spectrometry 250
Nocardicin 615
Nocardicin A 612, 615, 616
Noflo 673
Nolicin 673
Non-aromatic ring 590
Non-flavonoid polyphenolics 753
Non-heterocyclic alkaloids 397
Non-insulin-dependent diabetes mellitus (NIDDM) 765
Non-lactonic combranoid 700
Non-peptide marine biotoxin 710
Non-isoprenoid 216
Nor-ephedrine 407
Nor-pseudo-ephedrine 381
Nor-Y-ephedrine 407, 410
Noracin 673
Noradrenaline 481, 482
Noraxin 673
Nocardicin A 612
Norden 703
Norephedrine 409
Norepinephrine 481
Norfen 703
Norfloxacin 670, 671, 673
Norhalichondrin A 718, 720
Norkel 553
Norlupinane 399
Normal bovine collagen 784
Normal O-glycosides 134
Normicotine 459, 460
Norocin 673
Noroxin(e) 673
Norpseudoephedrine 405, 409, 410
Norxacin 673
Notannol 329
Notaral 627
Novamin 581
Novantrone 596
Novel biological targets 815
Novel furanocembranoids as antineoplastic agents 727

- Novel inhibitors of human thrombin 821
- Novel ionophoric marine substance acanthifolicin 719
- Novel polycyclic β -carboline-alkaloids 728
- Novel vaccines 63
- Novinose 663
- Novobiocin 663, 664
- Novobiocin monosodium salt 664
- Novocef 604
- Nuclear magnetic resonance (NMR) spectroscopy 822
- Nucleoside 201, 202, 214, 706, 708
- Nucleotide 201, 202
- Numerous software systems 816
- Nutmeg 295
- Nutraceutical 735, 736, 737, 764
- Nutrients of the future 737
- Nutropin (R) 80
- Nuts starch 91
- Nuvapen 633
- Nux vomica 516, 547
- Nystatin 689
- 
- O- and C-glucoside linkages 762
- O-19, 20-diacetyl-oxy-15 α -cyano-puupehenol 731
- O-19, 20-diacetyl-oxy-15 α -nitro-ethanepuupehenol (3) 733
- O-19, 20-diacetyl-oxy-15 α -nitro-methanepuupehenol 733
- O-19,20-diacetyl-oxy-15 α -cyano-puupehenol 29 732
- O-glycosides 124, 125, 135, 159
- O-glycosylation 59
- O-hydroxycinnamic acids 345
- O-primeverose 304
- O-vanillin 282
- O3-demethylthebaine 493
- Ocimene 25, 252, 253
- Ocimum 225
- Octacosanol 764
- Octacosyl alcohol 764
- Octamic acid 405
- Octopamine 703
- Odour 248
- Oflocet 674
- Ofloicin 674
- Ofloxacin 671, 673, 674
- Ofloxacin 674
- Oil of anise 249, 287, 292, 294
- Oil of balm 249
- Oil of basil 249
- Oil of bay 252
- Oil of bergamot 249, 269
- Oil of bitter almond 250
- Oil of caraway 286, 287
- Oil of chenopodium 299
- Oil of cherry laurel 249
- Oil of cinnamon 28, 249, 283
- Oil of citronella 267, 274, 301
- Oil of clove 28, 249
- Oil of eucalyptus 297
- Oil of fennel 292
- Oil of garlic 249
- Oil of geranium 267
- Oil of hops 252
- Oil of lavender 283
- Oil of lemon 249
- Oil of lemon grass 267, 272
- Oil of marjoram 288
- Oil of mentha 249
- Oil of neroli 268
- Oil of orange 249
- Oil of orris rhizome 283
- Oil of palmarose 267
- Oil of palmarosea 26, 249
- Oil of rose 26, 250, 267
- Oil of sasafras 249, 294
- Oil of savin 260
- Oil of summer savory 288
- Oil of thyme 249
- Oil of turpentine 252, 318, 320
- Oil of wintergreen 250, 301, 304
- Oil origanum 288
- Oil petit grain 268
- Oil pimenta 28
- Oils of cohobation 284
- Oils of nutmeg 28
- Oils of verbena 272
- Okadaic acid 718, 719, 720
- Okinawan marine sponge 728
- OKT3-monoclonal antibody 47
- Old maid 505, 506, 508, 512
- Old man 347, 348
- Older fluoroquinolones 676
- Oleaceae 168
- Oleandomycin 619, 623
- Oleandomycin hydrochloride 624
- Oleandomycin triacetate ester 624
- Oleandrose 85, 619
- Oleanol 187
- Oleanolic acid 187
- Oleo-gum resins 314
- Oleo-gum-resin 218, 328, 329, 330, 332, 333
- Oleoptene 249
- Oleoresin 247, 281, 314, 324, 326, 327, 328
- Olibanum 310
- Oligomerization 750
- Oligonucleotide primer 56
- Oligosaccharides 35, 85
- Ololiuqui 539
- Omega oil from salmon 736
- Omega-3 fatty acids 776
- Omega-3 fish oil 776
- Omnipen 633
- Omnipen-N 634
- Oncogenesis 748
- Oncoscint (R) CR/OV 42
- Oncovin 515
- Ondena 691
- Onion 735
- Ontogeny 23
- Ophelic acid 199
- Ophtagram 582
- Ophthalmokalixan 583
- Opiate alkaloids 814
- Opium 34, 7, 480
- Opium alkaloids 486, 810
- Opium latex 39
- Opium poppy 374, 375, 377, 486, 490, 491, 493, 495, 542
- Optical rotary dispersion (ORD) analysis 250
- Optical rotation 249
- Optimycin 653
- Optium 633
- Oracefal 602
- Oracilline 631
- Oral contraceptives 183
- Oral hypoglycemic drugs 831
- Orange 243
- Orange carotenoids 738
- Orange juice 736
- Oratren 631

- Orbenin 632
 Ordeal bean 532
 Orelox 609
 Organic additives 802
 Organogenesis 801
 Organosulphur phytochemicals 759
 Organum oil 287
 Orient ginseng 189
 Orientomycin 641
 Oriparine 497
 Oripavine 486, 493, 494, 495
 Oripavine hydrochloride 494
 Oripavine methiodide 494
P
 Orspray 651
 Orstanorm 539
 Orthoclone OKT3 79
 Orthocresol 298
 Orthodox medicaments 814
 Orvagil 669
 Osoapiole 296
 Ospamox 633
 Ossien 792
 Oswego tea oil 225
 Osyritin 749
 Osyritrin 749
 Otokalixin 583
 Otreon 609
 Ouabagenin 156
 Ouabain 156
 Ovaglobulin 790
 Ovalbumin 790
 Ovulation dipstick test 67
 Oxacepham 612
 Oxacillin 628, 631
 Oxaldin 674
 Oxamycin 641
 Oxapenam 611
 Oxidation 217, 219
 Oxidation of nicotine 458
 Oxidative products of terpenes 308
 Oxide volatile oils 251, 300
 Oxidising enzyme 783
 Oxiniacic acid 459
 Oxlopar 652
 Oxolinic acid 670, 671
 Oxomate 774
 Oxsoralen 351
 Oxybiocycline 652
 Oxychelidonic acid 34
 Oxycopaivic acid 310
 Oxycyclin oxy-dumocyclin 652
 Oxygenated acids 523
 Oxygenated NADPH 593
 Oxygenated pinane derivative 279
 Oxymycin 652
 Oxytetracid 652
 Oxytetracycline 649, 650, 651, 652, 653
 Oxytetracycline dihydrate 652
 Oxytetracycline hydrochloride 652
 Oxytetrin 652
 Oxytocin 428
 p 53 tumour suppressor 754
 p, p'-dimethoxystilbene 292
 p-hydroxy cinnamic acid 752
 p- α -dimethyl benzyl alcohol 332
 p-amino benzoic acid (PABA) 667
 p-aminophenyl 663
 p-anisaldehyde 280
 p-coumaric acid 178, 341, 369
 p-cymene 26, 28, 224, 253, 254, 257
 p-hydroxy-cinnamic acid 369
 p-hydroxy-cinnamoylferuloyl-methane 332
 p-hydroxy-phenylpyruvic acid 369
 p-menthane 218, 224, 254, 259, 260, 262
 p-methoxy benzoic 293
 p-nitrobenzoate 225
 Pacific nudibranch 721
 Pacific yew 8, 235
 Pacitaxel 8
 Pakistani ephedra 357
 Paldomycin 654
 Paliuroside 749
 Palm oil 771
 Palmidin A, B, and C 132, 133
 Palthe senna 139
 Palytoxin 709, 710, 711, 716
 Panaeolus 500
 Panaxadipol 188
 Panaxasides 188
 Panaxatriol 188
 Pancrealipase 783, 787
 Pancreatic enzymes 787
 Pancreatin 635, 783, 786, 787
 Pangram 582
 Panicillium AT 38
 Panmycin 650
 Pansies 161
 Papain 781, 783, 786
 Papaverine 382, 383, 388, 398
 Papaya 375
 Papillioideae alkaloids 544
 Papilloma 693
 Paprika 37, 323, 348, 397, 411, 418, 768
 Para-cymene 253
 Para-aminobenzoic acid (PABA) 666
 Para-aminosalicylic acid (PAS) 679
 Para-cinnamoyl CoA 354
 Para-coumaric acid 341, 371, 752
 Para-coumaroyl CoA 354
 Para-cymene 254, 273
 Para-dimethylaminobenzaldehyde 470
 Para-menthadienes 255
 Para-menthane 254, 255, 256, 259, 261
 Paraguay jaborandi 437
 Paraguay tea 2, 359
 Paralytic shell fish poisoning 711, 714
 Paramorphine 491
 Paramycin 580
 Parasicyonis action stoloides 705
 Parent plants 810
 Parenteral preparations 577
 Parfuran 677
 Paromomycin 576
 Parsley 159, 295
 Parsley oil 29
 Parsley seed oil 296
 Parsnips 350
 Parthenolide 231
 Partition 394
 Pasetocin 633
 Passiflorin 501
 Passion flower 501
 Pthalic anhydride 357
 Pathways for ajmalicine, preakummicine 510
 Pathways for coryanthe, aspidosperma and iboga 505

- Pavecef 603
Paveltia 427
Pavine clover 341, 342
PbTxS 713
PC-based automated bioreactors 813
PCR-technique 58
Pd-catalyzed hydrogenation 655
Peakummicine 510
Pectin 35, 108, 109, 133, 770
Pectin galacturonan 109
Pectinmethylesterase 782
Peegee 347
Pegamome 428
Peganine 428
Pelargonidin 167, 745
Pelletierine 374, 398
Pen-oral 631
Penam 611
Penamox 633
Penavlon V 631
Penbritin-S 634
Penem 611
Penicillanic 625
Penicillanic acid 1 639
Penicillanic acid sulfone 639
Penicillin 573, 596, 602, 625, 631, 633, 635, 637, 646, 696
Penicillin fermentation 572
Penicillin G 574, 577, 626
Penicillin G potassium 627
Penicillin N 596
Penicillin O 16
Penicillin production 573
Penicillin V 628, 631
Penicillin V potassium salt 631
Penicillin-resistant staphylococci 664
Penicillin-resistant strains 676
Penicillinase 600
Penicillinase-resistant penicillin 627, 628, 632
Penicillins G sodium 628
Penicillium 570
Penicillium O 38
Penicline 633
Penimox 633
Penitracin 645
Pentacyclic terpenoid 834
Pentacyclic triterpenes 821
Pentacyclic triterpenoid saponins 186, 213
Pentacyclic triterpenoids 186, 210, 236
Pentid 627
Pentosan 106
Pentosans 86
Pentrex 634
Peonidin 167, 745
Pepper spray 768
Peppercorns 768
Peppermint camphor 27
Peppermint oil 223, 285
Peppers 411, 418
Pepsin 780, 784, 787
Pepsinogen 780
Peptide alkaloids in ergot 541
Peptides 60
Peptidyl transferase 659
Peptones 787
Periciclina 652
Perilla 290
Perilla sugar 277
Perillaldehyde 276, 277
Perillartine 277
Periwinkle 505, 506, 507, 508, 512
Peroxidase enzyme 102
Peroxynitrite free-radical 754
Peru balsam 336
Peruresinotannol 312
Peruresinotannol cinnamate 336
Peruvian balsam 334, 335, 336
Peruvian bark 521
Peruvial 336
Petit grain 301
Petroleum plant 343, 349
Petunidin 167, 745
Peyocactine 478
Peyote 479
Phagocytic cells 46
Phallandral 279
Pharmaceutical adjuvants 91
Pharmaceutical chemistry 571
Pharmacobiotechnology 41, 66
Pharmacognosy 1, 4, 25
Pharmacophore moieties 819
Pharmaceutical industry 5
Pharmorubicin 592
Phase I enzymes 751
Phase II enzymes 751, 757, 774
Phases in fermentative process 572
Phasins 794
Phellandral 278
Phenanthrene 399
Phenethyl isothiocyanate 757
Phenol glycosides 127, 139, 211
Phenol volatile oils 251
Phenolic acids 751
Phenolic alkaloids 391
Phenolic bitter principles 547, 548
Phenolic ether volatile oils 251, 290, 291
Phenols 28
Phenoxyethylpenicillin (penicillin V) 631
Phenyl alanine 206
Phenyl ethanol 360, 361
Phenyl propanoids 340
Phenyl propyl cinnamate 338
Phenyl-pyruvic acid 208
Phenylacetamide 574
Phenylacetic acid 626
Phenylacetic acid derivative 574
Phenylalanine 204, 208, 209, 210, 340, 535, 789
Phenylalanine pathway 208
Phenylalanine-derived alkaloids 405
Phenylbenzopyrilium based constituents 35
Phenylethylamine 396, 398, 626
Phenylethylamine alkaloids 475, 477
Phenylethylene 338
Phenylethylisoquinoline alkaloids 475
Phenylpropanoids 207, 340, 361, 366, 556
Phenylpropanolamine 11
Phenylpropenes 340, 344
Phenylpropyl cinnamate 334
Phenylpyruvic acid 207, 369
Phenytol 690
Pheromones 217
Phloroglucin test 564
Phloroglucinol 164, 325
Phosphate buffer 572
Phospho-enolpyruvic acid 208
Phosphomolybdic acid 438
Phosphoprotein 793
Phosphoric acid 201

- Phosphoric acid-complex 225
Phosphorylases 781
Photoanethole 292, 293
Photosantonin 551
Photosynthetic pigments 239
Physalaemin 705
Physcion 132
Physical characteristics of volatile oils 248
Physical properties of resins 308
Physiomycine 653
Physostigma 532
Physostigmine 10, 381, 382, 389, 399, 532, 533, 534
Physostigmine salicylate 533
Physostigmine sulphate 533
Physostigmine sulphite 534
Phytic acid 752
Phytoalexins 217
Phytochemicals 737, 738
Phytochemistry 3, 4, 24, 33, 344
Phytoestrogen 748
Phytofoods 737
Phytohemagglutinin 794
Phytomedicines 775
Phytomelin 749
Phytonutrients 737, 747, 760, 774
Phytopharmaceuticals 202, 828
Phytosterols 760
Phytotoxicity 653
Picric acid 394, 643
Picrocrocin 277
Picrolonic acid 394
Picropodophyllin 319
Picrorhiza 196
Picrorhiza kurroa 196
Picroside 197
Picrotin 562
Picrotoxin 561, 562, 563
Picrotoxinin 552, 562
Pictet-spengler isoquinoline synthesis 484
Pierami 581
Pilocarpine 16
Pill bearing spurge 39
Pilocarpic acid 437, 438
Pilocarpine 380, 382, 389, 396, 397, 399, 437, 438, 439, 440, 441
Pilocarpine perchromate 439
Pilopic acid 438
Pilosine 437, 440, 441
Pimaric acid 310
Pimpinellin 556, 557
Pinaceae 307
Pinane 259, 260, 264, 265
Pine oil 33, 241
Pine tar 322
Pineapple 359
Pinene 27, 223, 253, 742
Pinene hydrochloride 219, 261
Pinetar 287
Pinque 511, 512
Pinus 219, 321
Pinus elliotth 321
Piperacillin 629, 635, 637
Piperacillin sodium 638
Piperidine 373, 398, 441, 464
Piperidine alkaloids 396, 442
Piperidine-6-carboxylic acid 455
Piperine 383, 389, 442, 447, 448, 768
Piperitone 28, 226
Piperonol 294
Piperoyl-CoA 448
Pipermint oil 243, 249, 255
Pipracil 638
Pipril pentacillin 638
Piraldina 679
Piramox 633
Pirilene 679
Pisatin 807
Pistachia galls 318
Pitayine 527
Pituri 397, 456, 457, 469
Plant families 397
Plant growth hormone 801, 807
Plant hormones 807
Plant phenolics 340
Plant poisons 809
Plant propagation 800
Plant secondary metabolites 806
Plant terpenoids 217
Plant tissue culture technologies 798, 813
Plant-cell cultures 798, 810
Plantago 106, 107
Plantago seeds 14, 106, 108
Plantain 343, 356
Plantain seed 14
Plaque 53
Plasma cells 75
Plasma kinins 816
Plasma proteins 788, 791
Plasma volume expander 92
Plasmid 48, 49, 52
Plasmid DNA 51
Plasmid vector 56
Plasmid-mediated resistance 671
Plasmin 788
Plasminogen 788
Platinum chloride test 448
Pletil 670
Pluropon 750
Pneumatic method 245
Podofilox 30
Podophyllic acid 319, 320
Podophyllotoxin 30, 319, 362, 363, 812
Podophyllum 309, 318, 319, 362
Podophyllum resin 318
Poinsettia 342
Point mutation 76
Policosanol 764, 765
Poliomyelitis 64
Pollination 745
Poly β -ketomethylene acid 203
Poly-peptides 709
Polyantus narcissus 356
Polyaromatic hydrocarbons 769
Polycillin-N 634
Polycyclic polyether metabolites 713
Polycycline 650
Polyenes 681
Polyether ionophores 713
Polyether macrolide 720
Polygalacturonase 782
Polygalitol 192, 193
Polygallic acid 192
Polygonaceae 127
Polygris 681
Polyhalo-3-butene-2-one 717
Polyketide 657, 682
Polymerase chain reaction (PCR) 46
Polymethylgalacturonase 782
Polymixin antibiotics 643
Polymixin B 642, 643, 644, 645
Polymox 633
Polymycin 644
Polymycin E 642, 644
Polymyrcene 334
Polymyxin B sulphate 643, 646

- Polymyxin E1 642, 645
Polymyxin E2 642, 645
Polymyxin hydrochloride 643
Polymyxins 641, 642
Polypeptide antibiotics 579, 641
Polypeptide protein 825
Polypeptides 641
Polyphenol extracts 744
Polyphenolics 744
Polyphenols 777
Polysaccharides 35, 85, 86
Polyterpene 251
Polytoxin (C51-55) hemiacetal 710
Polytrim 644
Polyunsaturated fatty acid (PUFA) 714
Polyurethane 817
Pomegranate bark 391
Pomegranate juice 777
Pomegranate seeds 748
Poncyl-FP 681
Poppy capsule 814
Poppy seed 7
Poppyseed poppy keshi 490, 491, 493
Populin 139, 143
Potassium-cupric tartrate test 563
Potato 342
Potato vaccine 82
Potency of pancrealipase 787
Potency of pancreatin 786
Potency of pepsin 787
Potent newer antineoplastic drug 825
Potential medicinal plants 233
Potential nucleophilic donors 733
Potential secondary plant metabolites 808
Poultry products 736
Powdered gum 105
Powdered opium 487
p,p-dihydroxy dicinnamoylmethane 332
Prairie huckleberry 416
Praticef 605
Preakuammicine 509, 519, 521
Pregnancy 69
Pregnancy dipstick test 67
Pregnane 185
Pregnenolone 14
Prepacifenol 717
Preparation of resins 309
Preparative-scale fermentation 728
Prepared resins 309
Prephenic acid 208, 369
Presenegenin 192
Prevailing practised therapies 815
Prezetamid 679
Prickly poppy 486, 490
Prilagin 689
Primaric acid 31
Primary flow 307
Primary metabolites 202
Primary-amide 646
Primaxin 614
Primbactam 613
Primeverose 141, 304
Primveroside 193
Principle stilbene 754
Proanthocyanidins 745, 751
Probe DNA 49
Probenecid 579
Procaine 12, 626
Procanthocyanidins 168
Prodrug approach 578
Prodrugs 598, 599, 600, 609
Production of penicillin 575
Proenzyme 780
Progesterone 13
Proleukin (R) 42
Proleukin (R) (Aldesleukin) 79
Proline 535
Propanethial-S-oxide 759
Propagation of plant-tissue cultures 802
Propoxyphene 11
Propranolol 234
Procentrum lima 711
Prostaglandins 10, 728, 696, 765,
Prostaphlin-A 632
Prostasal 760
Prostrate knotweed 367
Protanol 35
Protaverine 421
Proteases 781, 786
Protectins 60, 794
Protein kinase 791
Protein variants 790
Proteins 36, 60, 63
Proteolytic enzyme 783, 785, 794
Proteoses 782, 787
Proto-alkaloids 372
Protocatechuic acid 290, 333, 355
Protocrocin 277, 278
Protogonyaulax cartenella 711
Protogonyaulax cohorticula 711
Protophenicol 662
Protoveratrine A and B 389, 420, 421
Protoverine 420
Provatene 740
Provitamin A 240
Prumase 176
Prunasin 122, 175, 176, 177, 207, 209
Prunetrin 166
Prunus amygdalus 174
Pseudo ionone 273, 274
Pseudoconhydrine 404, 405, 406, 408, 409, 477
Pseudoephedrine oxalate 477
Pseudojervine 420, 424, 426
Pseudomembranous colitis 670
Pseudotannins 38
Psilocin 495, 498, 499, 500
Psilocybin 499
Psilocybe 500
Psilocybin 495, 499, 500
Psittacosis 661
Psolaridin 171
Psoralea 170, 171
Psoralen 171, 172, 350, 351, 556, 558
Psoralen coryfolia 350
Psoralenol 172
Psoriasis 351
Psyllium seed 14
PTX 716
Ptychodiscus brevis 711
Pudina oil 736
Puffer fish 712
Pulegone 223
Purgative 124
Purine 2, 3, 399, 201
Purine alkaloids 396, 401
Purity of antibiotic 577
Purple grape-juice 754
Putrecine 476
Putrescine 460, 471, 476
Puupehenone 730, 731, 732, 733
Pycnogenols 745, 750
Pyelonephritis 602

- Pyocianil 636
 Pyopen 636
 Pyrafat 679
 Pyran 157
 Pyrazinamide 679
 Pyrazinecarboxamide 679
 Pyrethrins 30
 Pyrethrin I 30
 Pyrethrin II 30
 Pyrethrins 809
 Pyridine alkaloids 454
 Pyridoxine hydrochloride 802
 Pyrimidine base 201
 Pyrindine 398
 Pyrocatechin 333
 Pyrocatechol 290
 Pyrodinium bahamense 711
 Pyrogallol 358, 368, 369
 Pyrones 157, 158
 Pyrrolidine 398, 461, 464
 Pyrrolidine alkaloids 396, 462
 Pyrrolidine ring system 461
 Pyrrolizidine 400, 476, 449, 457
 Pyrrolizidine alkaloids 396, 462, 471, 474, 475, 544, 810
 Pyrroloindole 532
 Pyrroloindole alkaloids 495, 531
 Pyrroloindole nucleus 532
 Pyruvate oxidation factor 771
 Pyruvic acid 208, 409, 485
- Q**
- Q fever 661
 Qinghaosu 6
 Quadracycln 650
 Quantitative estimation of volatile oils 247, 248
 Quassia 547
 Quassia wood 563, 564
 Quassin 237, 561, 563, 564
 Quaternary 380
 Quebrachine 511
 Quebrachol 760
 Queen of the meadow 304, 359
 Queensland 39
 Quercetin 140, 161, 162, 163, 319, 750
 Quercetin (F) 206
 Quercitin 162
 Quest for new antibiotics 569
 Quillaia sapotoxin 194
 Quillaic acid 187, 194
 Quillaja 194
 Quillaja bark 194
 Quillaja saponin 187
 Quinamm 524
 Quinate 524
 Quinazoline alkaloids 427
 Quinazolines 396
 Quinbisan 524
 Quindine hemipentahydrate 528
 Quine 522, 524
 Quinic acid 376
 Quinicardine 527
 Quinidex 527
 Quinidine 379, 398, 429, 430, 431, 432, 433, 521, 522, 523, 524, 527, 528, 530, 531
 Quinidine bisulphate 528
 Quinidine gluconate 528
 Quinidine hydrogen sulphate tetrahydrate 528
 Quinidine polygalacturonate 528
 Quinidine salt 528
 Quinidine sulphate dihydrate 527, 528
 Quiniduran 528
 Quinine 10, 12, 24, 216, 375, 376, 381, 382, 387, 388, 389, 396, 397, 398, 429, 430, 431, 432, 433, 434, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 544, 814
 Quinine bimuriate 524
 Quinine bisulphate heptahydrate 524
 Quinine dichloride 524
 Quinine dihydrochloride 524
 Quinine hydrochloride 380
 Quinine hydrochloride dihydrate 524
 Quinine sulphate 380, 387, 431
 Quinine sulphate dihydrate 524
 Quinine trihydrate 524
 Quinol 729
 Quinoline 398, 429
 Quinoline alkaloids 396, 429, 435, 495, 521, 811
 Quinoline-ring-system 434, 495
 Quinolinic acid 460
 Quinolizidine 449, 454
 Quinolizidine alkaloids 449
 Quinolizidine derivatives 543
 Quinolone antibacterial drugs 670
 Quinolones and fluoroquinolones 665, 670
 Quinones 29
 Quinonic acid 459
 Quinonoid system 158
 Quinora 527
 Quinsan 524
 Quinuclidine 429
- R**
- R-(–)-β-Citronellol 276
 Racemic isomeride 388
 Racephedrine hydrochloride 478
 Racephedrine sulphate 478
 Radiation therapy 746
 Radio-immunoassays (RIA) 67
 Radish 757
 Radix rhei 131
 Raisins 751
 Rancidity 241
 Raphanin 757
 Rasping machine 89
 Raspings 244
 Raspings process 244
 Rathimed 669
 Raubasine 505
 Rauwolfia 506, 511, 512, 513
 Rauwolfia serpentina 7
 Raymond test 147
 Raymond's reagent 147
 Reaction with ferric chloride 288
 Receptor 816
 Recombinant DNA 18, 51, 68, 74
 Recombinant DNA products 63
 Recombinant DNA technology 63, 75, 79
 Recombinant human interleukin-3 80
 Recombinant human platelet derived growth 80
 Recombinant organism 47
 Recombinant plasmids 52
 Recombinant vaccination vector 46
 Recombinant vaccines 64

- Recombinant vaccinia vector 47
Recombinate(R) 79
Recombivax HB(R) 79
Recovery through adsorption 576
Red algae 716, 717, 723
Red carotenoids 738
Red cinchona 521
Red clover 341, 342
Red kidney bean 794
Red sea sponge 721
Red snapper 715
Red squill 153
Red tide 711
Red tide organisms 711
Red tide toxins 711
Red wine 751, 754
Red-algae 115
Red-tide toxins 709
Redox enzyme 783
Reduced nicotinic adenine dinucleotide (NADH) 772
Reducing enzyme 783
Reducing sugars test 103
Reference compounds 820
Refobacin 582
Refractive index 249
Reimer-tiemann's reaction 282
Reinacke salt 643
Reineckate complex 394
Reineckate salt solution 385
REL P analysis 69
Remicyclin 650
Remijia alkaloid 433, 524
Remijia pedunculata 528
Remijia purdieana 528
Renin 786
Rennet casein 793
Rennet extract 793
Rennets 781
Rennin 784, 787, 793
Reparil 828
Replication 823
Reported 710
Rescinnamine 509, 512, 513, 518, 519
Resene resins 313
Resenes 308, 310, 312
Reserpine 19, 7, 318, 381, 382, 389, 399, 509, 512, 518, 519
Reserpine hydrochloride hydrate 513
Reserpine 513
Reserve transcriptase 56
Resin acids 306, 308, 309, 310, 324, 333
Resin alcohols 306, 308, 309, 311, 313, 329
Resin combinations 218
Resin esters 306, 308, 309, 310, 336, 337
Resin guaic 39, 315
Resin phenols 309
Resinates 310
Resinols 306, 309, 312
Resinotannols 306, 309, 312
Resins 129, 39, 218, 306, 307, 308, 314, 315
Resistomycin 583
Resorcinol 298
Respiratory enzyme 783
Restriction endonucleases 47
Restriction enzymes 47
Restriction length fragment polymorphism (REL P) 68, 69
Resveratrol 744, 751, 753, 754
Reticuline 486, 493, 793
Retinol 240
Retronecine 471, 474, 475, 476
Reverse transcriptase (RT) 44, 45
Rhabdomyosarcoma 692
Rhamnaceae 127
Rhamnol 760
Rhamnose 107, 162, 163
Rhamrose 85
Rhein 132, 138, 761, 823, 824
Rhein-8-glucosides 125, 138
Rhein-8-diglucoside 138
Rheinolic acid 133
Rheum 131
Rhubarb 131, 132, 133
Rhubarb rhizome 131
Rhubarbs 761
Ribbed gourd 236
Ribose 201
Ricinine 384, 398
Rickettsial diseases 661
Ridged 236
Rifa 658
Rifabrodin 658
Rifabutin 678
Rifadin(e) 658
Rifaldazine 658
Rifaldin 658
Rifampicin 658
Rifampin 658, 659, 678, 679
Rifamycin 658
Rifamycin SV 658
Rifamycins A-E 658
Rifoldin 658
Rimactan(e) 658
Rindex 653
Rio jaborandi 437
Rivea 536
Roasted coffee 342
Robamox 633
Robiocina 664
Rocefin 607
Roferon-A (R) 80
Roman chamomile 297, 346, 349
Roman chamomile 232
Romicil 623
Randomycin 653
Root bark of viburnum prunifolium 359
Rose oil 241, 248
Rosemary 297
Rosenthaler's reagent 386
Rosequin test 433
Rosin 39, 311
Rossed bark 177
Rotary thin-film evaporator 392, 393
Rotenoid 166, 167, 561
Rotenone 166, 559, 560, 561, 809
Rotenonone 560
RP-13057 690
Ruban 430
Rubber 313, 738
Rubiaceae 127
Rubiaceous alkaloids 397
Rubiaceae 168
Rubidomycin 690
Rubiervine 200, 201, 420
Rubreserine 533
Rue 350, 353, 435
Rue oil 350, 353
Russian liquorice 191
Rutacridone 435, 436
Rutin 22, 161, 162, 744, 749, 750
Rutinose 162, 164
Rutoside 749

S

- S-adenosylmethionine (SAM) 500
 S-allyl cysteine sulphoxide 759
 S-glycosides 124, 125
 S-phase 51
 Sabadine 421
 Sabinene 259, 260
 Saffron 297
 Safranal 277
 Safrole 291, 294, 295, 300
 Sago starch 91
 Sagrada 127
 Saint john's wort 763
 Salaspermic acid 834
 Salicin 122, 139, 142, 143, 356, 358, 359
 Salicylaldehyde 355
 Salicyldehyde 280
 Salicylic acid 25
 Saligenin 142, 143
 Saliretin 142
 Salosanine 185
 Salts of alkaloids 375
 Salts of penicillins 626
 Salutaridine 495, 496
 Salutaridinol 495, 496
 Sample-selection strategy 816
 San pedro aguacolli 479
 Sanclomycine 650
 Sandbox tree 343
 Sandhox tree 341
 Sandracolic acid 310
 Santonin 551, 552
 Sapogenins 183, 418
 Saponification test 241
 Saponin glycosides 127, 182, 183, 193, 213, 743
 Saponins 743
 Saprophytic 782
 Sarcophine 727
 Sarcophine metabolites 727
 Sarcophytol A (2) 727
 Sarmenose 85
 Sarpaganda 506, 511, 512, 513
 Sassafras 287, 295, 493
 Saturated aliphatic alcohols 266
 Saw palmetto 343, 359
 Sawacillin 633
 Saxitoxin 702, 703, 704, 712
 Scammony 201, 309
 Scarification process 243
 Schaer's reagent 386, 470
 Scheibler's reagent 385
 Schematic method of isolation of cinchona alkaloid 432
 Schiff base 484
 Schneider-weppen test 488
 Schoenteten's reaction 130
 SCID 78
 Scillabiose 151
 Scillaren A 150, 151, 152
 Scillaren B 150, 151, 152
 Scilliroside 153
 Sclerotia 535
 Sclerotium 535
 Scopalamine (Hyscyamine) 10
 Scopalamine and nicotine patches 14
 Scopoletin 178, 346, 348, 349
 Scopolia 397
 Scopolin 348, 349
 Scotch broom 450
 Scrophulariaceae 236
 Sea urchin 37
 Search for AIDS vaccines 65
 Sebercim 673
 Secologanim 505
 Secologanin 504, 525
 Second generation antibiotics 604
 Second generation cephalosporins 600, 603, 606
 Second-generation drugs 605
 Second-line agents 679
 Second-line drugs 678
 Secondary constituents 24, 119
 Secondary flow 307
 Secondary glycosides 146
 Secondary metabolites 202, 797, 798, 805, 806, 807, 810, 811
 Secondary modes of action 647
 Secondary plant metabolites (SPMs) 809
 Secondary-amine form 381
 Secoridoid and a terpenoid derivative 504
 Sedamine 447
 Sedral 602
 Seed gums 106
 Seed-dispersal 745
 Seglor 539
 Semi-synthetic antibiotic 633, 635, 656
 Semi-synthetic β -lactamase 640
 Semi-synthetic cephalosporin 602, 605
 Semi-synthetic cephalosporin antibiotic 600
 Semi-synthetic products 718
 Semisynthetic alkaloid 539
 Senecic acid 475
 Senecionine 396, 471, 474, 475
 Senedrine 478
 Seneecionine 476
 Senega 192
 Senega root 193
 Senegenic acid 192
 Senegenin 192
 Senegin II 192
 Senna 127, 137, 761
 Senna leaves 824
 Sennas 761
 Sennecionine 400
 Senneciphylline 400
 Sennoside A 138, 762
 Sennoside B 138
 Sennoside C 138
 Sennoside D 138
 Sennosides 138, 762
 Seopolia 342
 Separation and isolation of antibiotics 575
 Separation of citronellal from citral 275
 Septicid 672
 Septopal 582
 Seratiopeptidase 784, 787, 788
 Serine protease enzymes 821
 Seromycin 641
 Serotonin 495, 498, 499
 Serotonin antagonist 231
 Serotonin complex with creatinine sulphate monohyd 498
 Serotonin hydrochloride 498
 Serpentine 519
 Serum albumin 790
 Serum globulin 790
 Serum protein 791
 Sesquiterpene 251, 332
 Sesquiterpenoid lactone 228, 230, 231

- Sesquiterpenoids 215, 218, 229, 807
- Severe combined immunodeficiency (SCID) 77
- Sevinon 684
- Sex hormones 183
- Shatavari 186
- Shavegrass 342, 456
- Shell fishes 711
- Shellac 307, 320
- Shellolic acid 320, 321
- Shensi rhubarb 343
- Shikimic acid 203, 204, 206, 208, 210, 663
- Shikimic acid pathway 208, 364, 369, 370
- Shikimic acid—mevalonic acid mediators 203
- Shikimic acid-phenylpropanoid pathway 662
- Shikon 810
- Shiomarin 608
- Shogaols 327
- Short-acting sulphonamide 668
- Shotgun-cloning-experiments 17
- Siam benzoin 337, 338
- Siaresinotannol 312
- Sigadoxin 654
- Sigamopen 633
- Signal molecules 217
- Silamox 633
- Silarine 750
- Silepar 750
- Silicotungstic acid 438
- Silicristin 365
- Silidianin 365
- Silirex 750
- Silliver 750
- Silmar 750
- Silver morning glory 539
- Silybin 365
- Silymarin 365, 750
- Simple β -carboline alkaloids 495, 500
- Simple indole alkaloids 495
- Simple protein 791
- Simple tetrahydro iso-quinoline alkaloids 475, 481
- Simplotan 670
- Sinalbin 180, 182, 756
- Sinapic acid 341, 343, 344
- Sinapic acid choline ester 343
- Sinapine 182
- Sinapine acid sulphate 182
- Sinigrin 36, 125, 180, 181, 209, 246, 765
- Sinistrin 151
- Sinularin 697
- Sinusitis 622
- Sisomicin 585, 586
- Sito lande 760
- Sitoindoside VII and VIII 830
- Sitosterin 760
- Skimmetin 33, 347
- Snake weed 39
- Snakeplant 539
- Socotrine aloe 128
- Socotrine and zanzibar aloes 131
- Soda process 97
- Sodium acetate 209
- Sodium artesunate 231
- Sodium desoxycholate 683, 684
- Sodium nitrite test 489
- Sodium santoninate 551
- Sodium sulfisoxazole 667
- Sodium sulfisoxazole [Ionized Form] 667
- Soframycin 585
- Soft-gelatine 792
- Solanaceae 375, 376
- Solanaceous alkaloids 397
- Solaneceae 168
- Solanidine 200, 201, 377, 396, 400, 415, 417, 418, 419
- Solanine 200, 376, 377, 418
- Solanum alkaloids 415, 417
- Solanum khasianum 184
- Solar energised biochemical reactor 202
- Solasodine 185, 415, 416, 418, 419
- Solatene 740
- Solatubine 417
- Solid-agar medium 801
- Solubility 308
- Soluble CD4 66
- Soluble mucopeptides 785
- Somatic Cell 77
- Sonnenschein's reagent 385
- Sophisticated analytical instrumentations 733
- Sophisticated skillful antibiotic preparations 578
- Sophoretin 750
- Sophoricoside 166, 822
- Sophorin 749
- Sorghum 209
- Sorquetan 670
- Source of carbon nutrition 572
- Sources of yeast 795
- Southernwood 347, 348
- Soxhlet extraction process 392
- Soy 736
- Soy products 747
- Soy-protein hydrolyzates 802
- Soyabean 776
- Soyabean milk 735
- Soyabean yogurt 735
- Soyalecithin 36
- Soybean lipooxygenase 781
- Soyfood intake 776
- Soyfoods 776
- Spafloxacin 671
- Spanish fly 566
- Spanish foxglove 149
- Spanish licorice root 367
- Spanish savory 301
- Spara 675
- Sparfloxacin 675
- Sparteine 399, 449, 450, 451
- Sparteine sulphate pentahydrate 451
- Spearmint oil 243, 285
- Special tests 130
- Specific biological targets 815, 816, 820
- Specific DNA sequence 47
- Specific gravity 249
- Specific plasma proteins 788
- Specific protein kinase inhibitor 822
- Specific test 103
- Specificity of vaccines 65
- Spectrabid 635
- Spectrazole 604
- Spent broths of penicillium glaucum 358
- Spermatophyta 306
- Speromycin 663
- Spetamycin III diacetate 625
- Spiramycin I, II and III 624, 625
- Spiramycins 619, 624
- Spiran 682

- Spiroamine alkaloids 544
Spirofulvin 681
Spiroketal steroids 210
Spiruline 773
Splicing 53, 74
Splitter 111
Splitting enzyme 783
Sponge 716, 718, 719, 720
Sponge metabolites 729
Sponge method 243
Spongegourd 236
Spongosine 698, 706, 707
Spongouridine 698
Sporostatin 681
Spot test 241
Squalene 210, 686
Squaw weed 474
Squill 150, 151
Squill glycosides 152
Stabicilline 631
Stable radical 744
Stachydrine 398, 462, 463
Stachydrine acid oxalate 464
Stachydrine aurichloride 464
Stachydrine hydrochloride 464
Stachydrine platinichloride tetrahydrate 464
Standard analytical procedures 820
Standard operating procedures (SOPs) 575
Staphybiotic 632
Star anise 290, 294, 297, 356
Star anise fruit 292
Starch 21, 86, 87, 133
Starch purification centrifuges 88
Stas-Otto process 393
Stationary phase 798
Stearoptene 249
Stemmadenine 509
Sterculia gum 104
Stereo-configuration 123
Stereochemically different isomer 731
Stereochemistry 407
Stereoisomerism 256
Stereoselective synthesis 616
Steric hinderence 610
Steriodal spanonin 186
Steroid 400
Steroid biogenesis 204
Steroid glycosides 127, 144, 211
Steroidal 418
Steroidal alcohol 760
Steroidal alkaloidal glycosides 200, 214
Steroidal alkaloids 32, 185, 396, 415
Steroidal moiety 705
Steroidal saponins 183, 186, 209
Steroidal triterpene 188, 236
Steroidal triterpene aglycone 188
Steroidal-glycoside 32
Steroids 21, 183, 204, 738, 809, 32
Sterols 760
Stevacin 652
Stigmasterol 14
Stock culture 571
Stomach poison 457
Storax 310, 334, 335, 337
Storesinol 312, 335
Stramonium 465, 469
Strawberry clover 34
Streptamine 588, 589
Streptidine 201, 579, 587, 588, 589
Streptobiosamine 587, 588
Streptobrettin 587
Streptokinase 784, 788
Streptomyces 570, 589, 592
Streptomycetaceae 570
Streptomycin 16, 201, 576, 577, 580, 586, 587, 588, 678
Streptomycin hydrochloride 587
Streptomycin sesquisulphate 587
Streptomycin sulphate 587
Streptomycin trihydrochloride 587
Streptomycin-penicillin 588
Streptonivicin 663
Streptose 588, 589
Stretobiosamine 201
Strictosidine 509, 510, 519, 525, 530, 531
Strophanthidin 122, 154, 155
Strophanthin 155
Strophanthobiase 155
Strophanthotriose 155
Strophanthus 153
Strophanthus fruit 154
Strophanthus glycosides 155
Strophanthus plants 154
Structural analogues of erythromycin 621
Strychnine 4, 381, 382, 389, 397, 399, 509, 516, 517, 519, 521
Strychnine dehydrogeissoschizine 518
Strychnine derivatives 517
Strychnine gluconate pentahydrate 518
Strychnine glycerophosphate hexahydrate 518
Strychnine hydrochloride dihydrate 518
Strychnine N6-oxide 517
Strychnine nitrate 517
Strychnine phosphate 517
Strychnine sulphate 517
Stryctinine 521
STXs 713
Styaresin 335
Styracin 303, 334
Styrax 303, 356
Styrene 334
Styresinol 335
Styresinolic acid 335
Styrocamphene 335
Styrogenin 335
Suberin 21
Substance P 705
Substituted furocoumarins 558
Subunit vaccines 64, 65
Sugar 4, 124
Sugar kelp 35
Sulbactam 634, 638, 639
Sulbactam ampicillin 630, 640
Sulbactam sodium 640
Sulfisoxazole 668
Sulforaphane 757
Sulmycin 582
Sulphamethoxazole 667, 668
Sulphanilamide 667
Sulphate 433
Sulphate process 97
Sulphides 759
Sulphonamide 291, 666, 667
Sulphonamide drug 588
Sulphonamides and trimethoprim 665, 666
Sulphonamides for general infections, 666
Sulphonamides for intestinal infections 666

- Sulphonamides for local infection 666
 Sulphonamides for urinary infections, 666
 Sulphonate process 97
 Sulphonyl ureas 690
 Sulphoraphane 773
 Sulphurated glycosides 179
 Sulphuric acid test 354, 562
 Sulphuric acid-dichromate test 517
 Sumatra benzoin 303, 337, 338, 356
 Sumbul 347
 Sumox 633
 Superficial conjunctival infections 661
 Superoxide dismutase 748, 772
 Suppressor T-cells 46
 Supracyclin 654
 Supramycin 650
 Surface sterilization 803
 Surface sterilization of explants 803
 Surface sterilizing agents 803
 Surinam balsam 335
 Surinam quassia 357, 563
 Suspen 631
 Suspension cultures 804, 806, 812
 Swansonine 453, 454, 455
 Swansonine 453
 Sweat violet 244
 Sweet basil 292, 294
 Sweet birch 304
 Sweet birch oil 30
 Sweet clover 169
 Sweet flag 294
 Sweet orange 305
 Sweet peppers 323, 348, 397, 415
 Sweet-potato starch 91
 Swerlia 199
 Swertiamarin 565
 Sylvestrene 258
 Sylvic acid 311
 Symphytum 471
 Synclotin 601
 Synonym foromacidin B 625
 Synpenin 634
 Syntheses of pseudo- and α and β -ionones 273
 Synthetases 781
 Synthetic antimicrobial agents 665
 Synthetic camphor 219, 221, 261
 Synthetic gene 82
 Synthetic monocyclic b-lactam (monobactam) antibio 613
 Synthetic oligonucleotides 49
 Synthetic pathway 520
 Synthetic racemic menthol 223
 Synthetic seed technology 800
 Synthetic template 817
 Synthetic vaccines 73
 Syntometrine 536
 Syphilis 69
 Syrian rue 501
 Syringin 830
- T**
- T-lymphocytes 77, 78
 T-cell production 742
 T-cells 78, 742
 Tabac 457
 Tabazur 456
 Tabersonine 504, 505, 509, 510
 Tachykinins 705
 Taka-diastrase 143
 Tall oil rosin 314
 Tamoxifen 774
 Tan acid 614
 Tannic acid 367, 368
 Tannic-acid complex 394
 Tannin 133, 140
 Tannings of nutgalls 358
 Tannins 38, 366, 367, 747
 Tansy 297
 Tar 321, 322
 Targocid 648
 Tarivid 674
 Tarragon 346
 Taxanes 824
 Taxifolin 34
 Taxol 14, 19, 8, 235, 381, 811, 814, 824
 Taxol precursors 235
 Taxon 397
 Taxonomic classification 397
 Tazicef 607
 Tazidime 607
 Tazobactam 638, 640
 Tazobactam sodium 640
 Tazobactam-piperacillin 630, 640
 Tazocilline 640
 Tazocin 640
 Tea 2
 Teaberry 304, 342, 343
 Teaberry oil 30
 Tebrazid 679
 Tegopen 632
 Teichomycin A2 648
 Teichoplanin 641
 Teicoplanin 648
 Teicoplanins 648
 Telepathine 502
 Tenebrarius 589
 Tenebrimycin 589
 Tenemycin 589
 Teniposide 363
 Teonanacatl 498
 Tephrosin 166
 Terbinafine 681, 685
 Terbinafine hydrochloride 686
 Terebene 318
 Teriposide 362
 Teriterpene acids 338
 Terminal biological methylation 618
 Terpene aldehyde 271, 272
 Terpene hydrocarbons 250, 251
 Terpene limonoids 742
 Terpene phenols 288
 Terpene-derivatives 266
 Terpenes 215, 250, 258, 306, 738
 Terpenoid alkaloids 412
 Terpenoid indole alkaloids 495, 504, 501, 511
 Terpenoid peroxide 299
 Terpenoids 21, 31, 215, 217, 218, 228, 234, 358, 362, 400, 738
 Terpentine 257
 Terpin hydrate 221, 222, 257, 267, 270
 Terpinene 253
 Terramycin 652
 Terrestrial organisms 718
 Tertiary-amine form 381
 Test organism 570
 Test plates 570
 Tetrabromo-2-heptanone 717
 Tetrabromodihydromyrcene 252, 253
 Tetracyclic pretetramide nucleus 656

- Tetracyclic triterpenoid saponins 183, 213
Tetracyclic triterpenoid saponins 183
Tetracycline 16, 649, 650, 653, 654, 656,
Tetracycline hydrochloride 650
Tetracycline variants 650
Tetracyclines 579, 649, 654, 655
Tetradecin 650
Tetradotoxin 720
Tetradox 654
Tetrahydric phenol 29
Tetrahydro- β -carboline system 509
Tetrahydrocannabinol 23, 317
Tetrahydroharmine 504
Tetrahydroisoquinoline 485
Tetrahydroisoquinoline derivatives 481
Tetrahydropteroyl synthetase 667
Tetrahydrozoline 11
Tetralisal 656
Tetralysal 656
Tetramycin 650
Tetramyl 656
Tetran 652
Tetrandrine 485, 486
Tetranontriterpenoid 238
Tetrasulphate salt of quinine 431
Tetraterpenoids 216, 218, 238
Tetrodotoxin 712, 713
Thalleioquin 523
Thalleioquin Test 433, 523, 524
Thaumatococin 795, 796
Thaumatococins I and II 795
Thea 2
Theaflavins 747
Thearubigins 747
Thebaine 23, 486, 491, 492, 494, 495, 496, 497
Thebaine binoxalate monohydrate 492
Thebaine bitartrate monohydrate 493
Thebaine hydrochloride monohydrate 492
Thebaine oxalate hexahydrate 492
Thebaine poppy 490, 491, 494
Thebaine salicylate 492
Theobromine 2, 3
Theophylline 2, 3, 389, 675
Thiamine 373
Thiamine hydrochloride 37, 802
Thiamine monochloride 373
Thienamycin 612, 614
Thioctacid 771
Thioctan 771
Thioctic acid 771
Thiocyanate glycosides 179
Thioester-bound lysergic acid 540
Thioglycosides 127, 179, 180, 182, 209, 213
Thiosemicarbazone 280
Thiosulphinate 759
Third generation cephalosporins 600, 605, 606, 607, 608, 610
Thorn apple 462, 465, 469
Threonine 789
Thrombin 821
Thromboxane pathways 765
Thujane 259, 265
Thuzane 264
Thyme 287
Thyme camphor 225
Thyme oil 225
Thymidine 44, 47, 48
Thymol 28, 219, 223, 225, 226, 227, 254, 288, 289
Thymol iodide 227
Thymosin alpha 1 66
Thymus histones 790
Thyroglobulin 790
Tiacarcillin 629
Ticar 637
Tiacarcillin 579, 636, 637
Tiacarcillin disodium 637
Ticarpen 637
Ticillin 637
Tiecoplanin A2 648
Tiemann's method 275
Tienam 614
Tilden reagent 217
Timentin 639
Tinidazole 669, 670
Tinnevelley senna 137
Tio-mid 680
Tiocan 771
Tissue cells 804
Tissue culture propagation 803
Tissue culture technique (PTCT) 799
Tissue explants 805
Tissue plasminogen activator (TPA) 62
Tissue plasminogen activator 785
Tliltliltzen 539
TMP-SMZ 668
TNF- α 740
TNG-alpha 753
Tobacco 456, 457
Tobacco plant 161, 375
Toban 444
Tobracin 589
Tobradistin 589
Tobralax 589
Tobramycin 576, 580, 589, 590
Tobrex 589
Tocopherols 770
Tocosamine 451
Tocotrienols 770
Tokiocillin 629, 633
Tollen's reagent 147, 162, 273, 284, 354
Tollen's test 147
Tolu balsam 334, 336, 337
Toluene 356
Toluresinotalol 312
Toluresinotannol cinnamate 337
Tomatidine 415, 417, 418, 419
Tomatine 417
Tomato 200
Tonco seed 169
Tonga 346, 347
Tonka bean 346, 347
Tonka bean comphor 346
Tonopres 539
Toothpick ammi 553
Topicycline 650
Topioca starch 91
Topitracin 645
Topoisomerase 591, 823
Toricelecin 601
Totacef 602
Total alkaloids 15
Totaquine 530
Totazina 644
Totomycin 650
Toxins 695
Tracer technology 202
Tracix 614
Traditional chinese medicine (TCM) 777

- Traditional vaccines 64
Tragacanth 101, 103
Tragacanthin 104
Trans- β -ocimene 25
Trans-aconitic acid 376
Trans-anethole 292
Trans- β -ocimene 252, 253
Trans-diethylstilbestrol from 12
Trans-ferulic acid 329, 343, 347
Trans-isomer 256
Trans-lycopene 739
Trans-trans-cinnamyl cinnamate 303
Transcription 823
Transferring enzyme 783
Transformation 663
Trantoin 677
Traumatic acid 807
Traveler's joy 341
Tree tobacco 456, 457
Trescatyl 680
Tretinoin 240
Trevir 463
Triacetyl derivative 730
Triazole 687, 688
Trichazol 669
Trichomoniasis 669
Tricocet 669
Tricolam 670
Tricyclic diterpene 311
Tricyclic sesquiterpenoids 228
Triflucan 690
Trigenelline 398
Trigenolline 37
Triggering mechanism 573
Trigonella foenum 205
Trigonelline 37, 317
Trimethoprim 644, 668
Trimethyl gallic acid 178
Trimethylamine 37
Trimonase 670
Trimox 633
Trimyo-gen™ 778
Triocetin 624
Triptidolides 812
Triperpenoids 236
Tripple extract 245
Tripsin 781
Trisaccharides 167
Triterpene 251
Triterpenoid 237
Triterpenoid acids 187
Triterpenoid aglycone 193
Triterpenoid saponin glycosides 194
Triterpenoid saponins 188, 209
Triterpenoids 216, 218, 236, 238
Troleandomycin 619, 624
Tropane 398, 461, 464
Tropane alkaloids 396, 462, 464, 544
Trophophase 572, 574
Tropic acid ester with tropine 465
Tropine 473
Tropine tropate 465
Tropinone 473
Tropolone 400
Trovafloracin 671, 676
Trovafloracin hydrochloride 676
Trovafloracin mesylate 676
Trovafloracine monomethane-sulfonate 676
Trypiamine 500
Trypsin 780, 784, 788
Tryptamine 534
Tryptamine secologanin 510
Tryptophan 789
Tsiklomitsin 650
TTX 712, 726
Tuberculosis 69
Tubic acid 560
Tubic acid lactone 560
Tubocurarine 380, 389, 392, 398, 485, 486
Tubocurarine chloride 8, 382
Tulsi 736
Tulsi oil 225
Tumorigenesis 748
Tumour imaging 68
Tumour necrosis factor (TNF) 62, 77
Tupa 443
Turkish tannins 38
Turmeric 329, 331, 332
Turmeric yellow 753, 829
Turnsole 474
Turpentine 258, 262, 309, 323, 327, 328, 743
Turpentine oil 243, 261, 262, 328
Tuttomycin 585
Tx-1 713
Tx-2 713
Tyloses 307
Type II β -lactamases 638
Types of yeast 794
Tyramine 482
Tyrosine 208, 340
Tyrosine kinases 748
- U**
- U-19920 698
U-gencin 582
Ubiquinones 738, 771, 772
Udicil 698
UDP-glucose 756
Ultracef 602
Ultroxim 604
Umbellic acid 329, 347
Umbelliferae 33, 168, 225, 307, 330, 347
Umbelliferone 33, 329, 346, 347, 558
Umbelliferone test 330
Unacid 640
Unacil 654
Unacin 640
Unasyn 640
Undecylenic acid 681, 684
Undecylenic acid zinc salt 685
Unicin 650
Unilobin 444
Unimycin 652
Unipen 632
Unipyranamide 679
Uniquin 673
United Nations Convention on Biodiversity (CBD) 817
Unmatal 456, 462, 469
Unsaturated acyclic hydrocarbons 251
Unstable alkaloidal salts 383
Uracil arabinoside 698
Urantoin 677
Uric oxidase 782
Uricolytic enzyme 783
Uritone 677
Urizept 677
Uro-tablinen 677
Uro-tensins I and II 708
Urodin 677
Urokinase 784, 788

- Urolong 677
 Urotensins I and II 709
 Urotensins I and II (U I, U II) 709
 Urotropin 677
 Uroxacin 673
 Ursolic acid 140
 Usefulness of the secondary metabolites 808
 Uticillin VK 631
 Utimox 633
 Utinor 673
 UV 734
 UV-chamber 332
 UV-spectral data 820
 UV-visible spectroscopy 250
- V**
- V-cillin 631
 Vaccines 63, 72, 73
 Vaccines from live attenuated viruses 72
 Vaccinia virus 46
 Vagilen 669
 Valerian 342
 Valerian alkaloid 412
 Valine 789
 Van Urk reagent 537
 Van Urk test 537
 Vancocin 646
 Vancomycin 577, 646, 647, 648, 649
 Vancomycin hydrochloride 647, 648
 Vancomycin monohydrochloride 647
 Vancor 647
 Vanilla 195, 196, 303, 356, 359
 Vanilla bean 281
 Vanilla pod 195, 281
 Vanillic alcohol 196
 Vanillin 27, 195, 196, 210, 280, 281, 282, 334, 337, 338, 356, 359, 360, 411, 412
 Vanillin HCl test 563
 Vanilloside 195, 360
 Vanillylamine 411, 412
 Vanobid 641, 664
 Vantin 609
 Varatridine 422
 Varnish tree 320
 Vasicine 389, 427, 428, 429
 Vasicinone 429
 Vaxsyn[®] HIV-1 80
 VCR 515
 Vebe 514
 Vectacin 586
 Veetrin 655
 Vegetable calomel 318
 Vegetable lutein 31, 741
 Vegetable lutenol 31
 Vegetable luteol 741
 Vegetable vaccine 82
 Vegetables sponge 236
 Vehem-Sandoz 363
 Velban 514
 Velmonit 672
 Vendarcin 652
 Venezuelae 660
 Venzonate 302
 Vepen 631
 Vepesid 362
 Veracevine 420
 Verapamil binding sites 726
 Veratetrine 421
 Veratramine 4, 400, 420, 424, 425
 Veratridine 420, 421
 Veratrine 421
 Veratrosine 420
 Veratrum alkaloids 419, 420
 Verbena 367
 Verbenaceae 127
 Verladyn 539
 Verongid sponge 732
 Vervain 367
 Vetstrep 587
 Vibra-tabs 654
 Vibramycin 654
 Vibramycin hyclate 654
 Vicillin 634
 Vidarabine 721, 723
 Vinblastine 14, 19, 381, 389, 399, 509, 514, 515, 518, 519, 520
 Vinblastine dihydrochloride dihydrate 515
 Vinblastine sulphate 382, 514
 Vinca 9
 Vinca rosea 515
 Vincal leukoblastine 399, 514
 Vincosamine 646
 Vincoside 434, 515
 Vincrex 515
 Vincristine 14, 19, 382, 389, 509, 514, 515, 518, 519
 Vincristine sulphate 515
 Vindesine 381
 Vindoline 505, 507, 509, 510, 519, 520
 Vindoline hydrochloride 508
 Vinegar 1
 Vinodoline 508
 Vintage wines 753
 Violaquercitrin 749
 Viral vaccines 72
 Viral vector 60
 Virginia tobacco 456, 457
 Virus particulates 70
 Viscardan 553
 Viscose 98
 Visiren 674
 Visnacorin 555
 Visnagalin 553
 Visnagen 553
 Visnagin 170, 554, 555, 556
 Visokellina 553
 Vitali-morin colour reaction 470
 Vitallin 790
 Vitamin A 239, 240, 274, 741, 808
 Vitamin B 795
 Vitamin B complex 795
 Vitamin B₁ 37, 373, 802
 Vitamin B₃ 454
 Vitamin B₆ 802
 Vitamin BT 766
 Vitamin C 37, 736, 742, 748, 772, 808
 Vitamin D 808
 Vitamin E 741, 771, 772, 808
 Vitamin K 808
 Vitiligo 351
 Vitreous 128
 VLB; 29060-LE 514
 Vncoled 646
 Vodol 689
 Volatile oil 21, 241
 Volatile oil in a plant material 247
 Volatile oil of bitter almond 246
 Volatile oil of black mustard 246
 Volatile oils 39, 218, 240, 242
 Volatile terpenoids 807
 Voltage-sensitive Na-channel 713
 Volume of distribution 656
 Vulcamycin 663
 Vumon 363

W

w-CgTX 726
w-CgTXVIA 726
w-Cm TX 726
w-Conotoxins 725, 726
Wagner-meerwein rearrangements 415
Wagner's reagent 385, 438, 448
Walnuts 777
Warfarin 690
Warfarine 675
Water and steam distillation 243
Water distillation 243
Water hemlock 442
Wayenecomycin 617
Welfurin 677
Well-defined bioassays 815
Wheat bran 770
Wheat germ oil 764, 771
White dextrin 100
White mustard 180, 181, 182
White thyme 301, 302
Whole-wheat bread 735
Whooping cough vaccine 72
Wieland-Gumlich aldehyde 519
Wild cherry bark 177
Wild coleus 290
Wild gourd 341, 342
Wild lettuce 469, 564
Winter savory 301, 302
Wintergreen 304, 342, 343
Wonderberry 416
Wood rose 539
Woodruff 169
Wormseed 304
Wound hormone 807
Woundwort 342
Wytrion 624

X

X-ray diffraction analysis 250
Xanthan gum 116, 117
Xanthine 2, 3
Xanthoeriodictyol 315
Xanthophyll 31, 239, 741
Xanthoscellide 151
Xanthotoxin 351, 352, 558
Xanthotoxol 558
Xanthidrol reagent 147
Xanthidrol test 147
Xenobiotics 773
Xylose 107, 247

Y

y-ephedrine 407, 408
y-ionone 273, 274
Yageine 502
Yanogonin 322, 323
Yarrow 463
Yeast 680, 794
Yeast artificial chromosomes (YACs) 50
Yeast variants 795
Yellow carotenoids 738
Yellow cinchona 521
Yellow dextrin 100
Yellow dock 355
Yellow enzyme 783
Yellow gentian 566
Yellow lady-slipper 357
Yellow resin 39
Yellow sweetlover 341
Yellow-sponge 724
Yerba mate 359
Yersinia 672
Ylang-ylang 294, 302, 304, 305, 345, 356

Yocon 511
Yohimbe 397, 511
Yohimbine 8, 381, 399, 509, 511, 518, 519
Yohimbine hydrochloride 511
Yohimex 511
Yohidrol 511
Yope 355
Yupa 355

Z

Zacin 768
Zadorin 654
Zadstat 669
Zagam 675
Zamocillin 633
Zanathid coral 710
Zanzibar aloes 128
Zavedos 595
Zeatin 807
Zeaxanthin 738, 741
Zeaxanthol 741
Zetamicin 586
Zienam 614
Zinacef 604
Zinamide 679
Zinc undecylenate 685
Zingiberaceous starch grains 332
Zingiberene 229, 327, 332
Zolicef 602
Zonarol 716
Zonathus sp 724
Zone of lysis 53
Zoolobelin 444
Zoroxin 673
Zostrix 36, 323, 410, 768
Zosyn 640
Zutracin 645
Zwitterion 659
Zygophyllaceae resins 313
Zymase 795